

Title: Population-based estimates of the prevalence of children with congenital heart disease and associated comorbidities

Authors: Devin M. Parker, MS¹; Meagan E. Stabler, PhD²; Todd A MacKenzie, PhD³; Meghan S. Zimmerman, MD, MPH⁴; Xun Shi, PhD⁵; Allen D. Everett, MD⁶; Emily M. Bucholz, MD, MPH, PhD;⁷ Jeremiah R. Brown, PhD^{1,3,8}

¹The Dartmouth Institute for Health Policy and Clinical Practice, Geisel School of Medicine at Dartmouth, Hanover, NH;

²Department of Family and Community Medicine, Northern New England CO-OP Practice and Community Based Research Network, Dartmouth Hitchcock, Lebanon, NH;

³Department of Biomedical Data Science, Geisel School of Medicine at Dartmouth, Hanover, NH;

⁴Department of Pediatrics, Dartmouth-Hitchcock Medical Center, Lebanon, NH;

⁵Department of Geography, Dartmouth College, Hanover, NH;

⁶Department of Pediatrics, Johns Hopkins School of Medicine, Baltimore, MD;

⁷Department of Pediatrics, University of Colorado, Aurora, CO

⁸Department of Epidemiology, Geisel School of Medicine at Dartmouth, Hanover, NH

Correspondence to: Devin M. Parker, Geisel School of Medicine at Dartmouth, Hanover, New Hampshire. devin.m.parker.gr@dartmouth.edu

Abstract word count: 304

Manuscript word count: 3175

Abstract

Background

Congenital heart defects (CHD) are the most common birth defects and are estimated to affect almost 1% of births annually in the US. CHDs are the leading cause of infant mortality associated with birth defects and can result in chronic disability, morbidity and increased healthcare costs. Most CHD prevalence estimates rely on data from population-based birth defects surveillance systems and these estimates are inconsistent due to varied definitions. It is therefore important to derive high-quality, population-based estimates of the prevalence of CHD to help care for this vulnerable population.

Methods

We performed a descriptive, retrospective 8-year analysis using all payer claims data (APCD) from Colorado (CO) from 2012 - 2019. Children with CHD were identified by applying ICD-9 and ICD-10 diagnoses codes from the American Heart Association-American College of Cardiology (AHA-ACC) harmonized cardiac codes. We included children with CHD <18 years of age who resided in CO, had a documented zip code, and had at least one healthcare claim. CHD type was categorized as simple, moderately complex and complex disease. Association with comorbid conditions and genetic diagnoses were analyzed using chi-square test. We used direct standardization to calculate adjusted prevalence rates, controlling for age, sex, primary insurance provider and urban-rural residence.

Results

We identified 1,566,328 children receiving care in CO from 2012 – 2019. Of those, 30,512 children had at least one CHD diagnosis, comprising 1.9% (95% CI: 1.93 – 1.97) of the pediatric population. Over half of the children with CHD also had at least one complex chronic condition and 11% had a genetic diagnosis.

Conclusion

The current study is the first population-level analysis of pediatric CHD in the US. Our study found a higher CHD prevalence than in previously reported studies and a high comorbidity burden. These findings can inform initiatives to improve screening, treatment and planning of care for this complex population.

INTRODUCTION

Congenital heart defects (CHD) are the most common birth defects and are estimated to affect almost 1% of births annually in the US.^{1,2} CHDs are the leading cause of infant mortality associated with birth defects and can result in chronic disability, morbidity and increased healthcare costs.³

Previous prevalence estimates of children with CHD have been inconsistent due to varied definitions, included only a subset of CHD diagnoses, were geographically limited and were generated from a small number of sources.^{3,4} Many frequently cited prevalence estimates were derived from birth defect registries, where a diagnosis was made in a regional pediatric cardiology center and relied on patient referrals. Studies that actively screen for cases of CHD by echocardiography are not done routinely.^{4,5} A seminal report by Hoffman and colleagues in 1965 estimated prevalence of structural heart defects at birth in 6-11 cases per 1,000 births. While informative, their study excluded the most common CHD diagnoses - ventricular and atrial septal defects.^{6,7} Hoffman et al. concluded that much of the observed variability across studies reflected differences in case ascertainment.^{3,6}

Another important source of variation is the nomenclature used in CHD prevalence studies. Disparate approaches have been used to identify CHD, including definitions specifically developed for congenital registries or screening programs and studies limited to CHD requiring surgery. The most widely cited prevalence estimates were generated from the Metropolitan Atlanta Congenital Defects Program (MACDP), a population-based birth defects surveillance system from 1998 - 2005. Children with CHD were defined by the nomenclature developed from the International Paediatric and Congenital Cardiac Code, an initiative developed by members of the Society of Thoracic Surgeons (STS) and the European Association for Cardio-Thoracic Surgery.⁸ All identified cases were limited to surgical diagnoses used by the STS congenital

heart surgery database. This study by Reller et al reports an overall CHD prevalence of 8.14 per 1,000 live births.³

Population-based empirical data on the epidemiology and prevalence of children with CHD have been limited. Marelli and colleagues report contemporary prevalence estimates of CHD using Canadian health services databases.⁹ Infants were identified by administrative diagnosis and surgical procedural codes, reporting an observed prevalence rate of 8.21 per 1,000 from 1983 - 2010. A more recent model developed from the Global Burden of Diseases, Injuries and Risk Factors Study found a 4.2% increase in global CHD prevalence since 1990.¹⁰ We continue to lack robust and empirical U.S.-based population estimates.⁹

Advancements in medical care and surgical treatments have significantly increased the survival of CHD individuals.¹¹⁻¹³ Despite such progress, research continues to show that children with CHD are associated with both the highest mortality and the highest inpatient resource use in pediatric care.¹⁴ In 2004, congenital abnormalities accounted more than 139,000 hospitalizations, of which over 30% were for cardiac and circulatory anomalies.¹⁵ In 2009, hospital admissions for CHD cost an estimated \$5.6 billion annually, comprising 15% of costs for all pediatric care, while only accounting for 3.7% of pediatric admissions.^{14,16,17} It is now widely recognized that CHD is associated with frequent health care use and lifelong comorbidity.^{9,12} The impact of ongoing disease burden includes depression, learning disabilities, asthma, cancer, obesity, rheumatologic disease, hypertension, stroke, and heart failure. The true magnitude of comorbidity burden is not known.^{16,18}

Estimating prevalence of children with CHD by healthcare administrative claims data presents a novel opportunity to study longitudinal patterns of this prominent disease. All-payer claims databases (APCDs) cache public and private claims for health care services provided to insured

individuals within a state.¹⁹ Twenty-one states have established APCDs, and many others are currently in development.^{20,21} The governance of APCDs, their funding, and the authority of APCDs to require claims submissions varies by state.

In the current study we present contemporary, longitudinal, population-based estimates for children with CHD. High-quality, population-based prevalence estimates of CHD prevalence is critical to support the understanding of pediatric CHD epidemiology, as it may lead to better insight into its etiology and care practices.¹⁰ A comprehensive assessment of the CHD health burden can support appropriate allocation of resources for diagnosis, care coordination, care access and cost-effective treatment strategies.

METHODS

DATA SOURCE: CO APCD

The Center for Improving Value in Health Care (CIVHC) is a nonprofit authorized by the state to collect Colorado (CO) claims data. The APCD includes medical claims and dates of service from commercial health plans, Medicare and Colorado's Medicaid program. CO APCD includes hospital, clinic and claims files related to diagnosis, procedure, enrollment, member and provider data.

The current study is a retrospective study of children with CHD represented in the CO APCD. This is a longitudinal dataset spanning eight calendar years (January 1, 2012- December 31, 2019). Every individual was assigned a unique encrypted identifier on their first healthcare encounter and followed across all diagnoses, encounters, settings and years.

STUDY POPULATION

Our study population included all children aged 18 years or younger on Jan 1, 2012, with a documented CO zip-code and at least one healthcare encounter in CO (N= 1,535,816) during the study period. Individuals with CHD were identified by the American Heart Association-American College of Cardiology (AHA-ACC) harmonized cardiac codes.¹⁰ Using this algorithm, CHD is defined by the *International Classification of Diseases, Ninth Revision*, or *International Statistical Classification of Diseases and Health Related Problems, Tenth Revision*, (ICD-9/10) in the primary diagnosis field (ICD-9: 745.xx, 746.xx, 747.xx; ICD-10: Q20.x, Q21.x, Q22.x, Q23.x, Q24.x, Q25.x, Q26.x, Q27.x, Q28.x) (Appendix Table 1). Our study population included well-recognized chromosomal abnormalities that are commonly associated with CHD. Children with co-occurring genetic abnormalities have often been excluded in CHD prevalence studies due to their more complex disease. These genetic abnormalities include Down syndrome, Turner syndrome, DiGeorge syndrome, trisomy 13, trisomy 18, Alagille syndrome, Noonan

syndrome, Holt-Oram syndrome, Williams syndrome, Kabuki syndrome, VATER/VACTERL association, CHARGE association, Jacobsen syndrome and Williams-Beuren syndrome.²²⁻²⁴

Our cohort did not include diagnoses related to arrhythmias, cardiomyopathies, or transplants. We also excluded diagnoses that are specifically associated with adult disease as the particular anatomy and progression of disease for children is distinctively unique from adult diagnoses. (Appendix Table 1). To increase the specificity of the algorithm for our study population, manual audits of CHD diagnoses (N = 120) were independently adjudicated by two pediatric cardiologists specializing in congenital heart disease. The manual audit had 95% inter-rater agreement. Discrepancies in diagnoses were agreed upon through expert consensus.

Comorbidities and Disabilities

Individual comorbidities were defined by the Pediatric Complex Chronic Conditions (CCC) V2 classification system. CCCs are defined as “any medical condition that can be reasonably expected to last at least 12 months (unless death intervenes) and to involve either several different organ systems or 1 organ system severely enough to require specialty pediatric care and probably some period of hospitalization in a tertiary care center”. A comprehensive set of ICD-9-CM and ICD-10-CM codes were identified as CCC and subsequently bracketed into 12 categories (cardiovascular, premature and neonatal, neurologic or neuromuscular, metabolic, gastrointestinal, renal, respiratory, hematologic or immunologic, malignancy, technology dependency, transplantation, and other congenital or genetic condition).²⁵ In our evaluation, we omitted categories related to cardiovascular and other congenital or genetic complex conditions as those are known to occur simultaneously with CHD.

Individual disabilities were defined by the Children with Disabilities Algorithm (CWDA), a validated, claims-based tool for identifying children with disabilities.²⁶ CWDA classifies children

as having a disability as those “who have long-term physical, mental, intellectual, or sensory impairments which in interaction with various barriers may hinder their full and effective participation in society on an equal basis with others.” CWDA also accounts for intellectual and mental conditions that “limit children’s ability to fully participate in society”. There are 17 distinct categories within the CWDA based on the disability concepts and definitions articulated by the World Health Organization (WHO).

Spatial attributes

Individuals were categorized as urban-or rural residing by joining home zip-codes to 2010 Rural-Urban Commuting Area (RUCA) categorizations.²⁷ RUCA codes categorize U.S. census tracts on measures of population density, urbanization, and daily commuting. RUCA codes used in this study were based on data from the 2010 decennial census. Rural is defined as “towns with populations below 10,000 and surrounding commuter areas with more than one-hour driving distance to the closest city”.^{27 28} Median household income estimates were derived from American Community Survey (ACS), a publicly available U.S. Census Bureau survey that enables linkage of community characteristics with individual zip-code tabulation area (ZCTA) codes.²⁹

Severity categories

CHD severity categories were defined by the American College of Cardiology/ American Heart Association guidelines for the Management of Adults with Congenital Heart Disease (ACHD). These guidelines define CHD severity within an anatomic and physiological framework.³⁰ Category designations are clinically relevant and structured to facilitate incorporation of administrative data to inform prevalence levels and trends. These three severity categories are descriptively named as: simple, moderately complex and complex (Appendix Table 2). We included an additional fourth category ‘other’ to encompass remaining, uncategorized

diagnoses. For individuals that had multiple diagnoses across different severity categories, we ranked codes hierarchically so that the most severe lesion would be assigned with the primary diagnosis (i.e. 'complex' diagnosis of transposition of the great vessels diagnosis would be considered primary even if child also had a 'simple' septal defect). We further assigned CHD diagnoses into anatomically oriented and clinically relevant lesion categories (Appendix Table 3).

CHD diagnoses were analyzed by sociodemographic and geographic factors. Age at the time of first encounter was categorized as newborn (0 – 30 days), infant (31 – 365 days), child (366 – 10 years old), and youth (>10 – 18 years old). Sex was characterized as a dichotomous variable, with male as the referent. Primary insurance provider was assigned to the insurer associated with the most claims by individual and categorized by commercial/self-pay or Medicaid. Median household income was categorized by quintile (lowest quintile: \$24,427 – 51,461; highest quintile: \$95,050 – 147,421).

Statistical Analysis

Prevalence is defined as the proportion of individuals in a population who have a disease at a specific point in time.³¹ We calculated the observed prevalence of children with CHD as the number of patients diagnosed with CHD in a given year (numerator), divided by the CO pediatric population in that year (denominator). Prevalence was reported as the number of unique CHD diagnosed children per 1,000 individuals. We applied the Cochran-Mantel-Haenszel test and Cochran-Armitage test for nonzero correlation to test for yearly trends. Categorical variables are described as frequencies and proportions and compared through χ^2 tests.

We used direct standardization to calculate adjusted prevalence rates across the study period. The direct method of adjusting for differences among populations involves computing the overall

rates that would result if all populations had the same frequency distribution. Direct standardization adjusts for any distortion caused by different distributions between the general population and study population. In the initial model, we adjusted for age, sex, primary insurance provider, region of residence, urban-rural residence and median household income. We performed tests of collinearity to refine our model. In the final model, we adjusted for age, sex, primary insurance provider and urban-rural residence.

RESULTS

Figure 1. illustrates the derivation of the 1,565,387 children in the CO APCD health claims database during our longitudinal study period. We identified 30,512 unique children with CHD (received a CHD diagnosis at any point during the study period), comprising 1.9% (95% CI: 1.93 – 1.97) of the pediatric population.

Prevalence by demographics, insurance provider, disease severity and community characteristics are summarized in Table 1. In total, 48.1% were female, 56.3% had their first CHD encounter before the age of one, 88.4% resided in urban core areas, and 71.8% were primarily insured by Medicaid.

Most children with CHD were diagnosed with simple, non-complex cardiac lesions (65.4%) and 19.9% were diagnosed with a moderately complex CHD lesion. By anatomically oriented category, atrial septal defect was most commonly diagnosed, comprising nearly 25% of all diagnoses, ventricular septal defects accounted for 11% and patent ductus arterioses were diagnosed in 11% of all diagnoses (Table 2).

Among the children with CHD, 3,023 (11%) were diagnosed with a genetic disorder, and 405 (1.3%) had more than one genetic diagnosis. Trisomy 21 (Down's syndrome) was diagnosed in 1,557 children (5.1%) and 866 (2.8%) had Kabuki syndrome. Other genetic disorders frequently associated with CHD in our cohort included DiGeorge syndrome, Noonan syndrome, VATER/VACTERL, Turner syndrome, Jacobsen syndrome and Trisomy 18 (Table 3).

More than half of the children with CHD also experienced at least one concurrent chronic complex condition (55.3%), as identified by the CCC algorithm. Among those with CCCs, 7,626 (45.2%) had multiple CCC diagnoses in addition to CHD. The most common chronic complex

conditions were related to respiratory, metabolic, gastrointestinal, neurologic and neuromuscular conditions. (Table 4).

In our CHD cohort, 11,749 (38.5%) received at least one disability diagnosis during the study period, as defined by the CWDA. There were 5,484 (18.0%) children with CHD that were treated for more than one disability. Nervous system and mental health disorders were most common (23.1% and 16.7%, respectively). We also found that children with CHD were treated for respiratory, circulatory, digestive, and genitourinary diseases. (Table 5).

Geography

Figures 2 and 3 illustrate the population density of all CO residing children and children with CHD in our study, respectively. While we observe a higher density of non-CHD children residing in the western and mountain regions, we note strong similarities between both cohorts in population clusters along the Front Range region.

DISCUSSION

We present the first U.S. population-based study to longitudinally estimate the prevalence and comorbidity burden of children diagnosed with CHD in the U.S. During 2012 to 2019, there were 30,512 children identified with CHD in CO, comprising 1.9% of the pediatric population. In comparison with previously reported estimates, our higher reported prevalence may be explained by the inclusion of children with concurrent genetic abnormalities, which have not been previously accounted for, as well as diagnoses captured after infancy. Our study summarizes the significant and complex health burden that children with CHD frequently experience, with more than half having at least one comorbid condition and 45% have multiple.

Most children with CHD were diagnosed with simple, non-complex cardiac lesions, and less than one-fifth of the children were diagnosed with a moderately complex lesion. There was an increased prevalence in ventricular and atrial septal defect subcategories, possibly related to improved detection or diagnosis after infancy. Ventricular septal defects are the most common form of CHD and studies show that at least 25% of all ventricular septal defects spontaneously close.³² Retaining an initial diagnosis of ventricular or atrial septal defect across the study period may also explain the increased prevalence.

CHD occurs in association with other anomalies or as part of an identified syndrome in 25 – 40% cases.³³ We found that 11% of children with CHD were also diagnosed with a genetic abnormality. It has been reported that fifty-percent of individuals born with Trisomy 21 (Down's syndrome) also have atrial and ventricular septal defects, and approximately one-third of females with Turner syndrome have CHD.²³ In our study, we report a similar proportion of children diagnosed with both CHD and Trisomy 21.

Improved survival of children with CHD has led to increased attention to extracardiac complex conditions, often found across multiple body systems.^{16,34-36} We found that over half of the children diagnosed with CHD have at least one coexisting chronic complex condition and 2 in 5 children have a disability in addition to CHD. The most common comorbidities were respiratory, metabolic and neurologic conditions but a significant proportion of children also had diagnoses related to renal, immune and malignant conditions.

It has been reported that long-term neurodevelopmental and cognitive disabilities are the most common comorbidities for children with CHD.^{37,38} There are multiple theories to potentially explain the high prevalence of mental health disorders seen in this subset of children, such as abnormalities of in utero brain development and metabolism resulting in more saturated cerebral blood flow and underlying genetic syndromes, parental mental health disorders, prematurity, cardiac bypass surgery, and prolonged hospitalizations are posited as factors that could increase the risk of neurodevelopmental disabilities.³⁹⁻⁴² A study of youths with CHD reported 7 times higher odds of anxiety and attention deficit/hyperactivity disorder (ADHD), compared to those without CHD.³⁹ Applying validated algorithms, we found that over 25% of children with CHD also experience mental health disability. To date, we are the first to report the prevalence of mental health disabilities for children with CHD, both longitudinally and across healthcare settings.

Most studies elect to exclude children with chromosomal syndromes and genetic conditions, as it has been argued that this subset of children are uniquely complex.^{43,44} Proportion estimates of CHDs associated with a chromosomal abnormality vary widely from 9% to 18%, depending on study location, inclusion criteria for children with CHDs, and overall study population.⁴⁵ We are the first to report on the prevalence of other chromosomal abnormalities such as 22q11.2

deletion (DiGeorge Syndrome), trisomy 21 (Down syndrome), trisomy 18 and trisomy 13, that historically have been excluded in population estimates.⁴⁵

Our analysis has several notable advances compared with previous prevalence estimates. First, the prevalence analysis represent CHD children at the time of their first healthcare encounter, not at the time of their first CHD diagnosis. This approach allowed us to evaluate CHD longitudinally and include children who may have been diagnosed before the study or those who present after infancy.^{10,46,47} Many previously published studies identified children with CHD by active case-finding, or simply limited the population to newborns and infants. These methodologies may result in underestimation of less severe lesions more likely to manifest after the first year of life.^{48,49} Second, we included all CHD diagnoses – regardless of setting, lesion severity, or surgical intervention. Many frequently cited CHD prevalence studies restricted analyses to only simple lesions, only severe lesions, or only CHD diagnoses requiring surgery.^{1,3,6,9,49,50} Third, we incorporated many new sources of data and applied validated algorithms to estimate comorbidity burden. Finally, a fundamental strength of our data was the ability to longitudinally include children of all insurance types, throughout eight consecutive years.

The present study has several limitations. First, identification of children with CHD was made by administrative diagnoses codes ICD-9-CM and ICD-10-CM, thus both inadvertently excluding and including patients who were either misdiagnosed or miscoded. In addition, administrative codes often lack greater specificity compared with clinical nomenclature.³⁸ Second, and relatedly, although subtyping of CHD is expanded in our dataset, it has yet to capture the full spectrum of disease. This is particularly true for defects that have a wide variety of presentations and severities. Third, we did not modify our analysis based on assumptions of spontaneous regression or surgical repair (iatrogenic regression). While children with small or

simple CHD lesions may not require immediate or lifelong treatment, studies have shown small lesions are commonly associated with health complications requiring lifelong care.^{1,51-53}

CONCLUSION

Knowledge of prevalence patterns of CHDs is important for improving our understanding of the magnitude of the disease, in both the population and extent of care necessary for children with CHD. An accurate assessment of the absolute and relative burden of CHD, as well as related comorbidities, on a population-based scale will support appropriate allocation of resources for diagnosis, care coordination, care access and cost-effective treatment strategies.

Acknowledgments

None

Sources of Funding

American Heart Association Congenital Heart Defect Pre-Doctoral Fellowship Award

Disclosures

Authors have no disclosures to report.

References

1. Bjornard K, Riehle-Colarusso T, Gilboa SM, Correa A. Patterns in the prevalence of congenital heart defects, metropolitan Atlanta, 1978 to 2005. *Birth Defects Res A Clin Mol Teratol*. Feb 2013;97(2):87-94. doi:10.1002/bdra.23111
2. Krasuski RA, Bashore TM. Congenital Heart Disease Epidemiology in the United States: Blindly Feeling for the Charging Elephant. *Circulation*. Jul 12 2016;134(2):110-3. doi:10.1161/CIRCULATIONAHA.116.023370
3. Reller MD, Strickland MJ, Riehle-Colarusso T, Mahle WT, Correa A. Prevalence of congenital heart defects in metropolitan Atlanta, 1998-2005. *J Pediatr*. Dec 2008;153(6):807-13. doi:10.1016/j.jpeds.2008.05.059
4. van der Linde D, Konings EE, Slager MA, et al. Birth prevalence of congenital heart disease worldwide: a systematic review and meta-analysis. *J Am Coll Cardiol*. Nov 15 2011;58(21):2241-7. doi:10.1016/j.jacc.2011.08.025
5. van der Bom T, Zomer AC, Zwinderman AH, Meijboom FJ, Bouma BJ, Mulder BJ. The changing epidemiology of congenital heart disease. *Nat Rev Cardiol*. Jan 2011;8(1):50-60. doi:10.1038/nrcardio.2010.166
6. Hoffman JIE, Kaplan S. The incidence of congenital heart disease. *Journal of the American College of Cardiology*. 2002;39(12):1890-1900. doi:10.1016/s0735-1097(02)01886-7
7. Hoffman JI. Natural history of congenital heart disease. Problems in its assessment with special reference to ventricular septal defects. *Circulation*. Jan 1968;37(1):97-125. doi:10.1161/01.cir.37.1.97
8. Jacobs JP, Franklin RCG, Béland MJ, et al. Nomenclature for Pediatric and Congenital Cardiac Care: Unification of Clinical and Administrative Nomenclature - The 2021 International Paediatric and Congenital Cardiac Code (IPCCC) and the Eleventh Revision of the International Classification of Diseases (ICD-11). *World J Pediatr Congenit Heart Surg*. Sep 2021;12(5):E1-e18. doi:10.1177/21501351211032919
9. Marelli AJ, Ionescu-Iltu R, Mackie AS, Guo L, Dendukuri N, Kaouache M. Lifetime prevalence of congenital heart disease in the general population from 2000 to 2010. *Circulation*. Aug 26 2014;130(9):749-56. doi:10.1161/CIRCULATIONAHA.113.008396
10. Global, regional, and national burden of congenital heart disease, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet Child Adolesc Health*. Mar 2020;4(3):185-200. doi:10.1016/s2352-4642(19)30402-x
11. Warnes CA, Williams RG, Bashore TM, et al. ACC/AHA 2008 guidelines for the management of adults with congenital heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Develop Guidelines on the Management of Adults With Congenital Heart Disease). Developed in Collaboration With the American Society of Echocardiography, Heart Rhythm Society, International Society for Adult Congenital Heart Disease, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *J Am Coll Cardiol*. Dec 2 2008;52(23):e143-e263. doi:10.1016/j.jacc.2008.10.001
12. Oster ME, Riehle-Colarusso T, Simeone RM, et al. Public health science agenda for congenital heart defects: report from a Centers for Disease Control and Prevention

experts meeting. *Journal of the American Heart Association*. Aug 28

2013;2(5):e000256. doi:10.1161/JAHA.113.000256

13. Ionescu-Iltu R, Mackie AS, Abrahamowicz M, et al. Valvular operations in patients with congenital heart disease: increasing rates from 1988 to 2005. *Ann Thorac Surg*. Nov 2010;90(5):1563-9. doi:10.1016/j.athoracsur.2010.07.017

14. Simeone RM, Oster ME, Cassell CH, Armour BS, Gray DT, Honein MA. Pediatric inpatient hospital resource use for congenital heart defects. *Birth Defects Res A Clin Mol Teratol*. Dec 2014;100(12):934-43. doi:10.1002/bdra.23262

15. Russo CA, Elixhauser A. Hospitalizations for Birth Defects, 2004: Statistical Brief #24. *Healthcare Cost and Utilization Project (HCUP) Statistical Briefs*. Agency for Healthcare Research and Quality (US); 2006.

16. Chowdhury D, Johnson JN, Baker-Smith CM, et al. Health Care Policy and Congenital Heart Disease: 2020 Focus on Our 2030 Future. *Journal of the American Heart Association*. Oct 19 2021;10(20):e020605. doi:10.1161/JAHA.120.020605

17. Gilboa SM, Devine OJ, Kucik JE, et al. Congenital Heart Defects in the United States: Estimating the Magnitude of the Affected Population in 2010. *Circulation*. Jul 12 2016;134(2):101-9. doi:10.1161/CIRCULATIONAHA.115.019307

18. Oster ME, Riser AP, Andrews JG. Comorbidities among Young Adults with Congenital Heart Defects: Results from the Congenital Heart Survey to Recognize Outcomes, Needs and well-beinG - Arizona, Arkansas, and Metropolitan Atlanta, 2016 - 2019. *MMWR Morb Mortal Wkly Rep* 2021;70:197-201.

19. Perrailon MC, Liang R, Sabik LM, Lindrooth RC, Bradley CJ. The role of all-payer claims databases to expand central cancer registries: Experience from Colorado. *Health Serv Res*. Jun 2022;57(3):703-711. doi:10.1111/1475-6773.13901

20. Quality. AfHRa. All-payer claims databases.

18 States have, interest in developing an APCD.

21. Council APCD. Interactive State Report Map.

<https://www.apcdouncil.org/state/map>

22. Smith P. Primary care in children with congenital heart disease. *J Pediatr Nurs*. Oct 2001;16(5):308-19. doi:10.1053/jpdn.2001.26572

23. Richards AA, Garg V. Genetics of congenital heart disease. *Curr Cardiol Rev*. May 2010;6(2):91-7. doi:10.2174/157340310791162703

24. Edwards JJ, Gelb BD. Genetics of congenital heart disease. *Curr Opin Cardiol*. May 2016;31(3):235-41. doi:10.1097/HCO.0000000000000274

25. Feudtner C, Feinstein JA, Zhong W, Hall M, Dai D. Pediatric complex chronic conditions classification system version 2: updated for ICD-10 and complex medical technology dependence and transplantation. *BMC Pediatr*. Aug 8 2014;14:199. doi:10.1186/1471-2431-14-199

26. CEPQM. *Children with Disabilities Algorithm (CWDA): A Claims-based Tool for Identifying Children with Disabilities*. 2015.

27. Rural-Urban Commuting Area Codes.

28. Hailu A, Wasserman C. Guidelines for Using Rural-Urban Classification Systems for Community Health Assessment

<http://www.doh.wa.gov/DataandStatisticalReports/DataGuidelines>.

29. Bureau UC. American Community Survey. <https://www.census.gov/programs-surveys/acs/>

30. Stout KK, Daniels CJ, Aboulhosn JA, et al. 2018 AHA/ACC Guideline for the Management of Adults With Congenital Heart Disease: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. Apr 2 2019;73(12):1494-1563. doi:10.1016/j.jacc.2018.08.1028
31. Hennekens C. *Measures disease frequency and association*. Epidemiology in medicine. 1987.
32. Cresti A, Giordano R, Koestenberger M, et al. Incidence and natural history of neonatal isolated ventricular septal defects: Do we know everything? A 6-year single-center Italian experience follow-up. *Congenit Heart Dis*. Jan 2018;13(1):105-112. doi:10.1111/chd.12528
33. Bernstein D. Evaluation of the cardiovascular system. *Nelson textbook of pediatrics*. 2004;17:1479-99.
34. Lui GK, Saidi A, Bhatt AB, et al. Diagnosis and Management of Noncardiac Complications in Adults With Congenital Heart Disease: A Scientific Statement From the American Heart Association. *Circulation*. Nov 14 2017;136(20):e348-e392. doi:10.1161/cir.0000000000000535
35. Khairy P, Ionescu-Iltu R, Mackie AS, Abrahamowicz M, Pilote L, Marelli AJ. Changing mortality in congenital heart disease. *J Am Coll Cardiol*. Sep 28 2010;56(14):1149-57. doi:10.1016/j.jacc.2010.03.085
36. Nasr VG, Faraoni D, Valente AM, DiNardo JA. Outcomes and Costs of Cardiac Surgery in Adults with Congenital Heart Disease. *Pediatric cardiology*. Oct 2017;38(7):1359-1364. doi:10.1007/s00246-017-1669-7
37. Marino BS, Lipkin PH, Newburger JW, et al. Neurodevelopmental outcomes in children with congenital heart disease: evaluation and management: a scientific statement from the American Heart Association. *Circulation*. Aug 28 2012;126(9):1143-72. doi:10.1161/CIR.0b013e318265ee8a
38. Razzaghi H, Oster M, Reefhuis J. Long-term outcomes in children with congenital heart disease: National Health Interview Survey. *J Pediatr*. Jan 2015;166(1):119-24. doi:10.1016/j.jpeds.2014.09.006
39. Gonzalez VJ, Kimbro RT, Cutitta KE, et al. Mental Health Disorders in Children With Congenital Heart Disease. *Pediatrics*. Feb 2021;147(2)doi:10.1542/peds.2020-1693
40. Triedman JK, Newburger JW. Trends in Congenital Heart Disease: The Next Decade. *Circulation*. Jun 21 2016;133(25):2716-33. doi:10.1161/CIRCULATIONAHA.116.023544
41. Verrall CE, Blue GM, Loughran-Fowlds A, et al. 'Big issues' in neurodevelopment for children and adults with congenital heart disease. *Open Heart*. 2019;6(2):e000998. doi:10.1136/openhrt-2018-000998
42. Kovacs AH, Brouillette J, Ibeziako P, et al. Psychological Outcomes and Interventions for Individuals With Congenital Heart Disease: A Scientific Statement From the American Heart Association. *Circ Cardiovasc Qual Outcomes*. Aug 2022;15(8):e000110. doi:10.1161/HCQ.0000000000000110
43. Pierpont ME, Basson CT, Benson DW, Jr., et al. Genetic basis for congenital heart defects: current knowledge: a scientific statement from the American Heart Association Congenital Cardiac Defects Committee, Council on Cardiovascular Disease

- in the Young: endorsed by the American Academy of Pediatrics. *Circulation*. Jun 12 2007;115(23):3015-38. doi:10.1161/circulationaha.106.183056
44. Roskes EJ, Boughman JA, Schwartz S, Cohen MM. Congenital cardiovascular malformations (CCVM) and structural chromosome abnormalities: a report of 9 cases and literature review. *Clin Genet*. Sep 1990;38(3):198-210. doi:10.1111/j.1399-0004.1990.tb03571.x
45. Hartman RJ, Rasmussen SA, Botto LD, et al. The contribution of chromosomal abnormalities to congenital heart defects: a population-based study. *Pediatric cardiology*. Dec 2011;32(8):1147-57. doi:10.1007/s00246-011-0034-5
46. Grabitz RG, Joffres MR, Collins-Nakai RL. Congenital heart disease: incidence in the first year of life. The Alberta Heritage Pediatric Cardiology Program. *Am J Epidemiol*. Aug 1988;128(2):381-8. doi:10.1093/oxfordjournals.aje.a114978
47. Hoffman JI. *Incidence, mortality and natural history*. Paediatric Cardiology. Churchill Livingstone; 2001.
48. Riehle-Colarusso T, Strickland MJ, Reller MD, et al. Improving the quality of surveillance data on congenital heart defects in the metropolitan Atlanta congenital defects program. *Birth defects research Part A, Clinical and molecular teratology*. Nov 2007;79(11):743-53. doi:10.1002/bdra.20412
49. Marelli AJ, Mackie AS, Ionescu-Ittu R, Rahme E, Pilote L. Congenital heart disease in the general population: changing prevalence and age distribution. *Circulation*. Jan 16 2007;115(2):163-72. doi:10.1161/CIRCULATIONAHA.106.627224
50. Bound JP, Logan WF. Incidence of congenital heart disease in Blackpool 1957-1971. *Br Heart J*. 1977;39:445-50.
51. Wren C, O'Sullivan JJ. Survival with congenital heart disease and need for follow up in adult life. *Heart*. Apr 2001;85(4):438-43. doi:10.1136/heart.85.4.438
52. Botto LD, Correa A. Decreasing the burden of congenital heart anomalies: an epidemiologic evaluation of risk factors and survival. *Progress in Pediatric Cardiology*. 2003;18(2):111-121. doi:10.1016/s1058-9813(03)00084-5
53. Oyen N, Poulsen G, Boyd HA, Wohlfahrt J, Jensen PK, Melbye M. Recurrence of congenital heart defects in families. *Circulation*. Jul 28 2009;120(4):295-301. doi:10.1161/CIRCULATIONAHA.109.857987

Table 1. Characteristics of children with and without congenital heart disease in Colorado, 2012 – 2019

	CHD N= 30,512 (%)	NON-CHD N = 1,565,815 (%)	P value
Age ¹			
Newborn	12,527 (41.1)	301,746 (19.7)	
Infant	4,647 (15.2)	90,632 (5.9)	
Child	7,238 (23.7)	591,138 (38.5)	
Youth	6,100 (20.0)	552,299 (36.0)	<0.001
Sex ²			
Male	15,843 (51.9)	767,356 (49.9)	<0.001
Primary insurer			
Medicaid	21,915 (71.8)	897,634 (58.5)	
Commercial	8,597 (28.2)	638,181 (41.6)	<0.001
Rurality ³			
Urban	26,957 (88.4)	1,329,187 (86.6)	<0.001
Region			
Eastern	947 (3.1)	45,314 (3.0)	
Front Range	21,827 (71.5)	1,055,938 (68.8)	
Mountain	5,148 (16.9)	286,072 (18.6)	
Southern	1,427 (4.7)	65,602 (4.3)	
Western	1,163 (3.8)	82,889 (5.4)	<0.001
Median household income ⁴			
Q1	6,965 (22.8)	312,453 (20.3)	
Q2	5,579 (18.3)	268,015 (17.5)	
Q3	6,178 (20.3)	304,892 (19.9)	
Q4	6,035 (19.8)	316,775 (20.6)	
Q5	5,755 (18.9)	333,680 (21.7)	<0.001
Diagnosis severity category ⁵			
Simple	19,959 (65.4)		
Moderately Complex	6,068 (19.9)		
Complex	2,618 (8.6)		
Other	1,867 (6.1)	N/A	
Complex chronic conditions ⁶	16,878 (55.3)	N/A	
Genetic abnormalities ⁷	3,023 (10.0)	N/A	
Disability ⁸	11,749 (38.5)	N/A	

1. Age at first encounter categorized as: Newborn (0 – 30 days old); Infant (31 – 365 days old); Child (366 days – 10 years old); Youth (>10– 18 years old)
2. Gender categorized as male or female.
3. Rurality defined by the 2010 Rural-Urban Commuting Area Codes
4. Median household income defined by the American Community Survey (2019) U.S. Census. 1st Quintile: \$24,427 – 51,461; 2nd Quintile: \$51,484 – 63,318; 3rd Quintile: \$63,745 – 74,648; 4th Quintile: \$74,809 – 94,875; 5th Quintile: \$95,050 – 147,421.
5. Severity category defined by ACHD Anatomy and Physiology Classification
6. Coexisting chronic conditions defined by the Pediatric Complex Chronic Conditions (PCCC) algorithm
7. Genetic abnormalities: Down syndrome, trisomy 13, trisomy 18, Turner syndrome, DiGeorge syndrome, Alagille syndrome, Noonan syndrome, Holt-Oram syndrome, Williams syndrome, Kabuki syndrome, VATER/VACTERL association, CHARGE association, Jacobsen syndrome, and Williams-Beuren syndrome
8. Disabilities defined by the Children with Disabilities Algorithm (CWDA)

Table 2. Most common diagnoses for children with congenital heart disease, by lesion category

Lesion Category	N (%)
Atrial septal defect	24.4
Ventricular septal defect	11.3
Patent ductus arteriosus	10.6
Aortic insufficiency	10.0
Pulmonary artery abnormality	5.9
Other congenital anomalies of the heart	18.2

Table 3. Prevalence of children with congenital heart disease in Colorado that have a concurrent genetic abnormality

N = 3,480 (11%); 405 (with more than 1 genetic diagnosis)

Genetic abnormality	N (%)
	3,480 (11.4)
Trisomy 21 (Down syndrome)	1,557 (5.1)
Kabuki syndrome	866 (2.8)
DiGeorge syndrome (22q11.2 deletion)	206 (0.7)
Noonan	205 (0.7)
VATER/VACTERL	201 (0.7)
Turner syndrome	166 (0.5)
Jacobsen	119 (0.4)
Trisomy 18	52 (0.2)
Alagille syndrome	**
Williams-Beuren	**
Trisomy 13 (Patau's syndrome)	**
Holt-Oram syndrome	**
Williams	**
CHARGE association	**

** < 0.1% of CHD population

Table 4. Prevalence of children with congenital heart disease that have concurrent complex chronic conditions

Table 5 Prevalence of children with congenital heart disease that have concurrent disability

Disability	N (%)
	11,749 (38.5)
Diseases of the Nervous System and Sense Organs	7,040 (23.1)
Mental Health Disorders	5,085 (16.7)
Diseases of the Respiratory System	3,361 (11.0)
Congenital Anomalies	3,357 (11.0)
Certain Conditions Originating in the Perinatal Period	2,514 (8.2)
Diseases of the Circulatory System	1,873 (6.1)
Diseases of the Digestive System	1,863 (6.1)
Diseases of the Genitourinary System	1,393 (4.6)
Supplemental Classification of Factors Influencing Health Status	687 (2.3)
Systems, Signs and Ill-Defined Conditions	**
Endocrine, Nutritional and Metabolic Diseases, and Immunity Disorders	**
Injury and Poisoning	**
Neoplasms	**
Infectious and Parasitic Diseases	**


** < 1%

Figure 1. Derivation of cohort

Figure legend. Flow diagram of study subjects.

Figure 2. Population density map of children residing in CO, 2012 – 2019

Figure legend. The population density map illustrates the aggregate CO pediatric population at the zip-code tabulation area (ZCTA) level. The population is further categorized by quintile.



Supplementary
Table 1. CHD diagnoses and ICD-9/10 Mappings

ICD9	ICD9 DX Description	ICD10	ICD10 DX Description
394	Diseases of mitral valve	I34.0	Nonrheumatic mitral (valve) insufficiency
395	Diseases of aortic valve	I34.8	Other nonrheumatic mitral valve disorders
396.0	Mitral/aortic stenosis	I37.0	Nonrheumatic pulmonary valve stenosis
424.0	Mitral valve disorders	I37.8	Other nonrheumatic pulmonary valve disorders
424.0	Mitral valve disorders	Q21.3	Tetralogy of fallot
424.3	Pulmonary valve disorders	Q28.8	Other specified congenital malformations of circulatory system
745.2	Tetralogy of fallot	Q20.0	Common arterial trunk
746	Congenital anomalies of heart.	Q20.3	Discordant ventriculoarterial connection
747.89	Circulatory anomaly NEC	Q20.1	Double outlet right ventricle
745.0	Common truncus	Q20.5	Discordant atrioventricular connection
745.10	Complete transposition of great vessels	Q20.3	Discordant ventriculoarterial connection
745.11	Double outlet right ventricle	Q20.8	Other congenital malformations of cardiac chambers and connections
745.12	Corrected transposition of great vessels	Q20.4	Double inlet ventricle
745.19	Other transposition of great vessels	Q21.0	Ventricular septal defect
745.19	Other transposition of great vessels	Q21.1	Atrial septal defect
745.3	Common ventricle	Q21.2	Atrioventricular septal defect
745.4	Ventricular septal defect	Q21.2	Atrioventricular septal defect
745.5	Ostium secundum type atrial septal defect	Q21.2	Atrioventricular septal defect
745.60	Endocardial cushion defect, unspecified type	Q20.8	Other congenital malformations of cardiac chambers and connections
745.61	Ostium primum defect	Q21.8	Other congenital malformations of cardiac septa
745.69	Other endocardial cushion defects	Q21.9	Congenital malformation of cardiac septum, unspecified
745.8	Other bulbus cordis anomalies and anomalies of cardiac septal closure	Q22.3	Other congenital malformations of pulmonary valve
745.8	Other bulbus cordis anomalies and anomalies of cardiac septal closure	Q22.0	Pulmonary valve atresia
745.9	Unspecified defect of septal closure	Q22.1	Congenital pulmonary valve stenosis
746.00	Congenital pulmonary valve anomaly, unspecified	Q22.2	Congenital pulmonary valve insufficiency
746.01	Atresia of pulmonary valve, congenital	Q22.9	Congenital malformation of tricuspid valve, unspecified
746.02	Stenosis of pulmonary valve, congenital	Q22.5	Ebstein's anomaly
746.09	Other congenital anomalies of pulmonary valve	Q23.0	Congenital stenosis of aortic valve

746.1	Tricuspid atresia and stenosis, congenital	Q23.1	Congenital insufficiency of aortic valve
746.2	Ebstein's anomaly	I34.2	Nonrheumatic mitral (valve) stenosis
746.3	Congenital stenosis of aortic valve	Q23.3	Congenital mitral insufficiency
746.4	Congenital insufficiency of aortic valve	Q23.4	Hypoplastic left heart syndrome
746.5	Congenital mitral stenosis	Q24.4	Congenital subaortic stenosis
746.6	Congenital mitral insufficiency	Q24.2	Cor triatriatum
746.7	Hypoplastic left heart syndrome	Q24.3	Pulmonary infundibular stenosis
746.81	Subaortic stenosis	Q24.8	Other specified congenital malformations of heart
746.82	Cor triatriatum	Q24.5	Malformation of coronary vessels
746.83	Infundibular pulmonic stenosis	Q24.0	Dextrocardia
746.84	Obstructive anomalies of heart, not elsewhere classified	Q24.1	Levocardia
746.85	Coronary artery anomaly	Q24.8	Other specified congenital malformations of heart
746.87	Malposition of heart and cardiac apex	Q23.8	Other congenital malformations of aortic and mitral valves
746.87	Malposition of heart and cardiac apex	Q24.8	Other specified congenital malformations of heart
746.87	Malposition of heart and cardiac apex	Q20.9	Congenital malformations of cardiac chambers and connections, unspecified
746.89	Other specified congenital anomalies of heart	Q24.9	Congenital malformation of heart, unspecified
746.89	Other specified congenital anomalies of heart	Q25.0	Patent ductus arteriosus
746.9	Unspecified congenital anomaly of heart	Q25.1	Coarctation of aorta
746.9	Unspecified congenital anomaly of heart	Q25.40	Congenital malformation of aorta unspecified
747.0	Patent ductus arteriosus	Q25.45	Double aortic arch
747.10	Coarctation of aorta (preductal) (postductal)	Q25.46	Tortuous aortic arch
747.20	Anomaly of aorta, unspecified	Q25.47	Right aortic arch
747.21	Anomalies of aortic arch	Q25.29	Other atresia of aorta
747.21	Anomalies of aortic arch	Q25.3	Supravalvular aortic stenosis
747.21	Anomalies of aortic arch	Q25.41	Absence and aplasia of aorta
747.22	Atresia and stenosis of aorta	Q25.42	Hypoplasia of aorta
747.22	Atresia and stenosis of aorta	Q25.43	Congenital aneurysm of aorta
747.29	Other anomalies of aorta	Q25.44	Congenital dilation of aorta
747.29	Other anomalies of aorta	Q25.48	Anomalous origin of subclavian artery
747.29	Other anomalies of aorta	Q25.49	Other congenital malformations of aorta
747.29	Other anomalies of aorta	Q25.5	Atresia of pulmonary artery
747.29	Other anomalies of aorta	Q25.71	Coarctation of pulmonary artery

747.29	Other anomalies of aorta	Q25.72	Congenital pulmonary arteriovenous malformation
747.31	Pulmonary artery coarctation and atresia	Q25.6	Stenosis of pulmonary artery
747.31	Pulmonary artery coarctation and atresia	Q25.79	Other congenital malformations of pulmonary artery
747.32	Pulmonary arteriovenous malformation	Q26.9	Congenital malformation of great vein, unspecified
747.39	Other anomalies of pulmonary artery and pulmonary circulation	Q26.2	Total anomalous pulmonary venous connection
747.39	Other anomalies of pulmonary artery and pulmonary circulation	Q26.3	Partial anomalous pulmonary venous connection
747.40	Anomaly of great veins, unspecified	Q26.0	Congenital stenosis of vena cava
747.41	Total anomalous pulmonary venous connection	Q26.1	Persistent left superior vena
747.42	Partial anomalous pulmonary venous connection	Q26.8	Other congenital malformations of great veins
747.49	Other anomalies of great veins	Q28.8	Other specified congenital anomalies of circulatory system
747.49	Other anomalies of great veins	Z87.74	Personal history of (corrected) congenital malformations of heart and circulatory system
747.49	Other anomalies of great veins	Q25.21	Interruption of aortic arch
747.89	Other specified congenital anomalies of circulatory system		
V13.65	Personal history of (corrected) congenital malformations of heart and circulatory system		

Table 2. Disease Severity Categories and ICD-9/10 Mappings

ICD9	ICD-9 DESCRIPTION	ICD10	ICD-10 DESCRIPTION
SIMPLE			
746.0	Anomalies of pulmonary valve	Q22.1	Congenital pulmonary valve stenosis
746.00	Pulmonary valve anomaly, unspecified	Q25.0	Patent ductus arteriosus
746.02	Stenosis, congenital	Q21.0	Ventricular septal defect
747	Patent ductus arteriosus	Q21.1	Atrial septal defect
745.4	Ventricular septal defect	Q26.0	Congenital stenosis of vena cava
745.5	Ostium secundum type atrial septal defect	Q26.8	Other congenital malformations of great veins
747.4	Anomalies of great veins*	Q26.9	Congenital malformation of great vein, unspecified
747.40	Anomaly of great veins, unspecified*		
747.49	Other anomalies of great veins*		
424.3	Pulmonary valve disorder		
MODERATELY COMPLEX			
746.2	Ebstein's anomaly	Q22.2	Congenital pulmonary valve insufficiency
746.5	Congenital mitral stenosis	Q22.3	Other congenital malformations of pulmonary valve
746.6	Congenital mitral insufficiency	Q22.5	Ebstein's anomaly
746.8	Other specified anomalies of heart	Q22.8	Other congenital malformations of tricuspid valve
746.81	Subaortic stenosis	Q22.9	Congenital malformation of tricuspid valve, unspecified
746.83	Infundibular pulmonic stenosis	Q23.2	Congenital mitral stenosis
746.84	Obstructive anomalies of heart, NEC	Q23.3	Congenital mitral insufficiency
747.1	Coarctation of aorta	Q25.1	Coarctation of aorta
747.1	Coarctation of aorta (preductal) (postductal)	Q25.3	Supravalvular aortic stenosis
745.2	Tetralogy of Fallot	Q21.3	Tetralogy of Fallot
746.3	Congenital stenosis of aortic valve	Q23.0	Congenital stenosis of aortic valve
746.4	Congenital insufficiency of aortic valve	Q23.1	Congenital insufficiency of aortic valve
747.41	Total anomalous pulmonary venous connection	Q26.2	Total anomalous pulmonary venous connection
745.6	Endocardial cushion defects	Q21.2	Atrioventricular septal defect
745.60	Endocardial cushion defect, unspecified type	Q21.4	Aortopulmonary septal defect
745.61	Ostium primum defect	Q21.8	Other congenital malformations of cardiac septa
745.69	Other AV canal*	Q21.9	Congenital malformation of cardiac septum, unspecified
745.8	Other	Q26.3	Partial anomalous pulmonary venous connection
745.9	Unspecified defect of septal closure	Q24	Other congenital malformations of heart

747.42	Partial anomalous pulmonary venous connection	Q24.2	Cor triatriatum
746.82	Cor triatriatum	Q24.3	Pulmonary infundibular stenosis
746.85	Coronary artery anomaly	Q24.4	Congenital subaortic stenosis
746.86	Congenital heart block	Q24.5	Malformation of coronary vessels
746.87	Malposition of heart and cardiac apex	Q25.6	Stenosis of pulmonary artery
747.3	Anomalies of pulmonary artery	Q25.7	Other congenital malformations of pulmonary artery
424	Mitral valve disorder	Q25.71	Coarctation of pulmonary artery
		Q25.72	Congenital pulmonary arteriovenous malformation
		Q25.79	Other congenital malformations of pulmonary artery
		Q25.8	Other congenital malformations of other great arteries
		Q25.9	Congenital malformation of great arteries, unspecified
		Q26.4	Anomalous pulmonary venous connection, unspecified
COMPLEX			
747.2	Other anomalies of aorta	Q20.0	Common arterial trunk
747.20	Anomaly of aorta, unspecified	Q20.1	Double outlet right ventricle
747.21	Anomalies of aortic arch	Q20.2	Double outlet left ventricle
745.0	Common truncus	Q20.3	Discordant ventriculoarterial connection
745.1	Transposition of great vessels	Q22.0	Pulmonary valve atresia
745.10	Complete transposition of great vessels	Q23.8	Other congenital malformations of aortic and mitral valves
745.11	Double outlet right ventricle	Q23.9	Congenital malformation of aortic and mitral valves, unspecified
745.12	Corrected transposition of great vessels	Q25.2	Atresia of aorta
745.19	Transposition - Other*	Q20.4	Double inlet ventricle
746.01	Atresia, congenital	Q22.4	Congenital tricuspid stenosis
745.3	Common ventricle	Q22.6	Hypoplastic right heart syndrome
745.7	Cor biloculare	Q23.4	Hypoplastic left heart syndrome
746.1	Tricuspid atresia and stenosis, congenital	Q20.5	Discordant atrioventricular connection
746.7	Hypoplastic left heart syndrome	Q20.6	Isomerism of atrial appendages
747.11	Interruption of aortic arch	Q20.8	Other congenital malformations of cardiac chambers and connections
747.22	Atresia and stenosis of aorta	Q20.9	Congenital malformation of cardiac chambers and connections, unspecified
		Q25.21	Interruption of aortic arch
		Q25.29	Other atresia of aorta
		Q25.40	Congenital malformation of aorta unspecified
		Q25.41	Absence and aplasia of aorta

		Q25.42	Hypoplasia of aorta
		Q25.49	Other congenital malformations of aorta
UNCATEGORIZED / OTHER			
746.09	Other	Q24.8	Other specified congenital malformations of heart
746.89	Other	Q24.9	Congenital malformation of heart, unspecified
746.90	Unspecified anomaly of heart	I34.2	Eisenmenger's syndrome
747.00	Other congenital anomalies of circulatory system	Q24.1	Levocardia
		Q25.43	Congenital aneurysm of aorta
		Q25.44	Congenital dilation of aorta
		Q25.46	Tortuous aortic arch
		Q25.47	Right aortic arch
		Q25.48	Anomalous origin of subclavian artery

Appendix Table 3. Lesion categories and ICD-9/10 Mappings

	Diagnosis	Codes
1	Non-specific mitral valve disease	394, 424, I34.8
2	Combined mitral and aortic disease	396, Q23.8
3	Mitral insufficiency	I34.0, 746.6, Q23.3
4	Non-specific pulmonary valve disease	424.3, 746.0, 746.09, I37.8, Q22.3
5	Pulmonary stenosis	I37.0, 746.02, Q22.1
6	Tetralogy of Fallot	745.2, Q21.3
7	Truncus arteriosus	745, Q20.0
8	Transposition of the great vessels	745.10, 745.19, 745.3, Q20.3, Q20.4
9	Other single ventricle disease (DORV, DILV)	745.11, 745.3, Q20.1, Q20.4
10	Corrected transposition of the great vessels	745.12, Q20.5
11	Ventricular septal defect	745.4, 745.8, 745.9, Q21.0, Q21.8, Q21.9
12	Atrial septal defect	745.5, Q21.1
13	Atrioventricular septal defect	745.60, 745.61, 745.69, Q21.2
14	Pulmonary atresia	746.01, Q22.0, Q25.5,
15	Pulmonary insufficiency	Q22.2
16	Tricuspid atresia/stenosis	746.1, Q22.9
17	Ebstein's anomaly	746.2, Q22.5
18	Aortic valve stenosis	746.3, Q23.0
19	Aortic insufficiency	746.4, Q23.1
20	Mitral stenosis	746.5, I34.2
21	Non-specific aortic valve disease	395
22	Hypoplastic left heart syndrome	746.7, Q23.4
23	Subaortic stenosis	746.81, Q24.4
24	Cor triatriatum	746.82, Q24.2
25	Pulmonary infundibular stenosis	746.83, Q24.3
26	Coronary artery anomaly	746.85, Q24.5
27	Patent ductus arteriosus	747.0, Q25.0
28	Coarctation of the aorta	747.10, Q25.1
29	Other aortic arch anomaly	747.20, 747.21, 747.29, Q25.40, Q25.45, Q25.46, Q25.47, Q25.21, Q25.42, Q25.43, Q25.44, Q25.48, Q25.49
30	Aortic atresia	747.22, Q25.29, Q25.41
31	Supravalvar aortic stenosis	Q25.3
32	Pulmonary artery abnormality	747.31, 747.32, 747.39, Q25.71, Q25.72, Q25.6, Q25.79
33	Total anomalous pulmonary venous return	747.41, Q26.2

34	Partial anomalous pulmonary venous return	747.42, Q26.3
35	Other congenital anomalies of the heart	746, 747.89, 746.84, 746.87, 746.89, 746.9, 747.40, 747.49, 747.89, V13.65, Q28.8, Q20.8, Q20.9, Q24.8, Q24.0, Q24.1, Q24.8, Q24.9, Q26.9, Q26.0, Q26.1, Q26.8, Q28.8, Z87.74

Colorado APCD children cohort
N=1,953,612



Patients 18 years old or younger on Jan 1, 2012,
without missing DOB, with a Colorado home zip
code, zip code matched to RUCA and who were
alive on Jan 1, 2012
N=1,853,897



Excluded
N=99,715



Patients with at least one claim from
Jan 1, 2012 – Dec 31, 2019
N=1,535,816





Excluded
N=318,081



Patients with at least one CHD diagnosis code
N=30,512 (1.9%)

Denver Metropolitan area



Colorado, 2012 - 2019
Pediatric CHD Population

- 1 - 32
- 33 - 92
- 93 - 183
- 184 - 324
- 325 - 572