

1 **Epidemiology of cruciate ligament surgery in Japan: A retrospective cohort study**
2 **from 2014 to 2019**

3

4 Shota Uchino^{1,2*}, Masataka Taguri³

5

6

7 ¹ Department of Data Science, Graduate School of Data Science, Yokohama City University,
8 Yokohama, Japan

9 ² REHASAKU Co., Ltd, Tokyo, Japan

10 ³ Department of Health Data Science, Tokyo Medical University, Tokyo, Japan

11

12 *Corresponding author:

13 Email: ucchi.no.shota@gmail.com

14

15

16

17 **Abstract**

18 Understanding the incidence and trends of cruciate ligament (CL) surgeries in Japan is crucial for
19 providing effective healthcare services. This study aimed to use open data available from the
20 National Database of Health Insurance Claims and Specific Health Checkups of Japan (NDB) to
21 determine the changes over time in CL surgeries and to analyze the characteristics of the Japanese
22 population in terms of sex and age. We retrospectively identified CL surgeries of the knee joint
23 registered from April 2014 to March 2020 using the NDB open data. Data on sex, age, and practice
24 were extracted to determine the number of cases per 100,000 population. Trends in the annual
25 incidence of CL surgeries were evaluated using Poisson regression analysis. In total, 112,686 CL
26 surgeries were performed from 2014 to 2019. Arthroscopic ligament reconstruction accounted for
27 98% of all the cases. The number of surgeries performed had increased significantly from 16,975 in
28 2014 to 19,735 in 2019 ($P < 0.001$). Overall, CL surgeries were most common in the 15–19 years age
29 group, followed by the 20–29 years age group. However, these trends differed between males and
30 females. The incidence of CL surgery in Japan has increased, and its characteristics vary by gender
31 and age. These characteristics have been identified not only in younger patients but also in middle-
32 aged and older patients. It would be valuable to further investigate the general patterns of CL surgery
33 in Japan.

34

35 **Introduction**

36 In recent years, the availability of medical and national big data has made it possible to determine the
37 incidence of ligament injuries and surgeries [1–9]. These studies have revealed epidemiological
38 models that allow for the generalization of disease incidence. They have also been used to improve
39 treatment strategies and health care services to enhance patients' quality of life.

40 In Japan, an epidemiological study using insurance records of middle and high school athletes
41 reported 30,458 anterior cruciate ligament (ACL) injuries (0.81 per 1000 athlete-years) in these
42 growing athletes over a 10-years from 2005 to 2014. Furthermore, the incidence rate of ACL injuries
43 was reported to be 2.8 times higher in female athletes than in male athletes [10]. This previous study
44 determined the incidence of ACL injuries using a large and long-term insurance dataset; however, it
45 only focused on populations considered at high risk for ACL injury, and the incidence and the post-
46 injury response of other populations remain unknown.

47 To understand the incidence and trends of the disease in the Japanese population, a larger sample
48 size should be examined. Epidemiological studies using the National Database of Health Insurance
49 Claims and Specific Health Checkups of Japan (NDB) established by the Ministry of Health, Labour
50 and Welfare (MHLW) and NDB open data, which is a more generalized compilation of the NDB, have
51 been actively conducted in the field of locomotor system [11–13]. Epidemiological modeling from
52 these big data could lead to improved healthcare services.

53 This study aimed to elucidate trends in the number of cruciate ligament (CL) surgeries, sex, age
54 distribution, and characteristics over time using NDB open data from 2014 to 2019. We hypothesized
55 that the number of CL surgeries would increase over the years, and that there would be a higher
56 incidence of such surgeries among females.

57

58

59 **Materials and methods**

60 This population-based retrospective cohort study uses the NDB open data provided by the MHLW
61 from April 2014 to March 2020. These data are widely accessed for research purposes and have been
62 anonymized. Therefore, obtaining informed consent was not necessary.

63 We accessed the NDB open data website and obtained data on the "Number of calculations by sex,
64 and age group" for "operation (Code K)" [14–19]. There were four categories of the CL of the knee
65 joint: ligament tear suture (K074), arthroscopic ligament tear suture (K074-2), ligament reconstruction
66 (K079), and arthroscopic ligament reconstruction (K079-2). Data on the number of registered CL
67 surgeries were stratified by fiscal year, sex, and age (5-year interval). No detailed analyses were
68 performed if the number of registered CL surgeries had no specific values shown in <10 cases. The
69 sex and age data were obtained by calculating the number of surgeries per 100,000 population. The
70 calculation method utilized population estimates published by the Statistics Bureau of the Ministry of
71 Internal Affairs and Communications, based on population data as of October 1 of each year [20].

72 Changes in the number of surgeries on the CL of the knee joint over time were assessed using Poisson
73 regression, with the number of surgeries as the outcome variable and the year of surgery as the
74 explanatory variable. The supporting information file contains the details of the STROBE checklist
75 (S1 Table). The NDB open data is widely available for research purposes by the Ministry of Health,
76 Labour, and Welfare (MHLW). The data are compiled and published based on existing anonymized

77 NDB, and since there is no corresponding table, individuals cannot be identified. Therefore, ethical
78 approval was not required for this study.

79 Results

80 In total, 112,686 CL surgeries were performed from April 2014 to March 2020, with 109,925 (98%)
81 being arthroscopic ligament reconstruction. The remaining procedures included ligament
82 reconstruction and arthroscopic ligament tear suture, each accounting for approximately 1.0% of the
83 total number of surgeries performed, with very few cases of ligament tear suture (Table 1). The number
84 of arthroscopic ligament reconstruction increased significantly from 16,997 (18.3 cases/100,000
85 population) in 2014 to 19,774 (22.1 cases/100,000 population) in 2019, surpassing the number of all
86 other techniques ($P < 0.001$) (Fig 1).

87

88 **Table 1. Annual number of cruciate ligament surgeries in the NDB open data from 2014 to 2019.**

Category, year	2014	2015	2016	2017	2018	2019
Ligament tear suture	26	24	15	20	16	19
Arthroscopic ligament tear suture	225	204	228	267	213	227
Ligament reconstruction	289	229	230	209	169	151
Arthroscopic ligament reconstruction	16,997	17,507	18,129	18,241	19,277	19,774
Total	17,537	17,964	18,602	18,737	19,675	20,171

89

90 **Fig 1. Number of registered arthroscopic ligament reconstruction procedures per 100,000**

91 **population from 2014 to 2019.**

92

93 Due to low registration numbers, data for the 0–9 and >70 years age groups were incomplete. For the

94 10–69 years age group, the highest number of registrations was observed in the 15–19 years age group

95 (108.5 cases/100,000 population), followed by the 20–24 years age group (41.8 cases/100,000

96 population). We noted that registrations declined with age (Fig 2 and S2 and S3 Tables).

97 In terms of sex, females outnumbered males by a factor of 6.7 in the 10–14 years age group and by

98 1.6 in the 15–19 years age group. In contrast, males were 1.6 times more likely than females to be

99 enrolled in the 20–24 years age group and more than twice as likely as females to be enrolled in the

100 25–29 and 30–34 years age groups. Registration numbers were similar for other age groups (Fig 3 and

101 S3 Table).

102

103 **Fig 2. Number of registered arthroscopic ligament t reconstruction procedures per 100,000**

104 **population by age group from 2014 to 2019.**

105 **Fig 3. Number of registered arthroscopic ligament reconstruction procedures per 100,000**

106 **population.**

107

108 **Discussion**

109 This study investigated the number of registered CL surgeries, sex differences, and age characteristics
110 in Japan from 2014 to 2019 using the NDB open data. As hypothesized, there was a consistent increase
111 in CL surgery over the study period. Notably, most CL surgeries were arthroscopic ligament
112 reconstruction procedures, which constitutes a significant finding of this study. With the advancement
113 of medical technology, arthroscopic surgery, which enables a detailed examination of the interior of a
114 joint during surgery, has become increasingly prevalent [21–23]. This development may have led to
115 the rise in the number of arthroscopic procedures in Japan.

116 Arthroscopic ligament reconstruction was found to vary by sex and age. Overall, the incidence was
117 higher among males. Conversely, females often undergo surgery during their teenage years, potentially
118 due to the disparity in growth rates between males and females. Height growth in Japanese females
119 reaches its peak around 2 years earlier than in males [24]. Variations in growth rates may influence
120 the timing of surgery. However, it should be noted that CL surgery has been a subject of intense debate
121 regarding its potential impact on growth retardation [25,26]. Other factors, such as female's inability
122 to be active in sports at times due to childbirth, may be related to the sex difference. Another finding
123 was that the incidence of surgery among middle-aged and older adults had also increased. Injuries
124 resulting from sports activities, everyday sprains, and falls are common among middle-aged and older
125 adults [27]. This can be attributed to increased health awareness and higher physical activity levels.

126 Additionally, there is a higher risk of knee joint dysfunction and secondary injuries (such as meniscus
127 and cartilage damage) [28,29]. Good outcomes of CL surgery in middle-aged and older patients have
128 also been reported [30,31]. Consequently, many patients may choose surgery with the goal of
129 regaining their previous level of activity. In Japan, like in other countries, the indications for surgery
130 have likely expanded due to changes in individual lifestyles and advancements in medical technology.

131 This study had several limitations. First, it failed to provide details on the number of ACL and PCL
132 surgeries as a categorization of the CL surgeries. Second, tissue injuries other than those caused by
133 CL surgery were not identified, as they cannot be used for ligament and meniscus surgeries for health
134 claims. Third, it was impossible to analyze information other than medical practice, sex, and age.

135 Finally, traffic accidents and occupational injuries were not included in the present data.

136 In future studies, we aim to investigate the number of injuries and reconstruction surgeries separately
137 for ACL and PCL injuries. We would also like to clarify the characteristics of Japanese patients
138 through comparison with overseas reports. We believe organizing information on knee joint ligament
139 injuries will lead to novel suggestions for rehabilitation and prevention programs.

140

141 **Conclusion**

142 The present study identifies an increased incidence of CL surgery in Japan, along with gender and age
143 characteristics. This trend is not limited to younger age groups but is also observed in middle-aged
144 and older age groups. In the future, more detailed data should be extracted from medical databases to
145 identify insights for optimal treatment strategies for specific populations.

146

147 **Acknowledgments**

148 This work was supported by JSPS KAKENHI Grant Numbers JP21K02905. We would like to express
149 our gratitude to all the members of Taguri's laboratory for their valuable discussions and comments.
150 Additionally, we would like to thank Editage (<https://www.editage.jp/>) for providing English language
151 editing services.

152

153

154 **References**

- 155 1. Gianotti SM, Marshall SW, Hume PA, Bunt L. Incidence of anterior cruciate ligament injury
156 and other knee ligament injuries: A national population-based study. *Journal of Science and*
157 *Medicine in Sport*. 2009;12: 622–627. doi:10.1016/j.jsams.2008.07.005
- 158 2. Nordenvall R, Bahmanyar S, Adami J, Stenros C, Wredmark T, Felländer-Tsai L. A
159 Population-Based Nationwide Study of Cruciate Ligament Injury in Sweden, 2001-2009:
160 Incidence, Treatment, and Sex Differences. *Am J Sports Med*. 2012;40: 1808–1813.
161 doi:10.1177/0363546512449306
- 162 3. Longo UG, Nagai K, Salvatore G, Cella E, Candela V, Cappelli F, et al. Epidemiology of
163 Anterior Cruciate Ligament Reconstruction Surgery in Italy: A 15-Year Nationwide Registry
164 Study. *JCM*. 2021;10: 223. doi:10.3390/jcm10020223
- 165 4. Lyman S, Koulouvaris P, Sherman S, Do H, Mandl LA, Marx RG. Epidemiology of Anterior
166 Cruciate Ligament Reconstruction: Trends, Readmissions, and Subsequent Knee Surgery. *The*
167 *Journal of Bone and Joint Surgery-American Volume*. 2009;91: 2321–2328.
168 doi:10.2106/JBJS.H.00539
- 169 5. Janssen KW, Orchard JW, Driscoll TR, van Mechelen W. High incidence and costs for
170 anterior cruciate ligament reconstructions performed in Australia from 2003-2004 to 2007-

- 171 2008: time for an anterior cruciate ligament register by Scandinavian model?: Incidence of
172 ACL surgery in Australia. *Scand J Med Sci Sports*. 2012;22: 495–501. doi:10.1111/j.1600-
173 0838.2010.01253.x
- 174 6. Granan L-P, Forssblad M, Lind M, Engebretsen L. The Scandinavian ACL registries 2004–
175 2007: baseline epidemiology. *Acta Orthopaedica*. 2009;80: 563–567.
176 doi:10.3109/17453670903350107
- 177 7. Longo UG, Viganò M, Candela V, de Girolamo L, Cella E, Thiebat G, et al. Epidemiology of
178 Posterior Cruciate Ligament Reconstructions in Italy: A 15-Year Study. *JCM*. 2021;10: 499.
179 doi:10.3390/jcm10030499
- 180 8. Chung KS, Kim JH, Kong DH, Park I, Kim JG, Ha JK. An Increasing Trend in the Number of
181 Anterior Cruciate Ligament Reconstruction in Korea: A Nationwide Epidemiologic Study.
182 *Clin Orthop Surg*. 2022;14: 220. doi:10.4055/cios20276
- 183 9. Chung KS. An increasing trend of posterior cruciate ligament reconstruction in South Korea:
184 epidemiologic analysis using Korean National Health Insurance System Database. *Knee Surg
185 & Relat Res*. 2021;33: 44. doi:10.1186/s43019-021-00126-y
- 186 10. Takahashi S, Okuwaki T. Epidemiological survey of anterior cruciate ligament injury in

- 187 Japanese junior high school and high school athletes: cross-sectional study. *Research in Sports*
- 188 *Medicine*. 2017;25: 266–276. doi:10.1080/15438627.2017.1314290
- 189 11. Katano H, Ozeki N, Kohno Y, Nakagawa Y, Koga H, Watanabe T, et al. Trends in
- 190 arthroplasty in Japan by a complete survey, 2014–2017. *Journal of Orthopaedic Science*.
- 191 2021;26: 812–822. doi:10.1016/j.jos.2020.07.022
- 192 12. Katano H, Koga H, Ozeki N, Otabe K, Mizuno M, Tomita M, et al. Trends in isolated
- 193 meniscus repair and meniscectomy in Japan, 2011–2016. *Journal of Orthopaedic Science*.
- 194 2018;23: 676–681. doi:10.1016/j.jos.2018.04.003
- 195 13. Yamaguchi S, Kimura S, Akagi R, Yoshimura K, Kawasaki Y, Shiko Y, et al. Increase in
- 196 Achilles Tendon Rupture Surgery in Japan: Results From a Nationwide Health Care Database.
- 197 *Orthopaedic Journal of Sports Medicine*. 2021;9: 232596712110341.
- 198 doi:10.1177/23259671211034128
- 199 14. Ministry of Health, Labour and Welfare. 1st NDB open data Japan. 2014. Available:
- 200 <https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/0000139390.html>
- 201 15. Ministry of Health, Labour and Welfare. 2nd NDB open data Japan. 2015. Available:
- 202 <https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/0000177221.html>

- 203 16. Ministry of Health, Labour and Welfare. 3rd NDB open data Japan. 2016. Available:
204 https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/0000177221_00002.html
- 205 17. Ministry of Health, Labour and Welfare. 4th NDB open data Japan. 2017. Available:
206 https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/0000177221_00003.html
- 207 18. Ministry of Health, Labour and Welfare. 5th NDB open data Japan. 2018. Available:
208 https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/0000177221_00008.html
- 209 19. Ministry of Health, Labour and Welfare. 6th NDB open data Japan. 2019. Available:
210 https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/0000177221_00010.html
- 211 20. Japanese population. Statistics Bureau of the Ministry of Internal Affairs and Communications
212 Website. [cited 9 Mar 2023]. Available: <https://www.stat.go.jp/data/jinsui/2.html>
- 213 21. Mahapatra P, Horriat S, Anand BS. Anterior cruciate ligament repair – past, present and
214 future. *J EXP ORTOP*. 2018;5: 20. doi:10.1186/s40634-018-0136-6
- 215 22. Hopper GP, Heusdens CHW, Dossche L, Mackay GM. Posterior Cruciate Ligament Repair
216 With Suture Tape Augmentation. *Arthroscopy Techniques*. 2019;8: e7–e10.
217 doi:10.1016/j.eats.2018.08.022
- 218 23. Davarinos N, O'Neill BJ, Curtin W. A Brief History of Anterior Cruciate Ligament

- 219 Reconstruction. *Advances in Orthopedic Surgery*. 2014;2014: 1–6. doi:10.1155/2014/706042
- 220 24. Zheng W, Suzuki K, Yokomichi H, Sato M, Yamagata Z. Multilevel Longitudinal Analysis of
221 Sex Differences in Height Gain and Growth Rate Changes in Japanese School-Aged Children.
222 *Journal of Epidemiology*. 2013;23: 275–279. doi:10.2188/jea.JE20120164
- 223 25. Chotel F, Henry J, Seil R, Chouteau J, Moyen B, Bérard J. Growth disturbances without
224 growth arrest after ACL reconstruction in children. *Knee Surg Sports Traumatol Arthrosc*.
225 2010;18: 1496–1500. doi:10.1007/s00167-010-1069-5
- 226 26. Gicquel P, Geffroy L, Robert H, Sanchez M, Curado J, Chotel F, et al. MRI assessment of
227 growth disturbances after ACL reconstruction in children with open growth plates–Prospective
228 multicenter study of 100 patients. *Orthopaedics & Traumatology: Surgery & Research*.
229 2018;104: S175–S181. doi:10.1016/j.otsr.2018.09.002
- 230 27. Weng C-J, Yeh W-L, Hsu K-Y, Chiu C, Chang S-S, Chen AC-Y, et al. Clinical and Functional
231 Outcomes of Anterior Cruciate Ligament Reconstruction With Autologous Hamstring Tendon
232 in Patients Aged 50 Years or Older. *Arthroscopy: The Journal of Arthroscopic & Related
233 Surgery*. 2020;36: 558–562. doi:10.1016/j.arthro.2019.08.047
- 234 28. Filbay SR, Culvenor AG, Ackerman IN, Russell TG, Crossley KM. Quality of life in anterior

- 235 cruciate ligament-deficient individuals: a systematic review and meta-analysis. Br J Sports
236 Med. 2015;49: 1033–1041. doi:10.1136/bjsports-2015-094864
- 237 29. van Meer BL, Meuffels DE, van Eijsden WA, Verhaar JAN, Bierma-Zeinstra SMA, Reijman
238 M. Which determinants predict tibiofemoral and patellofemoral osteoarthritis after anterior
239 cruciate ligament injury? A systematic review. Br J Sports Med. 2015;49: 975–983.
240 doi:10.1136/bjsports-2013-093258
- 241 30. Toanen C, Demey G, Ntagiopoulos PG, Ferrua P, Dejour D. Is There Any Benefit in Anterior
242 Cruciate Ligament Reconstruction in Patients Older Than 60 Years? Am J Sports Med.
243 2017;45: 832–837. doi:10.1177/0363546516678723
- 244 31. Liu C-H, Chiu C-H, Chang S-S, Yeh W-L, Chen AC-Y, Hsu K-Y, et al. Clinical and
245 functional outcomes of isolated posterior cruciate ligament reconstruction in patients over the
246 age of 40 years. BMC Musculoskelet Disord. 2022;23: 210. doi:10.1186/s12891-022-05151-w
247
248

249 **Supporting information**

250 S1 Table. STROBE Statement—checklist of items that should be included in reports of observational
251 studies.

252 S2 Table. Annual registrations of arthroscopic ligament reconstruction according to age groups from
253 2014 to 2019. To avoid the identification of individuals, aggregate units that are <10 in principle are
254 not included.

255 S3 Table. Annual registrations of arthroscopic ligament reconstruction per 100,000 population
256 according to sex and age groups from 2014 to 2019. To avoid the identification of individuals,
257 aggregate units that are <10 in principle are not included.




Figure1




Figure2




Figure3