

1 **Proportion of paediatric admissions with any stage of noma at the** 2 **Anka General Hospital, northwest Nigeria**

3 Short title: Noma in paediatric admissions in northwest Nigeria

4

5 Elise Sarah Farley ¹, Miriam Njoki Karinja ², Abdulhakeem Mohammed Lawal ², Michael Olaleye ¹,

6 Sadiya Muhammad ², Maryam Umar ², Fatima Khalid Gaya ², Shirley Chioma Mbaeri ², Mark

7 Sherlock ³, Deogracia Wa Kabila ¹, Miriam Peters ², Joseph Samuel ¹, Guy Maloba ², Rabi Usman ⁴,

8 Saskia van der Kam ³, Koert Ritmeijer ³, Cono Ariti ⁵, Mohana Amirtharajah ³, Grégoire Falq ³

9

10 **Affiliations**

11 1 Médecins Sans Frontières, Noma Children's Hospital, Sokoto, Nigeria

12 2 Médecins Sans Frontières, Nigeria Mission, Zamfara and Abuja, Nigeria

13 3 Médecins Sans Frontières, Amsterdam, Netherlands

14 4 Zamfara Ministry of Health, Zamfara, Nigeria

15 5 London School of Hygiene and Tropical Medicine, London, United Kingdom

16

17

18 Key words: noma, cancrum oris, malnutrition

19

20

21

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

22 **Author contributions**

23 Elise Sarah Farley (0000-0003-4457-4068): Conceptualization, Data Curation, Formal Analysis,
24 Investigation, Methodology, Writing – Original Draft Preparation.

25 Miriam Njoki Karinja (0000-0003-0474-9492): Data Curation, Investigation, Project Administration,
26 Supervision, Writing – Review & Editing.

27 Abdulhakeem Mohammed Lawal: Data Curation, Investigation, Project Administration, Supervision,
28 Writing – Review & Editing.

29 Michael Olaleye (0009-0003-4101-8469): Data Curation, Investigation, Project Administration,
30 Supervision, Writing – Review & Editing.

31 Sadiya Muhammad: Investigation, Writing – Review & Editing.

32 Maryam Umar: Investigation, Writing – Review & Editing.

33 Fatima Khalid Gaya: Investigation, Writing – Review & Editing.

34 Shirley Chioma Mbaeri: Investigation, Writing – Review & Editing.

35 Mark Sherlock (0009-0005-8186-4365): Conceptualization, Funding Acquisition, Writing – Review
36 & Editing.

37 Deogracia Wa Kabila: Funding Acquisition, Writing – Review & Editing.

38 Miriam Peters: Funding Acquisition, Writing – Review & Editing.

39 Joseph Samuel: Conceptualization, Funding Acquisition, Writing – Review & Editing.

40 Guy Maloba: Funding Acquisition, Writing – Review & Editing.

41 Rabi Usman (0000-0001-7869-5357): Writing – Review & Editing.

42 Saskia van der Kam (0000-0002-3608-8054): Conceptualization, Writing – Review & Editing.

43 Koert Ritmeijer (0000-0001-5924-872X): Conceptualization, Writing – Review & Editing.

44 Cono Ariti (0000-0001-7615-0935): Conceptualization, Formal Analysis, Methodology, Supervision,
45 Writing – Review & Editing.

46 Mohana Amirtharajah (0000-0002-4779-0308): Conceptualization, Funding Acquisition,
47 Methodology, Supervision, Writing – Review & Editing.

48 Grégoire Falq (0009-0002-2116-6475): Formal Analysis, Supervision, Writing – Review & Editing.

49

50

51

53

54 **ABSTRACT**

55 **Introduction**

56 Noma is a rapidly spreading infection of the oral cavity which mainly affects young children, and
57 without early treatment, can have a high mortality rate. Simple gingivitis is a warning sign for noma,
58 and acute necrotizing gingivitis is the first stage of noma. The epidemiology of noma is not well
59 understood. We aimed to gather evidence on the epidemiology of noma and its association with
60 malnutrition for use in program planning.

61 **Methods**

62 We conducted a prospective observational study enrolling patients aged 0 to 12 years who were
63 admitted to the Anka General Hospital, Zamfara, northwest Nigeria. Consenting caregivers of
64 participants were interviewed at admission. Participants had anthropometric and oral exams at
65 admission and discharge.

66 **Findings**

67 Of the 2346 participants, 58 (2.5%) were diagnosed with simple gingivitis and six (n=0.3%) with acute
68 necrotizing gingivitis upon admission. Of those admitted to the Inpatient Therapeutic Feeding Centre
69 (ITFC), 3.4% (n=37, CI 2.5 - 4.7%) were diagnosed with simple gingivitis upon admission compared
70 to 1.7% of those not admitted to the ITFC (n=21, CI 1.1 - 2.6%) (p=0.008). Risk factors identified for
71 having simple gingivitis include being aged over two years (2 to 6yrs old, odds ratio (OR) 3.4, CI 1.77
72 - 6.5; 7 to 12 yrs OR 5.0, CI 1.7 - 14.6; p<0.001) and being admitted to the ITFC (OR 2.1; CI 1.22 -
73 3.62).

74 **Conclusion**

75 Our study showed a small proportion of those admitted to the Anka General Hospital had simple or
76 acute necrotizing gingivitis. Those admitted to the ITFC were more likely to have simple gingivitis.
77 The lack of access to and uptake of oral health care indicates a strong need for oral exams to be

78 included in routine health services. This provision could improve the oral status of the population and
79 decrease the chance of patients developing noma.

80

81

82

83

84

85

86

87

88

89 **AUTHOR SUMMARY**

90 Noma is a rapidly spreading infection of the oral cavity which mainly affects young children and
91 without early treatment, can have a high mortality rate. Simple gingivitis is a warning sign for noma,
92 and acute necrotizing gingivitis is the first stage of noma. We aimed to gather evidence on the
93 epidemiology of noma and its association with malnutrition by conducting a prospective observational
94 study enrolling 2346 patients aged 0 to 12 years who were admitted to the Anka General Hospital,
95 Zamfara, northwest Nigeria. Consenting caregivers of participants were interviewed at admission.
96 Participants had anthropometric and oral exams at admission and discharge. Our study showed a small
97 proportion of those admitted to the Anka General Hospital had simple or acute necrotizing gingivitis.
98 Those admitted to the ITFC were more likely to have simple gingivitis. The lack of access to and uptake
99 of oral health care indicates a strong need for oral exams to be included in routine health services. This
100 provision could improve the oral status of the population and decrease the chance of patients developing
101 noma.

102

103

104 INTRODUCTION

105 Noma is an infection of the oral cavity which can cause the disintegration of the cheek, nose and/or eye
106 in a few weeks [1]. If untreated in the early reversible stages, noma has a reported 90% mortality rate
107 [2] and mainly affects children aged between two and six years [3]. Deaths in noma patients are
108 primarily due to starvation, aspiration pneumonia, respiratory insufficiency or sepsis [4,5]. With timely
109 antibiotic treatment, wound debridement and nutritional support, morbidity and mortality from noma
110 greatly decrease [1]. For those who survive, noma often leads to severe facial disfigurement, functional
111 issues and stigmatization [1,2,6,7]. The WHO classifies noma into stages: Warning Sign: simple
112 gingivitis, Stage 1: acute necrotizing gingivitis, Stage 2: oedema, Stage 3: gangrene, Stage 4: scarring
113 and Stage 5: sequelae [2].

114 There is limited research on noma and as a result, many gaps in knowledge exist around the disease [1].
115 The aetiology of noma is unknown but thought to be multifactorial [1]. Risk factors for noma include
116 poverty, poor oral hygiene, poor access to routine childhood vaccinations, limited access to quality
117 health care and immunosuppression resulting from comorbidities such as malnutrition, measles and
118 HIV [1,8–16]. Malnutrition is frequently listed as a risk factor for noma, however, evidence supporting
119 this theory is largely based on case reports and case series [11,17–27] and a handful of primary studies
120 [8–10,28–30]. The epidemiology of the disease is also not well understood. The WHO estimates that
121 140, 000 children contract noma each year globally [31]. In 2003, a northwest Nigerian study estimated
122 the noma incidence was 6.4 per 1000 children from 1996 to 2001 [32]. A recent study estimated that
123 the period prevalence of noma from 2010-2018 was 1.6 per 100,000 population at risk in northern
124 Nigeria [33]. A 2019 prevalence study in Sokoto and Kebbi states, Nigeria, identified that in children
125 aged 0 to 15 years (n=7122), simple gingivitis was diagnosed in 3.1% (n=181; 95% confidence interval
126 (CI) 2.6-3.8), acute necrotizing gingivitis in 0.1% (n=10; CI 0.1-0.3), and oedema in 0.05% (n=3; CI
127 0.02-0.2), no late stage noma cases were identified [34].

128 We conducted a prospective observational study assessing the proportion of pediatric admissions at
129 the Anka General Hospital with any stage of noma, to add to the understanding of the epidemiology
130 of the disease and examine the links between noma and malnutrition.

131 **METHODS**

132 **STUDY DESIGN AND DATA COLLECTION**

133 From 1st June to 24th October 2021, we conducted a prospective observational study utilising face-to-
134 face interviews with caregivers of paediatric patients (aged under 12 years) admitted to the Anka
135 General Hospital, northwest Zamfara, Nigeria and physical examinations of the patients themselves.
136 All caregivers of patients admitted between 09h00 and 16h00 Monday to Sunday were approached on
137 the day of admission by the study nurse. Consenting individuals were enrolled (discussed in the Ethics
138 section below). A tablet-based RedCap data collection form was used to collect the study specific data
139 on admission (sociodemographic characteristics, infant and current feeding practices, oral health
140 practices, health care access, oral examination, mid upper arm circumference, weight, and height) and
141 at discharge (oral exam and access to routinely collected medical records).

142 **SAMPLE SIZE**

143 We aimed to have a sample size large enough to estimate the proportion of paediatric admissions with
144 any stage of noma admitted to the Inpatient Therapeutic Feeding Centre (ITFC) or other wards. This
145 calculation was based on the following assumptions:

- 146 - 6%- Inpatient department proportion of patients diagnosed with any stage of noma (as this
147 had a lower proportion of any stage of noma diagnosed than the ITFC [35])
- 148 - 95% confidence interval
- 149 - 1.5% margin of error (i.e., can estimate a proportion between 4.5 and 7.5% assuming a true
150 proportion of 6%)

151 The sample size calculation showed that it was necessary to include 963 patients in the ITFC and 963
152 patients in the other wards.

153 **ANALYSIS**

154 We performed a descriptive analysis of the data collected. Categorical variables are shown as
155 frequencies and percentages. Continuous variables were summarised using means and standard

156 deviations (SD) or medians and interquartile ranges (IQR) depending on normality. Missing data
157 numbers are recorded in each table.

158 Age was categorised into 0-1 years, 2-6 years and 7-12 years. This age categorisation was selected as
159 children aged between two and six years have previously been reported to be at higher risk of noma [3].

160 Malnutrition estimates are not reported for participants aged zero to five months as we did not have
161 access to the confirmed birthweights of these children [36].

162 For children aged six months to 5 years, using Mid-Upper Arm Circumference (MUAC) measurements
163 taken upon admission, we estimated the proportion of severe acute malnutrition (SAM, MUAC
164 <115mm), moderate acute malnutrition (MAM, between MUAC \geq 115 and <125mm) and global acute
165 malnutrition (GAM, MUAC <125mm) [36].

166 For those aged six to 12 years, we report the WHO recommended Body Mass Index (BMI)-for-age Z-
167 scores [36]. BMI was calculated based on the weight (in kg) divided by the square of the height (in m)
168 of the individual. BMI-for- age are presented as Z-scores based on the 2007 WHO Growth Reference
169 [36].

170 Wealth scores were calculated by assigning a value of one to each of the following items owned by the
171 family: cell phone, radio, motorbike, tractor, bicycle, car (these items were chosen based on consultation
172 with local researchers, knowledgeable about the context). The minimum wealth score was zero and the
173 maximum was six, all items weighed equally in the scoring.

174 We compared the proportion of admissions with each stage of noma between those admitted to the
175 ITFC and those admitted to other wards using a Pearson's chi-square test.

176 We calculated the number of admissions who had any stage of noma upon admission, and the outcome
177 of those patients (noma stage at discharge) for all those who had oral examinations on both admission
178 and discharge.

179 We assessed risk factors for simple gingivitis using univariate logistic regression (patient age, wealth
180 score, ITFC admission status, if the child was sick during the past 3 months, parent primary caretaker,

181 if child had been given colostrum at birth, if the child eats pap, if the child was vaccinated, SAM,
182 MAM and GAM status upon admission, measles, and malaria diagnoses). These variables were
183 selected as they are reported risk factors for noma [1,8–27]. We conducted an exploratory analysis of
184 risk factors for acute necrotizing gingivitis; proportions are reported.

185 All data analysis was conducted with Stata 17 (StataCorp LP, College Station, TX, USA).

186 **ETHICS**

187 The Médecins Sans Frontières Ethics Review Board (ERB) (2017), Nigerian Federal Ministry of Health
188 ERB (NHREC/01/01/2007-29/04/2021), Zamfara Ministry of Health ERB (ZSHREC01112020) and
189 the Usman Danfodiyo University Teaching Hospital Health Research and Ethics Committee in Nigeria
190 (NHREC/30/012/2019) approved the study protocol.

191 Caregivers of participants were asked to sign an informed consent form and participants aged between
192 seven and 12 years were informed about the study and asked to sign an assent form after reading/
193 hearing the information provided on the information sheet. For participants or caregivers who were
194 illiterate, we requested them to provide an initial and a thumb print as a representation of consent, with
195 additional participation from an independent literate witness who checked that the participant/ caregiver
196 understood the contents of the consent form. This witness co-signed the consent form. The information
197 sheet and the informed consent and assent forms were written in English and Hausa.

198 **RESULTS**

199 **SOCIODEMOGRAPHIC CHARACTERISTICS**

200 Of those caregivers approached (n=2356), six did not provide consent and four did not provide assent
201 to be part of the study; thus 2346 participants were enrolled and included in the analysis, 2340 of
202 these had a discharge assessment. Caregivers were mainly aged between 18 and 25 years (n=1036,
203 44.3%) and 26 to 35 years (n=946, 40.5%), female (n=2318, 99.2%) and the mother of the child
204 (n=2211, 94.6%). The main source of income of most families was agriculture (n=1285, 55.0%). Most
205 (n=1696, 72.3%) families owned zero to two items on the wealth score list. Half the child participants

206 were female (n=1181, 50.6%), most were aged between zero and one years (n=1097, 47.1%) or two to
 207 six years (n=1135, 48.8%) (Table 1).

208 **Table 1- Sociodemographic characteristics, noma study Anka General Hospital 2021**

Caretaker age	N=2346	%
18 - 25 years	1036	44.3%
26 - 35 years	946	40.5%
36 - 45 years	283	12.1%
> 46 years	71	3.1%
Missing	10	
Caretaker sex		
Female	2318	99.2%
Male	19	0.8%
Missing	9	
Caretaker education		
None	732	31.3%
Arabic studies	1499	64.1%
Primary school	54	2.3%
Secondary school	50	2.1%
University	2	0.1%
Missing	9	
Income		
Agriculture	1285	55.0%
Trader	745	31.9%
Mining	88	3.8%
Daily labourer	53	2.3%
Civil servant	43	1.8%
Livestock	37	1.6%
Other	85	3.6%
Missing	10	
Relationship to child		
Mother	2211	94.6%
Other	125	5.4%
Missing	10	
Wealth score (own: cell phone, radio, motorbike, tractor, bicycle, car)		
0-2 items	1696	72.3%
3-6 items	650	27.7%
Child gender		
Female	1181	50.6%
Male	1154	49.4%
Missing	11	
Child age (years)		
0-1	1097	47.1%
2-6	1135	48.8%
7-12	95	4.1%

Missing	19
---------	----

209

210 **HEALTH CARE ACCESS AND PRIMARY DIAGNOSES**

211 Caregivers reported several challenges accessing care including difficulties with transport (n=454,
 212 19.4%) and the hospital being far from their home (n=432, 18.4%). Half of the respondents (n=1176,
 213 50.4%) reported that the child had been sick in the prior three months (not including the illness that
 214 made them seek care on the day of enrolment). Of these, almost all (n=1069, 90.9%) sought care for
 215 that illness, many from a street pharmacist (n=424, 39.7%) (Table 2).

216 **Table 2- Health care access noma study Anka General Hospital 2021**

Challenges when accessing care	N= 2346	%
Transport difficult	454	19.4%
Far to get to hospital	432	18.4%
Difficult to leave other children	22	0.9%
Takes a long time to get to hospital	52	2.2%
Child very sick, difficult to travel	6	0.3%
Difficult to get permission to travel from family	6	0.3%
Cost of care too expensive	13	0.6%
No challenges	1505	64.2%
Has your child been sick in the past 3 month? (other than the sickness that made you seek care today)		
No	1157	49.6%
Yes	1176	50.4%
Missing	13	
If yes, did you seek care due to that sickness? (N=1176)		
No	104	8.8%
Yes	1069	90.9%
Where did you seek care? (N=1069)		
Hospital	297	27.8%
Clinic	327	30.6%
Traditional healer	62	5.8%
Street chemist, pharmacist	424	39.7%

217

218 Of those respondents who had teeth (n=1907, 81.7%), almost all (n= 1652, 86.6%) cleaned their teeth,
 219 mainly using water (n= 1399, 84.7%). The main reasons for children cleaning their teeth were to get
 220 clean, bright teeth (n=537, 32.5%) and to get rid of foul breath (n=944, 57.1%). The main reasons for
 221 children not cleaning their teeth (n=251), were not knowing the benefits of cleaning (n=18, 7.2%) and

222 forgetting to clean their teeth (n= 19, 7.6%). Most children (n=2069, 88.5%) had not had any dental
 223 issues in the three months prior to study enrolment. The most common oral health issues reported
 224 (n=265, 11.4%) were a sore in the mouth (n=149, 56.2%), bleeding gums when touched (n=88,
 225 33.2%) and sore gums (n=34, 12.8%). Almost all the respondents (n=2274, 97.4) had not had an oral
 226 health check in the past year (Table 3).

227 **Table 3- Oral hygiene and health care noma study Anka General Hospital 2021**

Does your child have any teeth?	N= 2346	%
No	428	18.3%
Yes	1907	81.7%
Missing	11	
Does your child clean their teeth? (N=1907)		
Don't know	4	0.2%
No	251	13.2%
Yes	1652	86.6%
Reason for not cleaning (N=251)		
No time for brushing	12	4.8%
No money for brush and toothpaste	1	0.4%
Nobody brushes their teeth in my family	2	0.8%
Don't know of any benefits from brushing	18	7.2%
Gums are bleeding when brushing	1	0.4%
Useless, good teeth are hereditary	6	2.4%
Always forget to brush my teeth	19	7.6%
Reason for cleaning (N=1652)		
Clean, bright teeth	537	32.5%
Prevention of caries	279	16.9%
Prevention of bleeding gums	15	0.9%
Prevention of oral ulcer	107	6.5%
To get rid of foul breath	944	57.1%
Cleaning utensil (N=1652)		
Toothbrush	171	10.4%
Stick	2	0.1%
Cloth	29	1.8%
Ash	39	2.4%
Salt and water	110	6.7%
Water	1399	84.7%
Does your child use toothpaste? (N=1652)		
No	1418	85.8%
Yes	216	13.1%
No times teeth clean/day (Median, IQR) (N=1652)		
	1	1,2
How long does your child clean their teeth for? (N=1652)		
Less than 3 minutes	1580	95.6%
3 minutes or more	61	3.7%

Child oral health issues in the past three months?	N= 2346	%
No	2069	88.6%
Yes	265	11.4%
Missing	12	
Oral issues past 3 months (N=265)		
Sore gums	34	12.8%
Bleeding gums when touched	88	33.2%
Spontaneously bleeding gums	12	4.5%
Sore in mouth	149	56.2%
Pus in mouth	4	1.5%
Sore jaw	8	3.0%
Tooth pain	10	3.8%
Has your child visited an oral health care provider for a check-up in the past year		
No	2274	97.4%
Yes	60	2.6%
Missing	12	
Oral health provider (N=60)		
Hospital	10	16.7%
Clinic	26	43.3%
Traditional healer	14	23.3%
Street chemist, pharmacist	13	21.7%

228

229 The most common primary diagnosis of patients admitted to the hospital were malaria (n= 1086,
 230 46.3%) and severe acute malnutrition (n=977, 41.6%). The median number of days patients spent in
 231 the hospital was three (IQR 2, 5) (Table 4).

232 **Table 4- Diagnosis noma study Anka General Hospital 2021**

Primary diagnosis	N= 2346	%
Malaria	1086	46.3%
Severe acute malnutrition	977	41.6%
Measles	114	4.9%
Other	56	2.4%
Lower respiratory tract infection	50	2.1%
Acute watery diarrhoea	36	1.5%
Other	15	0.5%
Length of stay in hospital (days) (Median, IQR)	3	2, 5
Length of stay in hospital (days) (range)	0	97

233

234 **GINGIVITIS AND NOMA DIAGNOSES**

235 Simple gingivitis was diagnosed in 58 patients (2.5%) and acute necrotizing gingivitis was diagnosed
236 in six patients (0.3%) at admission. At discharge simple gingivitis was diagnosed in nine patients
237 (0.5%) and acute necrotizing gingivitis was diagnosed in one patient (0.1%) (Table 5).

238 **Table 5- Noma diagnosis upon admission and discharge, noma study Anka General Hospital**
239 **2021**

Admission Noma Stage	N= 2346	%
None	2256	97.2%
Stage 0: Simple gingivitis	58	2.5%
Stage 1: Acute necrotizing gingivitis	6	0.3%
Missing	26	
Discharge Noma Stage		
None	1667	99.4%
Stage 0: Simple gingivitis	9	0.5%
Stage 1: Acute necrotizing gingivitis	1	0.1%
Missing	669	

240
241 Of the 34 participants who were diagnosed with simple gingivitis on admission and who had an oral
242 exam at discharge: 28 (82.4%) had resolved at discharge; five (14.7%) still had simple gingivitis at
243 discharge; and one (2.9%) had acute necrotizing gingivitis upon discharge. One of the two patients
244 who had acute necrotizing gingivitis at admission and had an oral exam at discharge, had no stage of
245 noma at discharge, the other had simple gingivitis at discharge.

246 **MALNUTRITION**

247 Of those patients enrolled in the study who were aged between 6 months and five years and who had a
248 MUAC measurement completed upon admission (n=2039), n=692 (34.0%) had SAM, n=399 (19.6%)
249 had MAM and n=1091 (53.5%) had GAM (Table 6).

250 Of the participants aged between six and 12 years who had weight and height measured upon
251 admission (n=100), 44.0% (n=44) were wasted (BMI/age <-2 z-score). The majority of those admitted
252 to the ITFC were wasted (n=18, 58.1%), and the majority of those not admitted to the ITFC were
253 classified with normal BMI z-scores (n=41, 59.2%) (Table 6).

254 **Table 6: Malnutrition (SAM, MAM, GAM) upon admission in those aged 6 months to 5 years**
 255 **(n=2039) and BMI-for-age Z-scores classification for six to 12 year olds [36]**

	TOTAL		ITFC		Not ITFC		Ch sq
MUAC	N=2039	%	N=956	%	N=1083	%	
Normal, MUAC>125mm	948	46.5%	133	13.9%	815	75.3%	<0.05
SAM, MUAC <115mm	692	33.9%	604	63.2%	88	8.1%	
MAM, between MUAC ≥115 and <125mm	399	19.6%	218	22.9%	180	16.6%	
GAM, MUAC <125mm	1091	53.5%	823	86.1%	268	24.8%	<0.05
BMI-for-age Z-score	N=100	%	N=31	%	N=69	%	
Severe thinness (<-3SD)	28	28.0%	16	51.6%	12	17.4%	<0.05
Thinness (≥ -3 SD & < -2 SD)	16	16.0%	2	6.5%	14	20.3%	
Normal (≥ -2 SD & < +1SD)	51	51.0%	10	32.3%	41	59.2%	
Overweight (≥+1SD & <+2 SD)	2	2.0%	1	3.2%	1	1.5%	
Obesity (≥+2SD)	3	3.0%	2	6.5%	1	1.5%	

256

257 **RISK FACTORS**

258 Being aged over two years (2-6yr old odds ratio (OR) 3.4, CI 1.77, 6.5; 7 to 12 yr OR 5.0, CI 1.7,
 259 14.6; p=<0.001) and being admitted to the ITFC (OR 2.1; CI 1.22, 3.62) were associated with having
 260 simple gingivitis. Those whose primary diagnosis was malaria, were less likely to have simple
 261 gingivitis upon admission compared to those admitted for other reasons (OR 0.29; CI 0.15, 0.55;
 262 p=<0.001). Of those diagnosed with acute necrotizing gingivitis (n=6) n=4/6 were aged 2-6 years,
 263 were in the lower wealth score (indicating lower socioeconomic status) and ate pap; n=5/6 children
 264 had been sick in the 3 months prior to admission (Table 7).

265

266

267

268 **Table 7: Risk factors for stage 0 and 1 on admission**

	Total		Simple gingivitis					Acute necrotizing gingivitis	
	N =2346	%	N=58	%	OR	95% CI	P	N=6	%
Patient age (years)							<0.001		
0 to 1	1097	47.1%	12	20.7%	1.00	(Ref)		0	0.0%
2 to 6	1135	48.8%	41	70.7%	3.38	(1.77, 6.47)		4	66.7%
7 to 12	95	4.1%	5	8.6%	5.03	(1.73, 14.61)		2	33.3%
	2327		58					6	
Wealth score (own: cell phone, radio, motorbike, tractor, bicycle, car)							0.728		
0-2 items	1696	72.3%	43	74.1%	1.00	(Ref)		4	66.7%
3-6 items	650	27.7%	15	25.9%	0.90	(0.50, 1.63)		2	33.3%
	2346		58					6	
ITFC -Child admitted							0.007		
No	1269	54.1%	21	36.2%	1.00	(Ref)		5	83.3%
Yes	1077	45.9%	37	63.8%	2.10	(1.22, 3.62)		1	16.7%
	2346		58					6	
Malnutrition upon admission in those aged 6 months to 5 years							0.143		
Normal, MUAC>125mm	948	46.5%	17	33.3%	1.00	(Ref)		3	75.0%
SAM, MUAC <115mm	692	33.9%	23	45.1%	1.88	(1.00, 3.55)		0	0.0%
MAM, between MUAC ≥115 and <125mm	399	19.6%	11	21.6%	1.56	(0.72, 3.36)		1	25.0%
	2039		51					4	
GAM upon admission in those aged 6 months to 5 years							0.059		
Normal, MUAC>=125mm	948	46.5%	17	33.3%	1.00	(Ref)		3	75.0%
GAM, MUAC <125mm	1091	53.5%	34	66.7%	1.76	(0.98, 3.18)		1	25.0%
	2039		51					4	
Child sick past 3 months							0.942		
No	1157	49.6%	29	50.0%	1.00	(Ref)		1	16.7%
Yes	1176	50.4%	29	50.0%	0.98	(0.58, 1.65)		5	83.3%

		2333		58					6	
Primary caretaker							0.817			
	No	266	11.4%	6	10.3%	1.00	(Ref)		2	33.3%
	Yes	2070	88.6%	52	89.7%	1.11	(0.47, 2.60)		4	66.7%
		2336		58					6	
Colostrum given at birth							0.201			
	No	669	28.7%	21	36.2%	1.00	(Ref)		2	33.3%
	Yes	1663	71.3%	37	63.8%	0.70	(0.41, 1.21)		4	66.7%
		2332		58					6	
Does your child eat pap?							0.163			
	No	589	25.2%	10	17.2%	1.00	(Ref)		2	33.3%
	Yes	1746	74.8%	48	82.8%	1.63	(0.82, 3.25)		4	66.7%
		2335		58					6	
Was your child vaccinated?							0.434			
	No	922	39.5%	20	34.5%	1.00	(Ref)		2	33.3%
	Yes	1410	60.5%	38	65.5%	1.24	(0.72, 2.15)		4	66.7%
		2332		58					6	
Malaria- primary diagnosis							<0.001			
	No	1246	53.5%	46	79.3%	1.00	(Ref)		3	50.0%
	Yes	1084	46.5%	12	20.7%	0.29	(0.15, 0.55)		3	50.0%
		2330		58					6	
Measles- primary diagnosis							0.191			
	No	2216	95.1%	53	91.4%	1.00	(Ref)		5	83.3%
	Yes	114	4.9%	5	8.6%	1.87	(0.73, 4.77)		1	16.7%
		2330		58					6	
OR = odds ratio; CI = confidence interval; P = p value from univariate logistic regression model										

270 DISCUSSION

271 A small proportion of those enrolled in this study had simple gingivitis (2.5%) or acute necrotizing
272 gingivitis (0.3%) at admission, and no later stages of noma were identified in the cohort. Oral health
273 practices were not optimal and access to oral health care was limited and underutilised. Those in the
274 ITFC were more likely to have simple gingivitis. Known risk factors for noma (malnutrition and
275 being aged between two and six years) were associated with simple gingivitis in our cohort.

276

277 Our findings were similar to a community-based noma study in northwest Nigeria in 2020 where
278 simple gingivitis was identified in 3.1% (n=181; 95% CI 2.6 to 3.8), acute necrotising gingivitis in
279 0.1% (n=10; CI 0.1 to 0.3) and oedema in 0.05% (n=3; CI 0.02 to 0.2) [34]. These similarities were
280 expected as the studies were conducted a similar region (northwest Nigeria). However, as our study
281 was conducted in a hospital setting with a large proportion of patients being treated for malnutrition, it
282 could have been expected that we would have seen more noma cases being diagnosed. One plausible
283 explanation for this not being the case, is that the age range of our survey included a significant
284 number of children aged under two years (47.1%), which is an age group less at risk of developing
285 noma.

286 A study conducted in Cameroon in 2015 including school children (n=2287) [37] found similar
287 proportions of cases with gingivitis as in our study (2.1% had moderate gingivitis which could reflect
288 our simple gingivitis cases and 0.6% had severe gingivitis which could reflect our acute necrotizing
289 gingivitis cases), however comparison is difficult as the study methods and definitions differed
290 significantly.

291 The caregivers in our cohort reported several challenges accessing care; these are similar to
292 challenges reported in other noma research work in this setting, including difficulties accessing
293 transport and the long distances patients and their caregivers need to travel in order to access care
294 [38,39]. These issues should be taken into consideration for programs aiming to alleviate barriers to
295 health care in this setting.

296 Many of the respondents who sought care sought care for a prior illness first visited a street
297 pharmacist (39.7%) showing that they would be important agents for promoting oral health and
298 awareness about noma. This finding is supported by another study in Northern Nigeria which showed
299 that engaging community pharmacists may help to reduce oral health disparities by increasing oral
300 health awareness and improving the quality of life via cost-effective delivery of pharmacy-based oral
301 health-care services [40].

302 Many respondents reported cleaning their teeth (86.6%), however the main utensil used was water
303 (84.7%), and only 10.4% reported using a toothbrush. The importance of tooth brushing and
304 mechanisms to encourage and facilitate tooth brushing have been discussed in other literature
305 [2,37,41–43]. In our cohort, the main reasons for children cleaning their teeth were to get clean, bright
306 teeth (32.5%) and to get rid of foul breath (57.1%). The main reasons for children not cleaning their
307 teeth, were not knowing the benefits of cleaning (7.2%) and forgetting to clean their teeth (7.6%).
308 These provide useful starting points for oral hygiene trainings which should focus on educating
309 children and caregivers on the benefits of tooth brushing, teaching mechanisms for instilling
310 toothbrushing into the daily routine, and explaining that toothbrushing will lead to having nice
311 smelling breath and clean, bright teeth.

312 The most common oral health issues patients reported having in the prior three months were a sore in
313 the mouth (56.2%), bleeding gums when touched (33.2%) and sore gums (12.8%). These issues are all
314 likely varying degrees of gingivitis and show that children living in the communities served by the
315 hospital show the clinical effects of poor oral hygiene and that most oral complaints are transient, and
316 not chronic or progressive.

317 Almost all the respondents (97.4%) had not had an oral health check in the past year. This low level of
318 access and uptake of oral care is similar to other studies in the setting [44–47]. The main reasons for
319 the lack of oral health care that have been reported are a lack of human resources, high costs of care,
320 sparse provision of services and low levels of knowledge about different treatment options for oral
321 conditions [44–47]. These barriers indicate the need to incorporate oral health into primary health care
322 in this setting, a call common in oral health literature [45–47]. In terms of noma, it would be worthwhile

323 encouraging the inclusion of oral examinations in primary health care provision, malnutrition surveys
324 and vaccination campaigns, particularly in vulnerable populations.

325 Those admitted to the ITFC were more likely to be diagnosed with simple gingivitis. This adds weight
326 to the link between noma and malnutrition [8–10,28–30]; however, caution is needed when
327 interpreting these findings as most cases of simple gingivitis do not lead to noma.

328 The most common primary diagnosis of patients admitted to the hospital were malaria (46.3%) and
329 severe acute malnutrition (41.6%). Both of these conditions have been reported as risk factors for the
330 development of noma [8–10,48]. However, our univariate analysis did not support this, possibly
331 because our analysis was looking at risk factors for simple gingivitis, whereas the references are
332 reporting risk factors for late stage noma. Another limitation is that our analysis was based on
333 diagnosis at the same time as assessment of any stage of noma, whereas the references discuss
334 diagnosis of comorbidities in the three months prior to onset of noma.

335 A further limitation was that we only recruited patients during the day time, this could have meant
336 that patients with serious, acute complications (such as late stage noma) admitted at night time, were
337 not included in our survey.

338 Our survey was conducted during the rainy season, which could have resulted in a possible bias, as
339 this time of year could have meant increased difficulties for sicker patients to travel to the hospital
340 which could have meant fewer later stage noma patients reached care. It could also have meant
341 increased food insecurity, or difference in the epidemiology of other morbidities in the area. Future
342 studies in other seasons or the year-round inclusion or oral screenings in routine activities could offer
343 insight into whether or not the season affects the number of simple and acute necrotizing gingivitis
344 cases identified in this context.

345 **CONCLUSION**

346 Our study showed a small proportion of those admitted to the Anka General Hospital, Zamfara, had
347 simple or acute necrotizing gingivitis, most of which had resolved upon discharge. Those admitted to
348 the ITFC were more likely to have simple gingivitis, providing evidence of the link between

349 malnutrition and the warning sign of noma. Many of those in our study cohort exhibited known risk
350 factors for noma such as being malnourished, having a comorbidity, low socioeconomic status, poor
351 oral hygiene practices, and a lack of access to healthcare, indicating there is a large population at risk
352 of developing later stage noma in this setting. The lack of access to and uptake of oral health care
353 indicates a strong need for oral exams to be included in routine health services. This provision could
354 improve the oral status of the population and decrease the chance of patients developing later stage
355 noma.
356

357 **AUTHOR STATEMENTS**

358 **Acknowledgements**

359 Thank you to the participants in this study. A big thank you to the study team for all their hard work
360 and dedication to this project. Thanks to Vincent Kimui Ndichu, Christian Mwemezi and Rahima
361 Shuaibu for their contribution to the study. We would also like to thank all the health care workers who
362 are advocating for and treating noma patients around the world.

363 **Conflict of interest**

364 The authors declare no conflict of interest.

365 **Funding**

366 This study was funded by MSF as part of the Noma Operational Research Agenda.

367 **Ethics**

368 The MSF Ethics Review Board (ERB) (2017), Nigerian Federal Ministry of Health ERB
369 (NHREC/01/01/2007-29/04/2021), Zamfara Ministry of Health ERB (ZSHREC01112020) and the
370 Usman Danfodiyo University Teaching Hospital Health Research and Ethics Committee in Nigeria
371 (NHREC/30/012/2019) approved the study protocol.

372 **Data Availability Statement**

373 MSF has a managed access system for data sharing that respects MSF's legal and ethical obligations to
374 its patients to collect, manage and protect their data responsibility. Ethical risks include but are not
375 limited to the nature of MSF operations and target populations being such that data collected often
376 involves highly sensitive data. The dataset supporting the conclusions of this article is available on
377 request in accordance with MSF's data sharing policy (available at: [http://](http://fieldresearch.msf.org/msf/handle/10144/306501)
378 fieldresearch.msf.org/msf/handle/10144/306501). Requests for access to data should be made to
379 data.sharing@msf.org.

380 **REFERENCES**

- 381 1. Ashok N, Tarakji B, Darwish S, Rodrigues JC, Altamimi MA. A Review on Noma: A Recent
382 Update. *Glob J Health Sci* [Internet]. 2016;8(4):53–9. Available from:
383 [http://www.ccsenet.org/journal/index.php/gjhs/article/view/48228doi: 10.5539/gjhs.v8n4p53](http://www.ccsenet.org/journal/index.php/gjhs/article/view/48228doi:10.5539/gjhs.v8n4p53)
- 384 2. World Health Organization Regional Office for Africa. Information brochure for early
385 detection and management of noma. 2017. [Internet]. Available from
386 <https://apps.who.int/iris/handle/10665/254579>; 2017. Available from:
387 <https://apps.who.int/iris/handle/10665/25457979%0AAdditional>
- 388 3. World Health Organisation (WHO). Noma [Internet]. [cited 2017 Nov 13]. Available from:
389 <http://www.who.int/topics/noma/en/>
- 390 4. Baratti-Mayer D, Pittet B, Montandon D, Bolivar I, Bornand JE, Jaquinet A, Schrenzel J, Pittet
391 D, Hugonnet S, Jaquinet A, Schrenzel J, Pittet D. Noma : an “ infectious ” disease of unknown
392 aetiology. *Lancet Infect*. 2003;3(7):419–31. doi: 10.1016/s1473-3099(03)00670-4
- 393 5. Silva K, Twaddell S, Powers D. A 40-year-old man with a perforated cheek. *Am J Med Sci*
394 [Internet]. 2011;341(5):399–403. Available from:
395 <http://dx.doi.org/10.1097/MAJ.0b013e3181e2ed3e>doi: 10.1097/maj.0b013e3181e2ed3e
- 396 6. Feller L, Altini M, Chandran R, Khammissa RAG, Masipa JN, Mohamed A, Lemmer J. Noma
397 (cancrum oris) in the South African context. *J Oral Pathol Med*. 2014;43(1):1–6. doi:
398 10.1111/jop.12079
- 399 7. Shaye D, Winters R, Rabbels J, Adentunje A, Magee A, Vo D. Noma Surgery. *Laryngoscope*.
400 2019;129:96–9. doi: 10.1002/lary.27230
- 401 8. Adeniyi S, Awosan K. Pattern of Noma (Cancrum Oris) and its risk factors in northwestern
402 Nigeria: a hospital-based retrospective study. *Ann Afr Med*. 2019;18(1):17–22. doi:
403 10.4103/aam.aam

- 404 9. Lazarus D. Cancrum oris - a 35-year retrospective study. *S Afr Med J*. 1997;87:1379–82.
- 405 10. Denloye O, Aderinokun G, Lawoyin J, Bankole O. Reviewing trends in the incidence of
406 cancrum oris in Ibadan, Nigeria. *West Afr J Med*. 2003;22(1):26–9.
- 407 11. Ghosh K. Use of penicillin in cancrum oris. *Indian J Pediatr*. 1947;14(4):171–172. doi:
408 <https://link.springer.com/article/10.1007%2fbf02812636?li=true>
- 409 12. Pedro K, Smit D, Morkel J. Cancrum Oris (noma) in an HIV-positive adult: a case report and
410 literature review. *SADJ*. 2016;71(6):248–52.
- 411 13. Koech K. Cancrum oris in an adult with human immunodeficiency virus infection: case report.
412 *East Afr Med J*. 2010;87(1):38–40.
- 413 14. Sykes L, Essop R. Combination intraoral and extraoral prosthesis used for rehabilitation of a
414 patient treated for cancrum oris: A clinical report. *J Prosthet Dent*. 2000;613–6.
- 415 15. Chidzonga M. HIV/AIDS orofacial lesions in 156 Zimbabwean patients at referral oral and
416 maxillofacial surgical clinics. *Oral Dis*. 2003;9:317–22.
- 417 16. Barrios T, Aria A, Brahney C. Cancrum oris in an HIV-positive patient. *J Oral Maxillofac*
418 *Surg*. 1995;53(7):851–5. doi: <https://www.ncbi.nlm.nih.gov/pubmed/7595806>
- 419 17. Fayrer J. A case of acute malarial poisoning, enteric fever ensuing, complicated by brain and
420 lung symptoms, also by extensive cancrum oris. *Lancet*. 1889(2):296–300.
- 421 18. Auluck A, Pai K. Noma: life cycle of a devastating sore - case report and literature review. *J*
422 *Can Dent Assoc*. 2005;71(10):757a-e.
- 423 19. Gupta P, Chakravarty N. Penicillin in Cancrum Oris Complicating Kala-Azar. *Ind Med Gaz*.
424 1945;542–5.
- 425 20. Usher S, Ross D. A case of cancrum oris following typhoid fever with plastic repair. *Can Med*

- 426 Assoc J. 1931;446–9.
- 427 21. Weledji E, Njong S. Cancrum Oris (noma): the role of nutrition in management. *J Am Coll*
428 *Clin Wound Spec* [Internet]. 2016;7:50–2. Available from:
429 <http://dx.doi.org/10.1016/j.jccw.2016.08.003>doi: 10.1016/j.jccw.2016.08.003
- 430 22. Chiandussi S, Luzzati R, Tirelli G, Lenarda R, Biasotto M. Cancrum oris in developed
431 countries. *Aging Clin Exp Res*. 2009;21(6):475–7.
- 432 23. Hefferman L. Cancrum oris treated by excision and subsequent tube grafting. *Ind Med Gaz*.
433 1923;434–7.
- 434 24. Jelliffe D. Infective gangrene of the mouth (Cancrum Oris). *Pediatrics*. 1952;9(544):1–9.
- 435 25. Barrera J, Connor M. Noma in an Afghani child: A case report. *Int J Pediatr Otorhinolaryngol*
436 [Internet]. 2012;76(5):742–4. Available from:
437 <http://dx.doi.org/10.1016/j.ijporl.2012.01.034>doi: 10.1016/j.ijporl.2012.01.034
- 438 26. Srour L, Watt B, Phengdy B, Khansoulivong K, Harris J, Bennett C, Strobel M, Dupuis C,
439 Newton P. Noma in Laos: Stigma of severe poverty in rural Asia. *Am J Trop Med Hyg*.
440 2008;78(4):539–42.
- 441 27. Maley A, Desai M, Parker S. Noma: A disease of poverty presenting at an urban hospital in the
442 United States. *JAAD Case Reports*. 2015;1(1):18–20. doi: 10.1016/j.jdcr.2014.10.001
- 443 28. Baratti-Mayer D, Gayet-Ageron A, Hugonnet S, François P, Pittet-Cuenod B, Huyghe A,
444 Bornand J, Gervaix A, Montandon D, Schrenzel J, Mombelli A, Pittet D. Risk factors for
445 noma disease: a 6-year, prospective, matched case-control study in Niger. *Lancet Glob Heal*.
446 2013;1(2):87–96. doi: 10.1016/s2214-109x(13)70015-9
- 447 29. Enwonwu C, Phillips R, Ferrell C. Temporal relationship between the occurrence of fresh
448 noma and the timing of linear growth retardation in Nigerian children. *Trop Med Int Heal*.

- 449 2005;10(1):65–73.
- 450 30. Oginni F, Oginni A, Ugboko V, Otuyemi O. A survey of cases of cancrum oris seen in Ile-Ife,
451 Nigeria. *Int J Paediatr Dent*. 1999;9:75–80.
- 452 31. World Health Organization, OMS. World Health Report Life in the 21st century A vision for
453 all Report of the Director-General. 51st World Heal Assem [Internet]. 1998;1–226. Available
454 from: https://www.who.int/whr/1998/en/whr98_en.pdf?ua=1
- 455 32. Fieger A, Marck KW, Busch R, Schmidt A. An estimation of the incidence of noma in north-
456 west Nigeria. *Trop Med Int Heal*. 2003;8(5):402–7. doi: 10.1046/j.1365-3156.2003.01036.x
- 457 33. Bello SA, Adeoye JA, Oketade I, Akadiri OA. Estimated incidence and prevalence of noma in
458 north central Nigeria, 2010–2018: A retrospective study. *PLoS Negl Trop Dis* [Internet].
459 2019;13(7):2010–8. Available from: <http://dx.doi.org/10.1371/journal.pntd.0007574>doi:
460 10.1371/journal.pntd.0007574
- 461 34. Farley E, Oyemakinde MJ, Schuurmans J, Ariti C, Saleh F, Uzoigwe G, Bil K, Oluyide B,
462 Fotso A, Amirtharajah M, Vyncke J, Brechard R, Adetunji AS, Ritmeijer K, Van Der Kam S,
463 Baratti-Mayer D, Mehta U, Isah S, Ihekweazu C, et al. The prevalence of noma in northwest
464 Nigeria. *BMJ Glob Heal*. 2020;5(4):1–15. doi: 10.1136/bmjgh-2019-002141
- 465 35. Médecins Sans Frontières. Risk factors for gingivitis in Anka General Hospital- internal report
466 from cross-sectional operational research study. Anka, Zamfara; 2018.
- 467 36. Emergency Nutrition Network Global Nutrition Cluster. The Harmonised Training Package
468 (HTP): Resource Material for Training on Nutrition in Emergencies, Version 2 [Internet].
469 NutritionWorks. 2011. Available from: [https://www.enonline.net/attachments/1051/m6-](https://www.enonline.net/attachments/1051/m6-measuring-malnutrition-individual-assessment-entire-module.pdf)
470 [measuring-malnutrition-individual-assessment-entire-module.pdf](https://www.enonline.net/attachments/1051/m6-measuring-malnutrition-individual-assessment-entire-module.pdf)
- 471 37. Azodo CC, Agbor AM. Gingival health and oral hygiene practices of schoolchildren in the
472 North West Region of Cameroon. *BMC Res Notes*. 2015;4–9. doi: 10.1186/s13104-015-1350-

473 2

474 38. Farley E. Noma in northwest Nigeria: a neglected disease in neglected populations [Internet].

475 University of Cape Town; 2020. Available from:

476 <https://open.uct.ac.za/handle/11427/32757?show=full>

477 39. Farley E, Mehta U, Srour ML, Lenglet A. Noma (cancrum oris): A scoping literature review of
478 a neglected disease (1843 to 2021). PLoS Negl Trop Dis [Internet]. 2021;15(12):e0009844.

479 Available from: <http://dx.doi.org/10.1371/journal.pntd.0009844>doi:

480 [10.1371/journal.pntd.0009844](https://doi.org/10.1371/journal.pntd.0009844)

481 40. Taiwo OO, Panas R. Roles of community pharmacists in improving oral health awareness in
482 Plateau State, Northern Nigeria. Int Dent J [Internet]. 2018;68(4):287–94. Available from:

483 <https://doi.org/10.1111/idj.12383>doi: [10.1111/idj.12383](https://doi.org/10.1111/idj.12383)

484 41. Trotter A, Cismas I. Noma & Human Rights Law – A Doctrinal Legal Analysis with Focus on
485 Burkina Faso, Niger and Laos, Background Study. 2020.

486 42. World Health Organisation (WHO) and Hilfsaktion Noma. Promoting Oral Health in Africa:
487 Prevention and control of oral diseases and noma as part of essential noncommunicable
488 disease interventions [Internet]. 2016. Available from:

489 <http://www.aho.afro.who.int/en/news/5236/official-launch-who-manual-“promoting-oral->
490 [health-africa”](http://www.aho.afro.who.int/en/news/5236/official-launch-who-manual-“promoting-oral-)

491 43. Vriese SD De. Moving from Daji towards Noma: Changing the perception of a spiritual
492 towards a treatable disease: A case study of Hilfsaktion Noma eV in Niger [Internet]. 2021.

493 Available from: [https://www.diva-](https://www.diva-portal.org/smash/record.jsf?pid=diva2:1573547%0Ahttps://www.diva-portal.org/smash/get/diva2:1573547/FULLTEXT01.pdf)

494 [portal.org/smash/record.jsf?pid=diva2:1573547%0Ahttps://www.diva-](https://www.diva-portal.org/smash/record.jsf?pid=diva2:1573547%0Ahttps://www.diva-portal.org/smash/get/diva2:1573547/FULLTEXT01.pdf)

495 [portal.org/smash/get/diva2:1573547/FULLTEXT01.pdf](https://www.diva-portal.org/smash/get/diva2:1573547/FULLTEXT01.pdf)

496 44. Agbelusi G, Jeboda S. Oral health status of 12-year old Nigerian children. WAJM.

497 2006;25(3):195–8.

498 45. Adeniyi AA, Sofola OO, Kalliecharan R V. An appraisal of the oral health care system in
499 Nigeria. *Int Dent J*. 2012 Dec;62(6):292–300. doi: 10.1111/j.1875-595x.2012.00122.x

500 46. Uguru N, Onwujekwe O, Ogu UU, Uguru C. Access to Oral health care: A focus on dental
501 caries treatment provision in Enugu Nigeria. *BMC Oral Health* [Internet]. 2020 May 19 [cited
502 2023 May 10];20(1):1–13. Available from:

503 <https://bmcoralhealth.biomedcentral.com/articles/10.1186/s12903-020-01135-1>doi:
504 10.1186/s12903-020-01135-1/figures/2

505 47. Uguru N, Onwujekwe O, Uguru C, Ogu U, Okwuosa C, Okeke C. Oral health-seeking
506 behavior among different population groups in Enugu Nigeria. *PLoS One* [Internet]. 2021 Feb
507 1 [cited 2023 May 10];16(2):e0246164. Available from:

508 <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0246164>doi:
509 10.1371/journal.pone.0246164

510 48. Konsem, T, Millogo, M, Assouan, C, Ouedraogo D. Evolving form of cancrum oris, about 55
511 cases collected at the Academic Hospital Yalgado Ouedraogo of Ouagadougou. *Bull Soc*
512 *Pathol Exot*. 2014;107:74–8. doi: 10.1007/s13149-014-0338-9

513