#### 1 COVID-19 vaccine information, misinformation, and vaccine uptake in Malawi.

- 2 John Songo<sup>1\*</sup>, Hannah S. Whitehead<sup>2\*</sup>, Khumbo Phiri<sup>1</sup>, Pericles Kalande<sup>1</sup>, Eric Lungu<sup>1</sup>, Sam Phiri<sup>1,3</sup>,
- 3 Joep J. van Oosterhout<sup>1,2</sup>, Agnes Moses<sup>1</sup>, Risa M. Hoffman<sup>2</sup>, Corrina Moucheraud<sup>4</sup>

#### 4 \* co-first authors

#### 5 Affiliations

- 6 <sup>1</sup>Partners in Hope, Lilongwe, Malawi;
- 7 <sup>2</sup>Department of Medicine, Division of Infectious Diseases, David Geffen School of Medicine, UCLA, Los
- 8 Angeles, California, USA;
- 9 <sup>3</sup>Department of Public Health and Family Medicine, Kamuzu University of Health Sciences, Lilongwe,
- 10 Malawi
- <sup>4</sup> Department of Health Policy and Management, Fielding School of Public Health, UCLA, Los Angeles,
- 12 California, USA
- 13 Corresponding Author: John Songo
- 14 Email: pemphosongo@gmail.com
- 15

16

17

#### 19 Abstract

20 Background: COVID-19 vaccine information – including source, content, and tone – may be an important 21 determinant of vaccination, but this dynamic is not well-understood in low-income countries where COVID-19 vaccine uptake remains low. We assessed the COVID-19 vaccine information environment in 22 23 Malawi, and its correlation with vaccine uptake. *Methods:* A survey was administered among 895 adult ( $\geq$ 18 years) clients at 32 Malawian health facilities 24 25 in mid-2022. Respondents reported their COVID-19 vaccination history, exposure to information about 26 the COVID-19 vaccine from different sources and its tone (positive, negative, or neutral/factual), and 27 whether they had heard of and believed in ten COVID-19 and COVID-19 vaccine conspiracy theories. 28 We described the COVID-19 vaccine information environment in Malawi and used logistic regression 29 analyses to assess the association of exposure to information sources and conspiracy theories with uptake of the COVID-19 vaccine. 30 31 Results: Respondents had received information about the COVID-19 vaccine most commonly from 32 friends and neighbors, healthcare workers, and radio (each reported by >90%). Men, urban residents, and 33 respondents with a higher education level were exposed to more COVID-19 vaccine information sources.

34 COVID-19 vaccine uptake was positively associated with exposure to a greater number of COVID-19

vaccine information sources (aOR 1.09, 95% CI 1.03-1.15), and more positive information (aOR 4.33,

36 95% CI 2.17-8.64) – and was negatively associated with believing COVID-19 vaccine conspiracy

37 theories to be true (OR 0.76, 95% CI 0.68-0.87).

*Conclusions:* Malawian adults were exposed to a variety of COVID-19 vaccine information sources, with
 less access to information among women, rural residents, and people with lower educational attainment.

40 Exposure to misinformation was common, though infrequently believed. Vaccination was associated with

41 exposure to high number of COVID-19 vaccine information sources, exposure to positive vaccine

42 information and endorsing fewer conspiracy theories. Vaccination programs should disseminate

43 communication with positive messaging, through multiple information sources, prioritizing the less

44 exposed groups we identified.

- 45
- 46
- 47
- 48 Key words: COVID-19, COVID-19 vaccine, vaccine up-take, COVID-19 vaccine information.

## 49 Background

50	During public health crises, information can shape individuals' knowledge of and/or attitudes about health
51	interventions, and ultimately inform their behaviors (1,2). The COVID-19 pandemic, which occurred
52	amidst widespread technology use and social media connectedness, was deemed an "infodemic" by the
53	World Health Organization – i.e., it was characterized by an "overabundance of information," both
54	accurate and misleading, about the virus, its origins, and control measures, including vaccines (3).
55	Misinformation about vaccines - i.e., incorrect or misleading information - can contribute to vaccine
56	hesitancy (4). The spread of vaccination misinformation accelerated during the COVID-19 pandemic (5)
57	and has been linked to decreased COVID-19 vaccination intent and behavior (6,7).
58	Two years into widespread availability of COVID-19 vaccines, most people in low-income countries have
59	still not been vaccinated (8); and the COVID-19 information and misinformation landscape in these
60	settings - and its association with vaccine uptake - remains poorly understood (9). In Malawi, a low-
61	income country in southeastern Africa that has recorded nearly 90,000 cumulative confirmed COVID-19
62	cases and close to 3,000 deaths (10), only approximately 1 in 4 people have received a COVID-19
63	vaccination (11). Misconceptions and conspiracy theories about the COVID-19 vaccine have been
64	documented in Malawi and throughout the region (12-14).
65	Globally, much of the literature on the COVID-19 "infodemic" focuses on the dissemination of
66	(mis)information online and via social media (15,16), but it is not well-understood whether the internet is
67	a major source of COVID-19 information in low-income countries. In countries like Malawi, access to
68	technology is rapidly changing and there is a need to understand what this means for the spread of
69	information during public health crises like the COVID-19 pandemic. Understanding what information
70	people are exposed to, the sources and tone of this information, and how this information influences
71	behaviors may help to inform vaccination promotion strategies in settings with low vaccine uptake. To
72	this end, we conducted a survey to describe the information environment about COVID-19 and the
73	COVID-19 vaccine in Malawi, and assessed how individuals' exposure to information (from different
74	sources, of differing tone) is correlated with vaccine uptake.

75

#### 76 METHODS

#### 77 Study design, site & participant selection

We conducted a cross-sectional survey of adults presenting at 32 purposefully-selected health facilities 78 79 supported by Partners In Hope (PIH), a Malawian non-governmental organization that assists with 80 implementation of the National HIV program across Malawi. Sites were selected to represent public and 81 faith-based health facilities in urban, peri-urban, and rural areas across all three of Malawi's regions 82 (Northern, Central, and Southern). Eligible clients were at least 18 years old and visited a selected facility 83 for care at an outpatient department (OPD), antiretroviral therapy (ART), or non-communicable disease 84 (NCD) clinic during a time when the COVID-19 vaccine was available to the general public in Malawi. 85 We used systematic random sampling (every second individual in queue to see a provider was approached) to invite individuals to participate in the survey. 86

#### 87 Survey Domains and Variable Definitions

88 The survey included questions asking whether respondents had been exposed to COVID-19 vaccine

89 information from different sources, and if so, whether the information was positive, negative, or

90 neutral/factual in tone. Respondents were also asked whether they had heard of ten common conspiracy

91 theories about COVID-19 and the COVID-19 vaccine and if they believed them to be true; these 10

92 pieces of misinformation were selected by study team members as relevant to and circulating in Malawi

from among those included in two prior COVID-19 surveys done in the UK and USA (17,18). We asked

94 respondents about their experiences with COVID-19 and with COVID-19 vaccines, including vaccination

95 details for those vaccinated (number of doses received, location, manufacturer, timing).

96 The survey instrument was developed in English and translated to Chichewa, the local language; it was

- 97 then reviewed by bilingual (English/Chichewa) study team members to ensure clarity and meaning.
- 98 Exposure to COVID-19 vaccine information was assessed by asking respondents if they had heard any
- 99 information about the COVID-19 vaccine from thirteen different potential sources of information,

100 spanning traditional media (newspaper; radio; television), health system and the government (health care 101 provider; government/Ministry of Health), community figures (church or religious leaders; traditional 102 medicine practitioner), personal relations (family member; friend; neighbor/community member), and 103 social media (posts on Facebook, Instagram or Twitter from a known person; posts from an organization, 104 company, or someone not known personally; conversations/groups in a messaging app such as WhatsApp 105 or Facebook Messenger). In analysis, "friend" and "neighbor/community member" were combined and 106 considered as a single information source. Exposure to information was quantified as the *count* of the 107 number of different information sources reported, as well as the *proportion* of respondents who had heard 108 information from a source.

109 *Tone of COVID-19 vaccine information* was measured as follows: for each source that a respondent

110 reported hearing COVID-19 vaccine information from, they were asked whether they felt the information

111 was overall positive, negative, neutral/factual, a mix of positive and negative, or that they did not know

the tone of the information. In analysis, these responses were combined into three categories: positive or

neutral, mixed tone or unknown, and negative. To quantify tone of information, for each respondent, we

114 calculated the *proportion* of the information sources they reported exposure to that were positive or

negative in tone (i.e., number of positive information sources divided by the total number of information

sources exposed to), as well as *counts* of the number of positive or negative information sources.

117 Respondents were also asked who they *trusted the most* to give them advice regarding COVID-19

118 vaccine.

*Vaccine uptake* was defined as having received any (1 or more) doses of any manufacturer's COVID-19
vaccine (Johnson & Johnson, AstraZeneca or Pfizer).

121 *Exposure to COVID-19 misinformation* was characterized by asking whether the respondents had heard

122 10 common conspiracy theories about COVID-19 and the COVID-19 vaccine (17,18) and quantified as

the *count* of heard conspiracy theories. For each conspiracy theory that the respondent had heard of, they

124 were asked whether they believed it to be true, false, or did not know whether it was true or false.

125 Endorsement of COVID-19 misinformation was quantified as both the proportion of heard conspiracies

- 126 believed to be true (number of conspiracies believed to be true divided by the number of conspiracy
- 127 theories heard), as well as the *count* of conspiracies believed to be true.
- 128

#### 129 Data Collection

- 130 The survey was administered face-to-face as an interviewer-administered survey, in a private area within
- the health facility by a trained research assistant. All respondents provided oral informed consent for
- anonymous data collection prior to commencing the survey, and all responses were recorded using the
- 133 SurveyCTO mobile data collection platform on Android tablets. All respondents received 4000 Malawi
- 134 kwacha (approximately US\$5) as compensation for opportunity costs.
- 135 Data were collected from 19 May to 30 June, 2022. At the time of data collection, all adults 18 years of

age or older were eligible to receive the COVID-19 vaccine in Malawi. Ethical approval for the study

137 was obtained from the National Health Sciences Research Committee in Malawi (#2883) and the

138 University of California Los Angeles Institutional Review Board (#22-000380).

139

#### 140 Data analysis

We described sample characteristics, exposure to, and tone of COVID-19 vaccine information, and 141 142 exposure to and endorsement of COVID-19 misinformation. Exposure to COVID-19 conspiracy theories 143 was also visualized with a heat map table, in which cells were color-coded ranging from bright green (smallest proportion of respondents had heard theory) to red (highest proportion of respondents had heard 144 theory). Chi-square tests, t-tests, and multivariable logistic regression models were used to assess whether 145 146 these variables were associated with various socio-demographic characteristics. We used multivariable 147 logistic regression models to assess whether COVID-19 vaccine uptake was associated with exposure to 148 and tone of COVID-19 vaccine information, or exposure to and endorsement of COVID-19 vaccine 149 misinformation. All multivariable logistic models were adjusted for gender, residence (urban/rural), age, 150 HIV status, and education. All analyses were conducted using Stata v17.

#### 151

#### 152 **RESULTS**

- 153 A total of 944 individuals were approached for participation in this survey; 46 declined to be screened or
- to consent to participation, and 3 were found to be ineligible (<18 years of age) during screening. The
- remaining 895 individuals (94.8% of those approached) completed the survey and are included in this
- analysis. Nearly half (43%) of respondents had received at least one dose of a COVID-19 vaccine. Over
- half of the respondents were female (57%) and three-quarters (75%) were married (Table 1). The majority
- 158 of the respondents were residing in rural areas (82%), were employed (68%) and identified as Christian
- 159 (91%).
- 160

Table 1: Sample characteristics (n=895)		
	n	%
Gender		
Male	382	42.7
Female	513	57.3
Age		
18-29	216	24.1
30-39	239	26.7
40-49	225	25.1
50-59	122	13.6
60+	93	10.4
Marital status		
Unmarried	226	25.3
Married	669	74.7
Place of residence		
Urban	158	17.7
Rural	737	82.3
Religious		
Christian	818	91.4
Other religion	50	5.6
Not religious	27	3.0

Educational attainment		
No school	250	27.9
Primary school	401	44.8
Secondary school	213	23.8
Beyond secondary	31	3.5
Employment status		
Employed	610	68.2
Not employed	285	31.8
Household income over past 12 months		
Insufficient	277	32.7
Just met expenses	443	52.3
Allowed for saving	127	15.0
COVID-19 vaccination status		
Unvaccinated	510	57.0
Vaccinated (1+ doses of any COVID-19 vaccine)	385	43.0

#### 162

#### 163 Information about COVID-19 vaccines

#### 164 *Exposure to information about COVID-19 vaccine*

165 All but one survey respondent had heard information about COVID-19 vaccines from at least one source.

166 Respondents had been exposed to information from a median of 7 sources (IQR 6-9). The most common

sources were friends and neighbors, health care workers, and radio – each reported by >90% of

respondents (Figure 1). Receiving vaccine information from a traditional medicine practitioner was least

169 common (3% of respondents), followed by social media, messaging apps, television and newspapers,

- each reported by 25-35% of respondents.
- 171 Figure 1: Exposure to information about COVID-19 vaccine, by source and tone.

- 173 Men, urban residents, and respondents with a higher education level were exposed to a higher mean
- number of COVID-19 vaccine information sources (men 8.2 vs. women 6.9; p<0.001; urban 8.6 vs. rural

175	7.2; p<0.001; secondary or higher education 9.3 vs. primary school or less 6.8; p<0.001). No differences
176	were seen in the mean number of information sources by age, HIV status, or marital status.
177	
178	Among specific information sources, men were significantly more likely than women to have heard
179	COVID-19 information from nearly all sources, as were people with more educational attainment (Figure
180	2). Significantly less rural respondents reported exposure to information from traditional media
181	(newspaper, television, radio) and social media than rural residents (OR 0.23, 95%CI: 0.079-0.609, p=
182	0.004)
183	
184	Figure 2: Odds of having received COVID-19 vaccine information from various sources, by respondent
185	characteristics.
186	
187	Tone of information about COVID-19 vaccine
188	Information about COVID-19 from health care workers, the government/Ministry of Health, religious
189	leaders, friends and family, and traditional media (newspapers, television, radio) was largely (>80%)
190	perceived as positive or neutral (Figure 1). Approximately one-fifth of people exposed to COVID-19
191	vaccine information on social media and messaging apps felt it was negative in tone and so did half of the
192	people exposed to COVID-19 vaccine information from traditional medicine practitioners.
193	
194	Women reported that a greater share of their information sources about COVID-19 vaccine was negative
195	in tone: on average, women considered 10.2% of their information sources about COVID-19 vaccines
196	negative in tone, versus 7.6% for men (p=0.015). Respondents with lower educational attainment reported
197	that a lower share of their information sources was negative in tone (29% for those with a primary school
198	education or less, versus 32% for those with secondary school education or higher, p=0.034). No
199	differences were seen by HIV status, area of residence, or marital status.
200	

#### 201 *Most trusted information sources*

202	Respondents were asked who they trust most for advice regarding the COVID-19 vaccine. A healthcare
203	provider was selected most often (69.4%), followed by a community health worker (22.1%); family,
204	friends, religious leaders, and celebrities were selected infrequently (0.2-5% of respondents each).
205	Unvaccinated individuals were twice as likely to most trust family, friends, or a religious leader for
206	COVID-19 vaccine information as individuals who had been vaccinated (10.6% versus 5.2%, p=0.017).
207	No significant differences in trusted information were seen by gender, HIV status, age, or education level.
208	
209	Association of COVID-19 vaccine information with vaccine uptake
210	Having received information about COVID-19 vaccines from more sources was positively associated
211	with COVID-19 vaccine uptake, with each additional source increasing vaccination probability by 9%
212	(OR 1.09, 95% CI 1.03-1.15). Among subgroups, this relationship was found primarily among men (OR:
213	1.13, 95% CI: 1.04-1.23), respondents 40 years and older (OR: 1.15, 95% CI: 1.06-1.24), people living
214	with HIV (OR: 1.17, 95% CI: 1.09-1.27), and those with a secondary school education or higher (OR:
215	1.15, 95% CI: 1.03-1.28) (Appendix 1). Adjusting for gender, residence (urban/rural), age, HIV status,
216	and education, the positive association between information exposure (number of reported sources) and
217	COVID-19 vaccination persisted (aOR 1.08, 95% CI 1.01-1.15) and the odds of vaccination was higher
218	when respondents reported a greater proportion of COVID-19 information being positive in tone (aOR
219	4.33, 95% CI 2.17-8.64).

220

#### 221 Misinformation about COVID-19 vaccines

222 Exposure to misinformation about COVID-19 and COVID-19 vaccines

223 Respondents reported having heard a median of 5 of the 10 surveyed conspiracy theories. The most

commonly heard were "The spread of COVID-19 is a deliberate attempt to reduce the size of the global

- population" (80.7% of respondents) and "The COVID-19 virus is a hoax" (74.1% of respondents) (Figure
- 226 3).

227

228 Figure 3: Heat map displaying the proportion of survey respondents (overall, by gender, and by place of 229 residence) who reported hearing of 10 common COVID-19 conspiracies. Color scale spans from bright 230 green (smallest proportion) to red (highest proportion). 231 Men, respondents with higher educational attainment, and urban respondents were exposed to a higher 232 233 mean number of conspiracy theories: men 5.5 versus women 4.8 (p=0.0001); urban residents 5.7 versus 234 rural residents 5.0 (p=0.002), and individuals with a secondary school education or higher heard 5.9 235 versus those with primary school or less 4.8 (p < 0.001). 236 Endorsing misinformation about COVID-19 vaccines 237 Most respondents (72%) did not believe any of the 10 conspiracy theories to be true. Overall, respondents 238 239 endorsed (believed to be true) a mean of 0.56 conspiracy theories; this increased to a mean of 1.96 among the 254 respondents who believed in at least one conspiracy theory. Younger age groups, urban 240 241 respondents and respondents with higher education attainment endorsed more conspiracy theories: 18–39-242 year-olds endorsed 0.66 conspiracy theories versus 0.45 among 40+ year olds (p = 0.007), urban 243 respondents endorsed 0.75 versus 0.51 among rural respondents (p=0.01), and individuals with a 244 secondary school education or higher endorsed 0.76 versus 0.48 among those with primary school or less 245 (p=0.001) 246 247 The most commonly endorsed myths were "The government is exaggerating the number of COVID-19 deaths" (22.7%), "Big pharmaceutical companies created COVID-19 to profit from vaccines" (17.3%), 248 249 "The spread of COVID-19 is a deliberate attempt to reduce the size of the global population" (16.9%) and 250 "Bill Gates has created COVID-19 in order to reduce the world population" (15.2%) (Table 2). 251

252 Association with COVID-19 vaccine uptake

254	Exposure to a greater number of COVID-19 conspiracy theories was associated with increased odds of
255	vaccine uptake (OR 1.10, 95% CI 1.05-1.16), while endorsing more conspiracy theories was associated
256	with lower odds of COVID-19 vaccine uptake (OR 0.76, 95% CI 0.68-0.87). This association between
257	conspiracy theory endorsement and vaccine uptake remained significant in univariate analyses among all
258	considered subgroups (by gender, HIV status, age, and education) except urban residents (Appendix
259	Table 2).

260

Table 2: Endorsement of 10 common conspiracy theories and association with COVID-19 vaccine uptake					
	% Believe true, among those who theory		of vaccination theory to be t	ation if believed to be true	
Conspiracy theory have heard of the conspiracy theory		OR	95% CI	p-value	
The government is exaggerating the number of COVID-19 deaths	22.7	0.75	0.50-1.10	0.14	
Big pharmaceutical companies created COVID-19 to profit from vaccines	17.3	0.44	0.26-0.75	0.00	
The spread of COVID-19 is a deliberate attempt to reduce the size of the global population	16.9	0.33	0.21-0.52	0.00	
Bill Gates has created COVID-19 in order to reduce the world population	15.2	0.4	0.20-0.80	0.01	
Deaths due to COVID-19 are being intentionally hidden by the government	10.7	0.71	0.34-1.50	0.37	
The COVID-19 vaccine can change your DNA	8.4	0.13	0.04-0.46	0.00	
You can get COVID-19 from the vaccine	5.5	0.23	0.07-0.82	0.02	
The COVID-19 vaccine will be used to carry out mass sterilization	4.1	0.27	0.09-0.82	0.01	
COVID-19 vaccines have been shown to cause infertility	3.1	0.23	0.06-0.81	0.02	
The COVID-19 virus is a hoax	2.9	0.22	0.06-0.75	0.01	

261

262

263 Respondents who believed that COVID-19 vaccines can change your DNA had the lowest odds of

vaccination (OR 0.13, 95% CI 0.04-0.46), followed by those who believed that the COVID-19 virus is a

hoax (OR 0.22, 95% CI 0.06-0.75) compared to those who believed these conspiracy theories to be false

or didn't know. Believing that the government is exaggerating the number of COVID-19 deaths and that

deaths due to COVID-19 are being intentionally hidden by government were not associated with vaccine

- uptake.
- 269

270 **DISCUSSION** 

In a survey of adults presenting at health facilities in Malawi conducted while the COVID-19 vaccine was
available to the general public, we found that people hear COVID-19 vaccine information from a variety
of sources, and exposure to information – including the source, content, and tone – is associated with
COVID-19 vaccine uptake. Exposure to more sources of information and to positive information were
strongly associated with vaccine uptake. Endorsement of common COVID-19 conspiracy theories –
though quite uncommon – was associated with decreased odds of vaccination.

277

278 Respondents heard COVID-19 vaccine information from a mean of 7 different sources. The 3 most 279 common sources were peers, health care providers, and the radio. Information from health care providers, the Ministry of Health, traditional media, and church or religious leaders was mostly perceived as 280 positive. Health care providers were reported as the most trusted source of COVID-19 vaccine 281 282 information, as has been reported in other settings in sub-Saharan Africa (19,20). Information from peers 283 (friends, neighbors, and community members) was largely mixed in tone, while information from social media, messaging apps, and traditional medicine practitioners was mostly perceived to be negative in 284 285 tone. We found that the odds of vaccine uptake increased as the proportion of information sources that 286 were positive in tone increased. Additionally, most people in our sample had heard several conspiracy 287 theories about COVID-19, and endorsement of misinformation, while uncommon, was associated with 288 decreased odds of vaccination. Negative vaccine information and misinformation have been reported in Malawi (21) and the harmful impacts on vaccine uptake have been well researched (5,15,16). Our 289 findings underline the need for greater monitoring and action against vaccine information that is negative 290 291 in tone as well as information that is intentionally misleading (15).

292

The degree of exposure to COVID-19 vaccine information varied across socio-demographic groups, with men, urban respondents, and individuals with more education tending to report exposure to a greater number of both vaccine information sources and conspiracy theories, likely reflecting greater literacy, internet/mobile penetration, and media access among these groups. Given the association between greater

297 information exposure and vaccine uptake, these findings bolster previous studies that have found higher 298 COVID-19 vaccine acceptance among males and individuals with more education (22). We also found 299 greater endorsement of conspiracy theories among younger, less educated and urban respondents, 300 possibly explaining findings from previous studies in sub-Saharan Africa that younger (23) and urban 301 (24) respondents are more likely to be vaccine hesitant or believe in certain COVID-19 conspiracy 302 theories, but conflicting with findings from Ghana that individuals with higher education are more likely 303 to be vaccine hesitant(24). Our findings highlight demographic groups that may need to be reached with 304 COVID-19 vaccine information, as well as where to target efforts at countering misinformation.

305

Our results point to important implications for programming to encourage COVID-19 vaccine uptake. 306 307 Our finding that exposure to more information sources and information that is positive in tone is 308 associated with increased vaccine uptake adds to an emerging literature on this topic (25,26) and is the 309 first of its kind in Africa. This positive relationship has direct implications for communication strategies 310 aimed at increasing vaccination. Campaigns should disseminate positive information through a variety of 311 sources, and aim to reach less-exposed populations (such as women, rural residents, and those with less 312 education) via multiple information sources, particularly those that have been shown to predominantly 313 disseminate positive vaccine information (e.g., radio, television, or health care workers). Trust in 314 healthcare workers should be leveraged by positioning them as spokespeople and vaccine "champions", and by training them to share accurate, factual information about COVID-19 vaccines during one-on-one 315 316 clinical encounters. While found to be low in our sample, endorsement of misinformation should be 317 closely monitored, and vaccination programs may need to counter misinformation and conspiracy theories through culturally appropriate messaging targeted at the groups most susceptible to misinformation (15). 318 319 Effective approaches would address and correct both negative and false information, particularly as 320 spread by community members and via social media. More research is needed to identify effective and 321 locally-relevant strategies for this in Malawi and similar contexts; one promising approach is a 322 WhatsApp-based counselling intervention for countering COVID-19 fake news trialed in Nigeria (27).

2	2	2
.5	2	.5
-	_	-

324	We note several limitations of this study. The study respondents were recruited at health facilities hence
325	our sample is very likely to over-represent people with stronger trust in health care services, including
326	COVID-19 vaccination. Vaccination status was ascertained based on self-report, which may have
327	overestimated coverage and underestimated due to social desirability bias.
328	
329	Conclusion:
330	This study represents one of the first in-depth explorations of the COVID-19 vaccine information
331	environment and how it affects vaccine uptake in sub-Saharan Africa. Our findings provide important
332	insights on how people's vaccination behavior may be shaped by their local vaccine information sources,
333	tone, and content. Investment is needed in targeted information interventions that leverage prominent and
334	trusted sources of COVID-19 vaccine information, are designed to reach all socio-demographic groups,
335	and combat misinformation.
336	

## 337 Acknowledgements

- 338 We are grateful for the support of the participating health facilities and their management teams for
- allowing us to conduct surveys at the health facilities. We are also thankful to the research assistants who
- 340 conducted the surveys.

### 342 APPENDIX

343 Appendix 1: Association between hearing more information sources about COVID-19 vaccine, and

344 COVID-19 vaccination, unadjusted odds ratios

Subgroup	COVID-19 vaccination (OR)	95% CI	p-value
Male	1.13	1.04-1.23	0.004
Female	1.05	0.98-1.13	0.136
Urban	1.12	1.00-1.25	0.054
Rural	1.08	1.02-1.15	0.009
HIV+	1.17	1.09-1.27	0.000
HIV-	1.01	0.94-1.09	0.724
18-39 years	1.05	0.98-1.13	0.169
40+ years	1.15	1.06-1.24	0.001
Primary school or less	1.06	0.99-1.13	0.111
Secondary school or higher	1.15	1.03-1.28	0.010

345

346 Appendix 2: Association between endorsing more conspiracy theories about COVID-19 and COVID-19

#### 347 vaccine uptake

Subgroup	OR for vax uptake	95% CI	p-value
Male	0.70	0.57-0.86	0.001
Female	0.82	0.68-0.98	0.026
Urban	0.86	0.67-1.09	0.209
Rural	0.72	0.61-0.85	< 0.001
HIV+	0.77	0.64-0.94	0.009
HIV-	0.75	0.62-0.91	0.004
18-39 years	0.82	0.68-0.99	0.034
40+ years	0.74	0.61-0.91	0.004
Primary school or less	0.75	0.63-0.90	0.002
Secondary school or	0.75	0.60-0.92	0.006
higher			

#### 349 **REFERENCE**

- Anwar A, Malik M, Raees V, Anwar A. Role of Mass Media and Public Health Communications in the COVID-19 Pandemic. Cureus [Internet]. 2020 Sep 14; Available from: https://www.cureus.com/articles/38293-role-of-mass-media-and-public-health-communications-
- 353 in-the-covid-19-pandemic
- Garfin DR, Silver RC, Holman EA. The novel coronavirus (COVID-2019) outbreak:
   Amplification of public health consequences by media exposure. Heal Psychol [Internet]. 2020 May;39(5):355–7. Available from: http://doi.apa.org/getdoi.cfm?doi=10.1037/hea0000875
- World Health Organization. The COVID-19 infodemic [Internet]. Available from:
   https://www.who.int/health-topics/infodemic/the-covid-19-infodemic#tab=tab
- 4. Carrieri V, Madio L, Principe F. Vaccine hesitancy and (fake) news: Quasi-experimental evidence from Italy. Health Econ [Internet]. 2019 Nov 20;28(11):1377–82. Available from:

361 https://onlinelibrary.wiley.com/doi/10.1002/hec.3937

- 362 5. Naeem S Bin, Bhatti R, Khan A. An exploration of how fake news is taking over social media and putting public health at risk. Heal Inf Libr J [Internet]. 2021 Jun 12;38(2):143–9. Available from: https://onlinelibrary.wiley.com/doi/10.1111/hir.12320
- 365
  6. Roozenbeek J, Schneider CR, Dryhurst S, Kerr J, Freeman ALJ, Recchia G, et al. Susceptibility to
  366 misinformation about COVID-19 around the world. R Soc Open Sci [Internet]. 2020 Oct
  367 14;7(10):201199. Available from: https://royalsocietypublishing.org/doi/10.1098/rsos.201199
- Loomba S, de Figueiredo A, Piatek SJ, de Graaf K, Larson HJ. Measuring the impact of COVIDvaccine misinformation on vaccination intent in the UK and USA. Nat Hum Behav [Internet].
  2021 Feb 5;5(3):337–48. Available from: https://www.nature.com/articles/s41562-021-01056-1
- 8. United Nations Development Programme (UNDP). Global Dashboard for Vaccine Equity
  372 [Internet]. Available from: https://data.undp.org/vaccine-equity/
- Safary E, Mtaita C. A qualitative exploration of perceptions of the COVID-19 vaccine in Malawi during the vaccine rollout phase. One Heal Implement Res [Internet]. 2022;2(2):79–87. Available from: https://ohirjournal.com/article/view/4965
- Ministry of Health Malawi. COVID-19 National Information Dashboard [Internet]. Available
   from: https://covid19.health.gov.mw/
- 378 11. OurWorldInData.org. Coronavirus (COVID-19) Vaccinations [Internet]. Available from:
   379 https://ourworldindata.org/covid-vaccinations
- Aron MB, Connolly E, Vrkljan K, Zaniku HR, Nyirongo R, Mailosi B, et al. Attitudes toward COVID-19 Vaccines among Patients with Complex Non-Communicable Disease and Their Caregivers in Rural Malawi. Vaccines [Internet]. 2022 May 17;10(5):792. Available from: https://www.mdpi.com/2076-393X/10/5/792
- Wonodi C, Obi-Jeff C, Adewumi F, Keluo-Udeke SC, Gur-Arie R, Krubiner C, et al. Conspiracy
  theories and misinformation about COVID-19 in Nigeria: Implications for vaccine demand
  generation communications. Vaccine [Internet]. 2022;40(13):2114–21. Available from:
  https://www.sciencedirect.com/science/article/pii/S0264410X22001268
- 388 14. Dereje N, Tesfaye A, Tamene B, Alemeshet D, Abe H, Tesfa N, et al. COVID-19 vaccine
  389 hesitancy in Addis Ababa, Ethiopia: a mixed-method study. BMJ Open [Internet]. 2022 May
  390 1;12(5):e052432. Available from: http://bmjopen.bmj.com/content/12/5/e052432.abstract
- 391 15. Gisondi MA, Barber R, Faust JS, Raja A, Strehlow MC, Westafer LM, et al. A Deadly Infodemic:
  392 Social Media and the Power of COVID-19 Misinformation. J Med Internet Res [Internet]. 2022
  393 Feb 1;24(2):e35552. Available from: https://www.jmir.org/2022/2/e35552
- Cinelli M, Quattrociocchi W, Galeazzi A, Valensise CM, Brugnoli E, Schmidt AL, et al. The
  COVID-19 social media infodemic. Sci Rep [Internet]. 2020 Oct 6;10(1):16598. Available from:
  https://www.nature.com/articles/s41598-020-73510-5
- Freeman D, Loe BS, Chadwick A, Vaccari C, Waite F, Rosebrock L, et al. COVID-19 vaccine
  hesitancy in the UK: the Oxford coronavirus explanations, attitudes, and narratives survey

399		(Oceans) II. Psychol Med [Internet]. 2022 Oct 11;52(14):3127–41. Available from:
400		https://www.cambridge.org/core/product/identifier/S0033291720005188/type/journal article
401	18.	Hamel, Liz; Lopes, Luna; Ashley, Kirzinger; Sparks, Grace; Broddie M. KFF COVID-19 Vaccine
402		Monitor: Media and Misinformation. KFF [Internet]. 2018; Available from:
403		https://www.kff.org/coronavirus-covid-19/poll-finding/kff-covid-19-vaccine-monitor-media-and-
404		misinformation/
405	19.	Katoto PDMC, Parker S, Coulson N, Pillay N, Cooper S, Jaca A, et al. Predictors of COVID-19
406		Vaccine Hesitancy in South African Local Communities: The VaxScenes Study. Vaccines
407		[Internet]. 2022 Feb 25;10(3):353. Available from: https://www.mdpi.com/2076-393X/10/3/353
408	20.	Solís Arce JS, Warren SS, Meriggi NF, Scacco A, McMurry N, Voors M, et al. COVID-19
409		vaccine acceptance and hesitancy in low- and middle-income countries. Nat Med [Internet]. 2021
410		Aug 16;27(8):1385–94. Available from: https://www.nature.com/articles/s41591-021-01454-y
411	21.	Chimatiro C, Hajison P, Jella C, Tshotetsi L, Mpachika-Mfipa F. Barriers affecting COVID-19
412		vaccination in Phalombe District, Malawi: A qualitative study. South African Med J [Internet].
413		2023 Mar 8; Available from: https://samajournals.co.za/index.php/samj/article/view/842
414	22.	Moola S, Gudi N, Nambiar D, Dumka N, Ahmed T, Sonawane IR, et al. A rapid review of
415		evidence on the determinants of and strategies for COVID-19 vaccine acceptance in low- and
416		middle-income countries. J Glob Health [Internet]. 2021 Nov 20;11:05027. Available from:
417		http://jogh.org/documents/2021/jogh-11-05027.pdf
418	23.	Ovenseri-Ogbomo G, Ishaya T, Osuagwu UL, Abu EK, Nwaeze O, Oloruntoba R, et al. Factors
419		associated with the myth about 5G network during COVID-19 pandemic in sub-Saharan Africa. J
420		Glob Heal Reports [Internet]. 2020 Nov 3;4. Available from: https://www.joghr.org/article/17606-
421		factors-associated-with-the-myth-about-5g-network-during-covid-19-pandemic-in-sub-saharan-
422		africa
423	24.	Brackstone K, Atengble K, Head M, Boateng L. COVID-19 vaccine hesitancy trends in Ghana: a
424		cross-sectional study exploring the roles of political allegiance, misinformation beliefs, and
425		sociodemographic factors. Pan Afr Med J [Internet]. 2022;43. Available from:
426		https://www.panafrican-med-journal.com/content/article/43/165/full
427	25.	Kamal A, Hodson A, Pearce JM. A Rapid Systematic Review of Factors Influencing COVID-19
428		Vaccination Uptake in Minority Ethnic Groups in the UK. Vaccines [Internet]. 2021;9(10).
429		Available from: https://www.mdpi.com/2076-393X/9/10/1121
430	26.	Umakanthan S, Patil S, Subramaniam N, Sharma R. COVID-19 Vaccine Hesitancy and Resistance
431		in India Explored through a Population-Based Longitudinal Survey. Vaccines [Internet].
432		2021;9(10). Available from: https://www.mdpi.com/2076-393X/9/10/1064
433	27.	Talabi FO, Ugbor IP, Talabi MJ, Ugwuoke JC, Oloyede D, Aiyesimoju AB, et al. Effect of a
434		social media-based counselling intervention in countering fake news on COVID-19 vaccine in
435		Nigeria. Health Promot Int [Internet]. 2022 Apr 29;37(2). Available from:
436		https://academic.oup.com/heapro/article/doi/10.1093/heapro/daab140/6369161
437		

	% of respondents reporting they have heard conspiracy theory				
Conspiracy Theory	Overall	Male	Female	Urban	Rural
The spread of COVID-19 is a deliberate attempt to reduce the size of the global population	80.7	85.0	77.4	82.9	80.2
The COVID-19 virus is a hoax	74.1	77.7	71.3	79.7	72.9
The government is exaggerating the number of COVID-19 deaths	65.9	70.9	62.2	77.2	63.5
COVID-19 vaccines have been shown to cause infertility	54.3	55.5	53.4	58.2	53.5
The COVID-19 vaccine will be used to carry out mass sterilization	51.4	52.9	50.3	51.3	51.4
Big pharmaceutical companies created COVID-19 to profit from vaccines	47.2	52.4	43.3	60.1	44.4
You can get COVID-19 from the vaccine	36.9	41.6	33.3	38.6	36.5
Deaths due to COVID-19 are being intentionally hidden by the government	34.5	35.1	34.1	35.4	34.3
The COVID-19 vaccine can change your DNA	33.3	39.5	28.7	36.1	32.7
Bill Gates has created COVID-19 in order to reduce the world population	30.1	40.3	22.4	50.0	25.8

Figure 3



\*Traditional media= newspapers, radio, tv; Traditional medicine practitioner/Religious leaders; social media = social media and messaging apps; Personal relations = family and friends; Healthcare worker/MoH = healthcare worker & Ministry of health

## Figure 2

	% have heard information from source	% positive or neutral	% mixed tone or unknown	% negative
Friend, neighbor, community member	94%	23%	66%	11%
Health care provider	91%	99%	1%	0%
Radio	90%	98%	2%	0%
Family member	83%	62%	25%	12%
Church or religious leader	80%	86%	5%	9%
Government/Ministry of Health	75%	99%	1%	0%
Television	34%	97%	3%	0%
Newspapers	29%	95%	5%	0%
Conversation/group in messaging app	28%	22%	55%	22%
Social media post from unknown person or organization	26%	30%	49%	21%
Social media post from friend or family	25%	27%	53%	20%
Traditional medicine practitioner	3%	52%	7%	41%

Created with Datawrapper

# Figure 1