

1

2 **Understanding the fine-scale heterogeneity and spatial drivers of**
3 **malaria transmission using model-based geostatistical methods in**
4 **Kenya**

5

6 [†]Donnie Mategula^{1,2,3}, [†]Judy Gichuki^{4,5}, [†]

7 [†]Shared first authors

8

9 **Institutional affiliations**

10 1. Malawi-Liverpool-Wellcome Programme, Blantyre, Malawi

11 2. Liverpool School of Tropical Medicine, Liverpool, UK

12 3. Kamuzu University of Health Sciences, Blantyre, Malawi

13 4. Strathmore University, Institute of Healthcare Management, Nairobi, Kenya

14 5. Nairobi City County Government, Health Services Department, Nairobi, Kenya

15

16

17 Corresponding Author: dategula@mlw.mw

18

19

20

21 **Abstract**

22 Malaria remains a public health concern. Monitoring the fine-scale heterogeneity of the malaria
23 burden enables more targeted control efforts. Although malaria indicator surveys (MIS) have been
24 crucial in evaluating the progress of malaria control interventions, they are only designed to provide
25 a cross-sectional national and regional malaria disease burden. Recent advances in geostatistical
26 methods allow us to interpolate national survey data to describe subnational disease burden that is
27 crucial in informing targeted control.

28 A binomial geostatistical model employing Markov chain Monte Carlo (MCMC) parameter estimation
29 methods is used to understand the spatial drivers of malaria risk in Kenya and to predict malaria risk
30 at a fine-scale resolution, including identifying hotspots.

31 A total of 11,549 children aged six months to 14 years from 207 clusters were sampled in this survey
32 and used in the present analysis. The national malaria prevalence based on the data was 8.4%, with
33 the highest in the lake endemic zone (18.1 %) and the lowest in the low-risk zone (<1 %). The
34 analysis shows that elevation, proportion of ITN distributed, rainfall, temperature and urbanization
35 covariates are all significant predictors of malaria transmission. The 5x5 Km resolution maps show
36 that malaria is heterogeneous in Kenya, with hotspot areas in the lake endemic area, the coastal
37 areas, and some parts of the shores of Lake Turkana and Kajiado.

38 The high-resolution malaria prevalence maps produced as part of the analysis have shown that
39 Kenya has additional malaria hotspots, especially in areas least expected. These findings call for a
40 rethinking of malaria burden classification in some regions for effective planning, implementation,
41 resource mobilization, monitoring, and evaluation of malaria interventions in the country.

42

43

44

45

46 **Background**

47 Malaria remains a public health concern and continues to be one of the most important tropical
48 diseases affecting human populations to date [1]. In 2020, an estimated 241 million malaria cases
49 occurred worldwide, 90% of which were in sub-Saharan Africa [2]. The disease is caused by protozoa
50 of the genus *Plasmodium* of which five known species, *Plasmodium falciparum*, *P. vivax*, *P. ovale*,
51 and *P. malariae*, more recently, *P. knowlesi*, are responsible for human infection [3,4]. The vector
52 responsible for human transmission is the female anopheles mosquito. In Sub-Saharan Africa,
53 malaria is one of the leading causes of morbidity and mortality, especially in children under five.
54 Other high-risk groups include pregnant women and immunologically naïve persons like travellers
55 coming from non-endemic places [5].

56 Malaria remains a significant public health problem in Kenya, accounting for an estimated 13% to
57 15% of outpatient cases, with nearly 70% of the population at risk for malaria [6]. Four out of the
58 five species of malaria parasite that cause human infection are present in Kenya, but the
59 *Plasmodium falciparum* parasite is the predominant cause of infection in the country. Over the past
60 decade, Kenya has substantially scaled up available malaria control tools, such as insecticide-treated
61 bed nets, indoor residual spraying and the use of artemisinin-based combination therapies [6].
62 Evidence of this massive scale-up of interventions is the observed decline in prevalence. Kenya has
63 experienced a decrease in the national prevalence of malaria among children ages six months to
64 14 years, from 13 per cent in 2010 to 8 per cent in 2015, and % in 2020 [6].

65 One of the key objectives of the Kenya Health Policy 2014–2030 is the elimination of communicable
66 diseases, including malaria. This is supported by the Kenya malaria strategy for 2019 to 2023, which
67 sets a vision of a malaria-free Kenya and targets to reduce malaria incidence and mortality by
68 seventy-five per cent by 2023, with 2016 as the baseline year[6].

69

70 Malaria transmission in Kenya varies geographically. This could be due to varied climatic conditions,
71 vector and parasite resistance, differences in intervention uptake across populations and other
72 unmeasured factors that are thought to be responsible for this increasing heterogeneity [6,7]. The
73 country is administratively divided into five malaria epidemiological zones based on risk profiles.
74 These zones include highland epidemic-prone areas, lake endemic areas, coast endemic, semi-arid
75 seasonal, and low-risk malaria areas. The endemic areas lie in the lake and coastal regions with
76 altitudes ranging from 0m to 1300 above sea level. These areas have perennial malaria transmission
77 due to rainfall, temperature, humidity and other critical factors that drive malaria transmission. The
78 semi-arid seasonal malaria transmission areas are in the country's northern, northeastern, and
79 southeastern parts. These areas experience short periods of intense malaria. The highland epidemic-
80 prone areas are located within the western highlands and have seasonal malaria transmission with
81 some yearly variation. The altitude in these zones is relatively higher than the other zones, lying
82 1500 meters above sea level. The malaria epidemics in the highland epidemic-prone zones are less
83 predictable. Lastly, the low-risk malaria areas cover Nairobi and the central highland. Temperatures
84 are usually too low to allow the completion of the sporogony cycle of the malaria parasite in the
85 vector in the low-risk zones[6] .

86 Malaria indicator surveys (MIS) measure progress on key malaria indicators in Kenya. The country
87 has conducted four MIS in 2007, 2010, 2015 and 2020. The MIS are nationally representative
88 household surveys that provide estimates of national and regional malaria indicators to assist
89 malaria control programs in tracking their progress and evaluating the impact of strategies and
90 interventions. The MIS follow a standard methodology recommended by the Roll Back Malaria
91 Monitoring and Evaluation working group guidelines [8]. Originally, MIS surveys were designed to
92 measure the blanket scale up of interventions like bed nets, using a classic two-stage sample design
93 and coverage indicators as the primary endpoints. Over time, as coverage increased, interest
94 expanded to the impact of parasite prevalence. While it is still a norm that the MIS traditionally
95 measures progress in these areas, the survey methodologies need to consider the underlying disease

96 heterogeneity [8]. Recent advances in statistical analyses, including geostatistics, have made it
97 possible to make fine-scale inferences of malaria transmission from survey data like the MIS, that is
98 not traditionally designed for such extrapolations. In this paper, we use the Kenya 2020 MIS data to
99 i) understand the relationship between malaria prevalence and several factors including
100 environmental factors and ii) understand the disease heterogeneity across the country's surface
101 including identification of hotspots.

102 **Methods**

103 **Country Profile**

104 Kenya is an East African country that covers an area of 582,550 km². It is bordered by Ethiopia to the
105 north, Tanzania to the south, Uganda to the west, South Sudan to the northwest, and Somalia to the
106 northeast. Approximately 80% of Kenya's land is arid and semi-arid, only 20% is arable, and only
107 1.9% of the total surface area is occupied by standing water. The great East African Rift Valley
108 extends from Lake Victoria to Lake Turkana and further southeast to the Indian Ocean [9]. The
109 country has a number of large rivers including the Tana, Galana, Turkwel and Nzoia [10]. Figure 1
110 below is a map of Kenya showing the five epidemiological zones as defined by the national malaria
111 program [6].

112 *Figure 1: Kenya epidemiological malaria zones*

113

114 **Data**

115 This secondary analysis used data from the Kenya MIS [6]. Access to the dataset was given to the
116 authors on Apr 28 2022. The datasets were de-identified. The IRB-approved procedures for DHS
117 public-use datasets do not in any way allow respondents, households, or sample communities to be
118 identified. Authors had no access to the names of individuals or household addresses in the data

119 files. Additionally, the geographic identifiers only go down to the regional level, which is hard to
120 identify individuals.

121

122 The 2020 MIS, the fourth conducted by the country, followed a similar design and set-up as the
123 former ones. It was conducted during the peak malaria season in November and December 2020. A
124 two-stage stratified sampling design was used, powered to give malaria parasite prevalence
125 estimates and other key malaria indicators at the national level (urban and rural areas) and for the
126 five epidemiological zones. The first stage sampling unit was a cluster developed from enumeration
127 areas (EAs). EAs are the smallest geographical areas created for purposes of census enumeration.
128 The EAs used were based on the 2019 Kenya population census. In the Kenya MIS, a cluster was
129 defined as either an EA or part of an EA. A total of 301 clusters (134 urban and 167 rural) were
130 sampled in this first stage using the probability proportional to size approach. The second stage
131 sampling unit was households. In each cluster, 30 households were selected from a line listing of the
132 sampled clusters using a systematic random sampling approach. A total of 7,952 households were
133 sampled. All women aged 15-49 in the selected households were eligible for individual interviews.
134 They were asked questions about preventing malaria during pregnancy and treating childhood
135 fevers. In addition, the survey included testing for anaemia and malaria among children aged six
136 months to 14 years using a finger- or heel-prick blood sample.

137 Permission to use the dataset was obtained from The Demographic and Health Surveys (DHS)
138 Program through the archiving office. The original study received ethical clearance from the
139 Kenyatta National Hospital/University of Nairobi Scientific and Ethics Review Committee. All
140 participants provided oral informed consent.

141

142 **Variables**

143 **Outcome variable**

144 In this analysis, the outcome variable was a binary outcome derived from the total number of
145 children tested and the total number testing positive. This was extrapolated to estimate the cluster-
146 level plasmodium falciparum malaria prevalence (*PfPR*).

147

148 **Explanatory variables**

149 The explanatory variables included cluster-level factors such as rainfall, temperature, elevation, and
150 urbanization and individual-level characteristics such as gender and age.

151 **Data cleaning**

152 The data cleaning and analysis were done in R[11]. Maps produced in the analysis were further
153 processed for better visualization in QGIS (Version 3.2). Relevant variables were extracted and
154 renamed to shorter names for ease of coding. Coordinates in the initial dataset were given the
155 longitude and latitude system, which were transformed into the universal coordinate system (UTM).
156 All distances were scaled to kilometres.

157 **Exploratory analysis**

158 The initial exploratory analysis was descriptive to understand the data and to explore the initial
159 relationships between the outcome variable of prevalence with the covariates in the data set.
160 Scatter plots with fitted linear regression lines were used for this step to observe the relationship
161 between prevalence and the explanatory variables. To further understand the variables, correlation
162 plots were used to understand the relationships between the variables to guide the decisions of
163 which covariates to include in the Model. The additional exploratory analysis involved plotting the
164 clusters on the Kenyan map's surface, showing the sampled cluster's distribution and the crude
165 malaria prevalence.

166 **Model fitting**

167 The first objective of the analysis was to understand the relationships between malaria prevalence
168 and several factors, including environmental factors. Several steps were followed:

- 169 1. Fitting a generalized linear model
- 170 2. Assessing evidence of residual correlation
- 171 3. Fitting a generalized linear mixed model
- 172 4. Reassessing evidence of residual correlations
- 173 5. Fitting a binominal geostatistical model and parameter estimation
- 174 6. Model validation

175 The model description for the generalized linear Model and the generalized linear mixed Model are
176 described in Supplementary file 1.

177

178 **Model description for the binominal geostatistical Model**

179 Let Y_i denote the number of individuals that test positive for plasmodium falciparum at survey
180 cluster location x_i

181 And that the survey team went to the sampled clusters given by x_i and sampled $m_i; i = 1 \dots n$
182 individuals at risk in the cluster and recoded the outcome of every person that tests positive and
183 negative for plasmodium falciparum malaria.

184 Then standard geostatistical Model assumes that:

$$185 \quad Y_i \sim \text{Binomial} (m_i, P(x_i))$$

186 Y_i is a Binomial distribution with m_i trials and probability of a positive test $P(x_i)$ specified in the
187 binomial geostatistical Model below:

188

$$189 \quad \log \left\{ \frac{P(x)}{1 - P(x)} \right\} = \alpha + d(x_i)^T \beta + S(x) + Z_i$$

190 Where α is the intercept parameter, $S(x)$ are the spatial random effects, representing the spatial
191 variation between the sampled clusters. Z_i are mutually independent zero-mean Gaussian random
192 variables with variance r and in this analysis represent the spatial variation within cluster variation,
193 measurement error or small-scale spatial variation.

194 $d(x_i)^T$ is a vector of observed spatially referenced explanatory variables associated with the
195 response Y_i , and β is a vector of spatial regression coefficients for the covariates.

196 The Matérn correlation function for the stationary Gaussian processes $S(x)$ used in this analysis, a
197 two-parameter family is given by:

198

$$199 \quad p(u, \varphi, k) = 2^{k-1} (u/\varphi)^k K_k + (u/\varphi)$$

200 Where:

- 201 • u denotes the distance between two locations x and x' ,
- 202 • $\varphi > 0$ is a scale parameter that determines the rate at which correlation decays to 0 as the
203 distance increases, and
- 204 • $k > 0$, is a smoothness parameter which determines the analytic smoothness of the underlying
205 process $S(x)$.

206 In the binomial geostatistical regression for this analysis, the Matérn shape parameter k was set to
207 0.5 variance parameters τ^2 to 0.

208 The covariates $d(x_i)^T$ used in the binomial geostatistical Model for prediction were obtained from
209 an exploratory analysis set to understand the relationship of the variables with the outcome variable
210 of malaria prevalence. This Model included the covariates: elevation, ITN usage, mean temperature,
211 rainfall, and cluster urbanization (urban vs rural). The Markov chain Monte Carlo (MCMC) methods
212 were used for parameter estimation in this Model. Confidence intervals of the estimates are
213 calculated on the log scale then transformed back to the non-log scale that is used to report the
214 results.

215 To test whether there was any evidence against spatial correlation in the data, empirical variogram
216 methods are used. A simulation of 1000 empirical variograms around the fitted Model is ran and
217 these are used to compute 95% confidence intervals at any given spatial distance of the variogram. A
218 conclusion is reached that that there is a spatial correlation in the data if the empirical variogram
219 obtained from the data falls outside the 95% tolerance bandwidth.

220 The second objective is to understand the disease heterogeneity across the surface of the country,
221 including identification of hotspots and the uncertainty attached to these hot spots. For this
222 purpose, a binomial geostastical model was used as described above but with covariates that were
223 available as raster. These included urbanization, temperature, and precipitation. The target for the
224 predictions was a prevalence of malaria over the 5 x 5 km regular grid surface covering the whole
225 surface of Kenya. A map of malaria prevalence was generated. Uncertainty of the prevalence was
226 addressed using *Exceedance Probabilities*, an approach that is more relevant to policy makers, than
227 the traditional approach of using confidence intervals. *Exceedance Probabilities (EP)* method sets
228 policy relevant thresholds. The *EP* can be formally expressed as:

$$229 \quad EP = \text{Probability} \{, P(x_i) > t \mid \text{data} \}$$

230 where t is the prevalence threshold, set to 10% in this analysis.

231 **Results**

232 A total of 11,549 children aged six months to 14 years were sampled. The analysis used 297 clusters.
233 The number of clusters per transmission zone is shown in Table 1 below. The lake endemic area had
234 the greatest number of clusters (98), while the coastal endemic area had the lowest number of
235 clusters (29). The national malaria prevalence based on the data was 8.4%, with the highest in the
236 lake endemic zone (18.1 %) and the lowest in the low-risk zone (<1 %).

237 *Table 1 Malaria Prevalence in Kenya across five epidemiological zones*

Epidemiological Zone	Total clusters	Number tested(N)	Number positive(n)	Prevalence (n/N)%
Coastal Endemic	29	1088	59	4.95
Highland Epidemic Prone	55	2122	33	1.56
Lake Endemic	97	4621	836	17.93
Low Risk	54	1307	1	0.08
Seasonal	56	2210	33	1.49

238

239 The map in Figure one below shows the sampled locations on the left and the cluster-level malaria
240 prevalence on the right. The lake endemic area has the highest number of clusters sampled and is
241 also the zone with the highest prevalence estimates at the cluster level.

242 *Figure 2 Map of Sampled locations(left) and Malaria Prevalence (Right)*

243 The weather pattern varies across the surface of Kenya. The maps in Figure 2 below show the
244 variation in temperature across space on the top, and the variation in annual precipitation for the
245 year 2020 on the bottom.

246 *Figure 3 Mean temperature (degrees Celsius) (top) and Annual precipitation (mm)in Kenya (2020)*
247 *(Bottom)*

248

249

250 **Binomial geostatistical model results**

251 The binomial geostatistical Model results indicate that the elevation, proportion of ITN distributed,
252 rainfall, temperature and urbanization covariates are all significant predictors of malaria
253 transmission (Table 2).

254 The odds of malaria transmission are less in the urban clusters compared to the rural ones. Urban
255 clusters have nearly 68% less malaria prevalence than rural ones (OR 0.32 CI: 0.26-0.39, P value
256 <0.0001). The higher the rainfall, the higher the risk of malaria transmission. Every mm increase in

257 the average rain increases malaria prevalence by 1.9 times (OR 1.91 CI 1.69- 2.15, P value <0.0001).
 258 Rise in mean temperature also increases the risk of malaria prevalence. Every degree increase in
 259 temperature increases the odds of malaria prevalence by 1.4 times (OR 1.37CI 1.28-1.47, P value
 260 <0.0001).

261 *Table 2: Binomial geostatistical Model*

262

	Estimate	Standard error	Lower Bound	Upper Bound	p-value
Intercept	0.00	3.54	0.00	0.00	<0.0001
elevation	1.86	1.05	1.70	2.02	<0.0001
INT coverage	3.15	1.07	2.75	3.61	<0.0001
Urban vs Rural	0.32	1.11	0.26	0.39	<0.0001
Rainfall(mm)	1.91	1.06	1.69	2.15	<0.0001
Temperature	1.37	1.04	1.28	1.47	<0.0001
Age in months (ref <12)					
12-23	1.93	1.54	0.83	4.48	<0.001
24-35	2.72	1.52	1.19	6.21	<0.01
36-47	3.56	1.51	1.59	7.96	<0.001
48-59	7.84	1.45	3.76	16.35	<0.001
Female vs male	1.20	1.08	1.03	1.40	<0.001
Sigma^2*	0.56	1.44	0.27	1.14	NA
Phi**	59.57	1.85	17.87	198.50	NA
Tau^2***	0.95	2.24	0.20	4.58	NA

263 **sigma2 is the variance of the Gaussian process, phi is the scale parameter of the spatial correlation**
 264 **and tau2 is the variance of the nugget effect**

265 **Model validation**

266 Using variogram-based techniques described above, the Model above was tested for evidence of
 267 spatial correlation. The results of this process are shown in figure 4 below. Since the empirical semi-
 268 variogram (solid line) falls within the 95% confidence interval (grey envelope), this shows that the
 269 Model is valid; the Model for malaria prevalence is compatible with the data.

270 *Figure 4 Model Validation*

271

272 **Prediction**

273 To understand the disease heterogeneity across the country's surface, including the identification of
274 hotspots, a 5 × 5 km resolution map for malaria prevalence in children six months to 14 years is
275 presented in Figure 5 below. Overall, malaria prevalence is low in most parts of the country.

276 Hotspots were notable in Western Kenya in the lake endemic areas around Lake Victoria, in the
277 endemic coastal regions along the Indian Ocean and three hotspot areas within the seasonal
278 epidemiological zone, one around the Lake Turkana region, one around the humid and sub-humid
279 belt in Meru County and the other in the semi-arid belt of Kajiado County.

280 *Figure 5 Malaria prevalence predictions among children six months- 14 years in Kenya*

281 Figure 6 below presents a map of malaria exceedance and probabilities, showing areas where $p(x) \geq$
282 10% with certainty on the colour gradient.

283 *Figure 6 Exceedance Probabilities.*

284

285 **Discussion**

286 Understanding the spatial distribution of malaria and the factors that drive its transmission are key
287 in malaria control. Given the heterogeneity of malaria transmission in Kenya, defining the malaria
288 burden at more localized locations is important to allow for targeted control activities. The national
289 malaria indicator surveys performed in the country are not designed to provide malaria prevalence
290 estimates at localized levels. This paper uses Model-Based geostatistical methods to understand
291 malaria transmission drivers in Kenya and map out malaria prevalence at a very high resolution (5 x 5
292 Km square grid).

293 In our analysis, we have found that several factors influence malaria transmission. These include
294 gender, age, temperature, rainfall, bed net coverage, elevation, and urbanization. This is consistent
295 with the well-known predictors of malaria transmission. Studies in the same area have previously
296 found a higher risk of malaria among males and an increasing risk with age compared to the first
297 year of life [11]. Both the natural environment and the artificial environment are known to affect
298 malaria transmission. Temperature, humidity, and rainfall all have interactions with mosquitos at
299 specific points in their life cycle. Temperature regulates the development of mosquitos at each
300 stage. The laying of eggs by mosquitos is reduced in temperature extremes, either too cold or too
301 hot [12]. Temperature also affects the mosquito stage transition, with the optimal temperature
302 being between 22 and 26 degrees Celsius[11].

303 Rainfall has been shown to be positively correlated with high malaria transmission. During the rainy
304 season, there is usually water logging in the ground, creating mosquito breeding grounds. This
305 analysis observed that prevalence nearly doubles for every mm increase in the annual rainfall. The
306 areas observed to have a higher prevalence of malaria in Kenya are known to have prolonged rainy
307 seasons[6].

308 The analysis also identified that malaria transmission is higher in rural areas compared to urban
309 areas. This finding is consistent with other studies in the same region. Urban areas may have better
310 housing and improved health services that are easier to access. These factors contribute to the lower
311 risk of malaria. Conversely, rural areas are primarily associated with favourable conditions for
312 malaria, including stagnant water, poor housing, inaccessible health services and agricultural
313 activities [13].

314 The finding of increasing malaria prevalence with higher bed net coverage can be explained through
315 reverse causality, which is often observed due to the higher distribution of bed nets in areas heavily
316 affected by malaria.

317 Malaria hotspot areas identified in the analysis include the entire lake and coastal regions classified
318 as malaria endemic [6]. This finding is in keeping with other previous analyses done for past time
319 points [18]. The climatic condition in these areas is known to support malaria transmission. We do
320 find additional hotspots, which highlights the strength in our analysis approach. Localized malaria
321 hotspots are identified in the county of Turkana. Though this area is classified as a seasonal malaria
322 transmission zone, a reactive case detection in the area conducted from 2018 to 2019 also detected
323 high malaria transmission with a prevalence as high as 33.6% [15]. Another study in a refugee camp
324 in the same region identified a malaria prevalence of 64.2% [15]. Evidence from a recent study
325 examining the contribution of *P. falciparum* parasite importation to local malaria transmission in
326 Central Turkana confirms that malaria in the area is rather endemic, with intense local transmission
327 as opposed to the importation of malaria [16]. Due to its malaria risk classification status, Turkana is
328 often left out of malaria control activities. This is an important finding where an area's transmission
329 is misclassified. The recent WHO malaria surveillance guide calls for countries to view malaria
330 transmission as a continuum in space and stratify the malaria burden for better targeting and
331 improved efficiency of malaria interventions[8]. As malaria transmission declines, it becomes
332 increasingly focal and prone to outbreaks. Understanding and predicting patterns of transmission
333 risk becomes an essential component of an effective elimination campaign, allowing limited
334 resources for control and elimination to be targeted cost-effectively. In this study, we also find
335 additional hotspots in the counties of Meru and Kajiado, areas with humid and arid weather
336 conditions, respectively. There is a need for more local surveillance in the area. These areas are also
337 characterized by low implementation of malaria control measures.

338

339 There are several strengths and limitations of the data used in the analysis. To the best of our
340 knowledge, this is the latest nationally representative data on malaria prevalence. With this, the
341 results of this study are generalizable to the entire population of Kenya. Use of the geostatistical
342 Model as opposed to the traditional non-spatial Model, is a key strength. It allows us to borrow

343 information from the sampled cluster to infer for the unsampled ones and at the same time, account
344 for predictors that influence malaria transmission. The major limitation in the analysis is the lack of
345 adequate environmental covariates to improve spatial predictions.

346 **Conclusion**

347 This analysis has shown that rainfall, urbanization, temperature, and bed net coverage are important
348 factors that affect malaria transmission. The high-resolution malaria prevalence maps produced as
349 part of the analysis are important in identifying hotspots which is an essential element in planning,
350 implementation, resource mobilization, monitoring, and evaluation of malaria interventions in the
351 country. We have also identified malaria hotspots in areas not traditionally classified as endemic,
352 highlighting the need to rethink the classification of malaria transmission epidemiology in Kenya.

353 **Acknowledgements**

354 Many thanks to the DHS programme team for allowing the authors to use the dataset and to the
355 participants that provided the data.

356 **References**

- 357 1. Murray CJL, Rosenfeld LC, Lim SS, Andrews KG, Foreman KJ, Haring D, et al. Global malaria
358 mortality between 1980 and 2010: a systematic analysis. *Lancet*. 2012 Feb 4;379(9814):413–
359 31.
- 360 2. World Health Organization. World malaria report 2021 [Internet]. 2021 [cited 2022 Jun 12].
361 Available from: [https://www.who.int/teams/global-malaria-programme/reports/world-](https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2021)
362 [malaria-report-2021](https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2021)
- 363 3. Millar SB, Cox-Singh J. Human infections with *Plasmodium knowlesi*—zoonotic malaria. *Clin*
364 *Microbiol Infect*. 2015 Jul 1;21(7):640–8.
- 365 4. White NJ. *Plasmodium knowlesi*: The Fifth Human Malaria Parasite. *Clin Infect Dis*. 2008 Jan
366 15;46(2):172–3.
- 367 5. Lalloo DG, Hill DR. Preventing malaria in travellers. *BMJ*. 2008 Jun;336(7657):1362–6.

- 368 6. Division of National Malaria Programme (DNMP) [Kenya], ICF. Kenya Malaria Indicator Survey
369 2020. 2021;158.
- 370 7. Mogeni P, Omedo I, Nyundo C, Kamau A, Noor A, Bejon P, et al. Effect of transmission
371 intensity on hotspots and micro-epidemiology of malaria in sub-Saharan Africa. *BMC Med*.
372 2017 Jun;15(1).
- 373 8. World Health Organization. Malaria surveillance, monitoring & evaluation: A reference
374 manual. Geneva; 2018.
- 375 9. Ingham K. Kenya | History, Map, Flag, Climate, Capital, & Facts | Britannica [Internet].
376 Britannica. 2019 [cited 2022 Jun 18]. Available from:
377 <https://www.britannica.com/place/Kenya>
- 378 10. KNBS. Population and housing census of Kenya, 2009 - Kenya Data Portal. 2010.
- 379 11. R Development Core Team. R: A Language and Environment for Statistical Computing.
380 Vienna, Austria; 2020.
- 381 12. Zhao X, Chen F, Feng Z, Li X, Zhou XH. Characterizing the effect of temperature fluctuation on
382 the incidence of malaria: An epidemiological study in south-west China using the varying
383 coefficient distributed lag non-linear model. *Malar J*. 2014 May 27;13(1):1–10.
- 384 13. Larson PS, Eisenberg JNS, Berrocal VJ, Mathanga DP, Wilson ML. An urban-to-rural continuum
385 of malaria risk: new analytic approaches characterize patterns in Malawi. *Malar J*. 2021
386 Dec;20(1):1–14.
- 387 14. Gopal S, Ma Y, Xin C, Pitts J, Were L. Characterizing the Spatial Determinants and Prevention
388 of Malaria in Kenya. *Int J Environ Res Public Heal* 2019, Vol 16, Page 5078. 2019
389 Dec;16(24):5078.
- 390 15. Meredith HR, Wesolowski A, Menya D, Esimit D, Lokoel G, Kipkoech J, et al. Epidemiology of
391 *Plasmodium falciparum* Infections in a Semi-Arid Rural African Setting: Evidence from
392 Reactive Case Detection in Northwestern Kenya. *Am J Trop Med Hyg*. 2021 Oct;105(4):1076–
393 84.
- 394 16. Markwalter CF, Menya D, Wesolowski A, Esimit D, Lokoel G, Kipkoech J, et al. *Plasmodium*
395 *falciparum* importation does not sustain malaria transmission in a semi-arid region of Kenya.
396 *PLOS Glob Public Heal*. 2022;2(8):e0000807.

397

398

399

400

401

402

403



Figure 1







Figure 2









Figure 4

Malaria Prevalence in Kenya




Figure 5




Figure 6