

1 **Uncertainty in serious illness:**
2 **A national interdisciplinary consensus exercise to identify clinical research priorities**
3

4
5 **Authors** Simon N Etkind^{1,2*}, Stephen Barclay¹, Anna Spathis^{1,2}, Sarah A Hopkins¹, Ben Bowers¹, Jonathan
6 Koffman³
7

8 **Affiliations**

- 9 1. Primary Care Unit, Department of Public Health and Primary Care, University of Cambridge
10 2. Cambridge University Hospitals NHS Foundation Trust
11 3. Hull York Medical School, University of Hull
12 * corresponding author
13
14

15 **Corresponding author contact details and address**

16 Simon Etkind

17 Sde23@medschl.cam.ac.uk

18 Primary Care Unit, Department of Public Health and Primary Care, East Forvie Building, Addenbrookes
19 Biomedical Campus, CB2 0SR
20
21

22 **Abstract:**

23 Background: Serious illness is characterised by uncertainty, particularly in older age groups. Uncertainty
24 may be experienced by patients, family carers, and health professionals about a broad variety of issues.
25 There are many evidence gaps regarding the experience and management of uncertainty.

26 Aim: We aimed to identify priority research areas concerning uncertainty in serious illness, to ensure that
27 future research better meets the needs of those affected by uncertainty and reduce research
28 inefficiencies.

29 Methods: Rapid prioritisation workshop comprising five focus groups to identify research areas, followed
30 by a ranking exercise to prioritise them. Participants were healthcare professionals caring for those with
31 serious illnesses including geriatrics, palliative care, intensive care; researchers; patient/carer
32 representatives, and policymakers. Descriptive analysis of ranking data and qualitative framework
33 analysis of focus group transcripts was undertaken.

34 Results: Thirty-four participants took part; 67% female, mean age 47 (range 33 – 67). The highest priority
35 was communication of uncertainty, ranked first by 15 participants (overall ranking score 1.59/3).
36 Subsequent priorities were: 2) How to cope with uncertainty; 3) healthcare professional
37 education/training; 4) Optimising clinical approaches to uncertainty; and 5) exploring in-depth
38 experiences of uncertainty. Research related to optimally managing uncertainty was given higher priority
39 than research focusing on experiences of uncertainty and its impact.

40 Conclusions: These co-produced, clinically-focused research priorities map out key evidence gaps
41 concerning uncertainty in serious illness. Managing uncertainty is the most pressing issue, and
42 researchers should prioritise how to optimally manage uncertainty in order to reduce distress, unlock
43 decision paralysis and improve illness and care experience.

44
45 **Key words**

46 Uncertainty; Communication; Serious illness; Palliative care; Qualitative research
47
48

49 **Key points**

- 50
- 51 • Uncertainty is ubiquitous and distressing in serious illness, and can paralyse decision making
 - 52 • In this consensus exercise, stakeholders identified research priorities for uncertainty in serious illness
 - 53 • Communication of uncertainty was the highest priority
 - 54 • Participants prioritised research concerning managing uncertainty above research to understand
55 experiences of uncertainty
- 56

57 **Background**

58 Uncertainty is ubiquitous in serious illnesses across all health settings, especially when living with long-
59 term conditions and frailty.[1-5] Encompassing “known unknowns”, uncertainty is characterised as an
60 inadequate understanding, a sense of incomplete, ambiguous or unreliable information, and conflicting
61 alternatives.[6, 7] It is inherently a complex concept and situations of uncertainty often result from
62 several inter-related factors.[8]

63

64 Irrespective of its origin, uncertainty matters because when suppressed and ignored, it can profoundly
65 negatively impact patients and their family.[9, 10] Older adults may be particularly affected as they
66 commonly experience complex and unpredictable illness, associated with irreducible uncertainties.
67 Uncertainty may precipitate extensive psychological and existential distress, potentially culminating in an
68 experience of ‘Total Uncertainty’ which may threaten an individual’s sense-of-self.[11]

69

70 If uncertainty is not addressed it may impact patient safety, adverse events, healthcare interactions and
71 relationships.[12, 13] One metric where this is recorded are complaints levelled at healthcare,
72 particularly in hospital settings.[14] Uncertainty can also limit patient participation in decision-making,
73 leading to ‘decision paralysis’,[15, 16] which may contribute to sub-optimal care and has repercussions
74 for the allocation of scarce health resources, including hospital admissions and longer inpatient stays.[17,
75 18]

76

77 It is not just patients who are affected by uncertainty. Despite uncertainty in medicine dating back to
78 Hippocrates, there exists a deeply rooted aversion to it in empirical medicine, where acknowledging
79 uncertainty can have connotations of failure.[18, 19] If poorly tolerated by health professionals,
80 uncertainty can adversely impact their confidence and competence, increasing the risk of moral injury,
81 burnout and depression.[20-23] This was particularly evident during the Covid-19 pandemic.[24, 25]

82

83 Uncertainty in illness is not necessarily distressing; it may be appraised negatively, neutrally, or in some
84 individuals, illnesses, and care circumstances, positively; for example holding onto prognostic uncertainty
85 can in some circumstances enable people to retain hope.[26-28] The distress caused by uncertainty, and
86 hence the block to decision making is therefore not inevitable; the negative impacts of uncertainty can
87 be at least partially ameliorated if it is addressed and communicated sensitively.[29-31]

88
89 There are innumerable possible situations of uncertainty, each of which may have its own optimal
90 approach. We still do not know how best to approach and address it in older patients living with long-
91 term or life-threatening illnesses in a way that best supports the individual and those involved in their
92 care;[32, 33] although previous work has explored how to support communication of uncertainty, [34]
93 uncertainty management,[12, 32] and shared decision making.[35] We do know that there can be no
94 one-size-fits-all approach to uncertainty, as it is experienced differently in different clinical contexts, by
95 different individuals.[6, 36] Uncertainties in some contexts may be more distressing than others and may
96 require different approaches.[37] Despite the importance of uncertainty for patients and clinicians alike
97 there has, to date, been no attempt to prioritise the most pressing areas to focus applied research on to
98 improve care.

99
100 We aimed to identify stakeholder priority research areas concerning uncertainty in serious illness, [38] to
101 enable future research to more effectively meet the needs of those affected by uncertainty and to
102 reduce research inefficiencies.[39, 40]

103 **Methods**

104 Design

105 National interdisciplinary one-day prioritisation workshop with a range of stakeholders: patient and carer
106 representatives, clinicians, researchers, and policymakers. Reporting according to REPRISE guidance (see
107 appendix).[41]

108

109 Patient and public involvement

110 A Patient and public involvement group supported development of the study aims and methods; 3 public
111 contributors participated in the workshop, contributing to focus groups and the ranking exercise, and
112 commented on the findings.

113

114 Participants and participant identification

115 Participants were clinicians, researchers, policymakers, people with lived experience of serious illness
116 and their informal carers. Researchers and policymakers were eligible if they were interested in the area.
117 Clinicians were from any profession or specialty with experience in providing care to people with serious
118 illnesses. The workshop invitation was disseminated widely through clinical and research networks and
119 social media, including: Applied Research Collaborative (ARC) East of England, UK uncertainty in serious
120 illness specialist interest group, the UK-wide Community Nursing Research Forum, regional and national
121 palliative care and gerontology contacts. Invitations were sent from December 2022, and registration
122 was open until 27th February 2023. The workshop was held on 28th February 2023. Workshop attendees
123 were informed of the research component in advance.

124

125 Workshop process

126 The workshop drew on existing approaches to prioritisation, incorporating idea generation,
127 consolidation, and ranking stages.[42] Written informed consent and self-reported demographic data

128 were collected from workshop participants at the start of the day, including age, gender, ethnicity,
129 participants' status as a researcher, clinician, policymaker, or patient representative, and details of their
130 field where relevant. To encourage debate, the workshop began with presentations on the "state of the
131 science" concerning uncertainty in serious illness. Models of uncertainty and evidence gaps from
132 relevant literature reviews were outlined.[11, 32, 34, 43, 44]

133 We then held focus groups with participants to explore their views on uncertainty and identify key areas
134 for future research. This approach mirrored the idea generation stage of the nominal group technique
135 and the first round of a Delphi process.[42, 45] The topic guide was developed by the research team and
136 was informed by literature review: it focused on experiences of uncertainty, views on desired outcomes
137 when addressing uncertainty, and ideas for research questions concerning uncertainty in serious illness.
138 Focus groups were led by clinicians and researchers with expertise in facilitation. Conversations were
139 audio recorded and a scribe within each group recorded the research questions identified by
140 participants.

141 Research areas from the focus groups were collated into a summary list during the day. In the final
142 session, this summary list was presented to participants who were invited to anonymously rank the top
143 three areas in order of priority using the online ranking tool "Slido" (© 2023 Cisco Systems, Inc.). Items
144 were presented in random order. See appendix for full workshop programme and topic guide.

145

146 Data analysis

147 *Analysis of focus group lists:* The lists generated by focus groups were reviewed by two researchers (SE &
148 JK) during the day, taking into account both researchers' existing knowledge of evidence gaps and
149 previously expressed areas for future research from literature reviews.(9, 28) The researchers combined
150 the lists produced by each focus group by removing duplicates and arranging similar questions under
151 "umbrella terms" to produce a single summary list of priority research areas for use in the subsequent
152 ranking exercise.

153

154 *Analysis of ranking exercise data:* Individual-level ranking scores were exported from Slido. We analysed
155 the scores descriptively and reported the number of participants ranking each item within their top
156 three. We calculated an average ranking score by allocating points to each item as follows: the item
157 ranked first received three points; the item ranked second received two points; the item ranked third
158 received one point; all other items received zero points. The average ranking score was calculated by
159 adding the points for each item and dividing by the number of participants. The maximum any item
160 could score was three if every participant ranked it as their top priority and the minimum was zero if no
161 participant ranked it in their top three.

162

163 *Analysis of focus group transcripts:* Following the workshop the focus group recordings were transcribed
164 verbatim, anonymised and analysed using a framework approach.[46] This stage aimed to identify
165 detailed research questions within the priority areas, as well as any additional areas discussed that were
166 relevant for future research. We used the research priority areas identified during the workshop as a
167 coding framework and one researcher (SE) coded text in the transcripts that described research
168 questions or participants' views about these areas. A second researcher (JK) independently reviewed one
169 focus group transcript. Coding was reviewed, and where there were differences, these issues were
170 reconsidered and debated by both researchers until consensus was achieved.[46] To avoid making
171 unwarranted claims about patterns and regularities in the data, we examined and coded unusual or non-
172 confirmatory views that did not fit easily into the original framework.[46] The framework was
173 condensed, summarised and discussed with the wider research team to refine it. Anonymised excerpts
174 from the transcripts are presented to illustrate themes and represent a range of views.

175

176 Ethical approval: This study received approval from the University of Cambridge Psychology Research
177 Ethics Committee [Reference:PRE.2022.125].

178 **Results**

179 Participant details

180 Thirty three participants took part in the focus groups, and 34 in the ranking survey, of whom 30
181 provided demographic information. The average age was 47 years (range 33 – 67), and 67% were female.
182 80% were of white ethnicity, 10% Asian, and 10% from mixed or multiple ethnic groups. 70% were
183 clinicians, 43% researchers, 10% patient or carer representatives, and 7% policymakers (participants
184 could state multiple roles). Of the clinician participants, twelve had a background in palliative care, three
185 geriatrics, two nursing and one each of intensive care, general practice, psychology, and physiotherapy.

186

187 Item generation and ranking

188 Five focus groups with six to seven participants in each were of 53 to 64 minutes' duration. The groups
189 generated 61 research questions, which we condensed to produce the 10 priority areas that were then
190 ranked by participants (Table 1). Communication of uncertainty was the highest-ranked item, scored first
191 by 15 participants. Participants ranked the next four priorities almost equally: coping with uncertainty;
192 training health professionals; optimising clinical approaches to uncertainty; understanding in-depth
193 experiences of uncertainty. The other areas received lower priority scores.

194

195 **Table 1. 10 priority areas for future uncertainty research in serious illness (n = 34):**

Overall Ranking	Priority	Number of participants ranking item in each top 3 positions (%)			Percentage of participants ranking items in top three	Ranking score*
		#1	#2	#3		
1	Communication of uncertainty	15 (44)	3 (9)	3 (9)	62	1.59
2	How to cope with uncertainty	6 (18)	5 (15)	3 (9)	41	0.91
3	Education/training of healthcare professionals	2 (6)	7 (21)	9 (26)	53	0.85
4	Optimising clinical approaches to uncertainty	6 (18)	2 (6)	4 (12)	35	0.76
5	Understanding patient/carer experiences of uncertainty in depth	2 (6)	7 (26)	5 (15)	41	0.74
6	Variation in experience/response to uncertainty between different individuals, groups, professions	1 (3)	4 (12)	1 (3)	18	0.35
7	Explore positive aspects of uncertainty	1 (3)	1 (3)	4 (12)	18	0.26
8	Impact of uncertainty on bereavement	1 (3)	2 (6)	1 (3)	12	0.24
9	Uncertainty in specific conditions/clinical situations	0 (0)	3 (9)	1 (3)	12	0.21
10	Factors associated with different uncertainty experiences	0 (0)	0 (0)	1 (3)	9	0.09

196 *Calculated as follows: item ranked first receives 3 points; item ranked 2nd receives 2 points; item ranked 3rd
 197 receives 1 point; all other items receive 0 points. The total points for each item are added and divided by the
 198 number of participants.

199

200 Detailed research priorities

201 We explored the 10 priority areas raised by workshop participants during qualitative analysis, and
 202 identified detailed sub-questions within each area (Table 2).

203

204 The highest priority research areas were related to managing uncertainty, by communication, supporting
205 individual coping mechanisms, or providing training. Participants particularly focused on how and when
206 to have conversations about uncertainty, and how to individualise communication to promote
207 psychological wellbeing:

208 *“Then there’s a really important research need around understanding more about how we*
209 *communicate uncertainty and the consequences of how we communicate uncertainty and how*
210 *we can talk about uncertainty in a way which is more patient-centred and supportive and*
211 *considers sort of psychological wellbeing.” Social scientist, focus group 3.*

212 Training of health professionals at varying stages was a recognised priority. This included training
213 professionals on toleration of their uncertainty as well as how to address the uncertainty experienced by
214 others. One participant highlighted that inadequate training of doctors to manage uncertainty has been
215 a longstanding issue and queried the best timing of such training:

216 *“We’ve been inadequate at teaching them [medical students] how to actually manage this*
217 *complexity for a long time, although you could argue that it’s quite difficult to teach until they*
218 *really get into the nitty-gritty of practising.” Geriatrician, focus group 1*

219 Participants recognised it is unlikely a single intervention can address all the nuanced multilevel aspects
220 of uncertainty, but nevertheless, some felt interventions could play a role in addressing uncertainty, as
221 long as they were situated in a broader societal context:

222 *“There are interventions that are being developed out there but there are no interventions that*
223 *deal with all the different types and layers of uncertainty and they can’t by nature. And I think*
224 *there is that kind of relational uncertainty, the kind of organisational uncertainties, there are*
225 *uncertainties on macro, meso, micro levels and you can deal with one component, but you can’t*
226 *deal with all the different components, and it’s how those interventions work within the broader*
227 *societal context of uncertainty.” Occupational therapist, focus group 4*

228 Older frail patients were seen as a priority for future research into how uncertainty can be managed.
229 Participants noted a lack of knowledge about approaching uncertainty in the context of frailty:

230 *“For old and frail elderly there’s no support you know, for that anxiety, fear, so it’s how do we*
231 *look at that, you know, and how do we maybe look at how we can support, supporting that*
232 *frailty and older.” Professor of end-of-life care, focus group 5*

233

234 Participants identified research questions concerning *experiences* of uncertainty, though these were
235 generally given lower priority than questions relating to *managing* uncertainty. Participants identified
236 illness contexts where uncertainty experiences could be explored further, including critical illness and
237 frailty. They also acknowledged it is important to explore how the experience of uncertainty varies
238 depending on perspective (patient, carer or health professional), or individual characteristics:

239 *“We’d want to look at different groups such as sort of learning disability, neurodiversity, sort of*
240 *hard-to-reach areas whether deprivation, LGBTQ+, sort of that kind of differences you might get.”*
241 *Palliative care consultant, focus group 1.*

242 Participants identified potential positive impacts of uncertainty and its utility in certain situations. They
243 suggested it was important to understand why some health professionals thrive when required to
244 manage uncertainty, whereas others exhibit a lower tolerance. Participants noted the relationship
245 between uncertainty and hope and reasoned that positive aspects of uncertainty should be explored
246 further:

247 *“I wonder....whether there’s something about learning to cope with uncertainty or to tolerate*
248 *uncertainty and whether people, patients and families can see that as a positive as well.”*
249 *Bereavement practitioner, focus group 4*

250

251 Additional areas for future research

252 Some questions identified from focus group transcripts were not identified as research priority areas by
253 participants. Additional areas included consideration of the legal and regulatory implications of
254 uncertainty and its management, the link between uncertainty and patient safety, how to prepare the
255 public for serious illness uncertainty, and the resource impacts of different levels of tolerance to
256 uncertainty:

257 *“Have we studied the use of resources around uncertainty? Because I’m sure loads of tests are*
258 *done completely unnecessarily because people just want to be sure.” Palliative care consultant,*
259 *group 1*

260 **Table 2. Detailed research priorities from framework analysis***

Theme	Priority area	Detail of priority area
<p style="text-align: center;">Uncertainty managemen t</p>	<p>Communication of uncertainty (research on how best to communicate about uncertainty)</p>	<ul style="list-style-type: none"> • Understand more about the detailed processes for uncertainty communication: who should be involved, optimum setting & timing, what phrases to use; how much uncertainty to share; how are such conversations received by patients • How to communicate uncertainty openly whilst maintaining a trusting relationship? • Interprofessional communication • How to individualise communication of uncertainty to different settings, situations, individual characteristics? • How does the communication of uncertainty affect decision-making and other outcomes?
	<p>How to cope with uncertainty (how can individuals best be supported to cope with their uncertainties)</p>	<ul style="list-style-type: none"> • How do health professionals cope when they feel uncertain and how do they achieve resilience/tolerance to it? • How can patients and carers be supported to cope with their uncertainty and develop resilience? • Who copes well and why? • How to cope with the long-term impact of decisions made under conditions of uncertainty? • What is the role of hope in coping with uncertainty?
	<p>Education/training of healthcare professionals (how can we train health professionals to approach uncertainty)</p>	<ul style="list-style-type: none"> • How to teach uncertainty management to medical students, what would be the goals of such training? • How to train HCPs at different levels to recognise/tolerate/hold their own uncertainty? • How to transfer expertise from areas where uncertainty is well managed? • Development of training interventions including psychological training
	<p>Optimising clinical approaches to uncertainty (how can the uncertainty of others best be managed and addressed, what are optimal approaches)</p>	<ul style="list-style-type: none"> • How to individualise the management of uncertainty depending on patient experiences, response to uncertainty, and how to identify how much uncertainty is tolerable to an individual? • Who should 'hold' uncertainty and how to find a balance in terms of information sharing and decision-making • How to maintain trust and manage expectations? • How to gauge the readiness of people to discuss uncertainty and time conversations? • How can technology be used to distil information and reduce/address uncertainty relating to complexity/information overload? • What is the role of uncertainty management interventions and models of care to address uncertainty, and how effective are these? • What are the key outcomes we should be aiming for when seeking to approach and manage uncertainty? • What are the barriers to addressing uncertainty?

		<ul style="list-style-type: none"> • How to manage uncertainty as an MDT and maintain continuity of approach
Uncertainty experiences	<p>Understanding patient/carer experiences of uncertainty in depth</p> <p>(what are patient and carer experiences of uncertainty)</p>	<ul style="list-style-type: none"> • How does the experience of uncertainty affect decision-making? • What is the lived experience of uncertainty amongst patients and families (what is helpful when faced with uncertainty and what is harmful? How do the multiple layers of uncertainty interact within an individual's experience (mapping uncertainty)? Are the experiences/impacts of different types of uncertainty different? How do past life or healthcare experiences affect current uncertainty experiences.)? • How is uncertainty transmissible i.e. how can HCP uncertainty be picked up by patients and vice versa and what is the effect of this?
	<p>Variation in experience/ response to uncertainty</p> <p>(how does the experience of uncertainty vary between different groups)</p>	<ul style="list-style-type: none"> • How does experience and response to uncertainty vary between different medical specialties and health professional groups? • How does experience and response to uncertainty change over time as the illness progresses. • How do individual characteristics e.g. culture, LGBTQ status, neurodiversity, age, affect experience and response to uncertainty? • How does career stage and knowledge/experience in a clinical role affect health professionals experience? • What are the long-term effects of uncertainty experiences in ITU on ITU survivors?
	<p>Explore positive aspects of uncertainty</p> <p>(what aspects of uncertainty are positive and how can these be promoted)</p>	<ul style="list-style-type: none"> • What is the relationship between sharing uncertainty and maintaining hope? • Explore positive utility of uncertainty e.g. as a way to promote a quest for knowledge or changing ways of thinking • Explore why some people thrive with uncertainty
	<p>Impact of uncertainty on bereavement</p> <p>(how does uncertainty in serious illness impact on bereavement experiences)</p>	<ul style="list-style-type: none"> • How does uncertainty in serious illness and at the end of life affect experiences and outcomes of bereavement?
	<p>Uncertainty in specific conditions/clinical situations</p> <p>(the uncertainties that are experienced in a particular clinical situation such as in ITU)</p>	<ul style="list-style-type: none"> • How do different clinical situations change how uncertainty is experienced and responded to? • Are there situations where the expression of uncertainty is not appropriate? • How to make good decisions in the context of uncertainty in the intensive care unit? • How is uncertainty experienced in children with complex neuro-disability? • How to approach uncertainty in adults with neurodegenerative disease and uncertain illness trajectory? • How to approach uncertainty in frailty with uncertain prognosis?

		<ul style="list-style-type: none"> • How to communicate uncertainty at transitions of care
	<p>Factors associated with different uncertainty experiences</p> <p>(what factors are associated with different experiences of or responses to uncertainty.)</p>	<ul style="list-style-type: none"> • What system factors underlie the challenges of uncertainty? • What factors are associated with uncertainty experiences and tolerance e.g. age, life experience, culture? • How did the COVID-19 pandemic affect experiences and response to uncertainty? • How does trust in clinicians affect uncertainty experiences?

261

*Priority areas are listed in the order they were ranked

262 **Discussion**

263 Summary of findings

264 This study co-produced clinically-focused research priorities to address known evidence gaps
265 concerning uncertainty in serious illness. Optimising communication of uncertainty was the top
266 research priority. Research into managing uncertainty was considered higher priority than research
267 investigating experiences of uncertainty.

268

269 Discussion of main findings

270 Communicating uncertainty was the top priority for participants, reflecting key evidence gaps and
271 recommendations in this field.[32, 47] In their narrative review of uncertainty communication,
272 Simpkin et al identified a number of evidence gaps, including identifying individuals' communication
273 preferences and tailoring communication to those preferences.[34] The question of how to maintain
274 hope whilst communicating uncertainty was noted as a priority; this has been explored in cancer
275 care,[48] but remains a key question in other serious illnesses. Additionally, participants raised
276 several sub-questions in terms of *how* to discuss uncertainty, reflecting the need for
277 implementation-focused communication research.

278 After communication, participants prioritised other aspects of managing uncertainty: identifying
279 how individuals can be supported to cope with their own uncertainty; investigating how we can
280 optimise clinical approaches to uncertainty; understanding more about how to equip health
281 professionals to deal with uncertainty through training. Though management of uncertainty has
282 been recognised as a core component of medical training for decades, curricula still make limited
283 reference to uncertainty, and filling this gap should be a priority.(2) Whilst we have an improving
284 understanding of how physicians manage uncertainty,[31] the evidence base for other professional
285 groups is very limited, yet nurses and allied healthcare professionals often lead the clinical care and
286 support of older people and their families. To date, uncertainty management and communication
287 interventions have had variable impact in serious illness,[32] and often prove challenging to

288 evaluate.[33, 49] There is scope for further intervention development,[50] especially work that
289 focuses on evaluation of uncertainty-focused interventions.

290 Whilst many of the questions identified by participants related to investigating experiences of
291 uncertainty, these were usually considered lower priority, perhaps because much is already known
292 about uncertainty experiences.[7, 51] For example, conceptual taxonomies of uncertainty are well
293 developed,(6) and there have been evidence syntheses of experience in some specific areas e.g.
294 multimorbidity.[11] However, there are still evidence gaps concerning how uncertainty affects
295 individuals in other clinical contexts, in particular older patients living with frailty. Given the rapidly
296 increasing complexity of the healthcare system and unpredictability of the frailty trajectory, this is an
297 area that warrants urgent exploration.[34] Understanding more about the impacts of uncertainty on
298 experiences of illness, care, and bereavement would enable us to develop interventions focused on
299 the real-world problems uncertainty can cause.

300

301 Strengths and limitations

302 The rapid prioritisation approach we used enabled a diverse group of interested individuals to
303 generate and rank research priorities in a single day. Those ranking the priorities had spent the
304 entire day considering uncertainty in serious illness and were well placed to express considered
305 views when ranking the list presented to them. By identifying evidence gaps before the workshop
306 and communicating these to participants during initial presentations, we were able to focus on areas
307 where more research is needed and incorporate the key stages of a traditional prioritisation
308 exercise. By additionally incorporating formal qualitative analysis we increased rigour and developed
309 a robust priority list. This approach was a more feasible and pragmatic alternative to lengthier
310 methodologies such as the Delphi process[52, 53]. However, the rapid nature of the prioritisation
311 process meant there was limited time to condense the findings of the focus group discussions, which
312 risked a loss of accuracy and ranking was limited to broad research areas. We ameliorated this by
313 subsequent analysis of focus group transcripts, which enabled us to identify detailed research

314 questions identified by participants, including areas that participants mentioned even when they
315 weren't identified as priorities at the time.

316 We incorporated patient and carer experiences, but most participants were healthcare professionals
317 or researchers, thus these groups were not equally represented in the ranking process. Whilst not
318 specifically excluded, social care professionals did not attend this workshop, which limits the findings
319 to healthcare. We recruited a UK-wide sample, but this was not an international study, and future
320 work should explore if these findings hold internationally, though literature from 17 countries
321 reported consistent findings on a similar topic.[11] The anonymous nature of the ranking means we
322 could not adjust for the background of participants when analysing ranking data. The largest group
323 of clinical participants were from palliative care backgrounds which may have shaped their views;
324 however, a broad range of health professionals and researchers were represented, and the degree
325 of agreement, particularly regarding the top priority of communication suggests the findings
326 represent true consensus.

327

328

329 **Conclusion**

330 Through a rapid prioritisation workshop, we have identified 10 ranked priority areas for clinically
331 focused research on uncertainty in serious illness. There was consensus that further research into
332 managing uncertainty, particularly communication, was of higher priority than research to
333 investigate experiences of uncertainty. Future targeted research could result in interventions to
334 reduce the distress associated with uncertainty, unlock decision paralysis and improve illness and
335 care experience.

336

337 **Acknowledgements**

338 Thank you to those who facilitated focus groups: Stephan Barclay, Jonathan Koffman, Anna Spathis,
339 Ben Bowers, Sarah Hopkins, Debbie Critoph, Markus Schichtel, Ikumi Okamoto

340 We would also like to thank our patient and public involvement representatives: Roberta Lovick,
341 Sarah Dixon, Rashmi Kumar

342 Many thanks to the workshop organiser Angela Harper & the professional services team at the
343 Primary Care Unit, University of Cambridge for ensuring the smooth running of this study

344 Thank you to Zoe Fritz for your advice during drafting of the manuscript

345

346 **Funding:**

347 This study and SB are supported by the National Institute for Health and Care Research (NIHR)
348 Applied Research Collaboration East of England (NIHR ARC EoE) at Cambridgeshire and Peterborough
349 NHS Foundation Trust. BB is supported by the Wellcome Trust [225577/Z/22/Z]. SAH is jointly
350 funded by The Dunhill Medical Trust and British Geriatrics Society [Grant ref. JBGS20\5].

351 The views expressed are those of the author(s) and not necessarily those of the NIHR or the
352 Department of Health and Social Care.'

353

354 **Contributions:**

355 Study design: SE, SB, AS, JK

356 Securing funding: SE, SB

357 Data collection: SE, SB, AS, SAH, BB, JK

358 Analysis: SE JK

359 Paper drafting: SE, SB, AS, SAH, BB, JK

360 Approval of final version: SE, SB, AS, SAH, BB, JK

361

362 **Conflicts of Interest:**

363 The authors declare that they have no conflicts of interest.

364

365

366 References

- 367 1. Fox RC. The evolution of medical uncertainty. *Milbank Memorial Fund Quarterly, Health and*
368 *Society*. 1980;58(1):1-49.
- 369 2. Etkind SN, Koffman J. Approaches to managing uncertainty in people with life-limiting
370 conditions: role of communication and palliative care. *Postgraduate Medical Journal*. 2016.
- 371 3. Etkind SN, Bristowe K, Bailey K, Selman LE, Murtagh FEM. How does uncertainty shape
372 patient experience in advanced illness? A secondary analysis of qualitative data. *Palliative Medicine*.
373 2016;31(2):171-80.
- 374 4. Cox CL, Miller BM, Kuhn I, Fritz Z. Diagnostic uncertainty in primary care: what is known
375 about its communication, and what are the associated ethical issues? *Family Practice*.
376 2021;38(5):654-68. doi: 10.1093/fampra/cmab023.
- 377 5. Higginson IJ, Rumble C, Shipman C, Koffman J, Sleeman KE, Morgan M, et al. The value of
378 uncertainty in critical illness? An ethnographic study of patterns and conflicts in care and decision-
379 making trajectories. *BMC Anesthesiol*. 2016;16:11. Epub 2016/02/11. doi: 10.1186/s12871-016-
380 0177-2. PubMed PMID: 26860461; PubMed Central PMCID: PMC4746769.
- 381 6. Han PK, Klein WM, Arora NK. Varieties of Uncertainty in Health Care A Conceptual
382 Taxonomy. *Medical Decision Making*. 2011;31(6):828-38.
- 383 7. Etkind SN. Uncertainty in multimorbidity: a shared experience we should recognise,
384 acknowledge and communicate. *British journal of community nursing*. 2022;27(11):540-4. doi:
385 10.12968/bjcn.2022.27.11.540.
- 386 8. Lipshitz R, Strauss O. Coping with uncertainty: A naturalistic decision-making analysis. *Organ*
387 *Behav Hum Decis Process* 1997;69:149–63.
- 388 9. Andersen HE, Hoeck B, Nielsen DS, Ryg J, Delmar C. A phenomenological-hermeneutic study
389 exploring caring responsibility for a chronically ill, older parent with frailty. *Nursing Open*.
390 2020;7(4):951-60. PubMed PMID: 32114136678.
- 391 10. Nanton V, Munday D, Dale J, Mason B, Kendall M, Murray S. The threatened self:
392 Considerations of time, place, and uncertainty in advanced illness. *British Journal of Health*
393 *Psychology*. 2015:n/a-n/a. doi: 10.1111/bjhp.12172.
- 394 11. Etkind SN, Li J, Louca J, Hopkins SA, Kuhn I, Spathis A, et al. Total Uncertainty: A systematic
395 review and thematic synthesis of experiences of uncertainty in older people with advanced
396 multimorbidity, their informal carers, and health professionals. *Age and Ageing* 2022;51(8).
- 397 12. Carey I, Shouls S, Bristowe K, Morris M, Briant L, Robinson C, et al. Improving care for
398 patients whose recovery is uncertain. The AMBER care bundle: design and implementation. *BMJ*
399 *supportive & palliative care*. 2015;5(4):405-11.
- 400 13. Yardley I, Yardley S, Williams H, Carson-Stevens A, Donaldson LJ. Patient safety in palliative
401 care: a mixed-methods study of reports to a national database of serious incidents. *Palliative*
402 *medicine*. 2018;32(8):1353-62.
- 403 14. Ombudsman PaHS. Dying without dignity Investigations by the Parliamentary and Health
404 Service Ombudsman into complaints about end of life care. London Parliamentary and Health
405 Service Ombudsman, 2015.
- 406 15. Sellars M, Chung O, Nolte L, Tong A, Pond D, Fetherstonhaugh D, et al. Perspectives of
407 people with dementia and carers on advance care planning and end-of-life care: A systematic review
408 and thematic synthesis of qualitative studies. *Palliative medicine*. 2019;33(3):274-90.
- 409 16. Oksavik; JD, Solbjør; M, Kirchhoff; R, Sogstad MKR. Games with uncertainty: the participation
410 of older patients with multimorbidity in care planning meetings – a qualitative study. *BMC Geriatrics*.
411 2021;21:242.
- 412 17. Bristowe K, Carey I, Hopper A, Shouls S, Prentice W, Caulkin R, et al. Patient and carer
413 experiences of clinical uncertainty and deterioration, in the face of limited reversibility: A
414 comparative observational study of the AMBER care bundle. *Palliat Med*. 2015;29(9):797-807. Epub
415 2015/04/02. doi: 10.1177/0269216315578990. PubMed PMID: 25829443; PubMed Central PMCID:
416 PMC4572938.

- 417 18. Simpkin AL, Schwartzstein RM. Tolerating Uncertainty - The Next Medical Revolution? The
418 New England journal of medicine. 2016;375(18):1713-5. Epub 2016/11/03. doi:
419 10.1056/NEJMp1606402. PubMed PMID: 27806221.
- 420 19. Gheihman G, Johnson M, Simpkin AL. Twelve tips for thriving in the face of clinical
421 uncertainty. Medical Teacher. 2020;42(5):493-9. doi: 10.1080/0142159X.2019.1579308.
- 422 20. Čartolovni A, Stolt M, Scott PA, Suhonen R. Moral injury in healthcare professionals: A
423 scoping review and discussion. Nursing ethics. 2021;28(5):590-602.
- 424 21. Di Trani M, Mariani R, Ferri R, De Berardinis D, Frigo MG. From resilience to burnout in
425 healthcare workers during the COVID-19 emergency: the role of the ability to tolerate uncertainty.
426 Frontiers in Psychology. 2021;12:646435.
- 427 22. van Iersel MB, Brantjes E, de Visser M, Looman N, Bazelmans E, van Asselt D. Tolerance of
428 clinical uncertainty by geriatric residents: a qualitative study. European Geriatric Medicine.
429 2019;10(3):517-22.
- 430 23. Begin AS, Hidrue M, Lehrhoff S, Del Carmen MG, Armstrong K, Wasfy JH. Factors Associated
431 with Physician Tolerance of Uncertainty: an Observational Study. Journal of general internal
432 medicine. 2022;37(6):1415-21. Epub 2021/04/28. doi: 10.1007/s11606-021-06776-8. PubMed PMID:
433 33904030; PubMed Central PMCID: PMCPCMC8074695.
- 434 24. Koffman J, Etkind SN, Gross J, Selman L. Uncertainty and Covid-19: How are we to respond?
435 Journal of the Royal Society of Medicine. 2020;1–6 doi: DOI: 10.1177/0141076820930665.
- 436 25. Davey Smith G, Blastland M, Munafò M. Covid-19's known unknowns. BMJ (Clinical research
437 ed). 2020;371:m3979. doi: 10.1136/bmj.m3979.
- 438 26. Mishel MH. Reconceptualization of the Uncertainty in Illness Theory. Image: the Journal of
439 Nursing Scholarship. 1990;22(4):256-62. doi: 10.1111/j.1547-5069.1990.tb00225.x.
- 440 27. Gough N, Ross JR, Riley J, Judson I, Koffman J. 'When something is this rare ... how do you
441 know bad really is bad...?'-views on prognostic discussions from patients with advanced soft tissue
442 sarcoma. BMJ Support Palliat Care. 2019;9(1):100-7. Epub 2015/12/31. doi: 10.1136/bmjspcare-
443 2015-000898. PubMed PMID: 26715566.
- 444 28. Brashers DE. Communication and Uncertainty Management. Journal of Communication.
445 2001;51(3):477-97. doi: 10.1111/j.1460-2466.2001.tb02892.x.
- 446 29. Krawczyk M, Gallagher R. Communicating prognostic uncertainty in potential end-of-life
447 contexts: experiences of family members. BMC palliative care. 2016;15(1):59. doi: 10.1186/s12904-
448 016-0133-4.
- 449 30. Meranius MS, Engstrom G. Experience of self-management of medications among older
450 people with multimorbidity. Journal of Clinical Nursing. 2015;24(19):2757-64. PubMed PMID:
451 rayyan-114139203.
- 452 31. Han PK, Strout TD, Gutheil C, Germann C, King B, Ofstad E, et al. How physicians manage
453 medical uncertainty: a qualitative study and conceptual taxonomy. Medical decision making.
454 2021;41(3):275-91.
- 455 32. Ellis-Smith C, Dawkins M, Gao W, Higginson IJ, Evans CJ. Managing clinical uncertainty in
456 older people towards the end of life: a systematic review of person-centred tools. BMC palliative
457 care. 2021;20(1):1-41.
- 458 33. Koffman J, Yorganci E, Yi D, Gao W, Murtagh F, Pickles A, et al. Managing uncertain recovery
459 for patients nearing the end of life in hospital: a mixed-methods feasibility cluster randomised
460 controlled trial of the AMBER care bundle. Trials. 2019;20(1):1-18.
- 461 34. Simpkin AL, Armstrong KA. Communicating Uncertainty: a Narrative Review and Framework
462 for Future Research. Journal of general internal medicine. 2019;34(11):2586-91. doi:
463 10.1007/s11606-019-04860-8.
- 464 35. Politi MC, Clark MA, Ombao H, Dizon D, Elwyn G. Communicating uncertainty can lead to less
465 decision satisfaction: a necessary cost of involving patients in shared decision making? Health
466 Expectations. 2011;14(1):84-91. doi: 10.1111/j.1369-7625.2010.00626.x.
- 467 36. Hill DL, Walter JK, Szymczak JE, DiDomenico C, Parikh S, Feudtner C. Seven Types of
468 Uncertainty When Clinicians Care for Pediatric Patients With Advanced Cancer. J Pain Symptom
469 Manage. 2020;59(1):86-94. Epub 2019/08/20. doi: 10.1016/j.jpainsymman.2019.08.010. PubMed
470 PMID: 31425822; PubMed Central PMCID: PMCPCMC6942218.

- 471 37. Ghosh AK, Joshi S. Tools to manage medical uncertainty. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*. 2020;14(5):1529-33. doi: <https://doi.org/10.1016/j.dsx.2020.07.055>.
- 472
- 473 38. Waggoner J, Carline JD, Durning SJ. Is There a Consensus on Consensus Methodology? Descriptions and Recommendations for Future Consensus Research. *Academic Medicine*. 2016;91(5).
- 474
- 475
- 476 39. Sibbald SL, Singer PA, Upshur R, Martin DK. Priority setting: what constitutes success? A conceptual framework for successful priority setting. *BMC Health Services Research*. 2009;9(1):43. doi: [10.1186/1472-6963-9-43](https://doi.org/10.1186/1472-6963-9-43).
- 477
- 478
- 479 40. El-Harakeh A, Lotfi T, Ahmad A, Morsi RZ, Fadlallah R, Bou-Karroum L, et al. The implementation of prioritization exercises in the development and update of health practice guidelines: A scoping review. *PLoS ONE*. 2020;15(3):e0229249. doi: [10.1371/journal.pone.0229249](https://doi.org/10.1371/journal.pone.0229249).
- 480
- 481
- 482 41. Tong A, Synnot A, Crowe S, Hill S, Matus A, Scholes-Robertson N, et al. Reporting guideline for priority setting of health research (REPRISE). *BMC Medical Research Methodology*. 2019;19(1):243. doi: [10.1186/s12874-019-0889-3](https://doi.org/10.1186/s12874-019-0889-3).
- 483
- 484
- 485 42. James Lind Alliance Guidebook 2021 [cited 2023 25th April]. Available from: <https://www.jla.nihr.ac.uk/jla-guidebook/>.
- 486
- 487 43. Lund S, Richardson A, May C. Barriers to Advance Care Planning at the End of Life: An Explanatory Systematic Review of Implementation Studies. *PLoS ONE*. 2015;10(2):e0116629. doi: [10.1371/journal.pone.0116629](https://doi.org/10.1371/journal.pone.0116629).
- 488
- 489
- 490 44. Tulskey JA, Beach MC, Butow PN, Hickman SE, Mack JW, Morrison RS, et al. A Research Agenda for Communication Between Health Care Professionals and Patients Living With Serious Illness. *JAMA Internal Medicine*. 2017;177(9):1361-6. doi: [10.1001/jamainternmed.2017.2005](https://doi.org/10.1001/jamainternmed.2017.2005).
- 491
- 492
- 493 45. Harvey N, Holmes CA. Nominal group technique: An effective method for obtaining group consensus. *International Journal of Nursing Practice*. 2012;18(2):188-94. doi: <https://doi.org/10.1111/j.1440-172X.2012.02017.x>.
- 494
- 495
- 496 46. Spencer L, Ritchie J, Lewis J. Quality in qualitative evaluation: a framework for assessing research evidence. London: 2003 2003. Report No.: Occasional Series No 2.
- 497
- 498 47. Berger Z. Navigating the unknown: shared decision-making in the face of uncertainty. *Journal of general internal medicine*. 2015;30:675-8.
- 499
- 500 48. van Vliet LM. Balancing explicit with general information and realism with hope. Communication at the transition to palliative breast cancer care: Netherlands Institute for Health Services Research; 2013.
- 501
- 502
- 503 49. Cox C, Fritz Z. What is in the toolkit (and what are the tools)? How to approach the study of doctor–patient communication. *Postgraduate Medical Journal*. 2022.
- 504
- 505 50. Tunnard I, Yi D, Ellis-Smith C, Dawkins M, Higginson IJ, Evans CJ. Preferences and priorities to manage clinical uncertainty for older people with frailty and multimorbidity: a discrete choice experiment and stakeholder consultations. *BMC Geriatrics*. 2021;21(1). doi: [10.1186/s12877-021-02480-8](https://doi.org/10.1186/s12877-021-02480-8).
- 506
- 507
- 508
- 509 51. Connolly T, Coats H, DeSanto K, Jones J. The experience of uncertainty for patients, families and healthcare providers in post-stroke palliative and end-of-life care: a qualitative meta-synthesis. *Age and Ageing*. 2020. doi: [10.1093/ageing/afaa229](https://doi.org/10.1093/ageing/afaa229).
- 510
- 511 52. Jünger S, Payne SA, Brine J, Radbruch L, Brearley SG. Guidance on Conducting and REporting DElphi Studies (CREDES) in palliative care: Recommendations based on a methodological systematic review. *Palliat Med*. 2017;31(8):684-706. Epub 2017/02/14. doi: [10.1177/0269216317690685](https://doi.org/10.1177/0269216317690685). PubMed PMID: 28190381.
- 512
- 513
- 514 53. Hasson F, Keeney S, McKenna H. Research guidelines for the Delphi survey technique. *J Adv Nurs*. 2000;32(4):1008-15. Epub 2000/11/30. PubMed PMID: 11095242.
- 515
- 516
- 517