

# The epidemiological investigation of hyperlipidemia in the northern Henan Province

Xing Lu<sup>1</sup>, Ziyang Lin<sup>2</sup>, Junzheng Yang<sup>2\*</sup>

<sup>1</sup>Cardiovascular medicine department IV of Xinxiang Central Hospital,  
Xinxiang Henan 453000, China.

<sup>2</sup>Guangdong Nephrotic Drug Engineering Technology Research Center, The R&D Center of Drug  
for Renal Diseases, Consun Pharmaceutical Group, Dongpeng Avenue, No.71, Guangzhou,  
Guangdong 510530, China.

\*Correspondence to Junzheng Yang (yangjunzheng606403@163.com), Guangdong Nephrotic  
Drug Engineering Technology Research Center, The R&D Center of Drug for Renal Diseases,  
Consun Pharmaceutical Group, Dongpeng Avenue, No.71, Guangzhou,  
Guangdong 510530, China.

## Abstract

**Background/purpose:** Hyperlipidemia is a kind of diseases with the characteristic of high level of blood lipids, it could directly result in several serious diseases including atherosclerosis and coronary heart disease to threat to human health. Therefore, to understand the epidemiological status and risk factors of hyperlipidemia is important for the prevention, diagnosis, and treatment of hyperlipidemia.

**Methods:** The clinical data in Xinxiang Central Hospital from 2019-2021 were collected, and those data were classified and analyzed according to the basic information of investigation subjects and risk factors.

**Results:** A total of 670 investigation subjects were collected in this investigation including 521 males and 149 females, accounting for 77.76% and 22.24% in the total investigation population, respectively; age range of investigation subjects was 20-97 year old; there were 390 people with a history of smoking and 346 people were never smoking, accounting for 58.21% and 51.64% in the total investigation population, respectively; in this investigation, the proportion of peasant farmers was the highest (353/670, 52.69%), followed by retire population (164/670, 24.48%); in terms of education level, the proportion of people graduating from middle school was the

highest (338/670, 50.45%), followed by those graduating from primary school (249/670, 37.16%); the statistical results found that the incident of hyperlipidemia were correlated to gender ( $P<0.01$ ), ethnicity ( $P<0.01$ ), age ( $P<0.05$ ), marital status ( $P<0.05$ ), education level ( $P<0.001$ ), hypertension ( $P<0.05$ ) and diabetes mellitus ( $P<0.001$ ), had no relation to occupation ( $P>0.05$ ), cerebral hemorrhage, angina pectoris, ischemic stroke, gastrointestinal ulcers, coronary heart disease, chronic renal insufficiency, myocardial infarction, heart failure, peripheral artery disease, chronic obstructive pulmonary disease ( $P>0.05$ ).

**Conclusion:** The epidemiological status of hyperlipidemia in northern Henan Province had the specificity, the incident of hyperlipidemia was correlated to gender, ethnicity, age, marital status, education level, hypertension, and diabetes mellitus, those results may provide basis for the prevention and treatment for hyperlipidemia in northern Henan Province.

**Keywords:** hyperlipidemia; epidemiology; prevention and treatment; risk factor; northern Henan Province.

## **Introduction**

Hyperlipidemia is a kind of disease that endangers human health caused by high levels of blood lipids (including triglyceride, total cholesterol, and low-density lipoprotein cholesterol increased, and high-density lipoprotein cholesterol decreased in plasma) in the human body, according to ‘the Prevention and Treatment guidelines for Dyslipidemia in Chinese adults (revised in 2016)’, hyperlipidemia could be divided into four types, including hypercholesterolemia, hypertriglyceridemia, mixed hyperlipidemia and low high-density lipoprotein, the clinical manifestations of hyperlipidemia include xanthoma caused by lipid deposition in the dermis and arteriosclerosis caused by lipid deposition in the vascular endothelium [1]. In the normal human body, most patients do not have obvious symptoms or abnormal signs until blood biochemical tests are conducted to detect an increase in plasma lipoprotein levels, which undoubtedly increase difficulty to the prevention and treatment of hyperlipidemia.

It is reported that the average serum level of TC, LDL-C, and TG of adults in

2018 in China were significantly increased compared with the data obtained from the national survey in 2015; moreover, the data demonstrated that the average levels of TC and non HDL-C in Chinese adults are approaching or exceeding the average level of some western countries [2, 3], and it is predicted that an increase in serum cholesterol levels in the Chinese population is expected to lead to an increase of approximately 9.2 million cardiovascular events in China from 2010 to 2030 [4], those evidences demonstrated that hyperlipidemia is becoming a chronic noninfectious disease diseases that threaten human life and health, to understand the epidemiological status and risk factors of hyperlipidemia is necessary for prevention and treatment of hyperlipidemia. For this purpose, we collected and analyzed the clinical data in Xinxiang Central Hospital from 2019-2021, hope to provide some useful information for prevention and treatment of hyperlipidemia in north Henan Province.

## **Methods**

A total of 670 clinical data in Xinxiang Central Hospital from 2019-2021 were collected, and those data were classified and analyzed according to the basic information of investigation subjects including gender, ethnicity, age, smoking, marital status, occupation, education level and risk factors including hyperlipidemia, diabetes mellitus, ischemic stroke, coronary heart disease, angina pectoris, myocardial infarction, heart failure, peripheral artery disease, chronic renal insufficiency, gastrointestinal ulcers, chronic obstructive pulmonary disease, cerebral hemorrhage.

## **Results**

### **1. The correlation between general information of investigation subjects and hyperlipidemia in North Henan Province**

The results demonstrated that there were 530 non-hyperlipidemia people and 130 hyperlipidemia people, accounting for 79.10% and 19.40% in the total investigation population, respectively; there were 96 male hyperlipidemia people and 44 female hyperlipidemia people, accounting for 68.57% and 31.43% in the total hyperlipidemia people, respectively; there were 664 Han ethnicity people and 6 Hui ethnicity people, accounting for 99.10% and 0.90% in the total investigation population, respectively;

the age range of investigation population was 20-97 year old , and the number of people aged between 61-80 year old was highest (340/670, 50.75%); there were 390 people with a history of smoking, accounting for 58.21% in the total investigation population, 346 people were never smoking, accounting for 51.64% in the total investigation population; in this investigation, the proportion of peasant farmers was the highest (353/670, 52.69%), followed by retire population (164/670, 24.48%); in terms of education level, the proportion of people graduating from middle school was the highest (338/670, 50.45%), followed by those graduating from primary school (249/670, 37.16%); the statistical results found that the incident of hyperlipidemia were correlated to gender ( $P<0.01$ ), ethnicity ( $P<0.01$ ), age ( $P<0.05$ ), marital status ( $P<0.05$ ), and education level ( $P<0.001$ ) (Table 1).

**Table 1.** The correlation between general information of investigation subjects and hyperlipidemia in North Henan Province

General information		Non-hyperlipidemia ( n=530)		Hyperlipidemia (n=140)		Significance /P value
		n	%	n	%	
Gender	Male	425	80.19%	96	68.57%	0.0043**
	Female	105	19.81%	44	31.43%	
Ethnicity	Han	524	98.87%	140	100%	0.0043**
	Hui	6	1.13%	0	0.00%	0.3528
Age	20-40	21	3.96%	2	1.43%	0.3528
	41-60	173	37.27%	47	28.53%	0.1933
	61-80	268	50.57%	72	51.43%	0.0171*
	≥81	68	12.83%	19	13.57%	0.9243
Smoking	Currently smoking	220	41.51%	48	34.29%	0.7792
	Quit smoking ≥ 1 years	33	6.23%	8	5.71%	0.1455
	Quit smoking ≥ 1 months	15	2.83%	0	0.00%	>0.9999
	Never smoking	262	49.43%	84	60.00%	0.05*

Marital status	Married (Remarried)	503	94.91%	135	96.43%	0.0288*
	Single	3	0.57%	0	0.00%	0.6553
	Bereave	19	3.58%	5	3.57%	>0.9999
	Divorce	5	0.94%	0	0.00%	>0.9999
Occupation	Peasant farmer	273	51.51%	80	57.14%	0.5984
	Worker	83	15.66%	19	13.57%	0.2541
	Civil servants/ Government officials	6	1.13%	2	1.43%	0.5984
	Retirees	129	24.34%	35	25.00%	0.6753
	Professional technical personnel	8	1.51%	0	0.00%	0.9121
	Individual business owners	6	1.13%	1	0.71%	>0.9999
Education level	Illiteracy	31	5.85%	2	1.43%	0.0282*
	Primary school	179	33.77%	70	50.00%	0.0006***
	Middle School/Vocational School	273	51.51%	65	46.43%	0.2971
	College/University	46	8.68%	2	1.43%	0.0014**
	Postgraduate	1	0.19%	1	0.71%	0.3745

\*P<0.05, \*\*P<0.01, \*\*\*P<0.0001.

## 2. The correlation between risk factors and hyperlipidemia in north Henan Province

And then we analyzed the correlation between risk factors and hyperlipidemia, found that the incident of hyperlipidemia was correlated to hypertension and diabetes mellitus, the P value were 0.0101 and 0.0001, respectively, had no relation to cerebral hemorrhage, angina pectoris, ischemic stroke, gastrointestinal ulcers, coronary heart disease, chronic renal insufficiency, myocardial infarction, heart failure, peripheral artery disease, chronic obstructive pulmonary disease (P>0.05) (Table 2).

**Table 2. The correlation between risk factors and hyperlipidemia in north Henan Province**

Risk factors	Non-hyperlipidemia ( n=530)		Hyperlipidemia (n=140)		Significance/ P value
	n	%	n	%	
Hypertension	255	48.11%	85	60.71%	0.0101*
Diabetes mellitus	85	16.04%	47	33.57%	<0.0001***
Ischemic Stroke	73	13.77%	28	20.00%	0.0834
Coronary heart disease	4	0.75%	1	0.71%	>0.9999
Angina pectoris	89	16.79%	15	10.71%	0.088
Myocardial infarction	39	7.36%	6	4.29%	0.2548
Heart Failure	8	1.51%	1	0.71%	0.6932
Coronary intervention	24	4.53%	2	1.43%	0.1362
Peripheral Artery Disease	1	0.19%	1	0.71%	0.3745
Chronic renal insufficiency	3	0.57%	1	0.71%	>0.9999
Gastrointestinal ulcers	15	2.83%	1	0.71%	0.2146
Chronic Obstructive Pulmonary Disease	7	1.32%	1	0.71%	>0.9999
Cerebral hemorrhage	6	1.13%	1	0.71%	>0.9999

\*P<0.05, \*\*P<0.01, \*\*\*P<0.0001.

## Discussion

Hyperlipidemia, also known as dyslipidemia, is usually associated with elevated triglycerides and total cholesterol in plasma, is developing into one of the main diseases that affect people's daily health. Hyperlipidemia can be divided into primary hyperlipidemia and secondary hyperlipidemia according to pathogenesis, it is reported that hyperlipidemia could occur at different ages and genders, the incidence rate of hyperlipidemia was increased with the age, and the highest incidence occurs between 50-69 years old; many factors including gender [5, 6], alcohol consumption [7, 8], smoking [9], diet [10], physical activity [11], and emotional activity [12] could affect the occurrence of hyperlipidemia. In our data, we also found that the incident of hyperlipidemia was correlated to gender (P<0.01), ethnicity (P<0.01), and age (P<0.05) in north Henan province, these results are consistent with the previously

reported results, taking regular exercise and dietary therapy may effectively decrease or alleviate the occurrence of hyperlipidemia; furthermore, our data also demonstrated that the marital status ( $P < 0.05$ ) and education level ( $P < 0.001$ ) also have the higher correlation to the incident of hyperlipidemia, there is a significant difference in the incidence of hyperlipidemia between married populations and unmarried populations, between high educated populations and low educated populations, the reason for this result may be because married populations and unmarried populations, high educated populations and low educated populations have the different economic situation and dietary habits, but the further research is needed.

Hyperlipidemia is closely related to hypertension and diabetes mellitus, in many cases, hypertension and diabetes mellitus also could result in hyperlipidemia, it is reported that hyperlipidemia is also the main risk factor of hypertension and diabetes mellitus [13, 14]. In our data, we found that the incident of hyperlipidemia was highly correlated to hypertension ( $P = 0.0101$ ) and diabetes mellitus ( $P = 0.0001$ ) in north Henan Province, these evidences demonstrated we should strengthen the medical care and management of people with hypertension and diabetes mellitus, simultaneously combined with taking regular exercise and dietary therapy to decrease the incidence of hyperlipidemia.

## **Conclusion**

The epidemiological status of hyperlipidemia in northern Henan Province had the specificity, the incident of hyperlipidemia was correlated to gender, ethnicity, age, marital status, education level, hypertension, and diabetes mellitus, those results may provide treatment or prevention reference guide for hyperlipidemia in northern Henan Province.

## **Acknowledgements**

None.

## **Conflict of interests**

There is no conflict of interest in this article.

## **Funding**

None.

## References

- [1] China Adult Dyslipidemia Prevention and Control Guide Revision Joint Committee Guidelines for the prevention and treatment of Dyslipidemia in Chinese adults (revised in 2016). *Chinese Journal of Cardiovascular Disease*, 2016, 44 (10): 833-853 (In Chinese).
- [2] Song PK, Man QQ, Li H, et al. Trends in lipids level and dyslipidemia among Chinese adults, 2002-2015. *Biomed Environ Sci.*, 2019, 32(8):559-570.
- [3] Yang W, Xiao J, Yang Z, et al. Serum lipids and lipoproteins in Chinese men and women. *Circulation*, 2012, 125(18): 2212-2221.
- [4] Moran A, Gu D, Zhao D, et al. Future cardiovascular disease in China: markov model and risk factor scenario projections from the coronary heart disease policy model-China. *Circ Cardiovasc Qual Outcomes*, 2010, 3(3): 243-252.
- [5] Lin CC, Nfor ON, Su CL, Hsu SY, et al. Interactive associations of sex and hyperlipidemia with calcific tendinitis of the shoulder in Taiwanese adults. *Medicine (Baltimore)*, 2020, 99(46):e23299.
- [6] Leemaqz SY, Kyinn M, Banks K, Sarkodie E, et al. Lipid profiles and hypertriglyceridemia among transgender and gender diverse adults on gender-affirming hormone therapy. *J Clin Lipidol.*, 2023, 17(1):103-111.
- [7] Chen EX, Tu Ya SQ, She ZF, et al. The clinical characteristic of alcohol-hyperlipidemia etiologically complex type of acute pancreatitis. *Eur Rev Med Pharmacol Sci.*, 2022, 26(19):7212-7218.
- [8] Huang X, Hui H, Zhu W, Chen N, et al. Effect of the interaction between alcohol and meat consumption on the hyperlipidaemia risk among elderly individuals: Evidence from Shanghai, China. *Front Nutr.*, 2022, 9:982626.
- [9] Zhenhan Li, Guoqi Zhu, Guo Chen, Mei Luo, et al. Distribution of lipid levels and prevalence of hyperlipidemia: data from the NHANES 2007-2018. *Lipids Health Dis.*, 2022, 21(1):111.
- [10] Zheng ZG, Zhu ST, Cheng HM, Zhang X, et al. Discovery of a potent SCAP degrader that ameliorates HFD-induced obesity, hyperlipidemia and insulin resistance



via an autophagy-independent lysosomal pathway. *Autophagy*, 2021, 17(7):1592-1613.

[11] He N, Ye H. Exercise and Hyperlipidemia. *Adv Exp Med Biol.*, 2020, 1228:79-90.

[12] Zhang Hai-Na, Yu Xu-Ben, Tang Cong-Rong, Cao Yan-Cheng, et al. Atorvastatin ameliorates depressive behaviors and neuroinflammatory in streptozotocin-induced diabetic mice. *Psychopharmacology (Berl)*, 2020, 237(3):695-705.

[13] Rutugandha Paranjpe, Hua Chen, Michael L Johnson, Kim Birtcher, et al. Adherence to concomitant diabetes, hypertension, and hyperlipidemia treatments among older patients. *J Am Pharm Assoc.*, 2022, 62(4):1351-1358.

[14] Tung YC, Lee MY, Wang YC, Chang CJ, et al. Effect modification of hyperlipidemia and hypertension on the association between type 2 diabetes and gout. *Prim Care Diabetes*, 2020, 14(6):654-662.