- Barriers influencing implementation of patient safety incident reporting and learning
- 2 guidelines in specialised care units, KwaZulu-Natal: A qualitative study
- 3 T.M.H Gqaleni <sup>1</sup>, Sipho. W. Mkhize <sup>1</sup>
- <sup>1</sup>School of Nursing and Public Health, College of Health Sciences, University of
- 5 KwaZulu-Natal, Durban, South Africa.
- 6 E-mail:gqalenit@ukzn.ac.za. (TMH)

7

8

#### Abstract

#### 10 **Background**

9

16

19

25

- Globally, increased occurrences of Patient Safety Incidents have become a public concern.
- The implementation of Patient Safety Incidents incident reporting and learning guidelines
- is fundamental to reducing preventable patient harm. To improve the implementation of
- these guidelines in specialised care units in KwaZulu- Natal, the views of healthcare
- professionals were unearthed.

#### Aim

- 17 This study explores the healthcare professional views toward the implementation of Patient
- Safety Incidents reporting and learning Guidelines in specialised care units.

#### Methods

- A descriptive, explorative qualitative approach, was used to collect qualitative data from
- 21 healthcare professionals working in specialised care units. The study was conducted in
- SCUs of three purposely selected public hospitals in two districts of KwaZulu-Natal. Group
- discussions and semi-structured interviews were conducted from March to May 2021.
- 24 Content data analysis was performed using Tesch's method of analysis process.

#### Results

- The main themes that emerged during data analysis were: Ineffective reporting system
- affecting the communication of PSI guidelines, Inadequate Institutional and management
- for the healthcare professionals, Insufficient education and training of healthcare
- 29 professionals, and poor human resources affecting the implementation of PSI guidelines.

The findings highlighted that there were more major barriers to the implementation of PSI reporting and learning guidelines.

#### Conclusion

This study confirmed that PSI reporting and learning guidelines are still not successfully implemented in the specialised care units and the barriers to implementation were highlighted. For rigorous implementation in South Africa, the study recommends revised PSI reporting and learning guidelines, designed in consultation with the frontline healthcare professionals, consisting of standardised, simple -user-friendly reporting process as well as a better implementation strategy to guide the healthcare professionals. Continuous professional development programmes may play an important role in the facilitation of the implementation process.

#### Introduction

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

Globally, increased occurrence of Patient Safety Incidents (PSIs) and near misses have become a public concern, which includes South Africa [1]. Near misses are incidents or situations that have the potential to cause harm but did not reach the patient due to timely intervention, whereas Patient Safety Incident (PSI) is harm caused by medical mismanagement, instead of the underlying disease[2]. Globally, unsafe care is one of the top 10 leading causes of death in the world and up to 83% of harm is avoidable [3]. The Institute of Medicine reveals death related to PSIs varies from 44 000 to 98 000 people each year with high-cost implications [4-6]. According to Flott, Fontana (3), over 420 million hospitalisations each year around the world resulted in nearly 43 million PSIs, in 2013. An analysis by Organisation for Economic Co-operation and Development (OECD) found that 15 percent of all hospital costs in OECD nations are due to patient harm from PSIs, with associated costs ranging between \$1.4 to \$1.6 trillion, each year [7]. According to Choi, Pyo (8), Lu, Ko (9), improving the hospital organization atmosphere is essential to facilitate useful communication of patient safety incidents. In Swedish and Brazilian health services, barriers and facilitators that influenced patient safety were identified and the relevance of a multifaceted system perspective on patient safety problems and solutions was pointed out [10, 11]. According to [3, 9], patient safety is a multifactorial concept, driven by a complicated array of technical, human and system factors. Several studies have identified barriers influencing the implementation of PSIs reporting in healthcare settings including specialised care units (SCUs). These includes underreporting, punitive culture, lack of standardized reporting system, staff shortages, lack of safety culture, poor processing of incident reports, inadequate engagement of healthcare professionals, limited institutional support of incident reporting systems including inadequate usage of evolving health information technology [1, 12-17].

In middle and low-income countries (LMICs), almost one-third of patients who suffered PSIs died and four out of five of those incidences were preventable [18]. Moreover, 134 million PSIs occur in hospitals due to unsafe care, resulting in 2.6 million deaths, each year (National Academies of Sciences and Medicine, 2018). Also, it is estimated that 75% of PSIs occur in the LMIC contexts, during hospital care and 10-15% of total deaths are as a result of poor quality care [3]. In South Africa, a study conducted in KwaZulu-Natal reveals high percentage of serious PSIs (47%) indicating poor implementation of PSI reporting guidelines and lack of improvement strategies [19]. While incidents reporting is considered a cornerstone to patient safety culture and quality improvement in healthcare [20, 21], underreporting in hospitals worldwide is still a challenge[1, 22, 23].

In South Africa, the implementation of Patient Safety Incident Reporting and Learning Systems guidelines, as recommended by KwaZulu-Natal Department of Health (KZN DoH), in line with National Department of Health (NDoH 2017) has provided no clear evidence of the reduction of PSIs and if these guidelines are adequately implemented in SCUs. Gqaleni and Mkhize (17), affirmed that healthcare professionals' perception of knowledge of PSI reporting and learning guidelines was good, however, the perception of the implementation was poor. This study further recommended a revised implementation to improve the quality and safety of healthcare in (SCUs). For SCUs, to heighten the positive performance and practices of PSIs reporting and learning guidelines, the views of healthcare professionals needed to be unearthed. Therefore, the aim of the study was to explore and describe the views of the senior healthcare professionals regarding the implementation of Patient Safety Incident reporting and learning guidelines in specialised care units of KwaZulu-Natal.

## **Materials and Methods**

## Study design and setting

A descriptive, explorative qualitative approach, was used to collect qualitative data from healthcare professionals working in SCUs. The study was conducted in SCUs of three purposely selected public hospitals in two districts of KwaZulu-Natal. Two hospitals (A and C), situated in the eThekwini district, provide both secondary and tertiary services, and the third tertiary hospital (B) is in the uMgungundlovu district. The settings consisted of specialised care units that included both Critical Care Units (CCUs), where the most critically ill patients were admitted as well as High Care Units, where the recovering critically ill patients were transferred but still needed specialised care.

#### Table 1: Inclusion and Exclusion criteria

Inclusion	Exclusion
Participants who were Operational Nurse Managers	Participants who were not Operational Nurse Managers
Participants who were Assistant Nurse Managers	Participants who were not Assistant Nurse Managers
Participants who were Consultant Medical doctors	Participants who were not Consultant Medical doctors
Participants who were Monitoring & Evaluation Managers	Participants who were not Monitoring & Evaluation Managers
Working experience of more than 10 years in specialised care units	Participants who did not work in these units and had less than 10 years of experience.
Willingness to participate	Not willing to participate

## Population, recruitment and sampling

Sampling is a process of selecting a portion of the population to represent the entire population, so that inferences about the population can be made [24]. The target population for this study consisted of senior healthcare professionals, who were involved in the implementation of PSI reporting and learning guidelines in the SCUs. Therefore, the participants that were purposely selected were had expertise and experience in SCUs environments and had vast experience in the handling of PSI reporting and learning guidelines. The focus group discussions sample was comprised of Operational Nurse Managers (OMs). Individual interviews included Assisting Nurse Managers (ANMs), Monitoring, and Evaluation Managers (M&E), and Consultant Medical doctors (CMD).

#### **Recruitment strategy**

Appointments were made by the researcher with the institutions' Senior Nurse Managers and Operational Nurse Managers and all the participants prior to data collection. The researcher had a briefing session with the participants, prior to data collection, explaining the aim of the study and its importance to research. Participants were informed that they were not forced to participate and that they had a right to withdraw from the study even if they had already given consent, without any negative consequences, thus ensuring voluntary participation.

## **Data collection process**

This study was conducted during COVID-19 and the researcher followed the protocols. The data were collected virtually to ensure social distancing, which was reinforced in the boardrooms, including the use of personal protective equipment by the participants. Data

was collected over a period of 3 months, from August to October 2021. Information guidelines guided the focus group discussions and individual interviews, which consisted of topics and probing questions to elicit more detailed information from the participants. The researcher prepared one essential question with a list of probing open-ended questions, although the participants were encouraged to speak freely about the topic, to obtain all the information required.

Focus groups discussions, which consisted of Operational Nurse Managers, working in the selected settings, as they formed homogeneous participants. These participants were also at the forefront of implementing the PSI reporting and learning guidelines. Individual interviews consisted of senior participants which were heterogenous, i.e. Assisting Nurse Managers, Monitoring, and Evaluation Managers, and Consultant Medical Doctors to solicit more in-depth information. The probing question were influenced by the participants' responses and the interviewer sought clarity to minimize bias and subjectivity. Data were collected until saturation was reached, that is, no new information will be emerging. Interviews were conducted and lasted between 45 minutes to 60 minutes, and this allowed the participants to freely elaborate on their opinions on the phenomenon. Participants gave consent to record all the interviews and the researcher transcribed the data verbatim.

#### **Ethical considerations**

The University of KwaZulu-Natal's Human Science Research Ethics Committee (HSSREC/00001651/2020), and the Directorate: Health Research and Knowledge Management Unit (NHRD Ref: Kz\_202010\_0240), granted ethical clearance and permission to conduct the study before data collection. Gatekeepers' permission was also obtained from the Chief Executive Officers, heads of department and Senior Nurse Managers. A verbal consent to participate and to the recording of the interview voices were

obtained from the participants prior to data collection. Grove, Burns (25) asserted that the researcher must comply with three principles as stated by the Belmont report, namely; the principle of beneficence, the principle of human dignity, and justice. The participants were informed of their rights to withdraw from the study at any time, without any negative repercussions.

### Data analysis

Content data analysis was conducted using Tesch's method of the analysis process [24]. Transcribed interviews were read several times and descriptive words were identified for the formulation of thematic statements and concepts that created meaning. Notes were made with ideas that came to mind. Each interview was read through and the underlying meaning behind the scripts was determined. Notes were made on transcripts during the reading process. A list of concepts was made and similar topics were clustered together. These clustered topics were then be mapped as major topics. Raw data comprising transcripts were reviewed again in the light of identified topics. The text was coded and organised. Descriptive words were identified in the text and used to describe themes or topics. Related topics were grouped together. A final decision regarding categories was made. Preliminary analysis was made and excerpts from the data supporting the identified categories and themes was coded and recoded. A reflective journal was used to form themes and subthemes.

#### **Trustworthiness**

During the research process, the researcher applied four criteria, to determine trustworthiness, namely, credibility, dependability, confirmability and transferability [26].

Credibility. According to Polit and Beck (27) credibility refers to the confidence in the truth of the data and its interpretations, that is, the data must reveal what it is searching for. The researcher ensured that this study was conducted in an authentic manner and the researcher had a prolonged engagement with the participants, during the data collection process, which assisted in gaining an in-depth understating of the phenomena under study.

**Dependability**. In this study the researcher provided detailed information of the research process, findings and recommendations, so that the other prospective researchers could replicate the study to their context. According to Polit and Beck (27), findings of an inquiry must be able to be repeated and replicated with the same participants in the same context.

Confirmability. Confirmability refers to objectivity, that is, the potential for congruence between two or more independent people about data's accuracy, relevance, or meaning [27]. In this study, the research report conveyed participants' experiences and not the perspectives of the researcher. Themes generated during data collection reflected the tone of the participants.

**Transferability**. Transferability refers to the extent to which findings can be generalized and applied in other settings or groups [27]. In this study the researcher was responsible for providing a detailed account of the research process, findings and recommendations, so that the other prospective researchers could evaluate the applicability to their context.

## **Results**

### Introduction

Data were collected from the purposively selected participants until data saturation was reached. Data analysis was conducted manually using the eight steps of Tesch's method of

analysis. The first section provides the participants' demographics, as illustrated in Table 2 and the second section outlines the findings of the study, thoughts and feelings expressed by the participants during the in-depth interviews. The names of the participating hospitals and the participants were not divulged to protect their identities.

**Table 2: Demographics of the participants** 

Hospital	Gender	Job level	
Focus group discussions			
Hospital A	7 Females; 3 Males	Operational managers	
Hospital B	8 Females; 2Males	Operational managers	
Hospitals C	6 Females; 2Males	Operational managers	
Individual interviews			
Hospital A	1 Female	Assistant Nurse Managers	
Hospital B	1 Male	Assistant Nurse Managers	
Hospitals C	1 Female	Assistant Nurse Managers	
Hospital A	1Female; 2 Males	Consultant Medical doctor	
Hospital B	1 Female; 2 Males	Consultant Medical doctor	
Hospitals C	1 Male	Consultant Medical doctor	
Hospital A	1 Female	Monitoring and evaluation Manager	
Hospital B	1 Male	Monitoring and evaluation Manager	
Hospitals C	1 Female	Monitoring and evaluation Manager	

Note: FGD- Focus group discussion

**OM-** Operational Nurse Manager

**ANM**-Assistant Nurse Manager

M&E- Monitoring and Evaluation managers

CMD- Consultant Medical doctor

A summary of themes and subthemes that emerged during data analysis is presented in Table 3.

### **Table 3: Themes and subthemes**

Theme	Description of a theme	<b>Sub-themes</b>
1.Ineffective reporting system	Communication factors are	1.1 Tedious, long and not simple
affecting communication of PSI	associated with the	
guidelines	communication of PSI reporting	1.2 Unclear PSI guidelines and
	and learning guidelines as	policies
	perceived by the healthcare	
	professionals	1.3 Lack of consultation at the
		forefront level
		1.4 Inadequate feedback to staff
		and family
2. Inadequate Institutional and	Institutional support relates to the	2.1 A hostile environment
management for the healthcare	support received by healthcare	characterized by hierarchical
professionals	workers within the institution	structure and red-tape
		2.2 Punitive culture
		2.3 Lack of management support to
		frontline healthcare workers
		2.4 Lack of teamwork
3. Insufficient education and	Education and training refer to	3.1 Inadequate training of
training of healthcare professionals	both qualification and expertise of	healthcare professionals on PSIs
	healthcare professionals	3.2 Lack of knowledge on how to
		construct the report properly
		3.3 Lack of formal training of
		healthcare professionals

3.4 Lack of formal training, inhouse training, and continued professional development programme.

4. Poor human resources affecting	Human resources refer to the	4.1 Shortage of staff
the implementation of PSI	factors that the healthcare	4.2 Inadequate competent a skill
guidelines	professionals perceived as	mix
	availability of competent staff to	4.3 High staff turnover and lack of
	implement PSI guidelines	staff retention
		4.4 Busy environment

The views of some of the participants are presented as direct quotations to support the identified themes and sub-themes.

# Theme 1: Ineffective reporting system affecting communication of PSI guidelines

Communication factors are associated with the communication of PSI reporting and learning guidelines as perceived by the healthcare professionals. Participants had a perception that communication was ineffective and not user-friendly to implement PSI reporting and learning guidelines. Most participants stated that the actual structure of the reporting system makes it difficult for the healthcare professionals to effectively implement the guidelines.

#### Subtheme 1.1: Tedious, long and not simple

The participants indicated that the reporting process was long and intense, tedious and laborious, hence other PSIs were not reported.

"It is a long form...the new form is very intense, apart from the report that you're going to write.... not simple at all." (ANM1, Hospital A, Female, from Individual interview)

220	"The time that it takes to fill in the PSI forms and then the possibility of further meetings
221	down the line, it is essentially inconvenience from the interview process, a very
222	laborious, inconvenient process." (CMD1, Hospital C, Male, from Individual
223	interview)
224	Subtheme 1.2: Unclear PSI guidelines and policies
225	The healthcare professionals also stated that the guidelines were not clear, confusing and
226	not standardised. This was considered a barrier to effective implementation of the
227	guidelines.
228	"The guidelines are not particularly clearer on when an adverse event can be
229	considered as just expected as a consequence of the disease for a particular patient
230	versus a true PSI. Our government is using their own terminology which is
231	unfortunately not based on international standards." (CMD2, Hospital A, Male, from
232	Individual interview)
233	Some participants verbalised that there was a mismatch of the guidelines with the actual
234	work environment
235	"Action plans and policies are nice and they are there, it's something that is there and
236	are nice to read, but when it comes to practicality of it guidelines may look good on
237	paper, but they are not effectively implemented." (OM2, Hospital A, Female, from
238	FGD)
239	Subtheme 1.3: Lack of consultation at the forefront level
240	Healthcare professionals also reported a lack of consultation at the forefront level and non-
241	clinical involvement during the development phase of PSI reporting and learning guidelines.

that the people that develop these guidelines are not clinical, they have no

understanding of what is happening on the ground, they have no medical training." 244 (CMD2, Hospital A, Male, from Individual interview) 245 246 "I didn't have any input, maybe from other managers. I just received a new form from the province informing us what is required." (ANM2, Hospital A, Male, from 247 Individual interview) 248 Subtheme 1.4: Inadequate feedback to staff and family 249 Inadequate feedback to staff and family was also identified by the participants as barrier to 250 the implementation of PSI reporting and learning guidelines. Lack of feedback was more 251 from the senior management and was more of a redress meeting. Partial family involvement 252 if a PSI has occurred, near misses are usually overlooked because it is perceived as harm 253 that did not reach the patient. 254 255 "So, people might say okay, what do we need to disclose that to the family because it was a near miss. It wasn't a miss. The patient didn't suffer any harm." (CMD1, Hospital 256 C, Male, from Individual interview) 257 "I've never had feedback. And I've done multiple incident reports reporting over the 258 years, never, ever, ever, ever had.... you would expect documents and feedback. You 259 260 don't ever get any feedback from them. It is a waste of time because you get nothing back. Then the next time something happens you go.... I'm not going to do it again

because what was the point?" (CMD1, Hospital B, Female, from Individual interview)

261

262

#### Theme 2: Inadequate Institutional and management for the healthcare

### professionals

Most participants expressed a lack of support, especially from the senior management, a disjuncture of the leadership from the challenges experienced by the frontline workers, resulting in poor of teamwork.

#### **Subtheme 2.1: Punitive culture**

Most participants reported that there is a punitive culture associated with the implementation of PSI reporting and learning guidelines. This was characterised by fear of the negative repercussions when healthcare professionals attempt to comply, therefore this led to underreporting of most PSIs.

"Sometimes this leads to disciplinary measures. Sometimes you say, I'm going to give you a warning. And the PN will complain that I committed myself by reporting, and now I must face the negative consequences. They even threaten to consult the unions."

(ANM1, Hospital B, Male, from Individual interview)

"Also, people have a view that it is punitive, in a way... so, I have seen different departments threaten each other with PSIs... eh... so if you call something a PSI there is still very much a mindset that this is an accusation... and it is even been recognised between departments to sort of thresh out their politics...... been used between departments as a sort as a blame. Even various people are using it almost punitively...aggressively" (CMD1, Hospital C, Male, from Individual interview)

#### **Subtheme 2.2: A hostile environment**

Participants described environment as hostile, characterized by hierarchical structure and red-tape resulting reluctance in the implementation of PSI guidelines by the healthcare professionals.

"registered nurses that fear, they don't want to report that that's why they are underreporting, because that fear you don't have support. It's always like a threatening environment. All they've learned is that they must keep quiet. And what if I get fired?"

#### (CMD3, Hospital A, Female, from Individual interview)

"To me it's like your call to the court where you're gonna sit and when they discuss this incident you feel that I am... I am on the wrong..." (OM1, Hospital B, Female, from FGD)

#### Subtheme: 2.3 Lack of management support to frontline healthcare workers

In addition, participants expressed a lack of guidance when trying to implement reporting and learning guidelines. Participants verbalised that an enabling environment is crucial, therefore inadequate organisational resources, patient safety committees that are not fully functional, lack of implementation strategies to guide healthcare professionals and learning opportunities were regarded as barriers.

"There is lack of support. Because... we are OMs, we work with staff, and we've got challenges. You need somebody to listen to you, if you say this and this is happening. Instead, you're blamed.... you are criticised... you... you ... you really feel bad." (OM2, Hospital B, Female, from FGD

"no one's learning from the previous mistakes And there's always another excuse. You just hit your head against the wall." (CMD1, Hospital B, Female, from Individual interview)

Participants also verbalised that patient safety committees were not existing, and those that were existing were at a development stage.

"But from what I hear they just getting it up and running and getting some processes in order. I think the first steps are happening there... I view it as the start of the process. but I don't know if we've quite got the systems at the unit level... we don't have the unit SOP or patient safety committees on how to deal with a PSI... it is more of an ad-hoc base ..." (CMD2, Hospital A, Female, from Individual interview)

"I'm not aware of any Patient Safety committee that exists. I don't even know what happens what the process is after I've handed the form as far as it goes from the ward to the ANM... that's the point of the SOP. I don't know what happens after that, which is bad as well, because we need to know the entire process... it's all... it's all very haphazard". (CMD2, Hospital C, Male, from Individual interview)

#### 2.4 Lack of teamwork

Participants reiterated that there should be a learning experience during the implementation of reporting and learning guidelines, not to blame someone. Participants reported that a teamwork approach between departments and collaboration with other institutions was lacking.

"We rely on nurses on the reporting of the PSI for formal reporting as such, because of a busy environment." (CMD2, Hospital A, Male, from Individual interview).

"Although we morbidity and mortality meetings where we discuss the PSIs as staff, there is still that lack with collaboration with other institutions, we hardly meet...I don't remember us meeting....so there is a gap there." (ANM1, Hospital B, Male, from Individual interview)

## Theme 3: Insufficient education and training of healthcare professionals

Participants reported that education and training associated with implementation of PSI reporting and learning guidelines, was multifaceted. The education and training challenges included the actual formal education of registered nurses, lack of training in specialisation of critical care, inadequate in-service training and workshops, to empower the participants on the guidelines. Furthermore, The PSI guidelines were not made, freely available and common knowledge to all the disciplines of departments.

#### **Subtheme 3.1: Lack of formal training of healthcare professionals**

Participants confirmed that some of the nurses that were allocated to the specialised units did not have the formal training and the skill that is required in this environment.

"In most cases we were not even sure of their training course, as they were training in these schools ...mushrooming everywhere, and now come in as ENAs and passed, they now want to be PNs, but still behaving like ENs...running the ward and there is no senior sister. You face these challenges, as they were the ENs recently and that affects the quality of nurses we employ." (ANM2, Hospital A, Male, from Individual interview)

"People don't know how to identify PSIs are? How do you report...There's a total lack of education here...? So, you can't blame the individuals if they weren't educated in

351	this and so I do think there's a major problem." (CMD 2, Hospital A, Female, from
352	Individual interview
353	Subtheme 3.2: Lack of knowledge on how to construct the report properly
354	Furthermore, the participants indicated that the reporting system is not user-friendly with
355	terminology that was not easy to understand. Although the participants knew about the
356	classification of PSIs, the reporting process was complex.
357	"I think the cause is a lack of knowledge of how to write a PSILack of knowledge on
358	how to construct properly how to report properly, the whole incident it has to be
359	conciseit has to be to the point, so if it is not written properly, I send it backso it's
360	more time needed." (ANM1, Hospital A, Female, from Individual interview)
361	"Then I mean, I'll say to the doctors you must fill out an incident reporting form and
362	they'll goWhat? that nobody knows what it is. What to do with it." CMD1, Hospital
363	A, Male, from Individual interview)
364	Subtheme 3.3: Lack of formal training, in-house training, and continued professional
365	development programme
366	Participants further indicated that there was insufficient continued professional development
367	programme to empower the healthcare professionals to construct the report properly.
368	Participants expected to have a common knowledge for all healthcare professionals and to
369	be familiar with the guidelines.
370	"The perception that the staff is not adequately trained to identify the PSIs and to
371	disclose to patients and relatives." (ANM2, Hospital A, Male, from Individual
372	interview)

"I think.... even the more senior clinician struggled to decide what is the PSI and what 373 isn't a PSI...ah... so, do you do label something a PSI" (CMD1, Hospital C, Male, from 374 Individual interview) 375 Subtheme 3.4: Inadequate training of healthcare professionals on PSIs 376 Some participants reported that some of the items that are in the PSI reporting and learning 377 guidelines were well understood. They explained that some of the PSIs were expected as 378 normal process of the underlying illness. 379 "Sometimes some of the items that are been included in those PSIs are not particularly 380 appropriate to the ICU, because its normal to follow the normal pathology in ICU." 381 (CMD2, Hospital B, Male, from Individual interview) 382 Theme 4: Poor human resources affecting the implementation of PSI 383 guidelines 384 Most participants confirmed that inadequate competent, knowledgeable and experienced 385 was a huge barrier to effective implementation of PSI reporting and learning guidelines. The 386 inadequate staff allocation was exacerbated during COVID-19 pandemic. 387 388 4.1 Subtheme: Shortage of staff 389 Specialised units are busy by nature, and normally a ratio of 1:1, that is comprised of 390 adequate skills mix. Participants stated that it was impossible to achieve this ratio, therefore 391 affecting the implementation of guidelines and compromised quality patient care. 392

393

394

"the... conditions that we working in are not allowing us to make sure can uhhh.... we

try and ensure, but there isn't adequate staff that staff is adequate, I mean people who

are who work here every day, and who knows what is happening here and not people that I will be orientating. But the situation is that instead of having staff in this ward to work, according to the guidelines, or the policy we still get people from all over who will come here not knowing exactly what to do. Shortage itself is a big issue (emphatically) You get somebody that has just qualified as PN, she has never worked in a ward, she's just new...new PN She hasn't seen a ventilator she's seeing the ventilator for the first time. She's seeing the monitor for the first time." (OM5, Hospital B, Female, from FGD)

#### Subtheme 4.2: Inadequate competent a skill mix

Participants also indicated that the staff that was allocated to their departments did not have specialised qualifications in SCU environment.

"You need staff that will understand what is happening.... we don't even have time to orientate them. It's just orientation... You need staff that will understand what is happening.... we don't even have time to orientate them. It's just orientation." (OM5, Hospital B, Female, from FGD)

410

411

412

413

414

415

416

417

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

#### Subtheme 4.3: High staff turnover and lack of staff retention

Moreover, some participants expressed their frustration on high staff turnover and lack of staff retention

"There is also an increase in the number of retiring staff and the management is not recruiting for the vacant posts... they are not doing anything new. The staffing challenge is there. It takes too long to replace staff and you must motivate why you need more staff. Although there may be trained and experienced ICU nurses, they are not enough

for the number of days and you find yourself not achieving that ratio of 1:" (ANM3, Hospital A, Female, Individual interview) **Subtheme 4.4: Busy environment** Participants indicated that they work in a very environment, looking after vulnerable critically ill patients that require individual attention. "It is very difficult for staff, especially in this busy critical care where there is staff shortages... to now follow up ... because somebody will be having a patient...remember here the ratio here is 1:1 or 1:2." (ANM3, Hospital A, Female, Individual interview) "we work with sort of skeleton staff. Every day you wonder how people are going to 

cope with more admissions." (OM6, Hospital B, Female, from FGD)

## **Discussion**

This study sought to explore the views of the healthcare professionals who were at senior level, regarding the barriers and facilitators to implementation PSI reporting and learning guidelines in SCUs. Participants highlighted that although there are stipulated guidelines, they are not effectively executed by the health care professionals. They reflected more barriers than facilitators which could be perceived as hindrance the implementation of these guidelines. All participants acknowledged the importance of revised improvement strategies to facilitate the rigorous implementation and improvement of quality patient care was essential.

## Theme 1: Ineffective reporting system affecting communication of PSI

### guidelines

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

In this study, the participants indicated that the difficulty was more on the process of communication of the PSI reporting and learning guidelines. This was at multi-level, amongst the individuals, departments, senior management and the way the whole system is structured. Also, an organisational patient safety cultures must be cultivated to create an atmosphere that is conducive to communication of these guidelines. Barriers that were mostly mentioned by the participants included a laborious inconveniencing process, unclear guidelines and policies, not simple and user-friendly, lack of consultation at the coalface and lack of feedback [1, 10, 11]. Several studies confirm that clear, standardised effective communication is key to effective implementation by the healthcare professionals. Although participants in Hospital B demonstrated a positive attitude to the implementation of PSI reporting and learning guidelines, as they stated that they were partially involved during the consultative process, this was not the case in hospital A and C, as they indicated that they were never consulted. The government plays a critical role in monitoring and evaluation of the implemented PSI guidelines, therefore the lack of involvement of Department of health in managing incident reporting means that they are not aware of patient safety issues in the hospitals in their regions [28] Lack of feedback to the staff and patients and uncertainty about what to report does not encourage the healthcare professionals to use the reporting system effectively, hence the participants indicated that there was a lot of undocumented PSIs [3, 11, 15]. Participants also indicated that there is no learning that takes place, where healthcare professionals can use the PSI as learning opportunity to prevent recurrences, as they learn from their mistakes. It is important that healthcare professional do not rely on reporting incidents only as reports

often do not lead to positive change [11]. Participants also stated that the feedback to patients and families is partial, only if it is a major event. Martos Algarra (29) affirms that patients become first victims and the professionals involved turn into second victims and all of them need adequate support.

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

## Theme 2: Inadequate Institutional and management for the healthcare professionals

Investing in patient safety culture can be viewed as building an organizational resource, which is beneficial for both improving the care quality and protecting staff well-being [9]. There is a need for more effective transformational leadership to facilitate a culture of patient safety incident reporting and effective implementation of PSI reporting and learning guidelines [22]. In this study, participants explained an atmosphere that is hostile, punitive and lack of support of the frontline healthcare workers from the management [11, 22]. Fear of negative repercussion and litigation were the main barriers of implementation of PSI reporting and learning guidelines [30]. According to Ontario (30), Lee, Wu (31), a nonpunitive, confidential, independent, expert analysis, timely, system-oriented, and responsive, was suggested, a hospital should particularly pay more attention to the confidentiality of case data in the system, when establishing an incident reporting system, to avoid disputes and enhance reporting intention. Lack of teamwork and collaboration with other departments and institutions was also mentioned by the participants. A similar study highlighted a lack of collaboration between physicians and other health care personnel functioned as a barrier to improved patient safety due to perceived hierarchical differences. [10]. Participants also stated that although they had risk management meeting to discuss PSIs, there was a lack of guidance in the form of patient safety committees, implementation strategy and standard operational procedure, that was of common knowledge. A similar

study highlighted that the availability and use of written protocols that provide structure for their work, such as guidelines and standardized care plans, positively influence patient safety [10]. According to Dhamanti, Leggat (28) Lack of commitment to and priority of patient safety, the complexity of the bureaucratic structure, and a lack of systematic partnership and collaboration are problems that need to be addressed by systematic improvement.

#### Theme 3: Insufficient education and training of healthcare professionals

Insufficient education and training that was expressed by participants was multifaced. Participants mentioned poorly qualified healthcare professionals with no skills of working in a specialised care environment. Insufficient skills could act as a barrier to enhanced patient safety. Being a newly graduated nurse was mentioned as a barrier, due to a general lack of experience and the need for learning concerning many new aspects of work [10]. Also, there was a lack on how to identify PSIs, how to construct a written report, inadequate in-service training. A similar study confirm that lack of knowledge about incident reporting systems, and lack of understanding about what constitutes an error was reported as common barrier [30]. The government plays a critical role in ensuring patient safety in health care services as they are actively involved in designing the guidelines and development of policies. Therefore, either the government or medical institute managers should collect and analyse the information in the system, reduce the reoccurrence of medical errors through education and training and improvement activities, and enhance the patient safety culture of the hospital [28]

## Theme 4: Poor Human resources affecting the implementation of PSI

### guidelines

Participants reported the lack of human resources was not only insufficient number of the staff allocated to the specialised care units, but skill mix was also lacking. The staff shortage was due to increased number of the staff at a retirement age, staff resigning for better employment renumeration, recruitment by other countries for competent staff. This brain drain could be associated with a burnout syndrome, therefore poor implementation of PSI reporting and learning guidelines. The specialised care units are characterised by a busy, stressful environment, with increased to work overload, working with a skeleton staff might lead to burnout [32]. Shortage of staff and increased workload was associated poor implementation of the PSI reporting guidelines [9, 16, 28]. Mossburg and Himmelfarb (32), affirm that healthcare managers can consider a simpler measure of this resource in ensuring adequate staffing levels across all departments of the institution.

## Strengths and limitations of the study

This study provided thought provoking findings as in-depth information was collected from the interviews from the focus group discussions and individual interviews from participants who were at the managerial positions and were actively involved in the implementation of PSI reporting and learning guidelines. In addition, the purposive sample from the focus group was homogenous, whereas the individual interviews were heterogenous, which provided a rich information of experiences from a various healthcare professional. To the researchers' knowledge this is the first study to enlighten the barriers to implementation of PSI reporting and learning guidelines by the healthcare professionals in South African SCUs. Furthermore, the study findings may inform the policy makers and healthcare

facilities to revise the PSI reporting guidelines, to be more relevant to the South African context and devise better implementation strategy, to improve patient safety care. In addition, the findings from this study have identified areas that need improvement in patient's safety, therefore its dissemination contributes to the cultivation of patient safety culture worldwide. The study participants were limited to three major government hospitals in KwaZulu-Natal, and this main limitation of the study, however the sample was representative, therefore enough to address the research question. Also, most of the literature that supported this study is more international, there is paucity of South African literature.

## Recommendations

This study was conducted in one province, therefore further research is recommended to be conducted in other provinces. The study research analysis and results can be used by global and country stake holders to improve the implementation of PSI reporting and learning guidelines, as per WHO recommendation. For rigorous implementation in South African SCUs, the study recommends revised PSI reporting and learning guidelines that consist of standardised, simple -user-friendly terminology as well as a better implementation strategy to guide the healthcare professionals to implement the guidelines effectively. These guidelines need to be designed in consultation with the frontline healthcare professionals by the policymakers. Continuous professional development programmes may play an important role in facilitating the implementation process. Education on PSIs reporting and learning guidelines should be incorporated early in the curriculum of healthcare professionals, so that it becomes common knowledge for all.

### Conclusion

This study confirmed that PSI reporting and learning guidelines are still not effectively implemented in the SCUs and the barriers of implementation were highlighted. There is a need to implement these guidelines in middle and low-income countries, including South Africa. The study revealed that ineffective reporting system, lack of institutional and managerial support, insufficient education and training of healthcare professionals and poor human resources were the main barriers to effective implementation of PSI and learning guidelines. To effectively improve patient safety, this study seeks to strengthen collaboration among the healthcare professionals, organisations, within the African continent and globally.

#### **Supporting information**

- S1 Appendix: Interview guide for the focus group discussion
- 566 S2 Appendix: Individual Interview guide

#### Acknowledgement

- The authors would like to acknowledge the gatekeepers and the healthcare professionals
- from the three hospitals, who assisted and participated in the study.

#### Author contributions.

- 572 Conceptualization: TMH Gqaleni, SW Mkhize
- **Study design, data collection**: TMH Ggaleni.
- Formal analysis: TMH Gqaleni.
- **Investigation**: TMH Gqaleni.
- **Methodology**: TMH Gqaleni.

577	Project administration: TMH Gqaleni.
578	Resources: TMH Gqaleni.
579	Supervision: SW Mkhize.
580	Validation: SW Mkhize.
581	Writing – original draft: TMH Gqaleni.
582	Writing – review & editing: TMH Gqaleni, SW Mkhize
583	
584	Conflict of interest: None
585	
586	
587	
588	
589	
590	
591	
592	
593	
594	
595	
596	
597	
598	
599	
600	

## References

602 603 604	1.	Yali G, Nzala SH. Healthcare Providers' Perspective on Barriers to Patient Safety Incident Reporting in Lusaka District. Journal of Preventive and Rehabilitative Medicine. 2022;4(1):44-52. https://doi: 10.21617/jprm2022.417
605 606	2.	WHO. National Guideline for Patient Safety Incident Reporting and Learning in the Public Health Sector of South Africa. World Health Organization; 2017.
607 608	3.	Flott K, Fontana G, Darzi A. The global state of patient safety. London: Imperial College London. 2019.
609 610 611	4.	James JT. A new, evidence-based estimate of patient harms associated with hospital care. Journal of patient safety. 2013;9(3):122-8. https://www.jstor.org/stable/26633011
612 613	5.	Makary MA, Daniel M. Medical error—the third leading cause of death in the US. Bmj. 2016;353. https://www.jstor.org/stable/26942667
614 615	6.	Lt K. To err is human: building a safer health system. Institute of Medicine, Committee on Quality of Health Care in America. 2000.
616 617	7.	National Academies of Sciences E, Medicine. Crossing the global quality chasm: Improving health care worldwide: National Academies Press; 2018.
618 619 620	8.	Choi EY, Pyo J, Ock M, Lee S-i. Nurses' perceptions regarding disclosure of patient safety incidents in Korea: a qualitative study. Asian Nursing Research. 2019;13(3):200-8. https://doi.org/10.1016/j.anr.2019.05.002
621 622 623 624	9.	Lu L, Ko Y-M, Chen H-Y, Chueh J-W, Chen P-Y, Cooper CL. Patient Safety and Staff Well-Being: Organizational Culture as a Resource. International Journal of Environmental Research and Public Health. 2022;19(6):3722. https://doi.org/10.3390/ijerph19063722
625 626 627	10.	Ridelberg M, Roback K, Nilsen P. Facilitators and barriers influencing patient safety in Swedish hospitals: a qualitative study of nurses' perceptions. BMC nursing. 2014;13:1-12. https://doi.org/10.1186/1472-6955-13-23
628 629 630 631	11.	Alves MdFT, Carvalho DSd, Albuquerque GSCd. Barriers to patient safety incident reporting by Brazilian health professionals: an integrative review. Ciencia & saude coletiva. 2019;24:2895-908. https://doi.org/10.1590/1413-81232018248.23912017
632 633 634 635	12.	Archer S, Hull L, Soukup T, Mayer E, Athanasiou T, Sevdalis N, et al. Development of a theoretical framework of factors affecting patient safety incident reporting: a theoretical review of the literature. BMJ open. 2017;7(12):e017155. https://doi: 10.1136/bmjopen-2017-017155

636 637 638	13.	Rashed A, Hamdan M. Physicians' and nurses' perceptions of and attitudes toward incident reporting in Palestinian hospitals. Journal of patient safety. 2019;15(3):212-7. https://doi: 10.1097/PTS.0000000000000218
639 640 641 642	14.	Khoshakhlagh AH, Khatooni E, Akbarzadeh I, Yazdanirad S, Sheidaei A. Analysis of affecting factors on patient safety culture in public and private hospitals in Iran. BMC health services research. 2019;19(1):1-14. https://doi.org/10.1186/s12913-019-4863-x
643 644 645	15.	Gqaleni TM, Bhengu BR. Adverse events reporting system as experienced by critical-care nurses in Kwazulu-Natal, South Africa. Africa Journal of Nursing and Midwifery. 2018;20(1):1-14. https://doi.org/10.25159/2520-5293/2456
646 647	16.	Martos Algarra C. The role of forgiveness in disclosure and victim suport after a patient safety incident. 2022. http://hdl.handle.net/10803/673273
648 649 650 651	17.	Gqaleni T, Mkhize S. Healthcare professionals' perception of knowledge and implementation of Patient Safety Incident Reporting and Learning guidelines in specialised care units, KwaZulu-Natal. Southern African Journal of Critical Care. 2023:25-30. https://doi.org/10.7196/SAJCC.2023.v39i1.559
652 653	18.	WHO. Patient safety tool kit: World Health Organization. Regional Office for the Eastern Mediterranean; 2015.
654 655 656 657	19.	Gqaleni TM, Bhengu BR. Analysis of Patient Safety Incident reporting system as an indicator of quality nursing in critical care units in KwaZulu-Natal, South Africa. Health SA Gesondheid. 2020;25(1):1-8. https://doi.org/10.4102/hsag.v25i0.1263
658 659 660 661	20.	Dhamanti I, Leggat S, Barraclough S, Tjahjono B. Patient safety incident reporting in Indonesia: an analysis using World Health Organization characteristics for successful reporting. Risk management and healthcare policy. 2019:331-8. http://doi.org/10.2147/RMHP.S222262
662 663 664	21.	ELMeneza S, AbuShady M. Anonymous reporting of medical errors from The Egyptian Neonatal Safety Training Network. Pediatrics & Neonatology. 2020;61(1):31-5. https://doi.org/10.1016/j.pedneo.2019.05.008
665 666 667	22.	Yusuf Y, Irwan AM. The influence of nurse leadership style on the culture of patient safety incident reporting: a systematic review. British Journal of Healthcare Management. 2021;27(6):1-7. https://doi.org/10.12968/bjhc.2020.0083
668 669 670	23.	Gong Y. Challenges and Opportunities of Patient Safety Event Reporting. Accident and Emergency Informatics. 2022:133-50. https://doi:10.3233/SHTI220014
671 672	24.	Burns N, Grove SK, Gray JR. Understanding nursing research-eBook: Building an evidence-based practice: Elsevier Health Sciences; 2015.
673 674	25.	Grove SK, Burns N, Gray J. The practice of nursing research: Appraisal, synthesis, and generation of evidence: Elsevier Health Sciences; 2012.

26. Shenton AK. Strategies for ensuring trustworthiness in qualitative research 675 projects. Education for information. 2004;22(2):63-75. DOI: 10.3233/EFI-2004-676 22201 677 Polit, Beck. Nursing research: Generating and assessing evidence for nursing 27. 678 679 practice. Philadelphia: Wolters Kluwer Lippincott Williams & Wilkins; 2017. 28. Dhamanti I, Leggat S, Barraclough S, Liao H-H. Comparison of Patient Safety 680 Incident Reporting Systems in Taiwan, Malaysia, and Indonesia. Journal of patient 681 safety. 2021;17(4):e299-e305. DOI: 10.1097/PTS.00000000000000622 682 29. Martos Algarra C. The role of forgiveness in disclosure and víctima suport after a 683 patient safety incident: Universitat Internacional de Catalunya; 2022. 684 http://hdl.handle.net/10803/673273 685 30. Ontario HQ. Patient safety learning systems: a systematic review and qualitative 686 synthesis. Ontario health technology assessment series. 2017;17(3):1. 687 PMID: 28326148 688 31. Lee Y-C, Wu H-H, Weng S-J, Huang Y-C, Hsieh W-L, Huang C-H. Application of 689 Hospital Information Systems-Construction of an Incident Reporting System. TEM 690 691 Journal. 2016;5(4). 32. Mossburg SE, Himmelfarb CD. The association between professional burnout and 692 engagement with patient safety culture and outcomes: a systematic review. Journal 693 of patient safety. 2021;17(8):e1307-e19. 694 695

696