

1 **Title page**

2

3 **Title**

4 Intronic *FGF14* GAA repeat expansions are a common cause of downbeat nystagmus syndromes: frequency,  
5 phenotypic profile, and 4-aminopyridine treatment response

6

7 **Authors**

8 David Pellerin<sup>1,2</sup>; Felix Heindl<sup>3</sup>; Carlo Wilke<sup>4,5</sup>; Matt C. Danzi<sup>6</sup>; Andreas Traschütz<sup>4,5</sup>; Catherine Ashton<sup>1,7</sup>; Marie-  
9 Josée Dicaire<sup>1</sup>; Alexanne Cuillierier<sup>8</sup>; Giulia Del Gobbo<sup>8</sup>; Kym M. Boycott<sup>8</sup>; Jens Claassen<sup>9,10</sup>; Dan Rujescu<sup>11</sup>;  
10 Annette M. Hartmann<sup>11</sup>; Stephan Zuchner<sup>6</sup>; Bernard Brais<sup>1,12,13</sup>; Michael Strupp<sup>3,\*</sup>; Matthis Synofzik<sup>4,5,\*</sup>

11

12 **Affiliations**

13 <sup>1</sup>Department of Neurology and Neurosurgery, Montreal Neurological Hospital and Institute, McGill University,  
14 Montreal, QC, Canada

15 <sup>2</sup>Department of Neuromuscular Diseases, UCL Queen Square Institute of Neurology and The National Hospital for  
16 Neurology and Neurosurgery, University College London, London, United Kingdom

17 <sup>3</sup>Department of Neurology and German Center for Vertigo and Balance Disorders, University Hospital, Ludwig-  
18 Maximilians University, Munich, Germany

19 <sup>4</sup>Division Translational Genomics of Neurodegenerative Diseases, Hertie-Institute for Clinical Brain Research and  
20 Center of Neurology, University of Tübingen, Tübingen, Germany

21 <sup>5</sup>German Center for Neurodegenerative Diseases (DZNE), Tübingen, Germany

22 <sup>6</sup>Dr. John T. Macdonald Foundation Department of Human Genetics and John P. Hussman Institute for Human  
23 Genomics, University of Miami Miller School of Medicine, Miami, FL, USA

24 <sup>7</sup>Department of Neurology, Royal Perth Hospital, Perth, WA, Australia

25 <sup>8</sup>Children's Hospital of Eastern Ontario Research Institute, University of Ottawa, Ottawa, ON, Canada

26 <sup>9</sup>Department of Neurology, University Hospital Essen, University Duisburg-Essen, Essen, Germany

27 <sup>10</sup>MediClin Klinik Reichshof, Reichshof-Eckenhagen, Germany

28 <sup>11</sup>Department of Psychiatry, Medical University of Vienna, Vienna, Austria

29 <sup>12</sup>Department of Human Genetics, McGill University, Montreal, QC, Canada

30 <sup>13</sup>Centre de Réadaptation Lucie-Bruneau, Montreal, QC, Canada

31 \*Shared last authors

32

33 **Corresponding author**

34 Matthis Synofzik, MD

35 Professor of Neurology

36 Division Translational Genomics of Neurodegenerative Diseases, Hertie-Institute for Clinical Brain Research and

37 Center of Neurology, University of Tübingen

38 Hoppe-Seyler-Str. 3, 72076 Tübingen, Germany

39 Phone: +49-7071-2982060

40 Fax: +49-7071-2925001

41 [matthis.synofzik@uni-tuebingen.de](mailto:matthis.synofzik@uni-tuebingen.de)

42

43 **Running title:** *FGF14* GAA expansions in downbeat nystagmus syndromes

## 44 **Abstract**

45 The cause of downbeat nystagmus (DBN) remains unknown in approximately 30% of patients (idiopathic DBN).  
46 Here, we hypothesized that: (i) *FGF14* (GAA)<sub>≥250</sub> repeat expansions represent a frequent genetic cause of idiopathic  
47 DBN syndromes, (ii) are treatable with 4-aminopyridine (4-AP), and (iii) *FGF14* (GAA)<sub>200-249</sub> alleles are potentially  
48 pathogenic.

49  
50 We conducted a multi-modal cohort study of 170 patients with idiopathic DBN that comprised: in-depth ocular  
51 motor, neurological, and disease evolution phenotyping; assessment of 4-AP treatment response, including re-  
52 analysis of placebo-controlled video-oculography treatment response data from a previous randomized double-blind  
53 4-AP trial; and genotyping of the *FGF14* repeat.

54  
55 Frequency of *FGF14* (GAA)<sub>≥250</sub> expansions was 48% (82/170) in the entire idiopathic DBN cohort. Additional  
56 cerebellar ocular motor signs were observed in 100% (82/82), cerebellar ataxia in 43% (35/82), and extracerebellar  
57 features in 21% (17/82) of (GAA)<sub>≥250</sub>-*FGF14* patients. Alleles of 200 to 249 GAA repeats were enriched in patients  
58 with DBN (12%; 20/170) compared to controls (0.87%; 19/2,191; OR, 15.20; 95% CI, 7.52-30.80;  $p=9.876e-14$ ).

59 The phenotype of (GAA)<sub>200-249</sub>-*FGF14* patients closely mirrored that of (GAA)<sub>≥250</sub>-*FGF14* patients. (GAA)<sub>≥250</sub>-  
60 *FGF14* and (GAA)<sub>200-249</sub>-*FGF14* patients had a significantly greater clinician-reported (80% vs 31%;  $p=0.0011$ ) and  
61 self-reported (59% vs 11%;  $p=0.0003$ ) response rate to 4-AP treatment compared to (GAA)<sub><200</sub>-*FGF14* patients.

62 This included a treatment response with high relevance to everyday living, as exemplified by an improvement of 2  
63 FARS stages in some cases. Placebo-controlled video-oculography data of four (GAA)<sub>≥250</sub>-*FGF14* patients  
64 previously enrolled in a 4-AP randomized double-blind trial showed a significant decrease in slow phase velocity of  
65 DBN with 4-AP, but not placebo.

66  
67 This study shows that *FGF14* GAA repeat expansions are a highly frequent genetic cause of DBN syndromes,  
68 especially when associated with additional cerebellar features. Moreover, they genetically stratify a subgroup of  
69 patients with DBN that appear to be highly responsive to 4-AP, thus paving the way for a “theranostics” approach in  
70 DBN syndromes.

71

72 **Keywords:** SCA27B; natural history; repeat expansion disorder; placebo-controlled trial; theranostics; genome-  
73 wide association study

## 74 **Introduction**

75 Downbeat nystagmus (DBN) is the most common form of acquired persisting nystagmus and is characterized by a  
76 pathological slow upward drift of the eyes followed by a corrective downward saccade.<sup>1,2</sup> In most cases it is caused  
77 by a hypofunction of the cerebellar flocculus/paraflocculus or their projections, which leads to disinhibition of the  
78 superior vestibular nuclei neurons, resulting in spontaneous slow upward eye drift.<sup>3-6</sup> DBN is a common feature of  
79 the allelic disorders spinocerebellar ataxia 6 and episodic ataxia type 2, while it is infrequently observed in other  
80 forms of spinocerebellar ataxia.<sup>7-10</sup> Despite the numerous known genetic and acquired etiologies,<sup>1,2</sup> the cause of  
81 DBN remains unknown in approximately 30% of cases (idiopathic DBN).<sup>2</sup>

82  
83 The recent identification of an association between a variation (rs72665334) in intron 1 of the fibroblast growth  
84 factor 14 (*FGF14*) gene and idiopathic DBN in a genome-wide association study (GWAS)<sup>11</sup> and of a dominantly  
85 inherited (GAA)<sub>≥250</sub> repeat expansion in intron 1 of *FGF14* as the cause of spinocerebellar ataxia 27B (SCA27B) /  
86 GAA-*FGF14* ataxia,<sup>12,13</sup> a late-onset slowly progressive cerebellar syndrome that is frequently associated with  
87 DBN,<sup>12,14</sup> provided evidence for a potential link between *FGF14* and DBN.<sup>12,14,15</sup> Furthermore, 4-aminopyridine (4-  
88 AP), a proven effective symptomatic treatment for DBN,<sup>16,17</sup> has shown promising benefits to reduce the frequency  
89 and severity of ataxic symptoms in previous small series of patients with GAA-*FGF14* ataxia.<sup>12,14,18</sup>

90  
91 Based on these findings, we hypothesized that *FGF14* (GAA)<sub>≥250</sub> repeat expansions are a frequent monogenic cause  
92 of idiopathic DBN syndromes, which are treatable with 4-AP, and are in disequilibrium with the genome-wide  
93 significant rs72665334 variant previously identified by GWAS. Moreover, we systematically studied the phenotype  
94 and 4-AP treatment response of patients with DBN carrying an *FGF14* (GAA)<sub>200-249</sub> allele, providing preliminary  
95 evidence that these so-called “intermediate” alleles might be pathogenic.

## 96 97 **Methods**

### 98 **Patient Enrollment**

99 We enrolled a consecutive series of 219 index patients with suspected DBN referred to the Department of Neurology  
100 or the German Center for Vertigo and Balance Disorders at the LMU Hospital in Munich, Germany, between 2012  
101 and 2020. Etiologic evaluation of DBN was performed through a detailed medical history, assessment of drug abuse,

102 comprehensive neurological examination, laboratory tests, and brain imaging by MRI or CT scan. Genetic screening  
103 for episodic ataxias and spinocerebellar ataxias was performed in select patients with a suggestive phenotype when  
104 deemed appropriate by the clinician. Patients were excluded from the study if: (i) no DNA was available for genetic  
105 screening ( $n=2$ ), (ii) DBN was not objectified on neuro-ophthalmological examination after review of the medical  
106 records ( $n=11$ ), or (iii) a competing etiology for DBN was identified ( $n=36$ ). The final study cohort comprised 170  
107 patients with a diagnosis of idiopathic DBN (**Fig. 1**). Part of this patient cohort ( $n=80$ ) has previously been reported  
108 in the idiopathic DBN GWAS.<sup>11</sup> All but three patients (one  $[GAA]_{\geq 250}$ -*FGF14* and two  $[GAA]_{<200}$ -*FGF14*) were of  
109 self-reported European descent. The three non-European patients were of Turkish descent. This study was approved  
110 by the ethics committees of the LMU Munich, the Montreal Neurological Hospital-Institute, and Clinical Trials  
111 Ontario and was carried out in accordance with the Declaration of Helsinki. We obtained written informed consent  
112 from all the participants in this study.

113

#### 114 **Deep Phenotyping**

115 All patients underwent at least one comprehensive neurological and ocular motor examination. Deep phenotyping  
116 was performed by systematically reassessing all medical records using a standardized data sheet while being blind to  
117 the *GAA-FGF14* genotype. Functional impairment was assessed in terms of the need for mobility aids and by means  
118 of the Friedreich Ataxia Rating Scale (FARS) functional stage (0=normal; 1=minimal signs on examination;  
119 2=minimal disability; 3=mild disability; 4=moderate disability, requires a walker; 5=severe disability, confined but  
120 can navigate a wheelchair; 6=total disability).<sup>19</sup> Results of routine brain MRI and nerve conduction studies were  
121 available for review in 70% (119/170) and 18% (31/170) of patients, respectively. Ancillary assessment of vestibular  
122 function was obtained in 94% (159/170) of patients using bithermal caloric stimulation ( $n=147$ ), video head impulse  
123 test (vHIT;  $n=100$ ), and rotatory chair test ( $n=1$ ). Posturography (Kistler platform, Kistler Instrumente AG) was  
124 obtained in 59% (100/170) of patients and results were analyzed with an artificial neural network.<sup>20</sup>

125

126 Patients were stratified in one of four phenotypic clusters based on the presence of additional cerebellar and/or  
127 extracerebellar signs: pure DBN, DBN plus additional isolated cerebellar ocular motor signs (DBN+COM), DBN  
128 plus cerebellar ataxia (DBN+CA), and DBN plus cerebellar and extracerebellar features (DBN+EC), which included  
129 bilateral vestibulopathy (BVP) and/or polyneuropathy. Along with the cerebellar system, the vestibular and sensory

130 systems appear to be preferentially impaired in *GAA-FGF14* ataxia.<sup>14,15</sup> Additional cerebellar ocular motor signs  
131 were defined by the presence of at least one of: saccadic pursuit, dysmetric saccades, gaze-evoked nystagmus,  
132 rebound nystagmus, or impaired visual fixation suppression of the vestibulo-ocular reflex (VOR). Cerebellar ataxia  
133 was defined by the presence of cerebellar dysarthria, dysdiadochokinesia, intention tremor, or, when available,  
134 evidence of cerebellar involvement on posturography. BVP was diagnosed as per the consensus criteria of the  
135 Bárány Society requiring the documentation of bilaterally reduced or absent angular VOR function by caloric  
136 stimulation, vHIT, or rotatory chair.<sup>21</sup> Polyneuropathy was diagnosed on nerve conduction studies (excluding focal  
137 entrapment neuropathies) or clinically defined by the combination of significantly decreased vibration sense at the  
138 ankles ( $\leq 3/8$  on the Rydel-Seiffer scale) and decreased or absent ankle reflexes.<sup>22</sup>

139

#### 140 **Assessment of Treatment Response to 4-aminopyridine**

141 Assessment of treatment response to 4-AP was performed by two independent approaches. (1) *Open-label real-*  
142 *world treatment response data.* Information on clinician-reported and patient-reported response to 4-AP treatment  
143 (fampridine 10mg twice a day) was collected (available for 83 patients). As all patients from this study who were  
144 treated with 4-AP received the drug before the discovery of *GAA-FGF14* ataxia, the assessment of treatment  
145 response was naturally blind to the underlying *GAA-FGF14* genotype. Patients were routinely evaluated 1 to 3  
146 months after initiating treatment with 4-AP. Patient-reported responses were assessed based on their global  
147 impressions of the impact of treatment on their neurological functioning. Clinician-reported responses were  
148 evaluated based on the clinician's global impression from neurological and neuro-ophthalmological examinations.  
149 (2) *Double-blind placebo-controlled clinical trial data.* Four patients with DBN had been part of an earlier placebo-  
150 controlled randomized double-blind trial assessing the efficacy of 4-AP in a cohort of 27 patients with DBN.<sup>16</sup> All  
151 four patients, now known via the current study to carry an *FGF14* (*GAA*) <sub>$\geq 250$</sub>  expansion, were genetically  
152 undiagnosed at the time of the original trial.<sup>16</sup> Due to unavailability of DNA, the genetic status of the remaining 23  
153 patients who took part in the 4-AP trial remains unknown. As part of the trial, patients had received 5 mg of 4-AP  
154 (or placebo) four times a day for 3 days and 10 mg of 4-AP (or placebo) four times a day for 4 days. Randomization  
155 resulted in all four patients receiving active treatment first, then placebo, separated by a 1-week washout.  
156 Assessments were done before the first, 60 min after the first, and 60 min after the last drug administration,  
157 measuring the slow phase velocity (SPV) of DBN by video-oculography (degrees/second) as primary outcome.

158

### 159 **Genetic Screening for *FGF14* Repeat Expansions**

160 The *FGF14* repeat locus was genotyped by capillary electrophoresis of fluorescent long-range PCR amplification  
161 products and bidirectional repeat-primed PCRs, as described previously.<sup>23</sup> Expansions of at least 250 GAA repeat  
162 units were considered pathogenic.<sup>12,13</sup> Alleles of 200 to 249 GAA repeat units were analyzed separately as their  
163 pathogenicity has recently been suggested.<sup>24</sup>

164

### 165 **Genotyping of the *FGF14* rs72665334 Variant**

166 The rs72665334 C>T variant in intron 1 of *FGF14* (GRCh38, chr13:102,150,076) was found to be associated with  
167 idiopathic DBN in 106 patients in a recent GWAS.<sup>11</sup> Genotyping data were previously generated using the  
168 HumanOmniExpress-24 array and data for the rs72665334 variant with call rate probability >70% were extracted for  
169 73 patients enrolled in this study.

170

171 To assess whether the rs72665334 variant is in disequilibrium with the *FGF14* GAA repeat expansion at a  
172 population level, we also genotyped the rs72665334 variant by Sanger sequencing in an independent and ethnically  
173 distinct cohort of 37 French-Canadian index patients with GAA-*FGF14* ataxia and DBN. PCR reactions were  
174 performed in a 24μL volume using the Qiagen Taq DNA polymerase kit (catalog no. 201209, Qiagen) with  
175 0.125mM dNTPs, 1μM of forward and reverse primers (forward primer: 5'-GCCCCTGTTCTAAAGCCTCT-3';  
176 reverse primer: 5'-GATCGTCCAGCCACATCTCT-3'), and 240ng of genomic DNA. Sanger sequencing of PCR  
177 amplification products was performed using the ABI 3730xl Analyzer (Applied Biosystems).

178

### 179 **Statistical Analysis**

180 We assessed differences between groups with the non-parametric Mann-Whitney U test for continuous variables and  
181 the Fisher's exact test for categorical variables. Effect sizes are reported as odds ratio (OR) with 95% confidence  
182 intervals (CIs). The non-parametric Kruskal-Wallis test was used for comparisons between multiple groups.  
183 Correlations were calculated using the Pearson's correlation coefficient or the Spearman's rank correlation  
184 coefficient. The Kaplan-Meier method was used to analyze disease-free survival. Cross-sectional FARS stages  
185 measured off 4-AP were analyzed by linear regression as a function of disease duration, age, and the expansion size.



186 To estimate the longitudinal progression of disability over the disease course, we used a linear mixed-effects model  
187 fitted by the restricted maximum likelihood method accounting for disease duration as fixed effect and with random  
188 intercepts and random slopes. We re-analyzed treatment effects of 4-AP and placebo by means of a mixed-effect  
189 analysis of repeated measures, with post-hoc comparison by uncorrected Fisher's least significant difference. We  
190 analyzed the data in R (version 4.3) and GraphPad Prism 9.3. *P* value of <0.05 was considered significant, using the  
191 Benjamini-Hochberg method to correct for multiple comparisons. All analyses were two-sided.

192

### 193 **Data Availability**

194 Individual deidentified patient data may be shared at the request of any qualified investigator upon reasonable  
195 request. No consent for open sharing has been obtained.

196

## 197 **Results**

### 198 **Frequency of *FGF14* GAA Repeat Expansions in Downbeat Nystagmus Syndromes**

199 A total of 170 index patients with a diagnosis of idiopathic DBN formed the final study cohort and were screened  
200 for *FGF14* GAA repeat expansions (**Figs. 1 and 2A**). We identified 82 patients (48%) who carried an *FGF14*  
201 (GAA)<sub>≥250</sub> expansion (median size of expansion, 324 repeat units; range, 250-516), including one patient carrying  
202 biallelic GAA expansions of 280 and 313 repeat units. Alleles of 200 to 249 repeat units were identified in 12% of  
203 patients (20/170; median size of allele, 234 repeat units; range, 207-249), compared to 0.87% in 2,191 previously  
204 reported control individuals<sup>25</sup> (19/2,191), suggesting an enrichment of this population of alleles in patients with  
205 DBN (OR, 15.20; 95% CI, 7.52-30.80; Fisher's exact test, *p*=9.876e-14). Two patients carried a likely non-  
206 pathogenic<sup>24,26</sup> (GAAGGA)<sub>n</sub> expansion of 319 and 335 triplet repeat units equivalent, respectively.

207

208 The frequency of *FGF14* (GAA)<sub>≥250</sub> expansions in DBN syndromes stratified by phenotype was 56% (42/75) for  
209 DBN plus additional isolated cerebellar ocular motor signs, 72% (23/32) for DBN plus cerebellar ataxia, 30%  
210 (17/56) for DBN plus cerebellar ocular motor signs and/or ataxia and extracerebellar features, which included BVP  
211 and/or polyneuropathy, and 0% (0/7) for pure DBN (**Fig. 2B**).

212

### 213 **Phenotype and Discriminative Features of (GAA)<sub>≥250</sub>-*FGF14*-related Downbeat Nystagmus Syndromes**

214 Table 1 summarizes the baseline characteristics and clinical features of patients carrying a (GAA)<sub>≥250</sub> expansion, a  
215 (GAA)<sub>200-249</sub> allele, and a (GAA)<sub><200</sub> allele.

216  
217 Symptoms in (GAA)<sub>≥250</sub>-*FGF14* patients with DBN started at a median age of 67 years (range, 30-81). The most  
218 common presenting symptom was gait unsteadiness (89%; 72/81), followed by visual disturbances (37%; 30/81),  
219 such as diplopia, oscillopsia, or visual blurring, vertigo and/or dizziness (20%; 16/81), and, rarely, dysarthria (1%;  
220 1/81) (**Fig. 2C**). We observed a weak inverse correlation between the age at onset and the size of the repeat  
221 expansion (74 patients; Pearson's  $r$ , -0.30;  $R^2=0.09$ ;  $p=0.01$ ) (**Fig. 2D**). Family history for DBN or ataxia was  
222 positive in 31 patients (38%), of whom 26 had evidence of autosomal dominant inheritance. In keeping with the  
223 likely reduced male transmission of the disease due to contraction of the *FGF14* repeat locus upon paternal  
224 transmission,<sup>12,25</sup> we observed that only 2 of 26 dominantly inherited cases (8%) were paternally inherited.

225  
226 The median age and disease duration at last examination were 73 years (range, 40-92) and 6 years (range, 0-26.5),  
227 respectively. Additional cerebellar ocular motor signs were observed in 100% (82/82), cerebellar ataxia in 43%  
228 (35/82), and extracerebellar features in 21% (17/82) of (GAA)<sub>≥250</sub>-*FGF14* patients (**Fig. 2E**). Cerebellar ocular  
229 motor signs comprised – in decreasing order of frequency – saccadic pursuit (99%; 80/81), impaired visual fixation  
230 suppression of the VOR (86%; 64/74 in patients without BVP), gaze-evoked nystagmus (70%; 57/82), dysmetric  
231 saccades (28%; 22/79), and rebound nystagmus (20%; 16/82). Signs of cerebellar ataxia included dysmetria of upper  
232 limbs (46%; 16/35), dysdiadochokinesia (47%; 15/32), dysarthria (35%; 12/34), and intention tremor (11%; 4/35).  
233 Cerebellar involvement reflected by ~3Hz titubation was identified in 18 of 55 patients (33%) on posturography.  
234 Brain MRI of eight patients showed cerebellar atrophy (13%; 8/60), which was limited to the vermis in three  
235 patients and extended to the hemispheres in five patients. Two patients with discrete pan-cerebellar atrophy showed  
236 no signs of ataxia on examination.

237  
238 The subgroup of (GAA)<sub>≥250</sub>-*FGF14* patients with DBN plus cerebellar ocular motor signs and/or ataxia and  
239 extracerebellar features comprised nine patients with polyneuropathy, six patients with BVP, and two patients with  
240 polyneuropathy plus BVP (**Fig. 2E**). Polyneuropathy and BVP occurred more frequently in patients with cerebellar  
241 ataxia (71%; 12/17) compared to patients exhibiting only cerebellar ocular motor signs (29%; 5/17) (**Fig. 2E**).

242 Polyneuropathy was diagnosed on nerve conduction studies in seven patients and clinically in four patients. Results  
243 of nerve conduction studies were consistent with length-dependent sensorimotor axonal neuropathy in six patients  
244 and sensory neuropathy in one patient. BVP was diagnosed in 8 of 79 patients (10%) by caloric stimulation ( $n=1$ )  
245 and vHIT ( $n=7$ ). Disease duration at time of vestibular assessment differed significantly between patients with and  
246 without BVP (median, 14 vs 4 years; Mann-Whitney U test,  $p=0.0018$ ).

247  
248 To identify discriminative features of DBN syndromes in  $(GAA)_{\geq 250}$ -*FGF14* patients, we compared their phenotypic  
249 features to that of  $(GAA)_{< 200}$ -*FGF14* patients (**Table 1**). Both groups did not differ significantly in terms of disease  
250 duration, baseline characteristics, functional impairment, and symptoms. Impairment of the visual fixation  
251 suppression of the VOR was strongly associated with  $(GAA)_{\geq 250}$ -*FGF14* DBN syndromes (86% vs 50%; OR, 6.29;  
252 95% CI, 2.45-17.24; Fisher's exact test,  $p=2.089e-05$ ,  $q=6.685e-04$ ). In keeping with the lower frequency of *FGF14*  
253 repeat expansions in patients with DBN plus extracerebellar features, impaired vibration at the ankles (14% vs 37%;  
254 OR, 0.28; 95% CI, 0.11-0.66; Fisher's exact test,  $p=0.002$ ,  $q=0.018$ ) and BVP (10% vs 32%; OR, 0.24; 95% CI,  
255 0.08-0.63; Fisher's exact test,  $p=0.001$ ,  $q=0.018$ ) were significantly less common in  $(GAA)_{\geq 250}$ -*FGF14* than in  
256  $(GAA)_{< 200}$ -*FGF14* patients. Furthermore, in the subset of patients in whom BVP was objectified on vHIT, the mean  
257 VOR gain was significantly higher (values of  $>0.6$  are considered normal) in  $(GAA)_{\geq 250}$ -*FGF14* compared to  
258  $(GAA)_{< 200}$ -*FGF14* patients ( $0.44 \pm 0.10$  vs  $0.18 \pm 0.17$ ; Mann-Whitney U test,  $p=0.006$ ,  $q=0.0496$ ), despite no  
259 significant difference in disease duration between the groups (median, 14.5 vs 8 years; Mann-Whitney U test,  
260  $p=0.1512$ ). These results suggest that BVP occurs late in the course of  $(GAA)_{\geq 250}$ -*FGF14*-related DBN syndromes  
261 and remains mild.

262  
263 **Disease Progression in  $(GAA)_{\geq 250}$ -*FGF14*-related Downbeat Nystagmus Syndromes**  
264 We used a Kaplan-Meier survival analysis to model the age at disease onset of  $(GAA)_{\geq 250}$ -*FGF14* patients,  
265 estimating their probability of remaining disease-free over time (**Fig. 3A**). Survival until disease onset in  $(GAA)_{\geq 250}$ -  
266 *FGF14* patients did not significantly differ between the three main phenotypic subgroups (median age at onset, 67  
267 years [33-80] for DBN+COM vs 67 years [36-81] for DBN+CA vs 66 years [30-78] for DBN+EC; Cox-Mantel test,  
268  $X^2(2)=0.1267$ ;  $p=0.939$ ). However, disease duration at last examination was significantly longer in patients with  
269 DBN plus cerebellar and extracerebellar features (median, 4.25 years [0-16] for DBN+COM vs 6 years [1-22] for

270 DBN+CA vs 9 years [3-26.5] for DBN+EC; Kruskal-Wallis test,  $p=0.014$ ) (**Fig. 3B**), suggesting that extracerebellar  
271 features may develop later in the disease course.

272  
273 We also studied the evolution of key symptoms in  $(GAA)_{\geq 250}$ -*FGF14* patients in relation to gait impairment (**Fig.**  
274 **3C**). Episodic symptoms (median, 0 year; range, -6-5) and oscillopsia (0.75 year; -8-5) manifested concurrently with  
275 gait impairment in most patients. However, episodic symptoms and oscillopsia preceded gait impairment in 10%  
276 (1/10) and 11% (3/28) of patients for whom information on age at onset of these features was available, respectively.  
277 In comparison, fine motor impairment (5 years; 0-8), speech impairment (4 years; 0-10), and falls (2 years; 0-12)  
278 developed later in the disease course, consistent with the findings from a recent progression study.<sup>14</sup> Sixteen patients  
279 (20%) eventually required a walking aid after a median disease duration of 4 years (range, 0-13), including three  
280 patients (4%) who became wheelchair dependent after a median disease duration of 8 years (range, 7-12). Further to  
281 this analysis, we observed that BVP developed at a significantly later stage in the disease compared to cerebellar  
282 atrophy and neuropathy (median disease duration at first occurrence, 4 years [1-7] for cerebellar atrophy vs 3 years  
283 [1-6] for polyneuropathy vs 10 years [3-21.5] for BVP; Kruskal-Wallis test,  $p=0.004$ ) (**Fig. 3D**).

284  
285 The progression of functional disability in  $(GAA)_{\geq 250}$ -*FGF14* patients was assessed by means of the FARS  
286 functional staging (**Fig. 4**). At time of last examination, the median FARS stage measured off 4-AP was 3 (range, 2-  
287 5), indicating a mild level of disability. We found no association between the FARS functional stage and disease  
288 duration (73 patients; Spearman's rho, 0.194;  $p=0.10$ ) (**Fig. 4A**). Neither age, disease duration, or expansion size  
289 was associated with the cross-sectional FARS stage in a multiple linear regression model ( $F(3,69)=2.602$ , adjusted  
290  $R^2=0.06$ ,  $p=0.059$ ; age,  $b=0.017$ ,  $p=0.111$ ; disease duration,  $b=0.027$ ,  $p=0.151$ ; expansion size,  $b=-0.001$ ,  $p=0.527$ ).  
291 Analysis of the cross-sectional and longitudinal FARS stage data revealed that half of patients (27/54) displayed at  
292 least mild disability within the first five years of disease onset. However, disability remained at most mild in the  
293 majority of patients (86%, 63/73) even after more than 25 years of disease duration. This observation was further  
294 supported by longitudinal data in 40 patients (148 observations) showing an overall slow intra-individual increase of  
295 0.10 FARS stage per year of disease (standard error, 0.02 by linear mixed-effects model) (**Fig. 4B**). While disease  
296 evolved relatively slowly in most patients, inspection of individual disability progression trajectories revealed inter-  
297 individual variability in rates of progression (**Fig. 4B**). Concurrent medical conditions substantially contributed to

298 disability burden in two patients. One patient became wheelchair-bound following a complicated medical admission.  
299 However, this patient also carried biallelic expansions (280 and 313 repeat units) which may have contributed to the  
300 faster disability progression. The second patient, carrying alleles of 218 and 270 repeat units, became wheelchair-  
301 bound following a hip fracture.

302

303 **(GAA)<sub>200-249</sub>-FGF14 Patients have a Similar Clinical Phenotype and Disease Evolution as (GAA)<sub>≥250</sub>-FGF14**

304 **Patients**

305 While previous studies have established a preliminary pathogenic threshold of at least 250 GAA repeat units,<sup>12,13</sup> the  
306 pathogenic potential of (GAA)<sub>200-249</sub> alleles has recently been suggested following the identification of a family with  
307 late-onset slowly progressive ataxia in which (GAA)<sub>200-249</sub> alleles and (GAA)<sub>≥250</sub> expanded alleles segregated with  
308 disease.<sup>24</sup> Alleles of 200 to 249 GAA repeat units were significantly enriched in patients with DBN compared to  
309 controls (12% vs 0.87%; OR, 15.20; 95% CI, 7.52-30.80; Fisher's exact test,  $p=9.876e-14$ ), thus providing further  
310 support for their potential pathogenicity.

311

312 We next aimed to characterize the phenotype of (GAA)<sub>200-249</sub>-FGF14 patients to assess whether it resembled that of  
313 (GAA)<sub>≥250</sub>-FGF14 patients (**Table 1**). (GAA)<sub>200-249</sub>-FGF14 patients did not significantly differ from (GAA)<sub>≥250</sub>-  
314 FGF14 patients in terms of baseline characteristics, symptoms, clinical signs, and progression of functional  
315 disability. In comparison, impairment of the visual fixation suppression of the VOR (indicating an involvement of  
316 the fixation system), which was strongly associated with (GAA)<sub>≥250</sub>-FGF14-related DBN syndromes, was  
317 significantly more common in (GAA)<sub>200-249</sub>-FGF14 patients compared to (GAA)<sub><200</sub>-FGF14 patients (94%, 16/17 vs  
318 50%, 23/46; OR, 15.45; 95% CI, 2.07-696.97; Fisher's exact test,  $p=0.001$ ;  $q=0.037$ ), but not different from  
319 (GAA)<sub>≥250</sub>-FGF14 patients (94%, 16/17 vs 86%, 64/74; Fisher's exact test,  $p=0.682$ ;  $q=1.000$ ).

320

321 **Association of the rs72665334 Variant with FGF14 (GAA)<sub>≥250</sub> Expansions in Patients with Downbeat**

322 **Nystagmus**

323 A recent GWAS has identified an association between DBN and the FGF14 rs72665334 C>T variant in 106 patients  
324 (**Fig. 5A**).<sup>11</sup> This same variant, which has an allele frequency of ~7% in the European population (1000 Genomes  
325 Project<sup>27</sup>), was also found to be part of a disease haplotype shared by three Australian patients with GAA-FGF14

326 ataxia in a previous study.<sup>13</sup> Given the frequent occurrence of DBN in GAA-*FGF14* ataxia, we therefore  
327 hypothesized that the rs72665334 variant might be in disequilibrium with the *FGF14* GAA repeat expansion, which  
328 is located ~11kb away (**Fig. 5A**). In the DBN cohort, we observed that a significantly greater proportion of patients  
329 with a C|T or T|T genotype carried an *FGF14* (GAA)<sub>≥250</sub> expansion compared to an *FGF14* (GAA)<sub><200</sub> allele (54%,  
330 19/35 vs 14%, 4/29; OR, 7.18; 95% CI, 1.91 to 34.43; Fisher's exact test  $p=0.0014$ ) (**Fig. 5B**). Similarly, we also  
331 found that 38% of patients (14/37) from an independent and ethnically distinct cohort of 37 French-Canadian index  
332 patients with GAA-*FGF14* ataxia and DBN had a C|T or T|T rs72665334 genotype. To assess whether the  
333 rs72665334 variant is in disequilibrium with the *FGF14* (GAA)<sub>≥250</sub> expansion at a population level, we compared  
334 the frequency of the C|T and T|T genotypes in 503 European controls from the 1000 Genomes Project and 72  
335 patients with GAA-*FGF14* ataxia and DBN (35 patients from this study and 37 French-Canadian patients). The C|T  
336 and T|T genotypes were significantly more frequent in patients with DBN carrying an *FGF14* (GAA)<sub>≥250</sub> expansion  
337 compared to controls (46%, 33/72 vs 12%, 60/503; OR, 6.22; 95% CI, 3.51 to 11.02; Fisher's exact test,  $p=1.063e-$   
338 10) (**Fig. 5C**). Together, these results show that the rs72665334 variant is in disequilibrium with the *FGF14* GAA  
339 repeat expansion. However, its absence in 54% of (GAA)<sub>≥250</sub>-*FGF14* patients suggests that *FGF14* GAA repeat  
340 expansions may arise on distinct haplotype backgrounds.

341

#### 342 **Treatment Response to 4-Aminopyridine**

343 *Open-label real-world treatment response data.* We assessed the response to 4-AP treatment in patients with DBN  
344 stratified by *FGF14* genotype (open-label treatment as part of routine clinical care). 81% of (GAA)<sub>≥250</sub>-*FGF14*  
345 patients (66/81), 61% of (GAA)<sub>200-249</sub>-*FGF14* patients (11/18), and 56% of (GAA)<sub><200</sub>-*FGF14* patients (37/66) had  
346 previously received 4-AP. Treatment response for all patients had been recorded before the discovery of GAA-  
347 *FGF14* ataxia and, therefore, both clinicians and patients were naturally blind to the underlying GAA-*FGF14*  
348 genotype ("real-world double-blind" to genotype). A clinician-reported treatment response was recorded for 81% of  
349 (GAA)<sub>≥250</sub>-*FGF14* patients (29/36; improvement of ocular motor signs, including DBN: 24 patients; and gait: 14  
350 patients), 80% of (GAA)<sub>200-249</sub>-*FGF14* (4/5; ocular motor signs: 2; gait: 3), and 31% of (GAA)<sub><200</sub>-*FGF14* patients  
351 (5/16; ocular motor signs: 3; gait: 2) (**Fig. 6A**). A patient-reported benefit was recorded for 59% of (GAA)<sub>≥250</sub>-  
352 *FGF14* (29/49; improvement of visual symptoms: 14 patients; and gait: 17 patients), 60% of (GAA)<sub>200-249</sub>-*FGF14*  
353 (3/5; visual symptoms: 2; gait: 2), and 11% of (GAA)<sub><200</sub>-*FGF14* patients (2/19; visual symptoms: 1; gait: 1) (**Fig.**

354 **6B**). ( $GAA_{\geq 250}$ -*FGF14* and ( $GAA_{200-249}$ -*FGF14* patients had a significantly greater clinician-reported (80% vs  
355 31%; OR, 8.63; 95% CI, 2.08 to 41.96; Fisher's exact test  $p=0.0011$ ) and patient-reported (59% vs 11%; OR, 11.96;  
356 95% CI, 2.45 to 117.27; Fisher's exact test  $p=0.0003$ ) response rate to 4-AP treatment compared to ( $GAA_{<200}$ -  
357 *FGF14* patients. To next evaluate the effect of 4-AP on disability, we examined the FARS disability stages assessed  
358 on and off 4-AP. Data were available for seven ( $GAA_{\geq 250}$ -*FGF14* and two ( $GAA_{200-249}$ -*FGF14* patients (**Fig. 7**).  
359 Remarkably, the FARS stages for all nine patients were invariably lower, by up to 2 stages, while on 4-AP  
360 compared to off 4-AP. The therapeutic benefit of 4-AP was further evidenced by the clear association between drug  
361 intake and improvement of disability in four patients in whom stopping the drug led to an increase in disability (**Fig.**  
362 **7**). Given the average progression of 0.10 FARS stage per year of disease (**Fig. 4B**), an improvement of 1 FARS  
363 stage on 4-AP treatment might be considered equivalent to "reversing" the disease progression by approximately 10  
364 years. These results, which need confirmation in randomized placebo-controlled trials, suggest that the *GAA-FGF14*  
365 genotype status stratifies a subgroup of patients with DBN who are high responders to 4-AP. The response to 4-AP,  
366 whether reported by the clinician or the patient, may also be a useful clinical sign to increase suspicion for *GAA-*  
367 *FGF14* disease, given its positive predictive value of 88% (95% CI, 74-96%) for ( $GAA_{\geq 250}$ -*FGF14* status and 89%  
368 (95% CI, 76-96%) for combined ( $GAA_{\geq 250}$ -*FGF14* and ( $GAA_{200-249}$ -*FGF14* statuses.

369  
370 *Double-blind placebo-controlled clinical trial data.* We next analyzed in a first preliminary probing approach  
371 whether the effects of 4-AP treatment in *GAA-FGF14*-related DBN could be validated not only in open-label  
372 assessments, but also in double-blind, placebo-controlled randomized treatment assessments. For this purpose, we  
373 re-analyzed placebo-controlled video-oculography data of four patients with DBN – now known to be ( $GAA_{\geq 250}$ -  
374 *FGF14* – from a previous randomized double-blind trial.<sup>16</sup> In addition to being double-blind, all trial assessments  
375 were blinded to the underlying *GAA-FGF14* genotype, which was not discovered until after the trial. The absolute  
376 value of the SPV decreased under treatment with 4-AP ( $F(2,6) = 6.31, p=0.033$ ), but not under placebo ( $F(2,5) =$   
377  $0.45, p=0.660$ ) (**Fig. 6C**). The decrease in SPV was significant 60 min after administration of a single dose of 5 mg  
378 of 4-AP compared to intra-individual baseline measurements (least squares mean difference: 2.63°/s,  $p=0.014$ ), and  
379 persisted under a stable dose of 40 mg of 4-AP daily (least squares mean difference: 1.97°/s,  $p=0.043$ ). This effect  
380 corresponds to relative reductions (improvements) of SPV by 38-109% after a single dose, and 32-99% under stable  
381 treatment with 4-AP (**Fig. 6D**).



382

## 383 Discussion

384 This study tested the hypothesis that *FGF14* (GAA)<sub>≥250</sub> repeat expansions are a recurrent cause of idiopathic DBN  
385 syndromes, where DBN may be the presenting phenotype of a wider GAA-*FGF14*-related neurological disease  
386 spectrum. Our findings identify *FGF14* (GAA)<sub>≥250</sub> repeat expansions as a highly frequent monogenic cause of DBN,  
387 particularly when associated with additional cerebellar features. Through systematic in-depth phenotyping, we  
388 characterized the disease evolution of (GAA)<sub>≥250</sub>-*FGF14*-related DBN and its treatment response to 4-AP.  
389 Additionally, we explored the phenotypic and 4-AP treatment response characteristics of patients with *FGF14*  
390 (GAA)<sub>200-249</sub> alleles, whose pathogenicity has recently been suggested.

391

392 The cause of DBN remains unknown in approximately 30% of cases (idiopathic DBN).<sup>1,2</sup> Identifying genetic causes  
393 for these cases has proven challenging. We now show that *FGF14* (GAA)<sub>≥250</sub> repeat expansions account for almost  
394 50% of cases of previously unexplained DBN in a large and strictly consecutive cohort of European patients.  
395 Dominantly inherited *FGF14* GAA repeat expansions have recently been shown to be a common cause of  
396 spinocerebellar ataxia (SCA27B), which is associated with DBN in up to 40-60% of patients.<sup>12,14</sup> However, the high  
397 frequency of *FGF14* (GAA)<sub>≥250</sub> repeat expansions in patients with a predominantly sporadic non-ataxic DBN  
398 presentation was unexpected. One explanation may be that DBN is a milder phenotypic presentation of GAA-  
399 *FGF14* disease, in which overt cerebellar ataxia and other multisystemic involvement can be absent (or, in some  
400 cases, develop later in the disease course). Our results further suggest that genetic testing for *FGF14* GAA repeat  
401 expansion should now become part of the diagnostic work-up of patients with idiopathic DBN.

402

403 By screening a cohort of patients with idiopathic DBN, our study closes a gap in the delineation of the phenotypic  
404 spectrum and evolution of GAA-*FGF14* disease. The identification of additional floccular/parafloccular cerebellar  
405 ocular motor signs in all (GAA)<sub>≥250</sub>-*FGF14* patients with DBN indicates that the basic dysfunction arises from this  
406 cerebellar region and that pure DBN is an uncommon manifestation in GAA-*FGF14* disease. Furthermore, the  
407 observation of DBN and ocular motor signs manifesting up to 8 years prior to the development of gait impairment  
408 suggests that such signs may present in isolation early in the disease course. However, the recurrent identification of  
409 *FGF14* expansions in patients with DBN plus additional isolated cerebellar ocular motor signs but without overt



410 ataxia despite disease duration of up to 16 years raises the possibility that *GAA-FGF14* disease may remain limited  
411 to the cerebellar ocular motor system without broader cerebellar involvement in a subset of patients. This indicates  
412 that cerebellar ataxia is not a universal and mandatory feature of *GAA-FGF14* disease and that ocular motor signs  
413 and ataxia may each represent different features along a continuum of variable cerebellar involvement. Future  
414 natural history studies will be needed to determine whether cerebellar ataxia eventually develops in all patients.  
415 Nevertheless, the identification of a more limited phenotype in a sizeable number of patients suggests that the  
416 overall frequency of *GAA-FGF14* disease may even be higher than previously estimated.

417  
418 While BVP has been suggested to be a recurrent feature of *GAA-FGF14* disease,<sup>15</sup> our study is the first to provide  
419 an estimate of the frequency and temporal evolution of BVP in a large and phenotypically unselected cohort of  
420 patients with *GAA-FGF14* disease. Ancillary vestibular assessment performed in more than 95% of  $(GAA)_{\geq 250}$ -  
421 *FGF14* patients documented BVP in 10% of them. BVP appears to be a late feature in *GAA-FGF14* disease  
422 compared to cerebellar ataxia, as it developed on average more than 10 years after disease onset. In addition to being  
423 significantly less frequent in  $(GAA)_{\geq 250}$ -*FGF14* compared to  $(GAA)_{< 200}$ -*FGF14* DBN syndromes, BVP tended to  
424 remain relatively mild in *GAA-FGF14*-related DBN syndromes despite prolonged disease duration. The origin of  
425 the vestibular pathology in *GAA-FGF14* disease is yet to be established, although a central etiology may be more  
426 likely given the absence of significant involvement of the peripheral nervous system in this disease.

427  
428 While previous studies of patients with SCA27B have shown rates of cerebellar atrophy ranging from 60 to  
429 100%,<sup>12,14</sup> only 13% of  $(GAA)_{\geq 250}$ -*FGF14* patients with DBN exhibited cerebellar atrophy (mainly of the vermis).  
430 This finding might stem from our cohort's main inclusion criterion being DBN, not cerebellar ataxia, resulting in  
431 over half of the patients displaying no overt ataxia. Moreover, the median disease duration at time of brain MRI in  
432 our DBN cohort was relatively short (4.5 years). Mild cerebellar vermis atrophy can also be challenging to identify  
433 on routine MRI and may have been missed.

434  
435 Leveraging in-depth phenotyping data, our study identified clinical features discriminating  $(GAA)_{\geq 250}$ -*FGF14* from  
436  $(GAA)_{< 200}$ -*FGF14* DBN syndromes. Of all clinical signs, impairment of the visual fixation suppression of the VOR  
437 was most strongly associated with and predictive of  $(GAA)_{\geq 250}$ -*FGF14* DBN, also indicating a dysfunction of the

438 cerebellar flocculus/paraflocculus.<sup>28</sup> While multiple areas of the brain are involved in gaze stabilization during head  
439 movement, the cerebellar flocculus/paraflocculus is critical in modulating the VOR.<sup>28</sup> Early dysfunction of these two  
440 structures of the vestibulocerebellum, which are also implicated in the pathophysiology of DBN,<sup>3-6</sup> provides a  
441 unifying theory for the frequent and early co-occurrence of DBN, impaired VOR cancellation and other eye  
442 movement abnormalities, and vertiginous symptoms in *GAA-FGF14* disease.<sup>29</sup>

443  
444 Functional impairment, as captured by the FARS functional disability stage and the need for walking aid, increased  
445 relatively slowly with disease duration in  $(GAA)_{\geq 250}$ -*FGF14* patients, confirming previous reports from patients  
446 *GAA-FGF14* ataxia.<sup>12,14</sup> The slow accrual of disability was evidenced by an increase of 0.10 FARS stage per year of  
447 disease and the need for mobility aids in 20% of patients. However, we observed inter-individual variability in the  
448 rate of disability progression among patients, which was at least in part accounted for by concurrent medical  
449 illnesses in some patients. These observations strengthen the previous hypothesis from patients with *GAA-FGF14*  
450 ataxia<sup>14</sup> that co-morbid medical illnesses, which are common in elderly populations, must be taken into account  
451 when analyzing patient data and planning treatment trials in late-onset diseases like *GAA-FGF14* disease as they  
452 may alter their natural progression.

453  
454 The pathogenic potential of  $(GAA)_{200-249}$  alleles has recently been suggested based on the observation of segregation  
455 of  $(GAA)_{200-249}$  alleles and  $(GAA)_{\geq 250}$  expanded alleles with the disease in an affected family with autosomal  
456 dominant late-onset slowly progressive cerebellar ataxia.<sup>24</sup> Extending this preliminary evidence derived from  
457 segregation analysis, we now provide quantitative gene burden evidence for this notion by showing a significant  
458 enrichment of  $(GAA)_{200-249}$  alleles in patients with DBN compared to controls. In addition, as a clinical-phenotypic  
459 line of evidence, we observed that impairment of the visual fixation suppression of the VOR and treatment response  
460 to 4-AP, both strongly associated with  $(GAA)_{\geq 250}$ -*FGF14* DBN syndromes, were significantly more common in  
461  $(GAA)_{200-249}$ -*FGF14* than  $(GAA)_{<200}$ -*FGF14* patients (but not different from  $[GAA]_{\geq 250}$ -*FGF14* patients). It might  
462 thus be conceivable that *FGF14*  $(GAA)_{200-249}$  alleles may cause a milder cerebellar phenotype more commonly  
463 manifesting with isolated DBN and cerebellar ocular motor signs. These findings highlight the need to conduct  
464 large-scale studies to re-evaluate the pathogenic threshold of *GAA-FGF14* disease.

465

466 Our study provides large scale evidence – including a first small set of placebo-controlled data – for the  
467 symptomatic benefit of 4-AP in patients with *GAA-FGF14* disease. Our findings showed that 4-AP treatment was  
468 associated with a clinician-reported treatment response in 80% and a patient-reported meaningful benefit in 59% of  
469  $(GAA)_{\geq 250}$ -*FGF14* and  $(GAA)_{200-249}$ -*FGF14* patients. In comparison, such treatment responses were significantly  
470 lower in  $(GAA)_{<200}$ -*FGF14* patients (31% and 11%, respectively). Although these data were collected without  
471 knowledge of the *GAA-FGF14* genotype but with open-label provision of treatment, the substantially lower  
472 response rate to 4-AP treatment in  $(GAA)_{<200}$ -*FGF14* patients diminishes the likelihood that the observed effects can  
473 be largely attributed to placebo. Response to 4-AP in fact represented a strong predictor of  $(GAA)_{\geq 250}$ -*FGF14* and  
474  $(GAA)_{200-249}$ -*FGF14* status in patients with idiopathic DBN (positive predictive value, 89%). The mechanisms by  
475 which 4-AP, a potassium channel blocker,<sup>30</sup> ameliorates symptoms in *GAA-FGF14* disease are yet to be established  
476 although it may involve restoration of cerebellar Purkinje cell rhythmic firing property, as shown in other forms of  
477 hereditary ataxia,<sup>30,31</sup> that is disrupted with the loss of *FGF14* function.<sup>32,33</sup>

478  
479 Albeit still preliminary due to the small sample set, we present the first double-blind (and genotype-blind), placebo-  
480 controlled randomized treatment data of 4-AP in patients with *GAA-FGF14* disease.<sup>16</sup> All four patients for whom  
481 DNA was available and who were found to carry an *FGF14*  $(GAA)_{\geq 250}$  repeat expansion showed an improvement of  
482 SPV of the DBN on 4-AP, but not placebo. This observation bolsters the existing open-label real-world evidence for  
483 the treatment efficacy of 4-AP in *GAA-FGF14* disease observed in our current cohort and in previous SCA27B  
484 cohorts.<sup>14,18</sup> A larger randomized controlled trial is, nonetheless, warranted to confirm these findings. Our study  
485 highlights the potential of video-oculographic assessment of DBN as a responsive outcome measure for treatment.  
486 This approach could complement a promising battery of quantitative digital-motor outcomes,<sup>18</sup> including sensor  
487 measures of gait and balance, in future trials of 4-AP for *GAA-FGF14* disease.

488  
489 The results of our study need to be interpreted in light of some limitations. First, multi-centre studies with patient  
490 cohorts with DBN from different ethnic backgrounds are needed to replicate our findings, which were drawn from a  
491 single-centre patient cohort of overwhelmingly European background. Such studies are warranted to assess the  
492 frequency of *FGF14* *GAA* repeat expansions in other populations as profiling of the *FGF14* repeat locus in controls  
493 of diverse ancestries has shown that larger *FGF14* alleles are less common in non-European persons.<sup>12,13</sup> Second,

494 future prospective natural history studies in patients with DBN and other non-ataxia predominant phenotypes will be  
495 required to accurately track the phenotypic evolution of *GAA-FGF14* disease from its earliest stage and to establish  
496 whether a subset of patients never develops overt cerebellar ataxia. Third, our findings of potential pathogenicity of  
497 *FGF14* (*GAA*)<sub>200-249</sub> alleles are largely preliminary and require validation through additional segregation studies,  
498 larger case-control series, and functional studies. Fourth, although we show a treatment effect of 4-AP in several  
499 outcomes, including clinician-reported and quantitative digital-motor outcomes, substantiating our real-world  
500 treatment response data, the benefits reported herein need validation in a larger randomized placebo-controlled trial.

501  
502 In conclusion, we showed that *FGF14* *GAA* repeat expansions are a highly frequent monogenic cause of DBN  
503 syndromes in the European population. Our study suggests that *GAA-FGF14* disease may present along a  
504 continuum of variable cerebellar involvement, with some patients exhibiting isolated cerebellar  
505 floccular/parafloccular ocular motor signs without overt ataxia, and that *FGF14* (*GAA*)<sub>200-249</sub> alleles may be  
506 associated with a disease phenotype. It also provides robust and large-scale evidence that the *GAA-FGF14* genotype  
507 status defines a subgroup of patients with DBN that appear to be highly responsive to 4-AP, further paving the way  
508 toward clinical trials.

509 **Author contributions**

510 Design or conceptualization of the study: DP, CW, BB, MSt, MSy.

511 Acquisition of data: DP, FH, CW, MCD, AT, CA, MJD, AC, GDG, KMB, JC, DR, AMH, SZ, BB, MSt, MSy.

512 Analysis or interpretation of the data: DP, FH, CW, MCD, AT, CA, MJD, BB, MSt, MSy.

513 Drafting or revising the manuscript for intellectual content: DP, FH, CW, MCD, AT, CA, MJD, AC, GDG, KMB,

514 JC, DR, AMH, SZ, BB, MSt, MSy.

515

516 **Declaration of interests**

517 DP reports no disclosures.

518 FH reports no disclosures.

519 CW reports no disclosures.

520 MCD reports no disclosures.

521 AT reports no disclosures.

522 CA reports no disclosures.

523 MJD reports no disclosures.

524 AC reports no disclosures.

525 GDG reports no disclosures.

526 KMB reports no disclosures.

527 JC reports no disclosures.

528 DR has received grant/research support from Janssen and Lundbeck; he has served as a consultant or on advisory

529 boards for AC Immune, Janssen, Roche and Rovi and he has served on speakers bureaus of Janssen and

530 Pharmagenetix. He also received honoraria from Gerot Lannacher, Janssen and Pharmagenetix, and travel support

531 from Angelini and Janssen, all unrelated to the present manuscript.

532 AMH reports no disclosures.

533 SZ has received consultancy honoraria from Neurogene, Aeglea BioTherapeutics, Applied Therapeutics, and is an

534 unpaid officer of the TGP foundation, all unrelated to the present manuscript.

535 BB reports no disclosures.

536 MSt is Joint Chief Editor of the Journal of Neurology, Editor in Chief of Frontiers of Neuro-otology and Section  
537 Editor of F1000. He has received speakers honoraria from Abbott, Auris Medical, Biogen, Eisai, Grünenthal,  
538 GSK, Henning Pharma, Interacoustics, J&J, MSD, NeuroUpdate, Otometrics, Pierre-Fabre, TEVA, UCB, and  
539 Viartis. He receives support for clinical studies from Decibel, U.S.A., Cure within Reach, U.S.A. and Heel,  
540 Germany. He distributes M-glasses and Positional vertigo App. He acts as a consultant for Abbott, AurisMedical,  
541 Bulbitech, Heel, IntraBio, Sensorion and Vertify. He is an investor and share-holder of IntraBio. All are unrelated to  
542 the present manuscript.

543 MSy has received consultancy honoraria from Janssen, Ionis, Orphazyme, Servier, Reata, Biohaven, Zevra, Lilly,  
544 GenOrph, and AviadoBio, all unrelated to the present manuscript. MSy is planning a treatment trial of 4-AP in  
545 GAA-*FGF14* disease together with Solaxa Inc. as a sponsor, but has not received any type of honoraria or funding  
546 from Solaxa.

547

## 548 **Acknowledgments**

549 We thank the patients and their families for participating in this study. We thank the Centre d'Expertise et de  
550 Services Génome Québec for assistance with Sanger sequencing. DP and GDG hold Fellowship awards from the  
551 Canadian Institutes of Health Research (CIHR).

552

## 553 **Study sponsorship and funding**

554 This work was supported by the Clinician Scientist program "PRECISE.net" funded by the Else Kröner-Fresenius-  
555 Stiftung (to CW, AT, and MSy), the grant 779257 "Solve-RD" from the European's Union Horizon 2020 research  
556 and innovation program (to MSy), and the grant 01EO 1401 by the German Federal Ministry of Education and  
557 Research (BMBF) (to MSt). This work was also supported by the Deutsche Forschungsgemeinschaft (DFG, German  
558 Research Foundation) N° 441409627, as part of the PROSPAX consortium under the frame of EJP RD, the  
559 European Joint Programme on Rare Diseases, under the EJP RD COFUND-EJP N° 825575 (to MSy, BB and – as  
560 associated partner – SZ), the NIH National Institutes of Neurological Disorders and Stroke (grant 2R01NS072248-  
561 11A1 to SZ), the Fondation Groupe Monaco (to BB), and the Montreal General Hospital Foundation (grant  
562 PT79418 to BB). The Care4Rare Canada Consortium is funded in part by Genome Canada and the Ontario  
563 Genomics Institute (OGI-147 to KMB), the Canadian Institutes of Health Research (CIHR GP1-155867 to KMB),

564 Ontario Research Fund, Genome Quebec, and the Children's Hospital of Eastern Ontario Foundation. The funders  
565 had no role in the conduct of this study.  
566

567 **References**

- 568 1. Wagner JN, Glaser M, Brandt T, Strupp M. Downbeat nystagmus: aetiology and comorbidity in 117  
569 patients. *J Neurol Neurosurg Psychiatry*. Jun 2008;79(6):672-7. doi:10.1136/jnnp.2007.126284
- 570 2. Tran TM, Lee MS, McClelland CM. Downbeat nystagmus: a clinical review of diagnosis and management.  
571 *Curr Opin Ophthalmol*. Nov 1 2021;32(6):504-514. doi:10.1097/icu.0000000000000802
- 572 3. Pierrot-Deseilligny C, Milea D. Vertical nystagmus: clinical facts and hypotheses. *Brain*. Jun 2005;128(Pt  
573 6):1237-46. doi:10.1093/brain/awh532
- 574 4. Kalla R, Deutschlander A, Hufner K, et al. Detection of floccular hypometabolism in downbeat nystagmus  
575 by fMRI. *Neurology*. Jan 24 2006;66(2):281-3. doi:10.1212/01.wnl.0000194242.28018.d9
- 576 5. Hufner K, Stephan T, Kalla R, et al. Structural and functional MRIs disclose cerebellar pathologies in  
577 idiopathic downbeat nystagmus. *Neurology*. Sep 11 2007;69(11):1128-35.  
578 doi:10.1212/01.wnl.0000276953.00969.48
- 579 6. Marti S, Straumann D, Büttner U, Glasauer S. A model-based theory on the origin of downbeat nystagmus.  
580 *Exp Brain Res*. Jul 2008;188(4):613-31. doi:10.1007/s00221-008-1396-7
- 581 7. Takahashi H, Ishikawa K, Tsutsumi T, et al. A clinical and genetic study in a large cohort of patients with  
582 spinocerebellar ataxia type 6. *J Hum Genet*. 2004;49(5):256-64. doi:10.1007/s10038-004-0142-7
- 583 8. Yabe I, Sasaki H, Takeichi N, et al. Positional vertigo and macroscopic downbeat positioning nystagmus in  
584 spinocerebellar ataxia type 6 (SCA6). *J Neurol*. Apr 2003;250(4):440-3. doi:10.1007/s00415-003-1020-5
- 585 9. Jen J, Kim GW, Baloh RW. Clinical spectrum of episodic ataxia type 2. *Neurology*. Jan 13 2004;62(1):17-  
586 22. doi:10.1212/01.wnl.0000101675.61074.50
- 587 10. Buttner N, Geschwind D, Jen JC, Perlman S, Pulst SM, Baloh RW. Oculomotor phenotypes in autosomal  
588 dominant ataxias. *Archives of neurology*. Oct 1998;55(10):1353-7. doi:10.1001/archneur.55.10.1353
- 589 11. Strupp M, Maul S, Konte B, et al. A Variation in FGF14 Is Associated with Downbeat Nystagmus in a  
590 Genome-Wide Association Study. *Cerebellum*. Jun 2020;19(3):348-357. doi:10.1007/s12311-020-01113-x
- 591 12. Pellerin D, Danzi MC, Wilke C, et al. Deep Intronic FGF14 GAA Repeat Expansion in Late-Onset  
592 Cerebellar Ataxia. *N Engl J Med*. Jan 12 2023;388(2):128-141. doi:10.1056/NEJMoa2207406



- 593 13. Rafehi H, Read J, Szmulewicz DJ, et al. An intronic GAA repeat expansion in FGF14 causes the  
594 autosomal-dominant adult-onset ataxia SCA27B/ATX-FGF14. *American journal of human genetics*. Jun 1  
595 2023;110(6):1018. doi:10.1016/j.ajhg.2023.05.005
- 596 14. Wilke C, Pellerin D, Mengel D, et al. GAA-FGF14 ataxia (SCA27B): phenotypic profile, natural history  
597 progression and 4-aminopyridine treatment response. *Brain*. May 11 2023;doi:10.1093/brain/awad157
- 598 15. Pellerin D, Wilke C, Traschütz A, et al. Intronic *FGF14* GAA repeat expansions are a common cause of  
599 ataxia syndromes with neuropathy and bilateral vestibulopathy. *J Neurol Neurosurg Psychiatry*. 2023;jnnp-2023-  
600 331490. doi:10.1136/jnnp-2023-331490
- 601 16. Claassen J, Spiegel R, Kalla R, et al. A randomised double-blind, cross-over trial of 4-aminopyridine for  
602 downbeat nystagmus--effects on slowphase eye velocity, postural stability, locomotion and symptoms. *J Neurol*  
603 *Neurosurg Psychiatry*. Dec 2013;84(12):1392-9. doi:10.1136/jnnp-2012-304736
- 604 17. Strupp M, Teufel J, Zwergal A, Schniepp R, Khodakhah K, Feil K. Aminopyridines for the treatment of  
605 neurologic disorders. *Neurol Clin Pract*. Feb 2017;7(1):65-76. doi:10.1212/cpj.0000000000000321
- 606 18. Seemann J, Traschütz A, Ilg W, Synofzik M. 4-Aminopyridine improves real-life gait performance in  
607 SCA27B on a single-subject level: a prospective n-of-1 treatment experience. *J Neurol*. 2023.
- 608 19. Subramony SH, May W, Lynch D, et al. Measuring Friedreich ataxia: Interrater reliability of a neurologic  
609 rating scale. *Neurology*. Apr 12 2005;64(7):1261-2. doi:10.1212/01.wnl.0000156802.15466.79
- 610 20. Krafczyk S, Tietze S, Swoboda W, Valkovic P, Brandt T. Artificial neural network: a new diagnostic  
611 posturographic tool for disorders of stance. *Clin Neurophysiol*. Aug 2006;117(8):1692-8.  
612 doi:10.1016/j.clinph.2006.04.022
- 613 21. Strupp M, Kim JS, Murofushi T, et al. Bilateral vestibulopathy: Diagnostic criteria Consensus document of  
614 the Classification Committee of the Bárány Society. *J Vestib Res*. 2017;27(4):177-189. doi:10.3233/ves-170619
- 615 22. England JD, Gronseth GS, Franklin G, et al. Distal symmetric polyneuropathy: a definition for clinical  
616 research: report of the American Academy of Neurology, the American Association of Electrodiagnostic Medicine,  
617 and the American Academy of Physical Medicine and Rehabilitation. *Neurology*. Jan 25 2005;64(2):199-207.  
618 doi:10.1212/01.wnl.0000149522.32823.ea
- 619 23. Bonnet C, Pellerin D, Roth V, et al. Optimized testing strategy for the diagnosis of GAA-FGF14  
620 ataxia/spinocerebellar ataxia 27B. *Scientific reports*. Jun 15 2023;13(1):9737. doi:10.1038/s41598-023-36654-8

- 621 24. Hengel H, Pellerin D, Wilke C, et al. As frequent as polyglutamine SCAs: SCA27B in a large German  
622 autosomal dominant ataxia cohort. *Movement Disorders*. 2023;In Press
- 623 25. Pellerin D, Gobbo GD, Couse M, et al. A common flanking variant is associated with enhanced meiotic  
624 stability of the *FGF14*-SCA27B locus. *bioRxiv*. 2023:2023.05.11.540430. doi:10.1101/2023.05.11.540430
- 625 26. Pellerin D, Iruzubieta P, Tekgül S, et al. Non-GAA Repeat Expansions in *FGF14* are Likely Not  
626 Pathogenic. *Movement Disorders*. 2023;In Press
- 627 27. Auton A, Brooks LD, Durbin RM, et al. A global reference for human genetic variation. *Nature*. Oct 1  
628 2015;526(7571):68-74. doi:10.1038/nature15393
- 629 28. Kheradmand A, Zee DS. Cerebellum and ocular motor control. *Frontiers in neurology*. 2011;2:53.  
630 doi:10.3389/fneur.2011.00053
- 631 29. Graves TD, Jen JC. Progressive Vestibulocerebellar Syndromes. In: Bronstein A, ed. *Oxford Textbook of*  
632 *Vertigo and Imbalance*. Oxford University Press; 2013:0.
- 633 30. Alviña K, Khodakhah K. The therapeutic mode of action of 4-aminopyridine in cerebellar ataxia. *J*  
634 *Neurosci*. May 26 2010;30(21):7258-68. doi:10.1523/jneurosci.3582-09.2010
- 635 31. Jayabal S, Chang HH, Cullen KE, Watt AJ. 4-aminopyridine reverses ataxia and cerebellar firing  
636 deficiency in a mouse model of spinocerebellar ataxia type 6. *Scientific reports*. Jul 6 2016;6:29489.  
637 doi:10.1038/srep29489
- 638 32. Bosch MK, Carrasquillo Y, Ransdell JL, Kanakamedala A, Ornitz DM, Nerbonne JM. Intracellular FGF14  
639 (iFGF14) Is Required for Spontaneous and Evoked Firing in Cerebellar Purkinje Neurons and for Motor  
640 Coordination and Balance. *J Neurosci*. Apr 29 2015;35(17):6752-69. doi:10.1523/jneurosci.2663-14.2015
- 641 33. Shakkottai VG, Xiao M, Xu L, et al. FGF14 regulates the intrinsic excitability of cerebellar Purkinje  
642 neurons. *Neurobiology of disease*. Jan 2009;33(1):81-8. doi:10.1016/j.nbd.2008.09.019
- 643

644 **Figure legends**

645 **Figure 1: Study flowchart of the recruitment of patients with idiopathic DBN**

646 DBN: downbeat nystagmus.

647

648 **Figure 2: Frequency of *FGF14* GAA repeat expansions in DBN syndromes**

649 (A) Allele distribution of the *FGF14* repeat locus in 170 patients with idiopathic DBN syndromes (340  
650 chromosomes). The density plot shows allele size frequencies, with higher densities indicating greater frequencies.  
651 The box-and-whisker plot above the graph shows the allele distribution. The box indicates the 25<sup>th</sup> percentile (first  
652 quartile), the median, and the 75<sup>th</sup> percentile (third quartile), and the whiskers indicate the 2.5<sup>th</sup> and 97.5<sup>th</sup>  
653 percentiles. Outliers are represented by black dots. The dashed gray line and the shaded gray area indicate the so-  
654 called “intermediate” allele range of (GAA)<sub>200-249</sub>, and the dashed red line represents the pathogenic threshold of  
655 (GAA)<sub>≥250</sub> repeat units. (B) Percentage of patients carrying an *FGF14* (GAA)<sub>≥250</sub> expansion (dark blue), an *FGF14*  
656 (GAA)<sub>200-249</sub> allele (light blue), and an *FGF14* (GAA)<sub><200</sub> allele (gray) in the subgroups with (1) pure DBN (0/7, 0/7,  
657 and 7/7 patients, respectively), (2) DBN plus additional isolated cerebellar ocular motor signs (DBN+COM) (42/75,  
658 10/75, and 23/75 patients), (3) DBN plus cerebellar ataxia (DBN+CA) (23/32, 2/32, and 7/32 patients), and (4) DBN  
659 plus cerebellar ocular motor signs and/or ataxia and extracerebellar features (DBN+EC) (17/56, 8/56, and 31/56  
660 patients). (C) Frequency of presenting symptoms in 81 patients with DBN carrying an *FGF14* (GAA)<sub>≥250</sub> expansion.  
661 Data on presenting symptoms were missing for one patient. Patients may present with multiple symptoms at disease  
662 onset. Visual disturbances include diplopia, oscillopsia, and visual blurring. (D) Weak inverse correlation between  
663 size of the repeat expansion and age at disease onset in 74 patients carrying an *FGF14* (GAA)<sub>≥250</sub> expansion for  
664 whom information on age at onset was available (Pearson’s  $r$ , -0.30;  $R^2=0.09$ ;  $p=0.01$ ). The gray area displays the  
665 95% confidence interval. (E) UpSet plot showing co-occurrence of cerebellar ocular motor signs (COM), cerebellar  
666 ataxia (CA), bilateral vestibulopathy (BVP), and neuropathy among 82 patients with DBN carrying an *FGF14*  
667 (GAA)<sub>≥250</sub> expansion.

668

669 **Figure 3: Phenotypic evolution in GAA-*FGF14*-related DBN syndromes**

670 (A) Kaplan-Meier curve showing the probability of being disease free as a function of the age (years) for 74 patients  
671 carrying an *FGF14* (GAA)<sub>≥250</sub> expansion for whom age at onset was available. Median survival before occurrence of

672 disease was 67 years. The shaded gray area displays the 95% confidence interval around the probability estimate.  
673 **(B)** Disease duration at last examination for patients carrying an *FGF14* (GAA)<sub>≥250</sub> expansion with DBN plus  
674 additional isolated cerebellar ocular motor signs (DBN+COM,  $n=36$ ; median, 4.25 years), DBN plus cerebellar  
675 ataxia (DBN+CA,  $n=21$ ; median, 6 years), and DBN plus cerebellar ocular motor signs and/or ataxia and  
676 extracerebellar features (DBN+EC,  $n=17$ ; median, 9 years). Disease duration was significantly longer for patients  
677 with extracerebellar features (Kruskal-Wallis test,  $p=0.014$ ). **(C)** Temporal evolution of select phenotypic features  
678 relative to the onset of gait impairment (dotted line). The solid black lines show the median time from onset of gait  
679 impairment for each individual feature. **(D)** Occurrence of cerebellar atrophy on brain MRI ( $n=7$ ), polyneuropathy  
680 on nerve conduction studies ( $n=7$ ), and BVP on caloric stimulation or vHIT ( $n=7$ ) in relation to disease onset. BVP  
681 occurred later in disease course compared to cerebellar ataxia and polyneuropathy (Kruskal-Wallis test,  $p=0.004$ ). In  
682 panels B and D, the individual between-group adjusted  $p$  values ( $q$  values) as per the Benjamini-Hochberg method  
683 are shown in the graphs. Adjusted  $p$  values  $< 0.05$  indicate statistically significant difference.

684

#### 685 **Figure 4: Progression of disability in GAA-*FGF14*-related DBN syndromes**

686 **(A)** Cross-sectional progression of the functional impairment as assessed by the FARS functional disability stage  
687 relative to disease duration ( $n=73$  patients carrying an *FGF14* [GAA]<sub>≥250</sub> expansion). The FARS functional stage  
688 was not significantly associated with disease duration (Spearman's rho, 0.194;  $p=0.10$ ). **(B)** Longitudinal intra-  
689 individual progression of functional impairment as assessed by the FARS functional disability stage relative to  
690 disease duration (148 observations from 40 patients carrying an *FGF14* [GAA]<sub>≥250</sub> expansion are shown).  
691 Observations from the same patient are connected by a dotted line. The solid black line shows the average  
692 progression of the FARS stage over disease duration across all patients as modelled by a linear mixed-effects model  
693 accounting for disease duration as fixed effect and with random intercepts and random slopes. In both panels, only  
694 FARS stages measured off 4-AP treatment are shown. Two patients who became wheelchair-dependent following a  
695 hip fracture and a prolonged medical admission, respectively, are represented by orange squares in panel A and by  
696 squares in panel B. These patients also carried alleles of 218 and 270 repeat units and 280 and 313 repeat units,  
697 respectively.

698

#### 699 **Figure 5: Association of the rs72665334 variant with *FGF14* GAA expansions**

700 (A) Diagram of the *FGF14* gene, isoform 1b showing the location of the rs72665334 variant (GRCh38,  
701 chr13:102,150,076) in relation to the (GAA)<sub>n</sub>•(TTC)<sub>n</sub> repeat locus in the first intron (GRCh38, chr13:102,161,575-  
702 102,161,726). The rs72665334 variant is located ~11kb 5' of the intronic GAA short tandem repeat. (B) Percentage  
703 of patients with DBN carrying an *FGF14* (GAA)<sub>≥250</sub> expansion (*n*=35; dark blue), an *FGF14* (GAA)<sub>200-249</sub> allele  
704 (*n*=9; light blue), and an *FGF14* (GAA)<sub><200</sub> allele (*n*=29; gray) with the C|C, C|T, and T|T rs72665334 genotypes.  
705 There was a significantly greater proportion of patients with a C|T or T|T genotype who carried an *FGF14*  
706 (GAA)<sub>≥250</sub> expansion compared to an *FGF14* (GAA)<sub><200</sub> allele (54% vs 14%; OR, 7.18; 95% CI, 1.91 to 34.43;  
707 Fisher's exact test *p*=0.0014). (C) Frequency of the C|C, C|T, and T|T rs72665334 genotypes in 503 European  
708 controls from the 1000 Genomes Project<sup>27</sup> and 72 patients with DBN carrying an *FGF14* (GAA)<sub>≥250</sub> expansion  
709 (*n*=35 from the idiopathic DBN cohort and *n*=37 from the French-Canadian cohort). The C|T and T|T genotypes  
710 were significantly more frequent in patients with DBN carrying an *FGF14* (GAA)<sub>≥250</sub> expansion compared to  
711 controls (46%, 33/72 vs 12%, 60/503; OR, 6.22; 95% CI, 3.51 to 11.02; Fisher's exact test, *p*=1.063e-10).

712

### 713 **Figure 6: Treatment response to 4-aminopyridine**

714 *Open-label real-world treatment response data.* Percentage of patients presenting (A) a clinician-reported response  
715 and (B) a patient-reported response to 4-AP treatment in the subgroups with an *FGF14* (GAA)<sub>≥250</sub> expansion (29/36  
716 and 29/49 patients, respectively), (GAA)<sub>200-249</sub> allele (4/5 and 3/5 patients), and (GAA)<sub><200</sub> allele (5/16 and 2/19  
717 patients). *Double-blind placebo-controlled clinical trial data.* (C) Effect of 4-AP on the slow phase velocity (SPV)  
718 of DBN in a randomized double-blind trial.<sup>16</sup> Compared to baseline, the absolute value of the SPV significantly  
719 decreased under 4-AP 60 min after the first dose and under daily treatment, but not under placebo. Note that all four  
720 patients were randomized to receive active treatment first (= 4-AP) and placebo second. SPV during all three  
721 measurements of the placebo phase was higher than at the baseline of the treatment phase. The improved values of  
722 the SPV under 4-AP were in the range of the placebo values. (D) Treatment effect of 4-AP, illustrated by relative  
723 reduction (improvement) of SPV as potential future trial outcome assessment. The genotype of the four patients was:  
724 patient 1, 65 and 298 repeat units; patient 2, 44 and 354 repeat units; patient 3, 16 and 265 repeat units; patient 4,  
725 280 and 313 repeat units. In panels C and D, the black line shows the mean value for all four patients at each time  
726 point.

727

728 **Figure 7: Effect of 4-aminopyridine on functional impairment**

729 Longitudinal intra-individual progression of functional impairment while on and off 4-AP treatment as assessed by  
730 the FARS functional disability stage relative to disease duration for **(A-G)** seven patients carrying an *FGF14*  
731  $(GAA)_{\geq 250}$  expansion and **(H-I)** two patients carrying an *FGF14*  $(GAA)_{200-249}$  allele. FARS stages were invariably  
732 lower while on 4-AP treatment for all nine patients, indicating an improvement in the level of disability. No patients  
733 carrying an *FGF14*  $(GAA)_{<200}$  allele had FARS stages recorded while on and off 4-AP treatment.

734

735

736 **Table**

	<i>FGF14</i> (GAA) <sub>≥250</sub> (n=82)	<i>FGF14</i> (GAA) <sub>200-249</sub> (n=20)	<i>FGF14</i> (GAA) <sub>&lt;200</sub> (n=68)	<i>FGF14</i> (GAA) <sub>≥250</sub> vs <i>FGF14</i> (GAA) <sub>&lt;200</sub>	
				p value	q value
Male sex	44 (54%)	11 (55%)	40 (59%)	-	-
Triplet repeat count of the larger allele*	324 (250-516)	234 (207-249)	37 (8-335)	-	-
Age at disease onset	67 (30-81)	72 (45-84)	66 (17-88)	0.611	0.760
Disease duration	6 (0-26.5)	4 (0-23)	4 (0-50)	0.180	0.407
Age at last examination	73 (40-92)	79 (58-86)	72 (21-91)	0.346	0.554
Positive family history	31/82 (38%)	4/20 (20%)	13/67 (19%)	0.019	0.097
FARS disability stage	3 (1.5-5)	3 (1.5-4)	2 (1-4)	0.115	0.307
History of falls	27/50 (54%)	8/14 (57%)	20/35 (57%)	0.827	0.882
Regular use of walking aid	16/81 (20%)	4/19 (21%)	15/66 (23%)	0.688	0.760
<b>Symptoms</b>					
Episodic symptoms	11/81 (14%)	0/19 (0%)	18/66 (27%)	0.059	0.173
Gait impairment	81/81 (100%)	20/20 (100%)	68/68 (100%)	1.000	1.000
Vertigo or dizziness	23/82 (28%)	10/20 (50%)	26/68 (38%)	0.222	0.442
Visual disturbances	47/82 (57%)	8/20 (40%)	30/68 (44%)	0.140	0.344
Fine motor impairment	12/80 (15%)	1/18 (6%)	12/66 (18%)	0.658	0.760
Speech impairment	13/81 (16%)	4/19 (21%)	13/66 (20%)	0.665	0.760
<b>Clinical signs</b>					
Impaired balance/gait	67/79 (85%)	13/19 (68%)	44/64 (69%)	0.027	0.108
<b>Cerebellar ocular motor signs</b>					
- Gaze-evoked nystagmus	57/82 (70%)	11/20 (55%)	42/67 (63%)	0.390	0.594
- Saccadic pursuit	80/81 (99%)	20/20 (100%)	59/67 (88%)	0.011	0.073
- Dysmetric saccades	22/79 (28%)	6/20 (30%)	22/67 (33%)	0.588	0.760
- Impaired VOR suppression**	64/74 (86%)	16/17 (94%)	23/46 (50%)	<0.001	<b>&lt;0.001</b>
<b>Cerebellar ataxia</b>					
- Ataxia of upper limbs	16/73 (22%)	1/14 (7%)	21/55 (38%)	0.051	0.163
- Dysidiadochokinesia	15/71 (21%)	1/16 (6%)	14/54 (26%)	0.531	0.760
- Dysarthria	12/80 (15%)	2/19 (11%)	12/63 (19%)	0.653	0.760
Tremor of upper limbs	12/79 (15%)	2/18 (11%)	6/66 (9%)	0.318	0.536
<b>Neuropathy</b>					
- Impaired vibration at ankle (≤3/8)	11/79 (14%)	6/18 (33%)	24/65 (37%)	0.002	<b>0.018</b>
- Ankle hyporeflexia	20/79 (25%)	4/18 (22%)	29/65 (45%)	0.021	0.097
Pyramidal tract signs	1/77 (1%)	0/18 (0%)	2/66 (3%)	0.595	0.760
<b>MRI</b>					
Vermis atrophy	8/60 (13%)	1/11 (9%)	10/47 (21%)	0.307	0.536
Cerebellar hemisphere atrophy	5/60 (8%)	0/11 (0%)	8/47 (17%)	0.235	0.442
Supratentorial atrophy	5/60 (8%)	2/10 (20%)	4/47 (9%)	1.000	1.000
Brainstem atrophy	0/60 (0%)	0/10 (0%)	2/47 (4%)	0.191	0.407
<b>Nerve conduction studies</b>					
Axonal neuropathy	7/15 (47%)	3/5 (60%)	10/11 (91%)	0.036	0.129
<b>Vestibular function evaluation – caloric stimulation, vHIT, rotatory chair</b>					
Bilateral vestibulopathy	8/79 (10%)	3/18 (17%)	20/62 (32%)	0.001	<b>0.018</b>
VOR gain on vHIT in BVP – mean (±SD)	0.44 (±0.10)	0.21	0.18 (±0.17)	0.006	<b>0.0496</b>

737

738 **Table 1: Characteristics and discriminative features of GAA-*FGF14*-related DBN syndromes**

739 Unless specified, data are reported as median (range). Bold values indicate statistically significant adjusted  $p$  values  
740 ( $q$  values) below the 0.05 threshold as per the Benjamini-Hochberg method. Data on age at onset were missing for  
741 eight patients in the  $(GAA)_{\geq 250}$  group, three patients in the  $(GAA)_{200-249}$  group, and 10 patients in the  $(GAA)_{<200}$   
742 group.

743  
744 \*The *FGF14*  $(GAA)_{\geq 250}$  group includes a patient carrying biallelic GAA repeat expansions (280 and 313 repeat  
745 units). The *FGF14*  $(GAA)_{<200}$  group includes two patients carrying a likely non-pathogenic  $(GAAGGA)_n$  expansion  
746 (335 and 319 triplet repeat units equivalent, respectively). \*\*In patients without bilateral vestibulopathy.

747  
748 Abbreviations: BVP: bilateral vestibulopathy; FARS: Friedreich Ataxia Rating Scale; vHIT: video head impulse  
749 test; VOR: vestibulo-ocular reflex.

750



**219** patients with suspected DBN

```
graph TD; A[219 patients with suspected DBN] --> B[2 no DNA available]; A --> C[11 no DBN]; A --> D[36 competing etiologies for DBN]; A --> E[170 idiopathic DBN screening cohort]; D --> D1[8 metabolic]; D --> D2[6 drug toxicity]; D --> D3[6 alcohol toxicity]; D --> D4[3 genetic disease]; D --> D5[3 neoplastic]; D --> D6[3 vascular]; D --> D7[2 auto-immune]; D --> D8[5 other];
```






**2** no DNA available





**11** no DBN

**36** competing etiologies for DBN


- **8** metabolic
- **6** drug toxicity
- **6** alcohol toxicity
- **3** genetic disease
- **3** neoplastic
- **3** vascular
- **2** auto-immune
- **5** other

**170** idiopathic DBN screening cohort







A




B




A
















B



C







▲ On 4-AP

● Off 4-AP