

1 **National Changes in Diabetes Care Practices during the COVID-19 Pandemic: Prospective**

2 **Study of US Adults**

3 Short title: Impact of COVID-19 Pandemic on Diabetes Care

4 Kushagra Vashist, MPH <sup>a,\*</sup>, Saria Hassan, MD <sup>a, b</sup>, Mary Beth Weber, PhD <sup>a</sup>, Rakale C. Quarells  
5 <sup>c</sup>, PhD, Shivani A. Patel, PhD <sup>a</sup>

6  
7 <sup>a</sup>Emory University Rollins School of Public Health, Atlanta, GA

8 <sup>b</sup>Emory University School of Medicine, Atlanta, GA

9 <sup>c</sup>Morehouse School of Medicine, Atlanta, GA

10  
11 \*For Correspondence:

12 Kushagra Vashist,

13 1518 Clifton Road NE

14 7<sup>th</sup> Floor R Randall Rollins Building

15 Atlanta GA 30322

16 Email: [Kushagra.vashist@emory.edu](mailto:Kushagra.vashist@emory.edu)

17

18

19 Current word count- 2,061

20 Number of tables- 2

21

22 **ABSTRACT**

23 **Background**

24 There is a lack of nationally representative prospective data on the impact of the COVID-19  
25 pandemic on diabetes care and management in adults with type 2 diabetes. We examined changes  
26 in diabetes care and management practices before and after the onset of the COVID-19 pandemic.

27 **Methods**

28 Using the National Health Interview Survey, we analyzed data from 870 adults living with type 2  
29 diabetes who were interviewed in 2019 and re-interviewed between August and December 2020.  
30 Exposure to the COVID-19 pandemic was defined by year of survey (2019, pre-pandemic; 2020,  
31 pandemic). We estimated percent change in past year blood sugar check by a health professional  
32 and current use of blood sugar lowering medication overall and by sociodemographic subgroups.

33 **Results**

34 Receiving an annual blood sugar test fell by -3.3 percentage points (pp) (95% CI -5.7, -1.0), from  
35 98.3% in 2019 to 95.0% in late 2020. The reduction in annual blood glucose testing was largely  
36 consistent across socio-demographic groups and was particularly pronounced among adults not  
37 working and adults aged 65 years and older. In the same time period, current use of diabetes  
38 medications increased by +3.8 pp (0.7, 6.9), from 85.9% to 89.7%. The increase in medication use  
39 was most pronounced among individuals aged 40-64-year old, employed, and those living in large  
40 central metropolitan areas.

41 **Conclusions**

42 Nationally, adults with Type 2 diabetes reported a reduction in annual blood glucose testing by a  
43 health professional and an increase in diabetes medication usage during the COVID-19 pandemic.

- 44 If sustained after the end of the COVID-19 public health emergency, these changes have
- 45 implications for national diabetes management and care.

## 46 **Introduction**

47           Diabetes is a complex chronic condition requiring routine monitoring and regular  
48 medication. At minimum, the American Diabetes Association (ADA) recommends having an  
49 annual blood sugar check <sup>1</sup> and when needed, pharmacologic treatments such as metformin or  
50 other agents, including combination therapy <sup>2</sup> in adults with diabetes. There has been much  
51 concern that the COVID-19 pandemic has impacted diabetes care. Disruptions to health services  
52 <sup>3,4</sup>, changes to routine care <sup>4</sup>, and fear of getting severely ill from the virus <sup>5</sup> may have prevented  
53 some individuals living with diabetes from getting the care they need. Lack of access to health  
54 services has been associated with diabetes-related complications such as cardiovascular disease,  
55 kidney disease, neuropathy, or blindness <sup>5</sup>. Conversely, access to healthcare has been associated  
56 with controlled levels of blood glucose, blood pressure, and blood lipids <sup>6</sup>, which collectively  
57 prevent the onset of diabetes-related complications <sup>7,8</sup>.

58           Several studies indicate that that people with diabetes experienced challenges with  
59 managing their disease during the pandemic. For example, an online community-based survey  
60 indicated that 25% of people needing an insulin pump or continuous glucose monitoring supplies  
61 had delays or difficulties in obtaining them and that 1 in 6 persons with diabetes had difficulty  
62 needing insulin, as of April 2020 <sup>9</sup>. However, these data do not provide a clear picture regarding  
63 how individuals' diabetes care and management have changed in response to the pandemic.

64           Identifying the indirect impact of the pandemic is critical to understanding the needs and  
65 requirements of those living with diabetes for future public health emergencies and the state of  
66 diabetes care as we move into post-public health emergency recovery. Using prospective data  
67 from a nationally representative sample of adults interviewed in 2019 and re-interviewed in  
68 2020, we examined changes in annual blood sugar checks and use of current medication to lower

69 blood sugar — two key diabetes care and management practices— before and after the onset of  
70 the COVID-19 pandemic among people living with type 2 diabetes.

## 71 **Materials and METHODS**

### 72 **Data and sample**

73 The National Health Interview Survey (NHIS) is a representative household interview  
74 survey which collects information on the health of non-institutionalized US population across the  
75 50 states through a complex multistage design. To understand the ramifications of the COVID-  
76 19 pandemic, NHIS re-interviewed a sample of adults in August-December 2020 from 2019.  
77 This longitudinal sample started with 21,161 adults from non-Medical Expenditure Panel Survey  
78 that completed or partially completed a 2019 Sample Adult Interview <sup>10</sup>. After satisfying the  
79 eligibility criteria, this number was reduced to 20,827. Of these eligible sample adults, 10,415  
80 adults interviewed in August-December 2020. We merged the publicly available 2019 NHIS  
81 adults sample data with their follow up data in 2020 to conduct analysis of change in diabetes  
82 outcomes <sup>10</sup>. The analysis was restricted to adults with self-reported type 2 diabetes. Of the 973  
83 longitudinal sample respondents living with type 2 diabetes aged 18-96 years, 10.6% were  
84 excluded from the analysis because of missing information on a study outcome or covariate,  
85 resulting in an unweighted analytic sample size of 870.

86 NHIS is approved by the Research Ethics Review Board of the National Center for  
87 Health Statistics and the U.S Office of Management and Budget. All NHIS respondents provided  
88 oral consent prior to participation <sup>11</sup>. This study used publicly available data containing no  
89 personal identifying information and is therefore deemed not human subjects research by the  
90 Emory IRB.

### 91 **Variables and description**

92 Our two outcomes included having blood sugar checked by a doctor, nurse, or other  
93 health professional within the past 12 months and taking medication to lower blood sugar for  
94 diabetes control (either oral/hypoglycemic agents or insulin).

#### 95 *Socio-demographic characteristics*

96 Demographics included sex (male/female), age (divided into three groups; 18-39, 40-64,  
97 and 65 and older), race/ethnicity (as non-Hispanic White, non-Hispanic Black, Hispanic, and  
98 Other), smoking status (former/never smoker and current everyday/someday smoker) and marital  
99 status (married or living with a partner). Socioeconomic characteristics included educational  
100 attainment (classified as having less than bachelor's degree or bachelor's degree or more),  
101 urbanicity (categorized as large central/fringe metropolitan [population > 1 million], medium and  
102 small metropolitan, and nonmetropolitan), employment status (full-time and/or part-time  
103 employed, retired, unemployed/not working), and insurance status (no insurance, private  
104 insurance, Medicare insurance, Medicaid insurance, and other public insurance). Types of  
105 insurance were mutually exclusive and individuals with multiple health insurance were assigned  
106 a single insurance in the following order: those with private insurance included individuals who  
107 only had private insurance or a combination with any other coverage; next, those with Medicare  
108 insurance included having Medicare alone or a combination with any other public insurance; and  
109 lastly, those having Medicaid insurance included persons who had Medicaid insurance alone or  
110 in combination with any non-Medicare, public insurance.

#### 111 **Statistical analysis**

112 We first described the distribution of socio-demographic characteristics of respondents at  
113 baseline in 2019.

114 We estimated the prevalence of annual blood sugar checked and taking medication to  
115 lower blood sugar for diabetes control in 2019. Next, we estimated the average within-person  
116 change in each outcome from 2019 to 2020. Changes were estimated for the total sample as well  
117 as by age, sex, race, smoking status, marital status, urbanicity, insurance type, education level,  
118 and employment status. We computed the unadjusted change in each stratum, as well as the  
119 marginally adjusted change accounting for other sociodemographic groups in the analysis.

120 All data were analyzed using SAS version 9.4 and SUDAAN version 11.0.1, accounting  
121 for the complex survey design.

## 122 **RESULTS**

123 The distribution of the sociodemographic characteristics of adults with type 2 diabetes at  
124 baseline (2019) are shown in Table 1.

125 Table 2 shows the prevalence of diabetes care practices in 2019 and the unadjusted and  
126 adjusted average percent change from 2019 to 2020. Nationally, the percentage receiving an  
127 annual blood sugar check fell by an average of 3.3 percentage points (pp) (95% Confidence  
128 Interval [CI] -5.7, -1.0) from 98.3% in 2019 to 95.0% in Aug-Dec 2020. The reduction in annual  
129 blood sugar checking was largely consistent across sociodemographic groups, and was  
130 particularly pronounced among males (-4.7 pp [CI: -8.4, -1.1]), those not working (-8.1 pp [CI:-  
131 14.4, -1.8]) and 65 years and older (-4.9 pp [CI: -9.2, -0.7]). The percentage of participants  
132 taking medication to lower blood sugar increased by +3.8 pp (CI: 0.7, 6.9), from 85.9% to 89.7%  
133 from 2019 to 2020. The increase in anti-glycemic medication was most pronounced among those  
134 living in large central metropolitan area (5.8 pp [CI: 0.6, 11.0]) and 40-64 years old (4.8 pp [CI:  
135 0.4, 9.2].

## 136 **DISCUSSION**

137 To our knowledge, this is the first study which uses nationally representative,  
138 community-based prospective data to examine within-person changes in diabetes care practices  
139 associated with the COVID-19 pandemic. Among adults with type 2 diabetes, having blood  
140 sugar checked by a health professional decreased during the pandemic. On the other hand, taking  
141 medications to lower blood sugar for diabetes control increased. The findings suggest that the  
142 pandemic had a negative impact on diabetes care but an unclear impact on diabetes medication.  
143 The increase in diabetes medication usage may suggest a positive impact on diabetes control. On  
144 the other hand, it may have been that blood sugar levels worsened during the pandemic, leading  
145 more individuals to take medication for diabetes control.

146 Reasons for decreased testing during the pandemic may have been due to disruption of  
147 health services <sup>3,4</sup>, changes in routine care <sup>4</sup>, and fear of getting ill from the COVID-19 infection  
148 when going to the doctor <sup>5</sup>. Although reductions in testing were consistent across socio-  
149 demographic subgroups, changes differed in magnitude for some. We saw a greater decrease in  
150 blood sugar checked in individuals 65 years and older compared to other age groups. Older  
151 individuals are at a higher risk of getting severely ill from COVID-19 <sup>12</sup>, which may have been a  
152 factor in deterring them from visiting a health professional for testing or health professionals  
153 might have encouraged telehealth only visits for this high risk group. Adults who were not  
154 working at the 2019 baseline reported the largest decrease in having blood sugar checked (-  
155 8.1%), compared to other employment groups. The not working employment group largely  
156 consisted of persons unemployed, unable to work for health reasons/disabled, and taking care of  
157 house or family, and it may have been that these individuals were restricted in their ability to  
158 seek healthcare for the very reasons they were classified as not working.

159 Respondents reported an average increase in use of medications to lower blood sugar in  
160 August-December 2020 compared to 2019. This finding is consistent with reports from a  
161 previous study using an online survey, which showed that among respondents with type 2  
162 diabetes who reported a change in the intake of diabetes medication, a greater proportion of  
163 individuals took medications more regularly during the pandemic compared to before the  
164 pandemic <sup>13</sup>. The increase in diabetes usage during the early pandemic could be explained by one  
165 of two scenarios: 1. Patients were previously prescribed a diabetes lowering medication but had  
166 not started it or not taking it for some reason; or 2. Patients previously not eligible for diabetes  
167 medication based on their blood sugar values, were now eligible for blood sugar lowering  
168 medication. Alternatively, persons may have been more diligently checking their blood sugar at  
169 home, identified rising blood sugars, contacted their provider, and received newly prescribed  
170 blood sugar lowering medication. It is noteworthy that individuals with diabetes were able to get  
171 refills on prescriptions without visiting the doctor in 2020, whereas they had to return to their  
172 doctor to get refills by 2021.

173 In our analysis, we found that those living in large central metropolitan areas and in the  
174 40-64 years old age group reported a higher increase in the intake of medication to lower blood  
175 sugar in 2020 compared to those living in less urban areas or younger/older age groups.  
176 Individuals living in urban areas may have better infrastructure to obtain medications <sup>14</sup>, better  
177 access to telehealth services <sup>15</sup>, and be able to better afford medications compared to rural places  
178 <sup>14</sup>. Individuals aged 40-64 reported an increase in medication to lower blood sugar. These  
179 individuals may also be more likely to have access to internet and telehealth services <sup>16</sup> to enable  
180 discussion with providers on the need for medication despite not being able to visit the provider.

181 Broadly, our findings regarding both medication and testing are consistent with other  
182 national studies examining diabetes testing and medication usage during the pandemic. A  
183 previous study using electronic medical records in the United States during the same time period  
184 found a decrease outpatient visits and HbA1c testing during the pandemic, without evidence of  
185 reduced medication fills or glucose control <sup>17</sup>. Considering serial cross-sectional data from NHIS  
186 respondents, the prevalence of blood glucose checking in US adults with diabetes was 96.8% in  
187 2019 compared with 94.2% in 2021, suggesting that the drop in glucose checking by health  
188 professionals was sustained even in the second year of the pandemic <sup>18</sup>. Similarly, reductions in  
189 diabetes testing have also been reported in England <sup>19</sup>. Reductions in HbA<sub>1C</sub> testing is of concern  
190 due the importance of glucose monitoring for clinical treatment decisions and feedback to  
191 patients on diabetes management <sup>19</sup>.

192 Our study has several strengths. First, we used nationally representative prospective data  
193 to quantify diabetes care practices among the same individuals with type 2 diabetes before  
194 (2019) and after the onset of the pandemic (2020). Second, we examined changes in taking  
195 medication for diabetes control and having annual blood sugar checked from 2019 to 2020 by  
196 key sociodemographic groups, highlighting significant changes. Several limitations of this study  
197 should be acknowledged. There may have been overlapping time periods between 2019 and 2020  
198 for responses to having blood sugar checked by a health professional in the past 12 months. We  
199 were not able to assess whether changes in diabetes testing and medication resulted from changes  
200 in diabetes control due to lack of laboratory data.

## 201 **CONCLUSIONS**

202 Using data from prospectively followed adults drawn from a nationally representative  
203 sample, we found a reduction in annual blood sugar checks by healthcare providers and an

204 increase in reporting of taking medication to lower blood sugar in 2020 compared to 2019,  
205 among adults with type 2 diabetes. Despite a reduction in testing, there was an increase in those  
206 taking medication for diabetes. The increase in medication warrants further examination.  
207 Whether and to what extent these changes were sustained after the first year and will set a “new  
208 normal” for diabetes care and management is not fully clear. Understanding the indirect impact  
209 of the pandemic on diabetes care is critical to equitably addressing the needs of those living with  
210 diabetes in future public health emergencies and their aftermath. The need to intervene on  
211 communities and individuals at greater risk is necessary to reduce the impact of public health  
212 emergencies and ensure equitable access to recommended diabetes care and management  
213 practices.  
214

215 **Acknowledgments**

216

217 **Funding and assistance**

218 KV, MBW, RCQ, and SAP were supported by grant number P30DK111024-05S2 from the  
219 National Institute of Health under the RADx® Underserved Populations (RADx-UP) Initiative,  
220 and MBW is supported by P30DK111024. SH is supported by a grant from the National Institute  
221 of Health/National Heart Lung and Blood Institute (K23HL152368).

222

223 **Conflict of interest**

224 The authors declare that there are no relevant conflicts of interest.

225

226 **Author contributions**

227

228 KV and SAP conceptualized the study. KV conducted the statistical analysis and wrote the first  
229 draft of the manuscript. SAP provided —supervision of the study and input into study design. All  
230 authors contributed to the interpretation of data. SH, MBW, RCQ, SAP reviewed and provided  
231 substantive revisions to the manuscript. All authors approved the final version of the manuscript.  
232 KV is the guarantor of this work and, as such, had full access to all the data in the study and takes  
233 responsibility for the integrity of the data and the accuracy of the data analysis.

## REFERENCES

1. ElSayed NA, Aleppo G, Aroda VR, et al. 6. Glycemic Targets: Standards of Care in Diabetes—2023. *Diabetes Care*. 2022;46(Supplement\_1):S97-S110. doi:10.2337/dc23-S006
2. ElSayed NA, Aleppo G, Aroda VR, et al. 9. Pharmacologic Approaches to Glycemic Treatment: Standards of Care in Diabetes—2023. *Diabetes Care*. 2022;46(Supplement\_1):S140-S157. doi:10.2337/dc23-S009
3. Gregg EW, Sophiea MK, Weldegiorgis M. Diabetes and COVID-19: Population Impact 18 Months Into the Pandemic. *Diabetes Care*. 2021;44(9):1916-1923. doi:10.2337/dci21-0001
4. Hartmann-Boyce J, Morris E, Goyder C, et al. Diabetes and COVID-19: Risks, Management, and Learnings From Other National Disasters. *Diabetes Care*. 2020;43(8):1695-1703. doi:10.2337/dc20-1192
5. Sujan MdSH, Tasnim R, Islam MdS, et al. COVID-19-specific diabetes worries amongst diabetic patients: The role of social support and other co-variates. *Prim Care Diabetes*. 2021;15(5):778-785. doi:10.1016/j.pcd.2021.06.009
6. Zhang X, Bullard KM, Gregg EW, et al. Access to Health Care and Control of ABCs of Diabetes. *Diabetes Care*. 2012;35(7):1566-1571. doi:10.2337/dc12-0081
7. Al Nozha OM. Diabetes care and control: the effect of frequent visits to diabetes care center. *Ann Saudi Med*. 2014;34(3):229-234. doi:10.5144/0256-4947.2014.229
8. Deshpande AD, Harris-Hayes M, Schootman M. Epidemiology of Diabetes and Diabetes-Related Complications. *Phys Ther*. 2008;88(11):1254-1264. doi:10.2522/ptj.20080020
9. Piante PD. Impact of COVID-19 on the Diabetes Community in the United States. dQ&A. Published April 17, 2020. Accessed June 1, 2023. <https://d-qa.com/impact-of-covid-19-on-the-usa-diabetes-community/>
10. Survey Description, National Health Interview Survey, 2020. Published online 2020.
11. NHIS - About the Survey page | NIOSH | CDC. Published December 8, 2022. Accessed July 27, 2023. <https://www.cdc.gov/niosh/topics/nhis/method.html>
12. COVID-19 Risks and Information for Older Adults. Centers for Disease Control and Prevention. Published February 22, 2023. Accessed July 5, 2023. <https://www.cdc.gov/aging/covid19/index.html>
13. Fisher L, Polonsky W, Asuni A, Jolly Y, Hessler D. The early impact of the COVID-19 pandemic on adults with type 1 or type 2 diabetes: A national cohort study. *Journal of Diabetes and its Complications*. 2020;34(12):107748. doi:10.1016/j.jdiacomp.2020.107748
14. Dugani SB, Mielke MM, Vella A. Burden and Management of Type 2 Diabetes Mellitus in Rural United States. *Diabetes Metab Res Rev*. 2021;37(5):e3410. doi:10.1002/dmrr.3410

15. Chen J, Amaize A, Barath D. Evaluating Telehealth Adoption and Related Barriers Among Hospitals Located in Rural and Urban Areas. *The Journal of Rural Health*. 2021;37(4):801-811. doi:10.1111/jrh.12534
16. Okoye SM, Mulcahy JF, Fabius CD, Burgdorf JG, Wolff JL. Neighborhood Broadband and Use of Telehealth Among Older Adults: Cross-sectional Study of National Survey Data Linked With Census Data. *Journal of Medical Internet Research*. 2021;23(6):e26242. doi:10.2196/26242
17. Patel SY, McCoy RG, Barnett ML, Shah ND, Mehrotra A. Diabetes Care and Glycemic Control During the COVID-19 Pandemic in the United States. *JAMA Internal Medicine*. 2021;181(10):1412-1414. doi:10.1001/jamainternmed.2021.3047
18. Casagrande SS, Lawrence JM. Blood glucose, blood pressure, and cholesterol testing among adults with diabetes before and during the COVID-19 pandemic, USA, 2019 vs 2021. *BMJ Open Diabetes Research and Care*. 2023;11(3):e003420. doi:10.1136/bmjdr-2023-003420
19. Carr MJ, Wright AK, Leelarathna L, et al. Impact of COVID-19 on diagnoses, monitoring, and mortality in people with type 2 diabetes in the UK. *The Lancet Diabetes & Endocrinology*. 2021;9(7):413-415. doi:10.1016/S2213-8587(21)00116-9

**Table 1. Distribution of socio-demographic characteristics among those living with type 2 diabetes, baseline interview, January 2019-December 2019**

Baseline interview distribution, Jan 2019-Dec 2019

	Unweighted n	Weighted % (95% CI)
<b>Characteristics</b>		
Overall	870	100.0 (-)
<b>Sex</b>		
Male	405	49.9 (45.4, 54.4)
Female	465	50.1 (45.6, 54.6)
<b>Age</b>		
18-39	20	*
40-64	365	50.7 (46.2, 55.3)
65 and older	485	45.1 (40.2, 50.0)
<b>Race/ethnicity</b>		
Non-Hispanic White	596	59.0 (53.9, 63.9)
Non-Hispanic Black	111	13.2 (10.2, 16.9)
Hispanic	95	16.7 (12.6, 21.7)
Other	68	11.1 (8.1, 15.0)
<b>Smoking status</b>		
Former/never smoker	772	89.6 (86.7, 92.0)
Current every day/some day smoker	98	10.4 (8.0, 13.3)
<b>Education</b>		
Less than 4 years of college	633	81.0 (77.8, 83.9)
4 years of college or more	237	19.0 (16.1, 22.2)
<b>Employment status</b>		
Employed	307	39.9 (35.6, 44.5)
Retired	403	39.0 (34.5, 43.7)
Not working†	160	21.0 (17.5, 25.1)
<b>Marital status</b>		
Married or living with a partner	423	63.0 (58.8, 67.1)
Unmarried	447	37.0 (32.9, 41.2)
<b>Insurance type‡</b>		

Not insured	32	6.6 (4.3, 10.0)
Private insurance	480	53.2 (48.5, 58.0)
Medicare	281	28.2 (24.2, 32.6)
Medicaid	62	10.0 (6.6, 14.8)
Other	15	*
<b>Urbanicity</b>		
Large central/fringe metropolitan	410	49.7 (43.8, 55.5)
Medium and small metropolitan	278	29.9 (24.9, 35.5)
Nonmetropolitan	182	20.4 (16.1, 25.5)

---

\*Percentages suppressed for n<30

†Consists of persons unemployed, unable to work for health reasons/disabled, taking care of house or family, going to school, and not working because of other reasons

\*Types of insurance coverage are mutually exclusive. Persons with multiple types of health insurance were assigned to the first appropriate category in the following order: privately-insured, Medicare-insured, and Medicaid-insured

**Table 2. Taking medication to lower blood sugar and having blood sugar checked, past 12 months at baseline and average percentage change at follow-up among those living with type 2 diabetes**

Characteristics	Blood sugar checked, past 12 months			Taking medication to lower blood sugar <sup>§</sup>		
	Baseline interview percentage, Jan 2019-Dec 2019 (95% CI)	Average percentage change from baseline during follow-up interview, Aug 2020-Dec 2020 (95% CI)		Baseline interview percentage, Jan 2019-Dec 2019 (95% CI)	Average percentage change from baseline during follow-up interview, Aug 2020-Dec 2020 (95% CI)	
		Unadjusted	Adjusted <sup>  </sup>		Unadjusted	Adjusted <sup>  </sup>
<b>Overall</b>	98.3 (96.9, 99.1)	<b>-3.3 (-5.6, -1.0)</b>	<b>-3.3 (-5.7, -1.0)</b>	85.9 (81.9, 89.1)	<b>+3.4 (0.7, 6.9)</b>	<b>+3.8 (0.7, 6.9)</b>
<b>Sex</b>						
Male	98.2 (95.4, 99.3)	<b>-4.5 (-8.3, -0.7)</b>	<b>-4.7 (-8.4, -1.1)</b>	87.1 (81.2, 91.3)	<b>+3.5 (0.1, 6.8)</b>	+2.9 (-0.5, 6.2)
Female	98.4 (96.8, 99.2)	-2.2 (-4.8, 0.4)	-1.9 (-4.6, 0.7)	84.7 (78.7, 89.2)	+4.2 (-1.2, 9.4)	+4.7 (-0.6, 10.0)
<b>Age</b>						
18-39	*	*	*	*	*	*
40-64	97.6 (94.9, 98.9)	-3.3 (-7.2, 0.6)	-1.9 (-5.8, 2.0)	85.4 (80.2, 89.4)	<b>+5.1 (1.5, 8.7)</b>	<b>+4.8 (0.4, 9.2)</b>
65 and older	99.0 (97.4, 99.6)	<b>-3.4 (-6.2, -0.7)</b>	<b>-4.9 (-9.2, -0.7)</b>	87.2 (80.6, 91.8)	+1.4 (-3.7, 6.5)	+1.3 (-4.7, 7.4)
<b>Race/ethnicity</b>						
Non-Hispanic White	98.1 (96.0, 99.1)	<b>-3.6 (-6.9, -0.3)</b>	-3.5 (-7.1, 0.0)	84.2 (79.2, 88.2)	<b>+3.3 (0.4, 6.1)</b>	<b>+4.5 (1.3, 7.8)</b>
Non-Hispanic Black	99.6 (96.8, 99.9)	-1.9 (-4.5, 0.6)	-2.9 (-6.3, 0.4)	86.3 (77.3, 92.1)	+1.8 (-6.0, 9.6)	+0.8 (-7.3, 8.8)
Hispanic	96.9 (90.6, 99.0)	-3.6 (-10.2, 3.0)	-2.5 (-10.6, 5.6)	93.1 (83.8, 97.3)	+3.6 (-3.1, 10.4)	+0.7 (-4.3, 5.8)
Other	100.0 (-)	-3.1 (-7.5, 1.3)	-3.9 (-8.5, 0.8)	83.4 (58.3, 94.7)	+9.4 (-8.8, 27.5)	+8.1 (-10.6, 26.8)
<b>Smoking status</b>						

Former/never smoker	98.3 (96.8, 99.2)	<b>-3.6 (-6.1, -1.0)</b>	<b>-3.4 (-5.9, -0.9)</b>	86.4 (82.1, 89.8)	<b>+3.5 (0.2, 6.8)</b>	<b>+3.8 (0.6, 7.0)</b>
Current every day/some day smoker	98.2 (92.5, 99.6)	-1.3 (-5.6, 3.0)	-2.9 (-8.3, 2.5)	81.4 (71.4, 88.5)	+6.0 (-3.3, 15.3)	+4.1 (-7.1, 15.2)
<b>Education</b>						
Less than 4 years of college	98.2 (96.4, 99.1)	<b>-3.5 (-6.2, -0.7)</b>	<b>-3.2 (-5.8, -0.7)</b>	87.2 (82.4, 90.8)	+3.6 (-0.0, 7.2)	<b>+3.9 (0.0, 7.8)</b>
4 years of college or more	99.0 (96.1, 99.8)	-2.8 (-5.8, 0.3)	<b>-3.7 (-7.1, -0.3)</b>	80.3 (72.8, 86.1)	+4.7 (-0.9, 10.4)	+3.4 (-2.6, 9.5)
<b>Employment status</b>						
Employed	97.4 (94.0, 98.9)	-3.6 (-8.6, 1.4)	-2.5 (-5.6, 0.7)	83.9 (78.4, 88.1)	<b>+4.8 (0.4, 9.2)</b>	+4.8 (-0.7, 10.3)
Retired	98.8 (97.0, 99.5)	<b>-3.8 (-6.8, -0.9)</b>	-1.6 (-5.5, 2.2)	87.8 (80.1, 92.8)	+1.6 (-4.0, 7.2)	+1.6 (-4.3, 7.4)
Not working†	99.1 (96.4, 99.8)	-1.7 (-4.6, 1.2)	<b>-8.1 (-14.4, -1.8)</b>	86.1 (76.0, 92.4)	<b>+6.8 (1.2, 12.4)</b>	+6.0 (-0.3, 12.4)
<b>Marital status</b>						
Married or living with a partner	97.9 (95.6, 99.0)	<b>-3.8 (-7.2, -0.4)</b>	<b>-3.7 (-6.5, -0.9)</b>	86.2 (80.5, 90.4)	<b>+5.1 (0.5, 9.7)</b>	+4.1 (-0.5, 8.8)
Unmarried	99.1 (97.7, 99.6)	-2.7 (-5.4, 0.1)	-2.7 (-5.8, 0.4)	85.4 (80.2, 89.4)	+1.9 (-1.6, 5.3)	+3.2 (-0.4, 6.8)
<b>Insurance type‡</b>						
Not insured	*	*	*	*	*	*
Private insurance	98.9 (96.3, 99.7)	-3.0 (-6.0, 0.1)	-2.8 (-5.6, 0.0)	87.2 (83.4, 90.3)	+4.6 (-0.3, 9.6)	+3.0 (-0.4, 6.4)
Medicare	98.6 (96.1, 99.5)	<b>-4.0 (-7.9, -0.1)</b>	-3.7 (-8.0, 0.6)	83.6 (73.6, 90.3)	+1.8 (-2.0, 5.5)	+7.0 (-0.3, 14.3)
Medicaid	97.2 (91.4, 99.1)	-3.9 (-14.6, 6.7)	1.9 (-2.6, 6.4)	90.3 (77.6, 96.2)	+3.1 (-3.8, 10.0)	-0.0 (-8.5, 8.5)
Other	*	*	*	*	*	*

**Urbanicity**

Large central/fringe metropolitan	98.6 (96.6, 99.5)	<b>-3.8 (-7.1, -0.5)</b>	<b>-3.9 (-7.3, -0.6)</b>	85.2 (78.8, 90.0)	<b>+5.8 (0.4, 11.2)</b>	<b>+5.8 (0.6, 11.0)</b>
Medium and small metropolitan	97.8 (93.6, 99.2)	-1.1 (-3.0, 0.9)	-1.4 (-3.8, 1.0)	87.5 (82.4, 91.2)	+2.1 (-1.4, 5.6)	+2.2 (-1.6, 6.0)
Nonmetropolitan	98.4 (94.8, 99.5)	-5.4 (-12.7, 1.9)	-4.7 (-10.8, 1.4)	85.1 (73.8, 92.1)	+1.5 (-4.1, 7.0)	+1.3 (-3.3, 5.9)

Bold: Significant at alpha=0.05

\*Percentages suppressed for n<30

†Consists of persons unemployed, unable to work for health reasons/disabled, taking care of house or family, going to school, and not working because of other reasons

\*Types of insurance coverage are mutually exclusive. Persons with multiple types of health insurance were assigned to the first appropriate category in the following order: Privately-insured, Medicare-insured, and Medicaid-insured

§Composite of either taking oral/hypoglycemic agents to lower blood sugar or taking insulin to lower blood sugar, or both

||Adjusted for all characteristics shown in the table