1 Running head: Human migration, city growth, and vector surveillance and emergence 2 Title: Migration and the Emergence of Chagas Disease Vectors in the Growing City of El Pedregal, 3 4 Peru 5 6 **Authors** Raquel Gonçalves¹, Kathryn P. Hacker^{2,3}, Carlos Condori¹, Sherrie Xie², Katty Borrini-Mayori¹, Lina 7 Mollesaca Riveros¹, Roger Quispe Apaza¹, Manuel Ysidro Arratea¹, Gustavo Nativio², Ricardo Castillo-8 Neyra^{1,2}, Valerie A. Paz-Soldan^{1,4}, Michael Z. Levy^{1,2}* 9 10 *Corresponding author: Michael Z. Levy: mzlevy@pennmedicine.upenn.edu 11 12 13 **Filiations:** ¹Zoonotic Disease Research Lab (LIEZ), One Health Unit, School of Public Health and Administration, 14 15 Universidad Peruana Cayetano Heredia, Lima, Peru ² Department of Biostatistics, Epidemiology and Informatics, University of Pennsylvania, Philadelphia, 16 Pennsylvania, United States of America 17 ³ Department of Epidemiology, University of Michigan, Ann Arbor, Michigan, United States of America 18 ⁴ Department of Tropical Medicine, Tulane University School of Public Health and Tropical Medicine, 19 20 New Orleans, Louisiana, United States of America 21 22 **Abstract** 23 The city of El Pedregal grew out of a desert, following an agricultural irrigation project in Southern Peru. 24 We conducted door-to-door entomological surveys to document the emergence of triatomines and bed 25 bugs into this new urban environment. We inspected 5,191 households for Triatoma infestans (known 26 locally as the Chirimacha); 21 (0.41%) were infested. These were extremely spatially clustered (Ripley's

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

these, living in poor housing conditions, and the constant migration flow; driven by seeking opportunities

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

3

to work, to buy a house, or to escape from adverse climate effects in other farming regions. Environmental and socioeconomic impacts of large-scale development disproportionately affect vulnerable populations. Design of surveillance systems that account for these contexts is urgently needed to support early detection and control of emergence of vector-borne diseases. Introduction Some great cities of the world have mythical origin stories; the origins of others are more mundane. In many cases the impetus for a *de novo* city is political—national capitals like Washington DC (1) and Brasilia (2) were placed in central locations, and the blank slates allowed for national architectural expression (1). In many other cases, cities grow de novo in response to the exploitation or cultivation of new economic resources, such as rubber (Manaus, Belem, Iquitos) (3), gold (San Francisco) (4), oil (Houston, Texas USA) (5), among many others. The rapid emergence of entirely new cities allows for the close study of the processes of arrival and spread of urban pests, including disease vectors. The foundation of new cities increases human migration, mainly as a workforce, and can act as a persistent social determinant in the processes of introduction and emergence of vector-borne diseases (6,7). Further landscape changes, such as expansion of agricultural frontiers, and changes in land use, provide broad opportunities for new contacts among hosts, vectors, and pathogens (8). Socioeconomic factors and the poorly planned occupation of spaces associated with poverty, are structural processes that can promote conditions for infestation by triatomine bugs, the vectors of the parasite Trypanosoma cruzi, causative agent of Chagas disease. Poor housing conditions, such as poor quality construction materials, cracks on the walls, unplastered surfaces, and presence of animals are factors associated with increased risk of triatomine infestation (9,10). In southern Peru, a new city, El Pedregal, was created de novo following an irrigation project that created numerous job opportunities attracting migrants from local towns as well as more distant regions. We

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

studied the early stages of emergence of the insect vector of Chagas disease, Triatoma infestans, in El Pedregal as well as bed bugs, Cimex sp., which are also competent vectors of Trypanosoma cruzi (11–13). We performed door-to-door entomological surveys throughout El Pedregal, and, in concert with targeted insecticide application, conducted in-depth interviews with affected residents and their neighbors to examine the economic and social forces that led to the rapid growth of the city in a barren desert. **Study site** The study was conducted in Villa El Pedregal (-16.32699, -72.19480), Distrito de Majes, Provincia de Caylloma, Departamento de Arequipa, Perú. Located in the Subtropical Coastal desert (14) the city sits at an altitude of around 500m, and boasts a year-round temperate climate (mean annual temperature of 19°C), but with very little precipitation (150mm annual mean) (15). In 2017, the Peruvian census for Majes district reported 60,108 inhabitants. In the 70's, the Peruvian Government initiated an irrigation project, 'Proyecto Especial Majes-Siguas', with the goal of providing a water source for the irrigation of 60,000 hectares destined to agroindustry in the 'Pampas de Majes'. The first phase of the project consisted of the construction of the Condoroma dam, and channelling (and in some cases tunnelling) water from the the Colca River to the Siguas River, and from there to Pampas de Majes. By the 1980s the project had supported the establishment of agroindustry and processing plants in El Pedregal. These in turn attracted subsistence farmers from different parts of Peru, who came to the region seeking work opportunities (16). Methods Entomological surveys In collaboration with the Ministry of Health of Arequipa we conducted at-the-door interviews and entomological inspections in El Pedregal between October, 2019 and April, 2022, with numerous disruptions due to the Covid-19 pandemic. These surveys were meant to assess the distribution of

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124


125

126

127

128

(Fig. 1).



and in-depth interviews conducted in El Pedregal.

Semi-structured interviews

We visited 97 houses (of 118) treated during the 2nd IRS application. Houses were visited up to three times, in different work hours and during weekends. A total of 65 householders were contacted, and individuals 18 years or older were invited to participate in semi-structured interviews. Sixty individuals agreed to participate, while 5 declined participation. Additionally, two community health workers and 2

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

Spatial Analysis

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

of Try. cruzi in these insects is in line with our observations in the city of Arequipa, where we have not

had parasite-infected insects in areas with very sparse vector infestation (22,23).

Semi-structured interviews Socio-demographic and migration patterns Table 1 presents participants demographic information, work at arrival in El Pedregal, and migration history. The majority of migrants (53.1%) reported repeated migration movements, living in at least one city different from their origin place before their move to El Pedregal.

Participants	Median age (IQR)	
Female (n=44)	39.5 (30.2; 48)	
Male (n=20)	40 (27; 58)	
Place of origin (Department level)	n (%)	
Apurimac	1 (1.56)	
Arequipa	25 (39.06)	
Arequipa (born in El Pedregal)	5 (7.81)	
Ayacucho	1 (1.56)	
Cusco	12 (18.75)	
Moquegua	1 (1.56)	
Huánuco	1 (1.56)	
Puno	18 (28.13)	
Education level	Female n (%)	Male n (%)
Uncompleted primary	4 (9.1)	2 (10)
Completed primary	3 (6.8)	2 (10)
Uncompleted secondary	3 (6.8)	3 (15)
Completed secondary	11 (25)	5 (25)
Institute/technical education	3 (6.8)	1 (5)
Graduated	4 (9.1)	3 (15)
Missing information	16 (36.4)	4 (20)
Work at arrival in El Pedregal	Female n (%)	Male n (%)
Agriculture	12 (27.3)	8 (40)
Agriculture and commerce	3 (6.8)	0
Agribusiness companies	4 (9.1)	3 (15)
Commerce and services	8 (18.2)	3 (15)
Housewives	8 (18.2)	0
Arrived as child	6 (13.6)	3 (15)
Born in El Pedregal	3 (6.8)	2 (10)
Missing information	0	1 (5)
Migration history		n (%)
Born in El Pedregal and has never left		5 (7.8)
Migrated from their place of origin straight to El Pedregal		21 (32.8)
Migrated through more than 1 local apart from their place of origin before moving to El Pedregal		34 (53.1)
Migrated from their place of origin to El Pedregal, left, and returned to El Pedregal (back-forth)		4 (6.3)

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

Reasons for migration to El Pedregal

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

"... Over there [Puno] we suffered from flooding... floods have taken all the crops, that is why we have come, we arrived looking for work... Now everything is being flooded over there..."

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

"... we bought the lot and they told us 'you have to live on the lot because if not we'll take it' ... So all the houses were just 4 esteras [Woven reed mats], there was nothing around the lot. Now people with a little money close their lot in" "At first we bought esteras [Woven reed mats] ... it was hard, I slept on a doubled-over estera, we folded an estera and my children slept below ... It has been hard... like everyone we started like that..." "... Not even the market was like that, it was just some fabrics, the health post also was made of esteras, it wasn't even concrete, just rustic, later they made it of wood." "... Modulo G was empty land, not a single house. Ours was a house of little esteras, with a little room... the school wasn't there, later bit by bit [the urbanization] formed, the rooms were of plywood and later they constructed them. Still people didn't live here, we didn't have any neighbors, we still don't..." Occurrence of dangerous insects and experiences regarding triatomine and bed bug infestation When asked about occurrences of dangerous insects in their community, participants recalled seeing triatomines at their homes among other insects and arachnids (e.g., spiders, scorpions, flies, mosquitoes, ticks). The oldest inhabitants reported triatomine infestation in different neighborhoods of El Pedregal (e.g., Pedregal Norte, Alto, Colina, Malvinas, and in the *parcelas*, particularly in the animal paddocks) and in the neighboring locality of Santa Rita de Siguas in the late 1980's/early 1990's. Frequent insecticide spraying campaigns are recalled having controlled triatomine infestations. Migrants demonstrated knowledge regarding risk factors for triatomine infestation, particularly related to poor housing conditions and breeding animals near human dwellings. Participants knew they had to report infestation at the health centre facility, and posters and flyers displaying pictures of triatomines were mentioned as sources of information about how to identify the insect and how to report infestation.

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

.... I'll tell you when, in 1986, 1987, there were tons. I lived in a parcel, I worked in a parcel, and when I arrived on the parcel itself there were tons and here in Pedregal the people complained a lot about Chirimachas. There was a fumigation company, they fumigated frequently, after that the bugs disappeared." "I knew of the Chirimacha because, when we lived in the valley, we used to see lots.... In the valley [The Valley of Siguas there are strong campaigns for Chagas disease, because there are tons of Chirimachas. The majority of the houses were of adobe and in the adobe lived the Chirimachas. The thing is, in the valley we had all our houses tied to the hillside, and well, the hills were of that red earth and there always were Chirimachas. Although in our house there weren't many because, as I raised roosters, I always fumigated. But one year they went to the neighbor who raised guinea pigs on the roof of her house. She made her house out of cement, supposedly more hygienic than my house that was of adobe, and then they fumigated her house in her guinea pig pens and waaaa how they came down! It looked like the procession of the Señor de los Milagros*, tons. We never saw so many Chirimachas as that day they fumigated the house of our neighbor." *the procession of the Señor de los Milagros is a religious procession that occurs in Peru each October "It was a big black thing, with a kind of swollen belly, I looked at it, I had never seen it before. I saw it on the wall when we turned on the light... What could it be? I didn't know, what animal could it be... As I often went with my daughter to the health post, I went there and I had seen a sign that said 'Chirimacha' and that it produced a disease, Chaga. So I left it there." "... I came to know it here because where I slept was made of esteras, while I slept, they walked all over me, on my face, and I'd turn on the light and see the Chirimachas..."

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

Participants reported having heard about bed bug infestations from their neighbors, friends, and health workers in door-to door prevention campaigns. Inhabitants who had their houses infested by bed bugs, reported use of different products and practices to control infestation, such as: use of "Sapolio® aerosol", "Bolfo®" (Propoxur- carbamates), chlorine, and cleaning activities (taking out the mattress to expose it to the sun). Some of those who experienced infestation believe that bedbugs may transmit diseases but did not know which one. "...I moved recently, a year ago, and bumps appeared... I washed, cleaned and there was nothing, you can't see them "G-d what is this? and a friend told me 'the bed bugs bite you that way in one spot, lots in one spot'. She asked me to show her the bumps and she told me that it was bed bugs, I was frightened, I asked her 'What do I have to do? Then she told me about a sign on a kiosk that said, 'Call when you have Chirimahcas'. I called then and I told them 'Please, I don't know what I have' so they came and inspected everything, the women took apart my bed and there they were, inside. I had to use bleach, boiling water; I didn't want to sleep there anymore". "...Because every little animal passes from neighbor to neighbor...If they had here those bugs I'd use Bolfo®, It doesn't cause harm, and put everything out in the sun, more than anything, put things in the sun, like for fleas, there used to be tons of fleas, we had to put the blankets out in the son, and as the sun warmed the covers the fleas walked off, they left to search for a place to hide themselves... because at night you can't see them... But if you spread the covers out in the sun they leave. They might come from the mountains in the things that travellers bring, they could come in clothing, in the bedding, in anything that they bring they could come and I understand that this little animal adapts itself to any environment, from cold to hot..." Although participants had their houses IRS treated, 79.3% (23/29) reported having used other insecticidal products, such as: Butox® (deltamethrin) against spiders and mosquitoes; Raid® against chicken

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

ectoparasites; Sapolio® against spiders, mosquitoes, flies, triatomines and bed bugs; Baygon® against spiders and flies. Two respondents reported using the agricultural pesticide Lannate® (Methomylcarbamate), applied indoors and outdoors (against triatomine infestations), and applied directly on a dog (for fleas). "... if there were a lot of [triatomines] we definitely have to fumigate...What we used was what we used on the fields, those sprayers the work by pressure, spraying with those, relying on a special product for Chirimachas, I dont remember which. There are various products that have various names, we used 'Lannate [Methomyl], which was a strong poison, we sprayed with that and we had to be outside the house for at least 12 hours." "In La Joya we sprayed, but I can't remember the name [of the insecticide]. We haven't sprayed here... A long time ago I had a dog... It got fleas and couldn't get rid of them... I tried everything until a neighbor told me to use Lannate, which they use on the farms, and then I could kill the fleas." Assessment of Household Development via Satellite Images Houses infested by T. infestans were constructed significantly earlier than control houses (Wilcoxon ranksum test: W=33, p=0.02). No statistical difference was observed between median years of construction when we compared houses infested by *Cimex* sp. and control houses (W=6.5, p=0.07). Spatial Analysis The difference in K functions (KD) indicated significant clustering of both triatomine and bed bug infestations at various spatial scales (Fig. 2). The Ripley's K functions of observed case and control data for both infestation types showed significant clustering across all distances considered (Fig. 2A -2B and 2D-2E), suggesting that the locations of inspected houses exhibit significant clustering at baseline. However, the observed KD functions were positive up to 1000m for triatomines and up to 600m for bed

bugs, which suggests that triatomine and bed bug infestation exhibit clustering over and above any clustering observed in non-infested homes (Fig. 2C and 2F).




Figure 2. Ripley's K functions measuring spatial clustering for triatomine cases (a) and controls (b) and bed bug cases (d) and controls (e), as well as the difference in K functions (KD) for cases and controls for triatomines (c) and bed bugs (f) for a distance of 0-1000m. Black lines indicate observed values, dotted red lines represent null hypothesis of no clustering, and grey shading represent 95% tolerance envelopes. Significant clustering for triatomine infestations is detected at all distances, while clustering for bed bug infestations is detected for distances less than ~600m.

Discussion

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

Throughout history, human populations have employed migration as a survival strategy in response to various challenges, including the impacts of climatic events (26–28), food insecurity (29,30), or pursuing improved living conditions (31). Governmental policies focused on the implementation of extensive infrastructure projects, extractive industries, and agribusiness activities, can generate a need for additional labor force (32), often leading to migration into newly urbanizing areas. The resulting, often haphazard, settlement promotes conditions conducive to disease transmission by connecting hosts, pathogens, and vectors (33). The Chirimacha, Triatoma infestans, successfully invaded urban and peri-urban areas in Arequipa, Peru (10,23); Cochabamba, Bolivia (34); São Paulo, Brazil (35); and Argentina (36,37); complicating regional efforts to eliminate vector-borne Chagas disease (23,37). Here, we describe the migration stories that contributed to population growth in El Pedregal, and we assess residents' perceptions regarding urban development, and their experiences regarding triatomine and bed bug infestations. The Majes irrigation project was designed to supply water resources for the establishment of agribusiness companies in El Pedregal (16). These companies had a huge impact in local farming practices and economy, by introducing new products and agriculture practices, fixing prices for milk, and eventually leading to a shift in the main products in this region: from cattle and dairy to fruits and vegetables. Their establishment required a permanent labor force, which attracted subsistence farmers from across the country. These populations declared different reasons for migrating to El Pedregal; while some of them were seeking opportunities to own a house or to improve their living conditions, others were only searching for a new place to keeping working the land. Subsistence farmers assumed the agribusiness companies would secure a permanent work offer, a different reality than they faced elsewhere, where their farming practices were vulnerable to seasonality, local trade dynamics, and to adverse climate events. Coming to a desert, these migrants arrived with few resources; only enough to live in shelters constructed

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

of woven reeds and other poor-quality materials. Over time, working in the fields with a fairly consistent income, they constructed more permanent houses. Most of the houses, however, continued to incorporate precarious materials, such as unmortared stone. The prolonged precarity of houses in the area provided stable environments for introduced triatomines to establish colonies. Through interviews and satellite images, we assessed house construction and the expansion of the settlement over space and time. We observed that not all constructed houses were immediately inhabited, as evidenced by the nearly one-to-one inhabitant-to-dwelling ratio in the 2017 Peruvian census for Majes district, which reported 60,108 inhabitants and 58,298 dwellings registered (38). This low ratio is likely due to the large number of dwellings that have been 'staked out' but not fully constructed. In our door-todoor surveys in the fairly established areas, we found that approximately 15% of houses were uninhabited. The long interval between the tenancy of space and its effective use was also mentioned by participants in the in-depth interviews. Uninhabited neighboring houses might act as a limiting factor for triatomine dispersion, as few sources of food are available in these structures (39). The uneven construction may thereby have restricted triatomines to a small number of households, spatially clustered, in the older sections of El Pedregal. Households infested with T. infestans tended to be older, in areas where migrants had come many years ago. In contrast, infestation by bed bugs presented a different pattern, more dispersed in comparison to those infested by T. infestans. Bed bug infestation was not associated with the age of the home, nor the level of construction of the structure. Although bed bug infested households were more spatially clustered than would occur by chance, they were much more widely dispersed across El Pedregal than the triatomines. Bed bugs are likely less dependent on active dispersal in El Pedregal, and, by being carried

passively in belongings, less affected by the barriers of uninhabited houses, barren vacant lots, and streets.

Awareness of bed bug infestation was amplified through communication amongst community members and health workers. Still, knowledge regarding infestation control is low, and indiscriminate use of

522

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

insecticides products were reported against domestic infestations of bed bugs, triatomines, and other synanthropic insects. Similar results were observed in peri-urban neighborhoods of the metropolitan Arequipa (40). Although householder application of insecticides was associated with lower risk of infestation by *T. infestans* in endemic area in Argentina (36), insecticide misuse might decrease susceptibility of vector populations to these and related products (41). Aerosol insecticide formulations, in particular, are easily available for purchase in Peru, and their use is widespread. These products often combine different insecticides and synergists agents and are unlikely to control triatomine infestations effectively. Use of aerosol products has shown to increase resistance to pyrethroids in populations of Aedes aegypti in Mexico (41). Health educational campaigns to improve community knowledge regarding methods to control synanthropic insects, particularly for bed bug infestations, could help to prevent misuse of insecticide products. However, until a safe and affordable alternative is made available for bed bugs, as it is for triatomines, it is hard to imagine the bed bug situation improving. We suggest, based on our synthesis of spatial entomological data and our interviews, that the *Chirimacha* populations we detected and eliminated arrived at El Pedregal rather recently, and were hindered from dispersing by the numerous uninhabited lots in the growing city. It is possible that the insects are remnants of much earlier infestation processes that our participants recounted, but we believe that alternative explanation is less likely, because it is hard to imagine the insects being constrained to such small foci of infestations for such a long time. Indeed, in the neighboring district of La Joya, *Chrimachas* redispersed following control in the 1990s to infest over a quarter of households (42). It is also possible that the actions of residents applying their own insecticides prevented the further dissemination of Chirimachas, though again it is hard to imagine those activities would be effective and uniform enough for long term control. If we are correct our results are concerning—they suggest that as El Pedregal develops further, and the empty lots are filled in, the new city will be highly susceptible to triatomine infestations.

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

The historic and current triatomine infestations, which occurred about 30 years apart, seem to differ in their dispersal and density and the settings of space occupation. The former is recalled in interviews as a widespread event, affecting different neighborhoods, houses, and animal corrals at high densities. Housing conditions in those days were described as improvised constructions, using poor quality materials. The infestations we observed in Pedregal, however, were extremely spatially clustered, of low insect densities and only in a few houses that were constructed many years ago. Still, those houses were infested at low densities, and infestation spread was limited, likely by the environmental barriers created by unoccupied, or very sparsely occupied, lots, interspersed with more permanent and, for the insects, habitable, dwellings. Housing conditions, particularly construction materials and crowding, which provide availability of refuges and source of food for triatomines, are limiting factors for domestic infestation by triatomines, density of its colonies, and infection by Try. cruzi across many contexts (9,10,43–45). In the interval between the two (re-)introduction events, changes in housing materials and conditions in El Pedregal may not have made houses entirely refractory to infestation, but they hampered infestation, dispersal, and density of the insects. Large government decisions involving water and changes in land use can have outsized impacts on the lives of farmers and other laborers, and these impacts can reverberate through society, leading to mass migration and, ultimately, increasing populations' vulnerability to disease (46–49). The cost of these decisions is ultimately borne by those with the least recourse to bear them (50,51). Migration has been also driven by adverse climate conditions, and this phenomenon is expected to increase in the following years. It is urgent to advocate to a shift in governance that contemplates both a responsible use of common natural resources, and a preparedness to ensure food security to populations affected and displaced by climatic events. Ironically for a city that grew around an irrigation project, many residents of El Pedregal lack access to clean water (16). Without running water many resort to storing water in barrels, which creates risk for the

573

574

575

576

577

578

579

580

581

582

583

584

585

586

587

588

589

590

591

592

593

594

595

emergence of Aedes aegypti and Dengue virus (52). Emergence of vectors highlight the importance that communities and health authorities to remain hypervigilant, as the complex forces of migration and poor housing conditions makes El Pedregal especially vulnerable to vector-borne disease. Acknowledgements The authors gratefully acknowledge the contributions of the Ministerio de Salud del Perú (MINSA), the Dirección General de Salud de las Personas (DGSP), the Estrategia Sanitaria Nacional de Prevención y Control de Enfermedades Metaxénicas y Otras Transmitidas por Vectores (ESNPCEMOTVS), the Dirección General de Salud Ambiental (DIGESA), the Gobierno Regional de Arequipa, the Gerencia Regional de Salud de Arequipa (GRSA), the Pan American Health Organization (PAHO/OPS), the Canadian International Development Agency (CIDA). We also thank Amparo Toledo, Claudia Arevalo Nieto and Laura Tamayo for their valuable contributions to these studies. References Stephenson, Richard W. A plan whol[1]y new : Pierre Charles L'Enfant's plan of the City of Washington. 1°. United States Govt Printing Office; 1993. Lispector, Clarice. Brasilia. In: Todos os Contos. 1º. Editora Rocco; 2016. p. 656. Barham BL, work(s): OTCR. Reinterpreting the Amazon Rubber Boom: Investment, the State, and Dutch Disease. Lat Am Res Rev. 1994;29(2):73–109. Available from: http://www.jstor.org/stable/2503594 St. Clair DJ. The Gold Rush and the Beginnings of California Industry. Calif Hist. 1998;77(4):185– 208. Available from: https://ch.ucpress.edu/cgi/doi/10.2307/25462514 5. Lee J. The regional economic impact of oil and gas extraction in Texas. Energy Policy. 2015; 87:60– 71. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0301421515300768

- 6. Barnett ED, Walker PF. Role of Immigrants and Migrants in Emerging Infectious Diseases. Med Clin 596 597 North Am. 2008;92(6):1447–58. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0025712508000941 598 599 Wilke ABB, Benelli G, Beier JC. Anthropogenic changes and associated impacts on vector-borne 600 diseases. Trends Parasitol. 2021;37(12):1027–30. Available from: 601 https://linkinghub.elsevier.com/retrieve/pii/S1471492221002385 602 Baker RE, Mahmud AS, Miller IF, Rajeev M, Rasambainarivo F, Rice BL, et al. Infectious disease in 603 an era of global change. Nat Rev Microbiol. 2022;20(4):193–205. Available from: 604 https://www.nature.com/articles/s41579-021-00639-z 605 9. Samuels AM, Clark EH, Galdos-Cardenas G, Wiegand RE, Ferrufino L, Menacho S, et al. 606 Epidemiology of and Impact of Insecticide Spraying on Chagas Disease in Communities in the Bolivian Chaco. McCall PJ, editor. PLoS Negl Trop Dis. 2013;7(8):e2358. Available from: 607 https://dx.plos.org/10.1371/journal.pntd.0002358 608 10. Levy MZ, Bowman NM, Kawai V, Waller LA, Cornejo del Carpio JG, Cordova Benzaquen E, et al. 609 Periurban Trypanosoma cruzi-infected Triatoma infestans, Arequipa, Peru. Emerg Infect Dis. 610 2006;12(7):1345–52. Available from: http://wwwnc.cdc.gov/eid/article/12/09/05-1662_article.htm 611 612 11. Salazar R, Castillo-Neyra R, Tustin AW, Borrini-Mayorí K, Náquira C, Levy MZ. Bed Bugs (Cimex 613 lectularius) as Vectors of Trypanosoma cruzi. Am J Trop Med Hyg. 2015;92(2):331-5. Available from: https://ajtmh.org/doi/10.4269/ajtmh.14-0483 614
- 12. Jorg ME, Natula CN. Cimex lectularius, L. (La chinche común de cama) transmisor de Trypanosoma
 cruzi. Prensa Médica Argent. 1982;69:528–33.

- 13. Jörg ME. Cimex lectularius, L. (la chinche comun de cama) trasmisor de Trypanosoma cruzi. Rev
- 618 Soc Bras Med Trop. 1992;25(4):277–8. Available from:
- 619 http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0037-
- 620 86821992000400012&lng=es&tlng=es
- 621 14. SINIA Sistema Nacional de Información Ambiental. Mapa Nacional de Ecosistemas del Perú.
- 622 Available from: https://sinia.minam.gob.pe/mapas/mapa-nacional-ecosistemas-peru
- 623 15. SENAMHI Sistema Nacional de Meteorología e Hidrología de Perú. Available from:
- https://www.senamhi.gob.pe/?p=mapa-climatico-del-peru
- 625 16. Erwin A, Ma Z, Popovici R, Salas O'Brien EP, Zanotti L, Silva CA, et al. Linking migration to
- 626 community resilience in the receiving basin of a large-scale water transfer project. Land Use Policy.
- 627 2022;114:105900. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0264837721006232
- 628 17. Gutfraind A, Peterson JK, Billig Rose E, Arevalo-Nieto C, Sheen J, Condori-Luna GF, et al.
- 629 Integrating evidence, models and maps to enhance Chagas disease vector surveillance. Basáñez MG,
- editor. PLoS Negl Trop Dis. 2018;12(11):e0006883. Available from:
- 631 https://dx.plos.org/10.1371/journal.pntd.0006883
- 632 18. Dedoose Software. Los Angeles, CA: SocioCultural Research Consultants, LLC; 2021. Available
- from: www.dedoose.com
- 19. Spatial Statistics. In: Spatial Statistics [Internet]. Hoboken, NJ, USA: John Wiley & Sons, Inc.; 2005.
- p. i–ix. (Wiley Series in Probability and Statistics). Available from:
- https://onlinelibrary.wiley.com/doi/10.1002/0471725218.fmatter
- 637 20. Baddeley A, Turner R. spatstat □: An R Package for Analyzing Spatial Point Patterns. J Stat Software.
- 638 2005;12(6). Available from: http://www.jstatsoft.org/v12/i06/

- 21. Joshua French. smacpod: : Statistical Methods for the Analysis of Case-Control Point Data. 2022. 639 22. Barbu CM, Buttenheim AM, Pumahuanca MLH, Calderón JEQ, Salazar R, Carrión M, et al. Residual 640 Infestation and Recolonization during Urban Triatoma infestans Bug Control Campaign, Peru1. 641 Emerg Infect Dis. 2014;20(12):2055-63. Available from: http://wwwnc.cdc.gov/eid/article/20/12/13-642 643 1820 article.htm 644 23. Delgado S, Ernst KC, Pumahuanca MLH, Yool SR, Comrie AC, Sterling CR, et al. A country bug in 645 the city: urban infestation by the Chagas disease vector Triatoma infestans in Arequipa, Peru. Int J Health Geogr. 2013;12(1):48. Available from: https://ij-646 647 healthgeographics.biomedcentral.com/articles/10.1186/1476-072X-12-48 24. Wilfredo Gonzales-Valero. Hazards to food caloric availability and coverage per capita due to climate 648 649 change in the Puno region, Peruvian Altiplano: Challenges in food security and sovereignty. Food 650 Energy Security. 2018;7(2). Available from: https://onlinelibrary.wiley.com/doi/full/10.1002/fes3.134 651 25. PubChem. Lannate. Available from: https://pubchem.ncbi.nlm.nih.gov/compound/5353758 652 26. McMichael C, Barnett J, McMichael AJ. An Ill Wind? Climate Change, Migration, and Health. Environ Health Perspect. 2012;120(5):646–54. Available from: 653 654 https://ehp.niehs.nih.gov/doi/10.1289/ehp.1104375 655 27. McLeman RA, Hunter LM. Migration in the context of vulnerability and adaptation to climate change: insights from analogues: Migration and adaptation to climate change. Wiley Interdiscip Rev 656
- 28. Louis N, Mathew TH, Shyleen C. Migration as a Determinant for Climate Change Adaptation:
 Implications on Rural Women in Muzarabani communities, Zimbabwe. In: Leal Filho W, Luetz J,

Clim Change. 2010;1(3):450–61. Available from: https://onlinelibrary.wiley.com/doi/10.1002/wcc.51

657

Ayal D, editors. Handbook of Climate Change Management: Research, Leadership, Transformation.

- 661 Cham: Springer International Publishing; 2020. p. 1–24. Available from: https://doi.org/10.1007/978-
- 662 3-030-22759-3 273-1
- 663 29. Sadiddin A, Cattaneo A, Cirillo M, Miller M. Food insecurity as a determinant of international
- 664 migration: evidence from Sub-Saharan Africa, Food Secur. 2019;11(3):515–30. Available from:
- https://link.springer.com/10.1007/s12571-019-00927-w
- 666 30. Carney MA, Krause KC. Immigration/migration and healthy publics: the threat of food insecurity.
- Palgrave Commun. 2020;6(1):93. Available from: https://www.nature.com/articles/s41599-020-0461-
- 668 0
- 31. Gustafsson C. "For a better life ... "A study on migration and health in Nicaragua'. Glob Health
- 670 Action. 2018;11(1):1428467. Available from:
- https://www.tandfonline.com/doi/full/10.1080/16549716.2018.1428467
- 672 32. Bebbington AJ, Humphreys Bebbington D, Sauls LA, Rogan J, Agrawal S, Gamboa C, et al.
- Resource extraction and infrastructure threaten forest cover and community rights. Proc Natl Acad
- 674 Sci. 2018;115(52):13164–73. Available from: https://pnas.org/doi/full/10.1073/pnas.1812505115
- 675 33. Alirol E, Getaz L, Stoll B, Chappuis F, Loutan L. Urbanisation and infectious diseases in a globalised
- world. Lancet Infect Dis. 2011;11(2):131–41. Available from:
- https://linkinghub.elsevier.com/retrieve/pii/S1473309910702231
- 678 34. Medrano-Mercado N, Ugarte-Fernandez R, Butrón V, Uber-Busek S, Guerra H, Araújo-Jorge TC de,
- et al. Urban transmission of Chagas disease in Cochabamba, Bolivia. Mem Inst Oswaldo Cruz.
- 680 2008;103(5):423–30. Available from: http://www.scielo.br/scielo.php?script=sci arttext&pid=S0074-
- 681 02762008000500003&lng=en&tlng=en

683

684

685

686

687

688

689

690

691

692

693

694

695

696

697

698

699

700

701

702

703

704

35. Ceretti-Junior W, Vendrami DP, de Matos MO, Rimoldi-Ribeiro A, Alvarez JV, Marques S, et al. Occurrences of triatomines (Hemiptera: Reduviidae) and first reports of Panstrongylus geniculatus in urban environments in the city of Sao Paulo, Brazil. Rev Inst Med Trop São Paulo, 2018; 60: e33. Available from: https://www.scielo.br/j/rimtsp/a/zzFKXqJCzFGSpnRfxMfnjSm/?lang=en 36. Gaspe MS, Fernández M del P, Cardinal MV, Enriquez GF, Rodríguez-Planes LI, Macchiaverna NP, et al. Urbanisation, risk stratification and house infestation with a major vector of Chagas disease in an endemic municipality of the Argentine Chaco. Parasit Vectors. 2020;13(1):316. Available from: https://parasitesandvectors.biomedcentral.com/articles/10.1186/s13071-020-04182-3 37. Provecho YM, Fernández M del P, Salvá L, Meli S, Cano F, Sartor P, et al. Urban infestation by Triatoma infestans (Hemiptera: Reduviidae), an overlooked phenomena for Chagas disease in Argentina. Mem Inst Oswaldo Cruz [Internet]. 2021 [cited 2023 Mar 31];116:e210056. Available from: http://www.scielo.br/scielo.php?script=sci arttext&pid=S0074-02762021000100402&tlng=en 38. Instituto Nacional de Estadistica e Informatica. Arequipa- resultados definitivos. 2018. Available from: https://www.inei.gob.pe/media/MenuRecursivo/publicaciones_digitales/Est/Lib1551/04TOMO_01.pd f 39. Levy MZ, Barbu CM, Castillo-Neyra R, Quispe-Machaca VR, Ancca-Juarez J, Escalante-Mejia P, et al. Urbanization, land tenure security and vector-borne Chagas disease. Proc R Soc B Biol Sci.2014; 281(1789):20141003. Available from: https://royalsocietypublishing.org/doi/10.1098/rspb.2014.1003 40. Larson AJ, Paz-Soldán VA, Arevalo-Nieto C, Brown J, Condori-Pino C, Levy MZ, et al. Misuse, perceived risk, and safety issues of household insecticides: Qualitative findings from focus groups in Arequipa, Peru. Hayden M, editor. PLoS Negl Trop Dis. 202;15(5):e0009251. Available from: https://dx.plos.org/10.1371/journal.pntd.0009251

- 705 41. Gray L, Florez SD, Barreiro AM, Vadillo-Sánchez J, González-Olvera G, Lenhart A, et al.
- Experimental evaluation of the impact of household aerosolized insecticides on pyrethroid resistant
- Aedes aegypti. Sci Rep. 2018;8(1):12535. Available from: https://www.nature.com/articles/s41598-
- 708 018-30968-8
- 42. Delgado S, Castillo Neyra R, Quispe Machaca VR, Ancca Juárez J, Chou Chu L, Verastegui MR, et
- al. A History of Chagas Disease Transmission, Control, and Re-Emergence in Peri-Rural La Joya,
- Peru. Gürtler RE, editor. PLoS Negl Trop Dis. 2011;5(2):e970. Available from:
- 712 https://dx.plos.org/10.1371/journal.pntd.0000970
- 43. Campbell-Lendrum D, Angulo V, Esteban L, Tarazona Z, Parra G, Restrepo M, et al. House-level
- 714 risk factors for triatomine infestation in Colombia. Int J Epidemiol.;36(4):866–72. Available from:
- 715 https://academic.oup.com/ije/article-lookup/doi/10.1093/ije/dym065
- 716 44. Cardona ÁS, Angulo V, Parra-Henao G, Quirós-Gómez O, Alexander N. House-Level Risk Factors
- 717 for Triatoma dimidiata Infestation in Colombia. Am J Trop Med Hyg.;92(1):193–200. Available
- 718 from: https://ajtmh.org/doi/10.4269/ajtmh.14-0273
- 719 45. Schofield CJ, Marsden PD. Efecto del revoque de las paredes sobre una población domestica de
- 720 Triatoma infestans. Bol Oficina Sanit Panam. 1982; 93(1). Available from:
- https://iris.paho.org/bitstream/handle/10665.2/15992/v93n1p3.pdf?sequence=1
- 46. Gates MC, Boston RC. Irrigation linked to a greater incidence of human and veterinary West Nile
- virus cases in the United States from 2004 to 2006. Prev Vet Med. 2009;89(1–2):134–7. Available
- from: https://linkinghub.elsevier.com/retrieve/pii/S0167587708002535
- 725 47. Kibret S, McCartney M, Lautze J, Nhamo L, Yan G. The impact of large and small dams on malaria
- 726 transmission in four basins in Africa. Sci Rep. 2021;11(1):13355. Available from:
- 727 https://www.nature.com/articles/s41598-021-92924-3

- 48. Lerer LB, Scudder T. Health impacts of large dams. Environ Impact Assess Rev.1999;19(2):113–23.
- Available from: https://linkinghub.elsevier.com/retrieve/pii/S0195925598000419
- 730 49. Rohr JR, Barrett CB, Civitello DJ, Craft ME, Delius B, DeLeo GA, et al. Emerging human infectious
- diseases and the links to global food production. Nat Sustain. 2019;2(6):445–56. Available from:
- 732 https://www.nature.com/articles/s41893-019-0293-3
- 733 50. Randell H, Klein P. Hydropower Development, Collective Action, and Environmental Justice in the
- 734 Brazilian Amazon. Soc Nat Resour. 2021;34(9):1232–49. Available from:
- 735 https://www.tandfonline.com/doi/full/10.1080/08941920.2021.1948649
- 736 51. Zhouri A, Pascoal WV. From the muddy banks of the Watu: The Krenak and the Rio Doce mining
- disaster in Brazil. In: Current Directions in Water Scarcity Research. Elsevier; 2022. p. 145–65.
- Available from: https://linkinghub.elsevier.com/retrieve/pii/B978012824538500008X
- 739 52. Heukelbach J, de Oliveira FAS, Kerr-Pontes LRS, Feldmeier H. Risk factors associated with an
- outbreak of dengue fever in a favela in Fortaleza, north-east Brazil. Trop Med Int Health.
- 741 2001;6(8):635–42. Available from: http://doi.wiley.com/10.1046/j.1365-3156.2001.00762.z