

# **Factors associated with exclusive breastfeeding among mothers of children under two years of age in Dalit community, Rajbiraj Municipality, Saptari, Nepal**

*Neha Kumari Das<sup>\*1</sup>, Nirmal Duwadi<sup>1</sup>, Ramchandra Sinha<sup>1</sup>, Alisha Dahal<sup>1</sup>*

<sup>1</sup>Department of Public Health, Nobel College, Pokhara University, Kathmandu, Bagmati Province, Nepal

\*Corresponding author

E-mail: [ndas03823@gmail.com](mailto:ndas03823@gmail.com)

## *Abstract*

The prevalence of exclusive breastfeeding (EBF) is suboptimal in Nepal and very low in Madhesh province. Dalits are commonly recognized for experiencing economic exploitation, a lack of political representation, social marginalization, educational disadvantage, being classified as untouchables, and enduring the denial of basic human dignity and social justice.

Objective of this study is to assess the proportion of mothers of children under 2 years of age, practicing exclusive breastfeeding and associated factors in the Dalit community of Rajbiraj Municipality, Saptari, Nepal.

The study utilized an analytical, cross-sectional design by using a semi-structured questionnaire to 156 Dalit mothers of children under 2 years of age in Rajbiraj Municipality, Saptari, Nepal. In the bivariate analysis, which focuses on exploring the connection between independent variables and dependent variables, chi-square statistics were utilized. Subsequently, the independent variables that displayed significance in the bivariate analysis were included in the multivariate logistic regression.

The estimated prevalence of exclusive breastfeeding in the population was 43.6%. Mother's occupation (adjusted odds ratio (AOR) = 4.459; CI = 1.444 -13.767), smoking habit (AOR = 2.755; CI = 1.120 – 6.774), colostrum milk feeding (AOR = 12.472; CI = 3.253 – 47.823), number of times visit the health center for ANC (AOR = 2.333; CI = 1.040 – 5.233) were positively associated with exclusive breastfeeding, whereas, sex of the child, type of family, knowledge about breastfeeding, counselling on EBF in ANC and/or PNC visit were also associated with exclusive breastfeeding. Among the respondents who did not practice exclusive breastfeeding, the common reasons cited were trouble initiating milk flow (23.7%), insufficient breast milk production (21.7%), breast milk not satisfying the baby (20.9%), domestic work burden (12%), difficulties with infant sucking or latching (6.4%), and the baby being unable to be breastfed owing to sickness (4.8%).

Maternal education, occupation, colostrum milk feeding, and antenatal care visits were identified as important influencers of exclusive breastfeeding. Healthcare professionals, policymakers, and community leaders can utilize these insights to formulate effective strategies and design intervention that encourage and support exclusive breastfeeding among Dalit mothers, ultimately improving the health and well-being of the infants of the marginalized community.

## *Introduction*

### *Background*

Nepal is a nation characterized by rich sociocultural diversity, is spread across seven provinces and three ecological regions: mountain, hills and Terai. Within this diverse tapestry, Nepal is home to 23 distinct caste and ethnic groups.(1)

However, amidst this diversity, one group, the Dalits faces significant challenges. In the hierarchical Hindu caste system, Dalits often endure economic exploitation, political marginalization, social degradation, educational disadvantages, and stigma of untouchables. These injustices compound to deny them basic human dignity and social justice, leading to a life marked by limited income opportunities, reliance on daily wages, livestock sales, and social security allowances. Despite constituting 13.6% of the population, Dalits face formidable barriers to achieving an equitable and prosperous life. (2, 3)

Despite ongoing social disparities, it's crucial to grasp the vital importance of breastfeeding—an inclusive practice with profound implications for infant health. Breast milk, often hailed as the healthiest nourishment for infants, stands as a precious gift from mothers, meeting all the energy and nutritional needs of a baby.(4) Recognizing the immense value of breastfeeding, the World Health Organization (WHO) advocates for exclusive breastfeeding, wherein infants receive only breast milk during the first six months of life, without any additional food or liquids. (5) Exception Only oral rehydration solution, authorized drugs, vitamins, and minerals for exclusively breastfed infants. (6) Exclusive breastfeeding (EBF) plays a vital role in reducing child morbidity and mortality, as well as contributing to the effective management of healthcare expenses in society. It serves as a significant public health strategy to improve the health of both mothers and children. (7)

The benefits of EBF are far – reaching, encompassing significant protection against various diseases, including infections, pneumonia, sudden infant death syndrome, malocclusion, diarrhea, diabetes, asthma, heart disease, and obesity. It significantly reduce the risk of infant mortality due to infections in the first two years of life. (8) Additionally, EBF supports healthy brain development and is associated with better intelligence tests results in children and adolescents. (5) It fosters a close bond between women and their children while reducing the risk of breast and ovarian cancer. (9) Despite these compelling benefits, the prevalence of exclusive breastfeeding in Nepal remains alarming low. (10-12)

To promote a child's survival, growth, and development, it is crucial to follow the right feeding and care practices. According to the annual report, the percentage of exclusively

breastfeeding infants in 2021 was 36.9%, which was significantly lower than the NDHS 2022 figure of 56.4% and the most recent MICS 2019 figure of 62.1%. According to a breakdown by province, Madhesh Province has the lowest percentage of infants aged 0 to 6 months who are exclusively breastfed, at 26%. Vulnerable communities, such as Dalits, are the least likely to exclusively breastfeed, and rates of exclusive breastfeeding fall far short of the national target due to humiliation and discrimination. (10) The Dalits of Nepal are the most disadvantaged caste group and have benefited least from the advances in maternal and child health services. (13) For example, ANMs, who are front-line healthcare providers from non-Dalit castes, avoid visiting Dalit homes and are hesitant to handle infants or pregnant women because it requires close physical contact.(14)

The purpose of this study is to measure the factors associated with exclusive breastfeeding among Dalit mothers with children under the age of two. We anticipate that the findings of this research will serve as a valuable guideline for policymakers, stakeholders, and individuals in planning feasible interventions and strengthening existing factors related to exclusive breastfeeding. Ultimately, these effects aim to help Nepal achieve its national exclusive breastfeeding targets, thereby aligning with the broader aspirations of the Sustainable Development Goals for 2030. This research contributes to improving nutrition status (SDG2), prevents child mortality and lowers the risk of non-communicable disease (SDG3), and supports cognitive development and education (SDG4). Additionally, it underscores the role of breastfeeding in ending poverty, promoting economic growth, and reducing inequalities. Through a comprehensive exploration of exclusive breastfeeding within the context of Dalit communities, this study holds the potential to transform the lives of mothers and children, fostering a brighter and healthier future for all.

## ***Materials and Methods***

### ***Study design***

The cross-sectional analytical study design was used. Data was collected from 23<sup>rd</sup> February 2023 to 8<sup>th</sup> March 2023.

### ***Study Settings***

The study was conducted in Dalit community of Rajbiraj Muicipality, Saptari District, Madhesh Province of Nepal.

A mid-sized municipality called Rajbiraj is found in the southern region of Nepal's Madhesh Province. Rajbiraj is the district headquarters of Saptari and the eighth-largest city in the province. The total population of Rajbiraj Municipality is 68396, which is a larger population in comparison to other municipalities in the Saptari district. (15) In this district, there is an all-Dalit population. As a result, this setting was ideal for conducting research.

### ***Study Population and Sampling Frame***

#### ***Study Population:***

The study population for this study was Dalit mothers of under 2 year children among Rajbiraj Municipality, Saptari, Nepal.

#### ***Sampling Frame:***

The list of mothers with children under 2 years was obtained from MIS the section of Rajbiraj Municipality.

## *Sample Size*

The Annual Report 2077/78 of Madhesh Province (DoHS, 2078) revealed that the overall prevalence of exclusive breastfeeding among children aged 0 to 6 months was 26%. While calculating the sample size, the allowable error was kept at 5% and the confidence level at 95%. The required sample size of the proposed study was calculated based on the aforesaid prevalence data.

Sample size was computed as

$$\text{Sample size} = (z^2 p q / d^2) \quad (16)$$

Where,

Z= 1.96 reliability coefficient at alpha 0.05.

Prevalence (P) = 26% = 0.26

q = 1-P = 1-0.26 = 0.74

Allowable error (d) = 5% = 0.05

Applying the formula,

$$n = \frac{(1.96)^2 (0.26 * 0.74)}{(0.05)^2} = 295.65 \sim 296$$

The required sample size for calculating the infinite population in calculated by considering the finite population,

Now, for a finite population,

Finite population (N) = 274 (which is the number of mothers of under 2 year children in Rajbiraj Municipality)

Sample size for finite population,


$$n = \frac{n_o}{1 + \frac{n_o}{N}}$$

$$n = 296/1 + 296/274 \\ = 142.30 \sim 142$$

Adding non-response rate of 10%, the required sample size = 156.2 ~ 156

### ***Sampling Techniques/Procedures***

Simple random sampling technique was used to select the study participants. The sample frame for this study is 274 populations.



***Figure 3: Sampling technique and procedure for the selection of the study participants***

### ***Data Collection Tools and Techniques***

The data was collected using a semi-structured questionnaire, validated by a thorough review of the literature relevant to the study's topic, and seeking the advice of a research expert. The research questionnaire was translated into Nepali and further translated into English. The questionnaire was divided into 2 parts. Information on sociodemographic

characteristics was included in the first section, while exclusive breastfeeding was covered in the second. The feasibility of the questionnaire was maintained by pre-testing in the population of the sample. The actual data was gathered after the questionnaire has been improved. The face-to-face interview was conducted with mothers of under two years' children in the Dalit community, Rajbiraj Municipality. Interviews were conducted by myself and an enumerator with a public health background. For illiterate participants, a witness was present during the consenting process, and the witness signed the informed consent form.

### ***Operational Definition:***

The following definitions were applied, consistent with WHO definitions:

**Exclusive Breastfeeding:** If an infant less than six months old took only breast milk and no additional food, water, or other liquids (with the exception of medicine and vitamins, if needed) continuing for the recommended 6 months.

**Pre-lacteal feeding:** Pre-lacteal feeding refers to the act of giving anything other than breast milk to an infant within the first three days after birth.

**Dalit Community:** Kalar, Kakaihiya, Kori, Khatik, Khatbe, Chamar, Chidimar, Dom, Tatma, Dusadh, Dhobi, Pattharkatta, Pasi, Bantar, Mushar, Mestar, and Sarvanga are Dalit castes of Madhesi origin who live in Rajbiraj municipality of Saptari District, Nepal. (3)

### ***Validity and Reliability of Tools***

The content validity of the instrument was established by consultation with the study advisor's subject experts, peers, and public health faculty research committee members at Nobel College. The questionnaire was modified as needed based on insightful opinions and ideas. Both English and Nepali was used in the design of the instrument. The original version of the tool was checked to ensure consistency in the meaning provided by the English and Nepali versions. After the interview is over, the completed questionnaire was carefully reviewed. A pretesting was conducted among 10% of the study sample before the main research to check feasibility and improve on the research tools. A pretesting was conducted among 16 randomly selected mothers of children under 2 years of age in Sambhunath municipality. Cronbach's alpha coefficient was calculated selecting 16 questionnaires and used to measure consistency of tool. The Cronbach's alpha of  $\alpha$  calculated was 0.806 which is greater than 0.7 and considered as standard and statistically acceptable.



## ***Data Management and Analysis***

A data management plan was developed prior to data collection to make sure that the data are collected in the right format, are effectively organized, and are better interpreted. The day after the data is collected, the acquired data was cleaned, coded, and correctly entered in MS Excel. Data was manually verified, entered, and exported from MS Excel to IBM SPSS 20 for additional analysis. Analyses was performed for descriptive statistics like frequency, percentage, measures of central tendency, and measures of dispersion. Frequency tables was used for categorical variables, while mean and standard deviation was calculated for continuous variables. For the analysis involving testing for the association of independent variables with dependent variables, bivariate analysis (chi-square statistics and a p-value at  $<0.05$ ) was used. Those that are significant in the bivariate analysis was further preceded by multivariate logistic regression. Independent variables with a P-value  $<0.05$  was entered into the multivariate analysis.

## ***Ethical Consideration***

Before collecting data, administrative approval was obtained from the concerned authority at Nobel College. Then, formal approval and ethical clearance was obtained from the Institutional Review Committee of Nobel College. Permission was obtained from the Rajbiraj Municipality Office for conducting the purposed study. Written Informed consent was obtained from the participants prior to data collection, assuring voluntary participation and that the individual's identity was not disclosed in the report. Confidentiality was maintained throughout the study period, and the collected data was used only for research purposes.

## *Results/Findings*

This chapter deals with the result tabulation and interpretation of the data obtained from 156 Dalit mothers of under 2 years' children who voluntarily participated in his study.

### *Univariate Analysis*

**Table 3: Sociodemographic variable (n=156)**

The study, encompassing 156 respondents, revealed that the mothers' ages ranged from 19 to 36 years, with a mean age of 24.59 years (confidence interval: 24.03 to 25.15). Over 60% fell within the 19 to 24-year age group. Children's ages ranged from 6 to 24 months, with a mean age of 11.39 months (confidence interval: 10.64 to 12.14). The majority of respondents (64.1%) had children in the 6 to 12-month age group. In terms of gender, 59.6% were male. Most respondents were homemakers (84%), with husbands predominantly working as laborers (57.7%). Most of mothers had completed secondary education (48.1%) while Husbands' education receiving secondary education (55.1%). Nearly 59% belonged to nuclear families, and the majority had two children (41.7%). Lifestyle-wise, 71.8% didn't smoke, and 85.9% abstained from alcohol.

<b>Variables</b>	<b>Frequency</b>	<b>Percent</b>
	<b>(n = 156)</b>	<b>(%)</b>
<b>The age group of Mother</b>		
19 to 24 year	95	60.9
25 to 30	43	27.6
31 to 36 year	18	11.5
<b>Mean: 24.59 SD: 3.541 (CI = 24.03 to 25.15)</b>		

### **Age of Child in Month**

6 to 12 Month	100	64.1
13 to 18 Months	45	28.8
19 to 24 Month	11	7.1

**Mean: 11.39 SD: 4.721**

**(CI= 10.64 to 12.14)**

### **Sex of Child**

Female	63	40.4
Male	93	59.6

### **Mothers Occupation**

Employed	25	16
Homemaker	131	84

### **Fathers Occupation**

Private or government worker	27	17.3
Labor worker	91	58.3

Un - Employed	16	10.3
Foreign Worker	22	14.1
<b>Mothers Education</b>		
Cannot read and write	24	15.4
Read and write only	20	12.8
Primary education (1-5)	37	23.7
Secondary Education ( 6 – 10 ) and Above	75	48.1
<b>Fathers Education</b>		
Cannot read and write	18	11.5
Read and write only	15	9.6
Primary education (1 – 5)	37	23.7
Secondary education (6 – 10) and above	86	55.1
<b>Family Type</b>		
Joint/extended	64	41
Nuclear	92	59

### **Number of Children**

One Child	64	41.0
Two Child	65	41.7
Three and above	27	17.3

### **Mother's Smoking habit**

Yes	44	28.2
No	112	71.8

### **Mother's Alcohol Drinking Habit**

Yes	22	14.1
No	134	85.9

---

### **Table 4: Knowledge about Exclusive breastfeeding**

Table 4 illustrates the distribution of the population's knowledge on breastfeeding. Among 156 respondents 57.1% had sufficient knowledge about breastfeeding, while 42.9% had no knowledge. Of the 89 respondents with knowledge, 78.7% were aware of the benefits, 73% knew about exclusive breastfeeding, but only 19% were informed about managing breast-related problems. Health workers were the primary source of information (83.1%), followed by friends and community (24.7%), social media (29.2%), and mothers' group meetings (29.2%).

<b>Variables</b>	<b>Frequency</b>	<b>Percent</b>
	<b>(n = 156)</b>	<b>(%)</b>
<b>Knowledge about Breastfeeding</b>		
Yes	89	57.1
No	67	42.9
<b>(n=89)</b>		
<b>Knowledge related information</b>		
Benefits of Breastfeeding	70	78.7
Exclusive Breastfeeding	65	73
Management of breast-related problems	19	19
<b>Source of information</b>		
Health worker	74	83.1
Friends and Community	22	24.7
Social Media	26	29.2
Meeting of Mothers' group	26	29.2

**Table 5: Child-related information**

Table 5 outlines the feeding practices observed among infants in the study, revealing that a significant majority (68.6%) were breastfed within one hour of birth, indicating a strong adherence to early breastfeeding initiation. Concerning pre-lacteal feeding, over four-fifths (82.7%) did not receive any pre-lacteal feed during their first three days of life. As for colostrum milk, nearly four-fifths (79.5%) were fed colostrum, while 20.5% were not. Regarding the introduction of solid or liquid food, 20.5% introduced them to their children between 0 to 2 months, 35.9% between 3 to 5 months, and the remaining 43.6% introduced complementary feeding at 6 months or later.

Variable	Frequency (n = 156)	Percent (%)
<b>Early initiation of breastfeeding right after delivery</b>		
Within one hour	107	68.6
After one hour	49	31.4
<b>Pre-lacteal feeding</b>		
Yes	27	17.3
No	129	82.7
<b>Colostrum milk feeding</b>		
Yes	124	79.5

No	32	20.5
<b>First, introduce solid or liquid food other than breast milk</b>		
0 to 2 Months	32	20.5
3 to 5 Month	56	35.9
6 Month and above	68	43.6

**Table 6: Mother-related information**

Table 6 displays mother-related information linked to exclusive breastfeeding, indicating that all 156 respondents attended health centers for ANC, with 59.6% visiting four or more times. Concerning postnatal care (PNC), 66.7% did not receive PNC services, and 33.3% did. About 47.4% received exclusive breastfeeding counseling during ANC and PNC visits, while 52.6% did not. For childbirth practices, 92.9% delivered at a health facility, predominantly through spontaneous vaginal delivery (87.2%), with 12.8% undergoing a cesarean section.

Variable	Frequency (n = 156)	Percent (%)
<b>At least one visit the health center for ANC</b>		
Yes	156	100



No	0	0
<b>Number of times visit the health center for ANC</b>		
Four and more times	93	59.6
Less than four times	63	40.4
<b>Visit the health center for PNC</b>		
Yes	52	33.3
No	104	66.7
<b>Mother receive counselling on exclusive breastfeeding through ANC, PNC visit</b>		
Yes	74	47.4
No	82	52.6
<b>Place of delivery child</b>		
Institutional delivery	145	92.9
Home delivery	11	7.1
<b>Type of delivery</b>		

Spontaneous vaginal delivery	136	87.2
Cesarean section	20	12.8

---

**Table 7: Exclusive breastfeeding-related information**

In Table 7, the majority (56.4%) of respondents did not practice exclusive breastfeeding, while 43.6% reported adhering to exclusive breastfeeding. Of the 68 respondents practicing exclusive breastfeeding, 42.6% were influenced by health workers, 35.3% made the decision independently, and 22.1% were influenced by family, friends, or the community.

Variable	Frequency	Percent
	(n = 156)	(%)
<b>Exclusive Breastfeeding for six months</b>		
Yes	68	43.6
No	88	56.4
<b>Source of information who influence feeding breastmilk exclusively (n=68)</b>		
Family/Friends/Community	15	22.1
Health Worker	29	42.6

My own decision	24	35.3
-----------------	----	------

---

**Table 8: Factors to discontinue exclusive breastfeeding**

Among the total respondents (88) not practicing exclusive breastfeeding, reasons for discontinuation were documented in Table 6. The primary reason, cited by 67% of respondents, was difficulty initiating milk flow. Additionally, 61.4% reported insufficient milk production by the mother, while 59.1% felt breast milk did not satisfy the baby. Other factors included excessive household duties (34.1%), maternal concerns about inadequate weight gain (29.5%), difficulties with the baby's sucking or latching (18.8%), and the baby falling ill and being unable to breastfeed (13.6%).

Variable	Frequency (n=88)	Percent (%)
I had trouble getting the milk flow to start	59	67
I didn't have enough milk	54	61.4
Breast milk does not satisfy my baby	52	59.1
I had too many household duties	30	34.1
I thought my baby was not gaining enough weight	26	29.5
My baby had trouble sucking or latching	16	18.8
My baby become sick and could not breastfeed	12	13.6

## *Bivariate Analysis*

**Table 9: Association between exclusive breastfeeding and selected sociodemographic variables**

Table 9 presents data on exclusive breastfeeding among 156 respondents, with 43.6% practicing it and 56.4% not. Within the exclusive breastfeeding group, age distribution shows 52.9% in the 19 to 24-year range, 32.4% in the 25 to 30-year range, and 14.7% in the 31 to 36-year range. No significant association with age was found ( $P = 0.079$ ). Sex of the child revealed a significant association ( $P = 0.033$ ), with 69.1% of exclusively breastfed being male. Mothers' occupation also showed significance ( $P = 0.031$ ), with 91.2% of homemakers exclusively breastfeeding. Mother Education level exhibited a significant association ( $P = 0.049$ ), with 36.7% of those completing secondary education and above practicing exclusive breastfeeding. Family type and smoking habits also revealed significant associations ( $P = 0.024$  and  $P = 0.027$ , respectively), while alcohol-drinking habits did not show an association ( $P = 0.230$ ).

Variable	Exclusive breastfeeding for six months			$\chi^2$	P-Value
	Yes	No	Total		
<b>Age of Mother</b>					
19 to 24 Year	36	59	95	3.083	0.079
25 to 30 Year	22	21	43		
31 to 36	10	8	18		

**Sex of child**

Female	21	42	63	4.521	<b>0.033</b>
Male	47	46	93		

**Mother Occupation**

Employed	6	19	25	4.646	<b>0.031</b>
Homemaker	62	69	131		

**Father occupation**

Private or government worker	12	15	27	4.921	0.178
Self Employed	5	11	16		
Labor	37	54	91		
Foreign worker	14	8	22		

**Mother Education**

Cannot read and write	15	9	24	7.841	<b>0.049</b>
Read and write only	11	9	20		

Primary education	17	20	37
-------------------	----	----	----

Secondary education and above	25	50	75
-------------------------------	----	----	----

**Father Education**

Cannot read and write	9	9	18	0.985	0.808
-----------------------	---	---	----	-------	-------

Read and write only	6	9	15
---------------------	---	---	----

Primary education	14	23	37
-------------------	----	----	----

Secondary education and above	39	47	86
-------------------------------	----	----	----

**Type of Family**

Joint	21	43	64	5.126	<b>0.024</b>
-------	----	----	----	-------	--------------

Nuclear	47	45	92
---------	----	----	----

**Number of children**

One	29	35	64	0.646	0.977
-----	----	----	----	-------	-------

Two	26	39	65
-----	----	----	----

Three and above	13	14	27		
<b>Mother's Smoking habit</b>					
Yes	13	31	44	4.916	<b>0.027</b>
No	55	57	112		
<b>Mother's Alcohol Drinking habit</b>					
				1.443	0.230
Yes	7	15	22		
No	61	73	134		

**Table 10: Association between other factors and exclusive breastfeeding**

Table 10 examines the association between exclusive breastfeeding and various factors among 156 respondents. Of the 68 (43.6%) exclusively breastfeeding respondents, 69.1% had knowledge about breastfeeding, showing a significant association ( $P = 0.007$ ). Among those with knowledge about breastfeeding, exclusive breastfeeding was significantly associated with knowledge about the benefits of breastfeeding ( $P = 0.04$ ) and knowledge about exclusive breastfeeding ( $P = 0.007$ ). The study found a significant association between exclusive breastfeeding and avoiding pre-lacteal feeding ( $P = 0.0001$ ) and practicing colostrum milk feeding ( $P = 0.0001$ ). Additionally, the number of ANC visits was significantly associated with exclusive breastfeeding ( $P = 0.033$ ), with 60.3% of exclusively breastfeeding mothers receiving counseling on the topic ( $P = 0.005$ ). Although no significant association was found, the type of delivery showed a trend towards significance ( $P = 0.073$ ).

Variable	Exclusive breastfeeding for six months			$\chi^2$	P-Value
	Yes	No	Total		
<b>Knowledge about BF</b>				7.162	<b>0.007</b>
Yes	47	42	89		
No	21	46	67		
<b>Knowledge related to the benefits of EBF</b>				4.224	<b>0.04</b>
Yes	33	37	70		
No	14	5	19		
<b>Knowledge related to EBF</b>				7.371	<b>0.007</b>
Yes	40	25	65		
No	7	17	24		
<b>Knowledge related to the management of Breast related problem</b>				1.111	0.292



Yes	8	11	19		
No	39	31	70		
<b>Initiation of breastfeeding right after delivery</b>				2.299	0.129
Within one hour	51	56	107		
After one hour	17	32	49		
<b>Pre-lacteal feeding</b>				25.230	<b>0.0001</b>
Yes	0	27	27		
No	68	61	129		
<b>Colostrum milk feeding</b>				19.166	<b>0.0001</b>
Yes	65	59	124		
No	3	29	32		
<b>Number of times visit the health center for ANC</b>				4.521	<b>0.033</b>
Four and more times	47	46	63		

Less than four times	21	42	93		
<b>Visit the health center for PNC</b>				0.639	0.424
Yes	25	27	52		
No	43	61	104		
<b>Mother receive counselling on EBF through ANC, PNC visit</b>				7.993	<b>0.005</b>
Yes	41	33	74		
No	27	55	82		
<b>Place of Delivery</b>				0.035	0.852
Institutional delivery	64	81	145		
Home Delivery	4	7	11		
<b>Type of Delivery</b>				3.224	0.073
Normal Vaginal Delivery	63	73	136		

Cesarean Section                      5                      15                      20

---

## *Multivariate Analysis*

**Table 11: Multiple logistic regression**

Table 11 presents logistic regression results exploring factors influencing exclusive breastfeeding among mothers of children under 2 years old. The logistic regression analysis identified several factors influencing exclusive breastfeeding practices among mothers of children under 2 years old. Initially, there appeared to be a gender disparity, with male children showing higher odds of exclusive breastfeeding, but this significance diminished after adjusting for other variables, indicating potential confounding. Homemaker mothers exhibited significantly higher odds of exclusive breastfeeding compared to employed mothers, a trend that strengthened after adjustment. Similarly, mothers living in nuclear families initially showed higher odds of exclusive breastfeeding, although this significance disappeared after adjusting for other variables. Non-smoking mothers were consistently more likely to exclusively breastfeed, even after adjustment. While factors like knowledge about breastfeeding and advice on exclusive breastfeeding showed significant associations in unadjusted analyses, they lost significance after adjustment, hinting at the influence of confounding variables. Notably, feeding colostrum milk and frequent ANC visits remained strongly associated with higher odds of exclusive breastfeeding even after adjustment. These findings underscore the multifaceted nature of exclusive breastfeeding determinants and the importance of tailored interventions to promote and support breastfeeding practices.

<b>Variable</b>	<b>Exclusive breastfeeding for six month</b>	<b>Unadjusted OR (95% CI)</b>	<b>P-Value</b>	<b>Adjusted OR (95% CI)</b>	<b>P-Value</b>
-----------------	--	-------------------------------	----------------	-----------------------------	----------------

	No	Yes				
<b>Sex of Child</b>						
Female	42	21	Ref			
Male	46	47	2.043	<b>0.035</b>	2.135	0.058
			(1.053 –		(0.974-4.679)	
			3.966)			
<b>Mothers occupation</b>						
Employed	19	6	Ref			
Homemaker	69	62	2.845	<b>0.036</b>	4.459	<b>0.009</b>
			(1.068 –		(1.444 –	
			7.580)		13.767)	
<b>Type of family</b>						
Joint/Extended	43	21	Ref			
Nuclear	45	47	2.139	<b>0.025</b>	1.634	0.218

				(1.102 –		(0.748 –	
				4.150)		3.570)	
<b>Smoking habit</b>							
Yes	31	13	Ref				
No	57	55	2.301	<b>0.029</b>	2.755	<b>0.027</b>	
				(1.091 –		(1.120 –	
				4.852)		6.774)	
<b>Knowledge about BF</b>							
Yes	42	47	2.451	<b>0.008</b>	1.364	0.661	
				(1.263 –		(0.340 –	
				4.757)		5.475)	
No	46	21	Ref				
<b>Colostrum milk feeding</b>							

Yes	59	65	10.650	<b>0.0001</b>	12.472 (3.253 – 47.823)	<b>0.0001</b>
			(3.082 – 36.795)			

No	29	3	Ref			
----	----	---	-----	--	--	--

**Number of times  
visit the health  
center for ANC**

Four and more times	46	47	2.043	<b>0.035</b>	2.333 (1.040 – 5.233)	<b>0.040</b>
			(1.053 – 3.966)			

Less than four times	42	21	Ref			
-------------------------	----	----	-----	--	--	--

**Mother receive  
Counselling on  
EBF in ANC or  
PNC visit**

Yes	33	41	2.531	<b>0.005</b>	1.810	0.396
-----	----	----	-------	--------------	-------	-------

			(1.322 -	(0.460 –
			4.847)	7.127)
No	55	27	Ref	

---

## *Discussion*

An analytical cross-sectional study design was used to assess the proportion of exclusive breastfeeding and associated factors among 156 Mothers of children under 2 years of age in the Dalit community. This study aimed to assess the proportion of mothers of children under 2 years of age, practicing exclusive breastfeeding and associated factors in the Dalit community of Rajbiraj Municipality, Saptari, Nepal.

The study revealed an exclusive breastfeeding prevalence of 43.6%, showing a positive trend compared to the national average of 36.9%. Regional variations were observed, with Madhesh province at a lower 26%. (17) Discrepancies were noted in national reports, with NDHS 2022 at 56.4% and MICS 2019 at 62.1%. (18, 19) Comparable rates were found in western hilly region of Nepal reported a prevalence of 47.6%, (20) which was relatively similar to the our study. For instance, study conducted in the mid-western and eastern regions of Nepal found a rate of EBF 23.2%, (21) while Acharya et al. reported a prevalence of 53.0% in rural Terai of Nepal. (22) Dahal reported a prevalence of 49.1% in slum areas of Kathmandu Valley, (23) while Panthi found a rate of 71.5% among mothers in central Nepal. (24) (25) Sharma and Kafle reported that approximately nearly half (50.5%) in Pokhara Slum area, falling within our study's range but precluding a direct comparison. (26) Globally, this findings align with the breastfeeding prevalence of (44%) but lower than the South-Asia region (57%). (27, 28) Comparing with neighboring countries show that present study has higher rate of EBF compared to India's Bihar (27.6%), (29) and China (37%). The study presents encouraging trends in EBF prevalence within the Dalit community of Rajbiraj Municipality.

The current study observed that the age of the mothers ranged from 19 to 36, with a mean age of 24.59 years aligning with previous research in mid-eastern and western regions of Nepal and Western Nepal, where mean ages of 25.38 and 24.9 years were reported, respectively. (21, 30) Notably, studies conducted in the Slum of Pokhara and Nepalgunj emphasized a majority of participants being below the age of 30 year old. (29, 31) These studies complemented our findings, as their age ranges also matched those observed in our study. The association between the age of the mothers and exclusive breastfeeding practices was assessed using chi-square statistics, which revealed no significant association ( $P = 0.079$ ) in the current study. This result is consistent with the findings of study conducted in Rural Southern Nepal, another study conducted in Central Nepal and study in China reported no significant association between age and exclusive breastfeeding ( $P = 0.545$ ,  $P = 0.12$ , and  $P = 0.73$ , respectively). (22, 24, 25) These consistent findings across different regions and countries suggest that age may not be a determining factor influencing exclusive breastfeeding practices among mothers in various contexts.



The findings of this study indicate that health workers have a substantial impact on promoting and influencing exclusive breastfeeding practices among mothers. The higher percentage of respondents being influenced by health workers (42.6%) compared to personal decision – making (35.3%) and influence from family/friends/community (22.1%) underscores the significance of health workers in EBF promotion. The comparison with the study conducted in Pokhara, further support this understanding, highlighting that (50.5%) of mothers receiving information on EBF from health workers. (32) the higher percentage of mothers influenced by health workers suggests their significant contribution to maternal decision-making, as they are trusted source of information capable of offering evidence based guidance. This emphasizes the importance of health worker engagement and education in promoting optimal infant feeding practices within communities.

The current study found that the child’s age in months ranged from 6 to 24 months, with a mean age of 11.39 months and a confidence interval of 10.64 to 12.14. The majority (64.1%) of the respondents had children in the 6 to 12-month age group, while 28.8% of the children were in the 13 to 18-month age group, and 7.1% were in the 19 to 24- month age group. Comparing these findings with the previous studies conducted in mid-western and eastern regions of Nepal, where the mean age of infants was  $9.98 \pm 2.26$  months, aligns with the child’s age range in the study. (21) A study in Central Nepal reported a median age of infants as 10 months (ranging from 4 to 14 months), falling within the observed age range (24) In the Slum of Pokhara, the distribution of children in the 6 to 12-month age group closely corresponds to the current study’s results. Additionally, also reported the distribution of children as 51.3% in the 6 to 12-month age group and 48.8% in the 13 to 24-month age group, which corresponds closely to the findings of the current study. (26) Similarly, a study in urban slums in Bihar, India, reported consistent distribution across age groups, albeit with slight variation in percentages, further supporting the reliability of the current study’s findings. (29)

The study, comprising 156 respondents, revealed that (59.6%) were male, and 40.4% were female. indicating a significant association with exclusive breastfeeding (P-Value = 0.003). specifically, male children had two-fold odds (OR=2.043) of being exclusively breastfed compared to female children, without adjusting for other factors (OR = 2.043; CI = 1.053 – 3.966). Comparing with previous studies, findings varied; rural southern Nepal reported 52.3% female children with no significant association with exclusive breastfeeding,(22) while study in Nepal and Dhulikhel Municipality found a slightly higher proportion of male infants, aligning with our results. (31, 33) In Central Nepal, the distribution was 58.8% male and 41.2% female, suffering from our study. (24) the slum of Pokhara reported 47% Male and 53% female infants, contrary to our findings. (26) In Pokhara more than half (57.4%) were male, aligning with our results, but they reported female infants more likely to be EBF (51.2%), contrary to our findings. (32) In urban slums, Bihar, India, the distribution was similar to our study, and they found that male infants had a higher chance

of EBF for six months, supporting our findings. (29) these comparisons highlight the variability in gender distribution and its association with exclusive breastfeeding across different regions and studies. Our findings suggest a significant association between gender and EBF practices, emphasizing the need for further research to explore underlying factors influencing this relationship.

In the current study, out of the total 156 respondents, the majority (84%) were homemakers, while 16% were employed. The chi-square analysis revealed a significant association between exclusive breastfeeding and mothers' occupation (0.031). After adjusting for other variables, homemaker mothers had 5.724 times higher odds of practicing exclusive breastfeeding compared to employed mothers (OR = 2.845; CI = 1.068 – 7.580). These findings align with similar research conducted in Nepal. For instance, a study found that the majority (76.4%) of the mothers in their study were not formally employed, which aligns with the current study's findings of a high proportion of homemakers. (33) Study conducted in central Nepal reported that a vast majority (74.3%) of the mothers in their study were homemakers, similar to the current study's results. (24) Study conducted in Slum of Pokhara found that the occupation of mothers in theirs in their study was predominantly housewife (86.5%), which aligns with the current study's results. (26) Study conducted in Urban Slum of India reported that almost 87.2% of the mothers in their study were housewives, which is consistent with the current study's findings. However, they did not assess the association between mothers' occupation and exclusive breastfeeding. (29) These consistent findings across different studies highlight the strong association between maternal occupation (homemaker vs. employed) and exclusive breastfeeding practices. The high prevalence of homemakers practicing exclusive breastfeeding underscores the potential impact of maternal employment status on infant feeding behaviors. Future research could further explore the underlying reasons for this association and identify strategies to support employed mothers in maintaining exclusive breastfeeding practices.

In the current study, approximately 68.6% of infants were fed breast milk within one hour right after delivery, while the remaining 31.4% were fed breast milk after one hour. The chi-square analysis indicated no significant association between exclusive breastfeeding and the initiation of breastfeeding right after delivery ( $P = 0.129$ ). Comparing these findings with National Data Annual Report (2077/78), Fifty-five percent (55%) of children age 0–23 months engaged in early initiation of breastfeeding. (17) The present study indicating a positive trend. Comparing these findings with the previous studies, there are mixed results regarding the association between the initiation of breastfeeding and exclusive breastfeeding. Some studies have reported a significant association, suggesting that early initiation of breastfeeding is associated with a higher likelihood of exclusive breastfeeding. (25, 26, 29) However, other studies, including the current study, did not find a significant association. (18, 24) It is important to note that the percentage of infants who were breastfed within one hour of delivery varied across the studies. While the current

study reported 68.6%, other studies reported percentages ranging from 41.8% to 96.3%. (21, 31, 33) These variation could be due to differences in study populations, cultural practices, healthcare settings and other contextual factors. The timing of initiation of breastfeeding is an essential factor that can influence exclusive breastfeeding practices. While early initiation of breastfeeding is recommended and associated with positive outcomes, its direct impact on exclusive breastfeeding may vary depending on various factors unique to each study setting. Further research is needed to better understand the complex interplay between early initiation of breastfeeding and exclusive breastfeeding practices in diverse populations and settings.

The findings of the current study indicate that approximately four-fifths (79.5%) of the infants were fed colostrum milk, while 20.5% did not receive colostrum milk. The chi-square analysis revealed a significant association between exclusive breastfeeding and colostrum milk feeding in the study population ( $P = 0.0001$ ). Furthermore, the study found that mothers who breastfed exclusively and fed colostrum milk had higher odds compared to mothers who did not feed colostrum milk, even after adjusting for other variables. Comparing these findings with the referenced data, it is evident that the prevalence of colostrum milk feeding varies across different studies. For instance, (21) reported a lower rate of infants not receiving colostrum (2.3%), while (33) found that 83.5% of infants were fed colostrum. (26) reported that 5.5% of mothers discarded colostrum milk before feeding breastmilk. (29) noted that the majority of mothers (78.5%) reported their babies receiving colostrum immediately after birth, while the rest were asked to discard it due to cultural beliefs. The relatively high prevalence of colostrum milk feeding in the study population (79.5%) is encouraging, as it indicates a positive trend toward optimal breastfeeding practice. However, the 20.5% of infants who did not receive colostrum milk highlights the need for targeted interventions and support to address this issue. Understanding the reasons behind this practice, such as cultural beliefs or lack of awareness, can help inform strategies to promote colostrum feeding and improve breastfeeding outcomes. Efforts to educate and empower mothers and families about the importance of colostrum milk and dispel myths or misconceptions surrounding its benefits are essential for promoting early and optimal breastfeeding practices.

The findings of the current study indicate that all respondents visited the health center for at least one-time antenatal care (ANC). Comparing with Annual Report 2021, Madhesh Province has the highest number 100% of adolescents who received first ANC services and First ANC visit, which aligns with this study. (17) NDHS (2022) Ninety-four percent (94%) of women reported receiving antenatal care from a skilled provider. (17, 18) These comparisons highlight the positive trend in ANC utilization observed in our study population, where all respondents sought antenatal care services at least once during their pregnancy. The high coverage of ANC visits is essential for promoting maternal and child health, ensuring timely interventions, and reducing maternal and neonatal mortality rates.

Among the total of 156 respondents, 59.6% visited the health center for ANC four or more times, while 40.4% visited the health center less than four times. The chi square analysis revealed a significant association between exclusive breastfeeding and the number of times visiting the health center for ANC in the study population ( $P = 0.033$ ). The odds of mothers who breastfeeding exclusively and visited a health facility for ANC four or more times were two-fold higher compared to those who visited less than four times ( $OR = 2.043$ ;  $CI = 1.053 - 3.966$ ). After adjusting for other variables, the odds increased to 2.333 ( $OR = 2.333$ ;  $CI = 1.040 - 5.233$ ). Comparing these findings with Annual report 2021, Four in five women (81%) had at least four ANC visits, this contradicts the findings. (17) Study conducted in Western Hilly region of Nepal reported that nearly half (49.7%) of the participants followed the recommended guidelines of attending ANC four or more times, which aligns with this study. (20) Study conducted in Pokhara observed that mothers who had ANC visits for four or more times had lower exclusive breastfeeding rates (47.6%) compared to those who visited for less than four times (57.9%). This contradicts the findings. (32) Urban slums, Bihar highlighted that completing three or more ANC visits was associated with prolonged exclusive breastfeeding ( $OR 151$ , 95%  $CI 21 - 537$ ). (29) These comparisons highlight the variability in the association between ANC visits and exclusive breastfeeding across different studies and settings. While our study found a positive association between more ANC visits and higher rates of exclusive breastfeeding, other studies have reported contrasting results. Factors such as regional differences, cultural practices, and healthcare settings may contribute to these variations. Further research is needed to better understand the underlying mechanisms linking ANC utilization to exclusive breastfeeding outcomes and to tailor interventions accordingly to promote optimal infant feeding practices.

The findings of this study states that significant proportion of respondents in the study population did not receive post-natal care (PNC) services, with two thirds (66.7%) not accessing (PNC) services, and only one-third (33.3%) visited the health center for PNC services. The Annual Report FY 2077/78 (2021 years) reported that, proportion of mothers attending three PNC visits 25.1 percent and in Madhesh province 14.5% . The present study indicating a positive trend. (17) NDHS (2022) reported that, 70% of women received a postnatal check within the 2 days after delivery. this contradict the findings. (18) Additionally, study conducted in Western hilly region of Nepal reported that only 37.5% of participants went for postnatal services two or more times. This finding aligns with this study, where a significant proportion of respondents did not receive PNC services, indicating a potential gap in postnatal care provision. (20)

The current study indicates that almost half (47.4%) of the respondents received counselling related to EBF through ANC or PNC visits, while the remaining 52.6 % did not receive such counselling. The chi-square statistics demonstrated a significant association between exclusive breastfeeding and mothers receiving counselling on

exclusive breastfeeding in the study population ( $P = 0.005$ ). Moreover, the unadjusted analysis revealed that mothers who breastfed exclusively and received counselling on exclusive breastfeeding through ANC or PNC visits had odds 2.531 times higher than those who did not receive such counselling ( $OR = 2.531$ ;  $CI = 1.322 - 4.847$ ). These findings align with previous research conducted in Nepalgunj Medical college, which indicated that only 59% of women received advice on breastfeeding during the antenatal period. (30) The study emphasized the importance of providing support and counselling routinely during ANC to prepare mothers for successful exclusive breastfeeding. Additionally, a study by Urban slum, Bihar reported that receiving counselling at the hospital was associated with higher odds of exclusive breastfeeding. This aligns with the present study's results, highlighting the significance of receiving counselling on exclusive breastfeeding in promoting and establishing exclusive breastfeeding practices. (29) To enhance EBF rates, healthcare providers should prioritize the provision of evidence – based counselling and support throughout the antenatal, perinatal, and postnatal periods. This includes integrating counselling and education on exclusive breastfeeding into routine ANC, ensuring support is available during childbirth, and offering continuous guidance in the postnatal period to facilitate sustained exclusive breastfeeding. Strengthening health systems to prioritize and deliver quality counseling and support services is essential for improving exclusive breastfeeding practices and overall maternal and child health outcomes.

The findings of the study provided that among the total 156 respondents, 87.2% had a spontaneous vaginal delivery, while 12.8% underwent a cesarean section. The chi-square statistics showed no significant association between exclusive breastfeeding and the type of delivery in the study population, although it approached significance at the 10% level ( $P = 0.073$ ). In the study conducted in Western hilly region of Nepal, the regression analysis revealed that exclusive breastfeeding was more likely among mothers who delivered through normal vaginal delivery. (20) Similarly, study conducted in Nepalgunj medical college found that mothers who had a normal delivery were two times more likely to practice exclusive breastfeeding compared to those who had an assisted delivery. (26) On the other hand, study conducted in Central Nepal reported that cesarean section was the most frequent cause (88.6%) for the late initiation of breastfeeding. (24) This suggests that cesarean section may have an impact on the timing of breastfeeding initiation, but its influence on exclusive breastfeeding practices may not be significant within the study population. The studies in Pokhara, Urban slum of Bihar, and China also assessed the association between mode or type of delivery and exclusive breastfeeding. (25, 29, 32) Study conducted in Pokhara found a significant association between mode of delivery and exclusive breastfeeding ( $p < 0.001$ ). (32) Urban slum of Bihar reported that mothers who delivered normally had two times higher odds of practicing exclusive breastfeeding compared to those who had a cesarean section. (29) In the present study, although the association between exclusive breastfeeding and type of delivery did not reach statistical significance, there is a trend suggesting a potential relationship. Further research with

larger sample sizes and more detailed analysis may help clarify the impact of mode of delivery on exclusive breastfeeding practices in different populations and settings. Understanding these associations is crucial for developing targeted interventions to promote and support exclusive breastfeeding among mothers, regardless of their mode of delivery.

A significant proportion of respondents cited issues related to milk production, including trouble getting the milk flow to start, not having enough milk, or perceiving inadequate secretion of breast milk. These reasons were reported by multiple studies, including study conducted in Central Nepal, Nepalgunj medical college, and mid-western and eastern regions of Nepal. (21, 24, 30) Another common reason mentioned by respondents was the perception that breast milk did not satisfy the baby. This finding was reported in central Nepal (2022) and Nepalgunj medical college (2017). (24, 30) Some respondents reported specific difficulties with breastfeeding, such as painful breasts, difficulty in sucking and problems with latching. These challenges were highlighted in the studies in Central Nepal, and mid-western and eastern regions of Nepal. (21, 24) Several studies, including Nepalgunj medical college and Urban slum Bihar Mentioned household duties and work related problems as barriers to exclusive breastfeeding. (29, 30) Mothers may face challenges in balancing their household responsibilities and the time commitment required for breastfeeding. This can lead to decreased breastfeeding duration or early introduction of other foods or milk substituted. These findings highlight the multifaceted challenges faced by mothers in maintaining exclusive breastfeeding practices. Addressing these barriers requires comprehensive support systems, including targeted education and counseling, improved access to lactation support services, and workplace policies that accommodate breastfeeding mothers. Efforts to address these challenges are crucial for promoting and sustaining exclusive breastfeeding practices, which have significant benefits for maternal and child health outcomes.

## ***Conclusion***

This study sheds light on the demographic characteristics, knowledge and breastfeeding practices among the Mothers of children under 2 years of age in Dalit community of Rajbiraj Municipality. The findings emphasize the need for targeted interventions and support to promote exclusive breastfeeding practices among mothers in the study population. Factors such as the sex of the child, the mothers' occupation and education level and maternal knowledge were found to significantly influence exclusive breastfeeding. Adjusted analysis revealed that being a homemaker mother, completing formal education, feeding colostrum milk, and attending ANC visits frequently were significantly associated with higher odds of exclusive breastfeeding. However certain associations initially observed lost significance after adjusting for confounding variables, highlighting the importance of considering various factors and potential confounders in promoting exclusive breastfeeding.

## ***Recommendation***

To promote exclusive breastfeeding among mothers and improve maternal and child health outcomes, the following recommendations are proposed:

1. Targeted interventions should focus on supporting breastfeeding practices especially among mothers who are employed or have lower levels of education.
2. Healthcare providers, particularly during antenatal care visits, should offer comprehensive and accurate information on exclusive breastfeeding, providing counselling and support to expectant and new mothers.
3. Identify and address common barriers and challenges encountered by mothers during exclusive breastfeeding, such as milk flow issues or concerns about infant satisfaction. Offer appropriate guidance, support, and interventions to overcome these challenges.

## ***Limitations***

1. The study's findings may not be generalizable to the entire population as it focused on a specific region or community.
2. Another limitation of this study was recall bias due to the retrospective nature of the data collection, possibly resulting in over or underestimation of actual feeding practices. Although recall biases cannot be avoided, the researcher conducted all interviews by asking probing questions to gather exact information.

## Reference

1. Poudel B, Kattel S. Social Changes in Musahar Community: A Case Study of Dhanusa District of Nepal. *Nepal Journal of Multidisciplinary Research*. 2019;2(4):9-16.
2. Kareriya BR. Socio-Economic Status of Dalits of the Western Tarai Villages, Nepal. *Geographical Journal of Nepal*. 2011;9(0):13-22.
3. NPHC. National Population and Housing Census 2011. 2011.
4. Breastfeeding [Internet]. 2022. Available from: [https://www.who.int/health-topics/breastfeeding#tab=tab\\_1](https://www.who.int/health-topics/breastfeeding#tab=tab_1).
5. UNICEF. Breastfeeding: A Mother's Gift for Every Child. 2018.
6. Ekubay M, Berhe A, Yisma E. Initiation of breastfeeding within one hour of birth among mothers with infants younger than or equal to 6 months of age attending public health institutions in Addis Ababa, Ethiopia. *International Breastfeeding Journal*. 2018;13(1):4.
7. Shitie A, Tilahun A, Olijira L. Exclusive breastfeeding practice and associated factors among mothers of infants age 6 to 12 months in Somali region of Ethiopia. *Sci Rep*. 2022;12(1):19102.
8. Bernardo H, Cesar V, World Health O. Long-term effects of breastfeeding: a systematic review. Geneva: World Health Organization; 2013 2013.
9. What's In Breast Milk? [Internet]. 2023. Available from: <https://americanpregnancy.org/healthy-pregnancy/first-year-of-life/whats-in-breastmilk/>.
10. Neifert M, Bunik M. Overcoming clinical barriers to exclusive breastfeeding. *Pediatr Clin North Am*. 2013;60(1):115-45.
11. Agho KE, Dibley MJ, Odiase JI, Ogbonmwan SM. Determinants of exclusive breastfeeding in Nigeria. *BMC Pregnancy and Childbirth*. 2011;11(1):2.
12. Mogre V, Dery M, Gaa PK. Knowledge, attitudes and determinants of exclusive breastfeeding practice among Ghanaian rural lactating mothers. *International Breastfeeding Journal*. 2016;11(1):12.
13. Chaurasiya SP, Pravana NK, Khanal V, Giri D. Two thirds of the most disadvantaged Dalit population of Nepal still do not deliver in health facilities despite impressive success in maternal health. *PLoS One*. 2019;14(6):e0217337.
14. Goswami S, Pinto EP. Employing innovative evidence-backed community processes for maternal health services by Dalit women. *Int J Equity Health*. 2022;21(Suppl 1):166.
15. CBS. Statistical District Profile of Saptari 2075. 2075.
16. Cochran WG. *Sampling Techniques* 1977.
17. DoHS. Annual Report. Kathmandu; 2078.
18. NDHS. Nepal Demographic and Health Survey. 2022 November 2022.
19. NMICS. Multiple Indicator Cluster Survey. 2019.



20. Adhikari N, Acharya K, Upadhya DP, Pathak S, Pokharel S, Pradhan PMS. Infant and young child feeding practices and its associated factors among mothers of under two years children in a western hilly region of Nepal. *PLoS one*. 2021;16(12):e0261301.
21. Dharel D, Dhungana R, Basnet S, Gautam S, Dhungana A, Dudani R, et al. Breastfeeding practices within the first six months of age in mid-western and eastern regions of Nepal: a health facility-based cross-sectional study. *BMC Pregnancy and Childbirth*. 2020;20(1):59.
22. Acharya D, Singh JK, Kandel R, Park J-H, Yoo S-J, Lee K. Maternal Factors and the Utilization of Maternal Care Services Associated with Infant Feeding Practices among Mothers in Rural Southern Nepal. *International Journal of Environmental Research and Public Health*. 2019;16(11):1887.
23. Dahal M. Knowledge and practice of Breastfeeding and its associated factors among mothers having 6-24 months of children in slum areas of Kathmandu valley. 2019.
24. Panthi PK, Prerna bansal, Meera Prasai, Kamal Khadka, Ranjeeta Phuyal, Ayasha Shrestha. Prevalence of Exclusive Breastfeeding and Factors Influencing Infant Feeding Practices among Mothers of Central Nepal. *Journal of College of Medical Sciences-Nepal*. 2022;18.
25. Li J, Zhao C, Wang Y, Wang YP, Chen CY, Huang Y, et al. Factors associated with exclusive breastfeeding practice among mothers in nine community health centres in Nanning city, China: a cross-sectional study. *International Breastfeeding Journal*. 2021;16(1):71.
26. Sharma D, Kafle R. Exclusive Breastfeeding and Complementary Feeding Practices among Children in Slum of Pokhara. *Journal of College of Medical Sciences-Nepal*. 2020;16(2):93-8.
27. Wasti SP, Shrestha A, Dhakal P, Gc VS. The prevalence of exclusive breastfeeding practice in the first six months of life and its associated factors in Nepal: A systematic review and meta-analysis. *Sexual & Reproductive Healthcare*. 2023;37:100863.
28. WHO. Babies and mothers worldwide failed by lack of investment in breastfeeding. 2017.
29. Kazmi S, Akparibo R, Ahmed D, Faizi N. Prevalence and predictors of exclusive breastfeeding in urban slums, Bihar. *J Family Med Prim Care*. 2021;10(3):1301-7.
30. Parajuli J, Mishra P, Thapa N. Knowledge and Practice Regarding Breastfeeding among Mothers Attending Immunization Clinic in Nepalgunj Medical College Teaching Hospital. *Journal of Nepalgunj Medical College*. 2017;13(1):32-9.
31. Bhandari N, Prajapati R. Prevalence of exclusive breast feeding and its associated factors among mothers. *Kathmandu Univ Med J KUMJ*. 2018;16:166-70.
32. Joshi B, Timilsina A. Factors Influencing Exclusive Breast Feeding Practice among the Mothers of Infants in Pokhara. *JHAS*. 2019;9.

33. Bhandari S, Thorne-Lyman AL, Shrestha B, Neupane S, Nonyane BAS, Manohar S, et al. Determinants of infant breastfeeding practices in Nepal: a national study. *International Breastfeeding Journal*. 2019;14(1):14.