

1 **Whole-genome mapping of APOBEC mutagenesis in metastatic**  
2 **urothelial carcinoma identifies driver hotspot mutations and a**  
3 **novel mutational signature**

4 J. Alberto Nakauma-González<sup>1,2,3,\*</sup>, Maud Rijnders<sup>3</sup>, Minouk T. W. Noordsij<sup>1</sup>, John W. M. Martens<sup>3</sup>,  
5 Astrid A.M. van der Veldt<sup>3,4</sup>, Martijn P. J. Lolkema<sup>3,6</sup>, Joost L. Boormans<sup>2</sup>, Harmen J. G. van de  
6 Werken<sup>1,2,5,7,\*</sup>

7 <sup>1</sup>Cancer Computational Biology Center, Erasmus MC Cancer Institute, University Medical Center Rotterdam,  
8 Rotterdam 3015 GD, the Netherlands

9 <sup>2</sup>Department of Urology, Erasmus MC Cancer Institute, University Medical Center Rotterdam, Rotterdam 3015  
10 GD, the Netherlands

11 <sup>3</sup>Department of Medical Oncology, Erasmus MC Cancer Institute, University Medical Center Rotterdam,  
12 Rotterdam 3015 GD, the Netherlands

13 <sup>4</sup>Department of Radiology & Nuclear Medicine, Erasmus MC, University Medical Center Rotterdam, Rotterdam  
14 3015 GD, the Netherlands

15 <sup>5</sup>Department of Immunology, Erasmus MC Cancer Institute, University Medical Center Rotterdam, Rotterdam  
16 3015 GD, the Netherlands

17 <sup>6</sup>Currently employed by Amgen Europe BV, Breda 4817 ZK, the Netherlands

18 <sup>7</sup>Lead contact

19  
20 \*Correspondence: [j.nakaumagonzalez@erasmusmc.nl](mailto:j.nakaumagonzalez@erasmusmc.nl) (JAN), [h.vandewerken@erasmusmc.nl](mailto:h.vandewerken@erasmusmc.nl) (HJGvdW), Cancer  
21 Computational Biology Center & Department of Immunology, Department of Urology, Erasmus MC Cancer  
22 Institute, University Medical Center; Rotterdam, the Netherlands. PO Box 2040, 3000 CA, NA-1218.  
23

## 24 **SUMMARY**

25 APOBEC enzymes mutate specific DNA sequences and hairpin-loop structures, challenging the  
26 distinction between passenger and driver hotspot mutations. Here, we characterized 115 whole-  
27 genomes of metastatic urothelial carcinoma (mUC) to identify APOBEC mutagenic hotspot drivers.  
28 APOBEC-associated mutations were detected in 92% of mUC and were equally distributed across the  
29 genome, while APOBEC hotspot mutations (ApoHM) were enriched in open chromatin. Hairpin-loops  
30 were frequent targets of didymi (twins in Greek), two hotspot mutations characterized by the APOBEC  
31 mutational signature SBS2, in conjunction with an uncharacterized mutational context (Ap[C>T]), which  
32 was associated with DNA mismatch. Next, we developed a statistical framework that identified 0.40%  
33 of ApoHM as drivers of mUC, which affected known driver genes and non-coding regions near exons of  
34 potential novel driver genes. Our results and statistical framework were validated in independent cohorts  
35 of 23 non-metastatic UC and 3744 samples of 17 metastatic cancers, identifying cancer-type-specific  
36 drivers. Our study highlights the role of APOBEC in cancer development and may contribute to  
37 developing novel targeted therapy options for APOBEC-driven mUC.

38

## 39 **KEYWORDS**

40 APOBEC; breast cancer; didymi; driver mutations; hairpin-loops; hotspot mutations; pan-cancer;  
41 mutational signature; twin mutations; urothelial carcinoma

42

43

## 44 INTRODUCTION

45 Cancer genomes accumulate somatic mutations via different mutagenic processes and one of the most  
46 common is attributed to the apolipoprotein B mRNA-editing enzyme catalytic polypeptide-like (APOBEC)  
47 family<sup>1</sup>. APOBEC has a specific mutational signature, which is characterized by C>T/G mutations in the  
48 TpC context and is captured in the SBS2 and SBS13 signatures, as defined by the *Catalogue Of Somatic*  
49 *Mutations In Cancer* (COSMIC)<sup>2</sup>. In some tumor types with high APOBEC activity, the contribution to  
50 the tumor mutational burden is substantial, which increases the neoantigen load favoring response to  
51 immune checkpoint inhibitors<sup>3,4</sup>. However, APOBEC is also responsible for the emergence of driver  
52 mutations that contribute to cancer development as shown in mouse models<sup>5</sup>. Discriminating driver  
53 events from passenger events is essential to reconstruct the evolutionary history of cancers and identify  
54 effective novel drug targets in APOBEC-driven tumors.


55 The mutational process of APOBEC has been extensively studied, revealing its preference for single-  
56 stranded DNA structures that form hairpin-loops<sup>6</sup>. This characteristic of APOBEC could result in identical  
57 somatic mutations in tumors from multiple patients, so-called hotspot mutations or hotspots. Due to their  
58 high prevalence, these hotspots can erroneously be assigned as driver mutations, especially in the non-  
59 coding area of the genome. However, the vast majority of mutations are passengers and do not  
60 contribute to cancer development, and the same principle may also apply to hotspot mutations<sup>7</sup>.  
61 Although bioinformatic strategies to identify driver hotspot mutations have been developed<sup>8,9</sup>, the unique  
62 characteristics of the APOBEC mutagenic process require specific considerations to account for all co-  
63 variables accurately.

64 APOBEC-derived mutations are a dominant contributor to the mutational landscape in urothelial  
65 carcinoma (UC). Therefore, we analyzed whole-genome DNA-sequencing data of 115 metastatic UC  
66 (mUC) and matched blood samples<sup>10</sup> to identify driver hotspot mutations in the context of APOBEC  
67 mutagenesis. The comprehensive characterization of APOBEC-enriched tumors identified a novel  
68 mutational signature associated with DNA mismatch repair as well as genomic co-variables associated  
69 with APOBEC-derived hotspot mutations (ApoHM), which we used to develop a statistical framework  
70 and identify driver ApoHM. Furthermore, our findings were validated in whole-genomes of an  
71 independent cohort of 23 non-metastatic UC, and the analysis was extended to include 442 metastatic  
72 breast cancer (mBC) and 3302 samples 16 other metastatic cancer types.

## 73 RESULTS

### 74 APOBEC mutagenesis dominates the mutational landscape of urothelial carcinoma

75 The analysis of whole-genome sequencing (WGS) data of mUC and matched blood samples revealed  
76 a median of 20,667 (Q1=14,304, Q3=31,411) single nucleotide variants (SNVs) per tumor. mUC with a  
77 significant enrichment (E) for C>T mutations in TCW (W = A or T) context were considered APOBEC  
78 positive (92%). These tumors were further stratified according to APOBEC enrichment as APOBEC-  
79 high (41%;  $E > 3$ ), APOBEC-medium (33%;  $2 < E < 3$ ), and APOBEC-low (18%;  $E > 1$ ) (**Figure 1A**). The  
80 median contribution of APOBEC COSMIC signatures (SBS2+SBS13) in APOBEC-high, -medium and -  
81 low tumors was 61%, 37% and 15%, respectively. For the remaining 8% of tumors lacking APOBEC  
82 mutations, the median APOBEC signature was <2%, potentially reflecting the noise of the mutational  
83 signature calling. We associated the APOBEC stratification with multiple factors to better assess the  
84 different APOBEC subtypes. Tumor purity, for instance, declined with increasing APOBEC mutagenesis  
85 (**Figure S1**). Moreover, age was associated with the enrichment of APOBEC mutations (**Figure S1**).  
86 The median clonal fraction of SNVs was lower in tumors with APOBEC mutations than in non-APOBEC  
87 tumors, suggesting higher tumor heterogeneity in APOBEC-enriched tumors (**Figure S1**). Localized  
88 hypermutation events (kataegis) strongly correlated with APOBEC enrichment (spearman  $r=0.80$ ,  
89  $p < 0.001$ ). Homologous recombination (HR) deficiency ( $n=3$ ) was only present in APOBEC-low tumors  
90 (Fisher's exact  $p=0.005$ ), while none of the patients with microsatellite instability (MSI;  $n=4$ ) had  
91 evidence of APOBEC mutagenesis (Fisher's exact  $p < 0.001$ ). Structural variants were more frequent in  
92 APOBEC tumors than in non-APOBEC tumors (**Figure S1**). Additionally, APOBEC tumors had a higher  
93 ploidy (median ploidy = 3) and a higher number of genes affected by copy number alterations (CNA)  
94 than non-APOBEC tumors (**Figure S1**), suggesting genomic instability in APOBEC-driven mUC tumors.  
95 APOBEC mutagenesis was not associated with sex, the primary origin of mUC (upper tract versus  
96 bladder) or chromothripsis.



97

98 **Figure 1. Genomic landscape and APOBEC activity of metastatic urothelial carcinoma (n=115)**  
 99 **stratified by APOBEC-enrichment**

100 (A) Whole-genome sequencing data of metastatic urothelial carcinoma (mUC) were classified  
 101 according to the enrichment of APOBEC-associated mutations as having high, medium, low or  
 102 no APOBEC enrichment. The genomic features are displayed from top to bottom as follows:  
 103 APOBEC mutagenesis; genomic subtype (GenS1-5) as previously described<sup>10</sup>; genome-wide  
 104 tumor mutational burden (TMB); mutational signatures grouped by etiology, while both  
 105 APOBEC signatures are shown separately; absolute frequency of structural variants (SV);  
 106 relative frequency of SV categories; clonal fraction; ploidy; tumor purity; microsatellite  
 107 instability (MSI) status; homologous recombination (HR) deficiency status; samples with at  
 108 least one chromothripsis event; frequency of kataegis events; female patients and; primary  
 109 origin of mUC (upper tract versus bladder).

110 (B) Expression of *APOBEC* and *AICDA* genes in 90 samples with available RNA-sequencing  
 111 data.

112 (C) Pearson correlation of RNA expression of *APOBEC3A* and *APOBEC3B*.

113 (D) Fold enrichment of C>T and C>G alterations in YTCA (related to APOBEC3A) and RTCA  
114 (related to APOBEC3B) context.  
115 (E) APOBEC score (normalized expression of *APOBEC3A*+*APOBEC3B*).  
116 (F) Percentage of mRNA C>U mutations in *DDOST* at position chr1:20981977. In (B), (E) and (F),  
117 the Wilcoxon rank-sum test was applied to compare APOBEC tumors vs. non-APOBEC  
118 tumors. P-values were Benjamini-Hochberg corrected in (B).  
119 See also Figures S1-S5.

120

### 121 **APOBEC mutagenesis is an ongoing process in metastatic lesions of urothelial carcinoma**

122 Next, we analyzed RNA-sequencing data of 90 matched samples of mUC. Pathway activity based on  
123 downstream gene expression, such as cell cycle or p53 was similar between the APOBEC groups  
124 (**Figure S2**). Similarly, analysis of APOBEC expression of all genes of the APOBEC family (*APOBEC1*  
125 was not expressed) revealed no significant differences between APOBEC and non-APOBEC tumors  
126 (**Figure 1B**). We detected a weak positive correlation between the expression of *APOBEC3A* and  
127 *APOBEC3B* (**Figure 1C**). To further investigate the mutagenic activity of both enzymes, the fold  
128 enrichment of C>T and C>G mutations, at DNA level, in the tetra-base YTCA (related to APOBEC3A; Y  
129 are pyrimidine bases) and RTCA (related to APOBEC3B; R are purine bases) context was calculated<sup>11</sup>  
130 (**Figure 1D**). In both cases, YTCA and RTCA mutations did not correlate with expression of *APOBEC3A*  
131 or *APOBEC3B* (**Figure S3**). The lack of correlation might be linked to the heterogeneous expression of  
132 APOBEC enzymes that oscillate throughout the cell cycle<sup>12,13</sup>. Furthermore, we detected that both,  
133 APOBEC3A and APOBEC3B, contributed to APOBEC-associated mutations (fold enrichment is above  
134 1.0). Nevertheless, APOBEC3A appeared to be the main contributor as suggested in primary  
135 cancers<sup>11,14</sup>. Considering the mRNA expression of both *APOBEC3A* and *APOBEC3B* enzymes, we  
136 calculated the APOBEC expression score (sum of the normalized expressions of *APOBEC3A* and  
137 *APOBEC3B*). It appeared that the level of APOBEC enrichment correlated with the APOBEC expression  
138 score (**Figure 1E**). This analysis confirmed the link between APOBEC RNA expression at the time of  
139 biopsy and the historical accumulation of APOBEC-associated mutations in mUC that others have  
140 reported in primary urothelial carcinoma<sup>15,16</sup>.

141 Recently, it was proposed that edited *DDOST* mRNA can be used to measure ongoing APOBEC  
142 mutagenesis<sup>17</sup>. We found that the frequency of C>U alterations in the *DDOST* mRNA at position


143 chr1:20981977 was enriched in tumors with APOBEC mutagenesis, with up to 15% of mRNA reads  
144 edited in one single sample (**Figure 1F**). Additionally, we analyzed ongoing APOBEC mutagenesis in  
145 mUC, by WGS of eight tumors from patients who had undergone serial biopsies of metastatic lesions  
146 (**Figure S4**). We observed that the APOBEC mutational signature was present in private mutations of  
147 the second biopsy of these patients, suggesting that APOBEC mutagenesis could be active in the period  
148 between the first and second biopsy (**Figure S4A**). A lower cancer cell fraction (normalized allele  
149 frequency by copy number and purity, see Methods) in private SNVs of the second biopsy compared to  
150 shared (trunk) mutations confirms that these mutations were acquired more recently as they are only  
151 present in a subpopulation of cancer cells (**Figure S4B**). This result, together with the APOBEC  
152 signature detected in the subclonal mutations (**Figure S5**), suggests that the presence of subclonal  
153 populations due to APOBEC mutagenesis may contribute to the ongoing evolution of UC in the  
154 metastatic setting.

155

#### 156 **APOBEC-associated hotspot mutations are enriched in highly accessible genomic regions**

157 The high resolution achieved by WGS, allowed us to investigate the enrichment of APOBEC and non-  
158 APOBEC mutations (non-TpC context) in specific genomic regions. We found that the number of non-  
159 APOBEC-associated SNVs, for instance, varied across the genome (**Figure 2A**). When this distribution  
160 overlapped with DNA accessibility and overall gene expression level, the frequency of non-APOBEC  
161 mutations decreased in open chromatin (highly accessible regions) and highly transcribed regions  
162 (**Figure 2B**). In contrast, the frequency of APOBEC-associated mutations was nearly constant across  
163 the genome. When restricting the analysis to APOBEC tetra-base mutations, we found that the  
164 difference in the distribution of Y/RTCA mutations across genomic regions decreases with the level of  
165 APOBEC enrichment (**Figure 2C**). Interestingly, in APOBEC-high tumors, YTCA mutations were evenly  
166 distributed. In contrast, RTCA mutations were enriched in low DNA accessible and low transcribed  
167 regions, although this enrichment was considerably less compared to tumors with lower levels of  
168 APOBEC mutations.





169

170 **Figure 2. Distribution of APOBEC-associated mutations across genomic regions of metastatic**  
 171 **urothelial carcinoma**

172 (A) Whole-genome sequencing data (n = 115) were analyzed to estimate the mean number of  
 173 single nucleotide variants (SNVs) in windows of one mega base pair (Mbp) across the entire  
 174 genome. The circos plot shows from outer to inner circles: the genomics ideogram from  
 175 chromosome 1 to X where the centrosomes are indicated in red; the mutational load of  
 176 APOBEC and non-APOBEC associated mutations (mutations in TpC or non-TpC context,  
 177 respectively); The density of kataegis events; Average RNA counts (expression) from tumors  
 178 with available RNA-sequencing data (n = 90); DNA accessibility estimation from different



179           ChIPseq experiments of multiple histone marks from normal urothelial samples derived from  
180           ENCODE<sup>45</sup>. Peaks represent highly accessible DNA.

181       (B) Linear regression of the mutational load for APOBEC- and non-APOBEC-associated  
182           mutations as well as the density of kataegis events across the genome with DNA accessibility  
183           and expression data.

184       (C) Relative distribution of APOBEC YTCA and RTCA mutations across DNA-accessible and RNA  
185           expression regions. Samples were stratified according to the level of APOBEC mutagenesis.

186           The Wilcoxon signed-rank test was applied and p-values were Benjamini-Hochberg corrected.

187       (D) Frequency of hotspot mutations grouped according to APOBEC and non-APOBEC-associated  
188           mutations, and DNA accessibility or RNA expression level.

189


190   Because of the high correlation between kataegis and APOBEC enrichment, we analyzed the  
191   distribution of kataegis loci across the genome. Contrary to the overall distribution of SNVs, our data  
192   suggest that kataegis events are more likely to occur in regions with high DNA accessibility and high  
193   transcriptional activity (**Figure 2B**). Moreover, we also evaluated the genome-wide distribution of all  
194   hotspot mutations (two or more mutations in a specific genomic position), representing 0.35% of all  
195   mutated genomic positions. We found that the frequency of highly recurrent ( $n \geq 4$ ) ApoHM were enriched  
196   in high DNA accessibility and high transcriptional active regions (**Figure 2D**). Thus, while general  
197   APOBEC mutagenesis seemed to occur uniformly across the genome, kataegis and ApoHM seemed to  
198   occur more frequently in open chromatin and highly transcribed loci.

199

## 200   **Recurrent hotspot mutations correlate with APOBEC mutagenesis**

201   Next, we investigated the genomic consequence of hotspot mutations and found that the most frequent  
202   hotspot mutations in mUC occurred in non-coding regions of the genome (**Figure 3A**). Hotspot  
203   mutations in the *TERT* promoter were present in 62% of tumors. In line with previous reports<sup>18,19</sup>, *TERT*  
204   expression did not differ between tumors with hotspot mutations and those being *wildtype* (**Figure S6A**).  
205   However, differential gene expression analysis showed that tumors with hotspot mutations in the *TERT*  
206   promoter had high expression of genes related to the biological oxidation pathway (**Figure S6B, Table**  
207   **S1**). Besides *TERT*, other frequent hotspot mutations were identified in the non-coding regions near  
208   *ADGRG6* (40%), *PLEKHS1* (33%), *LEPROTL1* (20%), and *TBC1D12* (15%). Similarly, these hotspot

209 mutations did not affect the expression of these genes but were associated with transcriptomic effects  
 210 in several genes (**Figure S6**) and biological pathways (**Table S1**). These hotspot mutations strongly  
 211 correlated with enrichment for APOBEC-associated mutations, suggesting their origin in APOBEC  
 212 mutagenesis (**Figure 3B**).



213  
 214 **Figure 3. Recurrent hotspot mutations of metastatic urothelial carcinoma correlate with APOBEC**  
 215 **mutagenesis**

216 (A) Overview of recurrent hotspot mutations present in at least five samples, including all  
 217 substitutions occurring in the same genomic position. Hotspot mutations occurring in the TpC  
 218 context are highlighted in red.

219 (B) The association of hotspot mutations and APOBEC fold enrichment (continuous values) was  
 220 interrogated with a logistic regression analysis applying the Wald test. P-values were

221 corrected using the Benjamini-Hochberg method and ordered accordingly. Bars above the  
222 dashed line ( $-\log_{10}(0.05)$ ) are statistically significant and are indicated in red.

223 See also Figure S6 and Table S1.


224

225 All frequent hotspot mutations in the coding region have been previously described and affected known  
226 driver genes: *FGFR3* S249C/R248C (8%, 4%), *PIK3CA* E54K (7%), *RXRA* S427F/Y (7%) and *TP53*  
227 E285K\* (6%). Comparing the expression of these known driver genes affected by hotspot mutations  
228 vs. the wildtype, only *FGFR3* hotspot mutations significantly affected the expression of this gene (**Figure**  
229 **S6A**).

230

### 231 **Hairpin-loops are targets of twin hotspot mutations called Didymi**

232 We noticed that the hotspot mutations near *ADGRG6*, *PLEKHS1*, *LEPROTL1* and *TBC1D12* are located  
233 within DNA hairpin-loop structures. It is known that DNA hairpin-loops are targets of APOBEC3A (**Figure**  
234 **4A**)<sup>20,21</sup>, therefore, we predicted DNA hairpin-loops for all mutated genomic positions (see Methods).  
235 The predicted hairpin-loops near *ADGRG6*, *PLEKHS1*, *LEPROTL1* and *TBC1D12* are each affected by  
236 two hotspot mutations, which are referred to as twin mutations<sup>21</sup>. Moreover, we noticed that the twin  
237 mutations were not mutually exclusive, which differs from the hotspot mutations in the *TERT* promoter  
238 (mutual exclusivity test,  $p < 0.001$ ). Only the twin hotspot mutations in *TBC1D12* co-occurred more  
239 frequently than expected among APOBEC-high tumors ( $p = 0.02$ ). Further analysis of co-occurred twin  
240 hotspot mutations revealed that very few had identical variant allele frequencies, suggesting that most  
241 twin mutations in the same tumor occurred in independent events as they were also found on different  
242 alleles (**Figure S7**).



243

244 **Figure 4. Genomic characteristics of twin hotspot mutations in metastatic urothelial carcinoma**

245 (A) Hairpin-loop structures affected by frequent hotspot mutations in *ADGRG6*, *TBC1D12*,  
 246 *PLEKHS1* and *LEPROTL1*. The positions of hotspot mutations are marked in red for TpC  
 247 context and blue otherwise.

248 (B) Distribution of twin mutations according to the distance between twin mutations and loop size.

249 (C) Mutational signatures (COSMIC v3.3) of twin mutations according to their frequency. The  
 250 stability of the signature call was tested by applying 1,000 bootstrap iterations. Only SBS2 was  
 251 very stable in highly frequent (n ≥ 5) twin mutations.

252 (D) Frequency distribution of hairpin loops affected by twin mutations according to APOBEC  
 253 mutagenesis,

254 (E) DNA accessibility,  
255 (F) number of mutations in TpC context within a loop and  
256 (G) DNA sequence between twin mutations. Wilcoxon rank-sum test was applied to compare  
257 APOBEC vs non-APOBEC tumors and p-values were Benjamini-Hochberg corrected.  
258 See also Figures S7-S9 and Tables S2 and S3.

259

260 Next, we investigated the properties and origin of these twin mutations. A comprehensive analysis of all  
261 DNA hairpin-loop structures in the human genome affected by two mutations in their loops revealed  
262 2,387 twin mutations (4,774 altered genomic positions), representing 0.16% of all mutated genomic  
263 positions. The distance between twin mutations varied, but when the frequency of mutations increased,  
264 the distance decreased to mainly one or two bases and the loop sizes to mainly three to four bases  
265 (**Figure 4B**). Additional examination of the 96 tri-nucleotide contexts of all twin mutations revealed that  
266 both APOBEC COSMIC signatures, SBS2 and SBS13, were dominant (**Figure 4C**). However, at higher  
267 mutational frequencies ( $n \geq 5$ ), only signature SBS2 remained. We also observed a secondary signature  
268 of C>T mutations in the ApC context that does not resemble any known COSMIC signature (**Figure S8**,  
269 **Table S2**). The absolute contribution of this signature was similar across all APOBEC tumors and its  
270 prevalence in the mUC cohort correlated with spontaneous deamination (SBS1) and defective DNA  
271 mismatch repair signatures (**Figure S9**; SBS6, SBS15, SBS20 and SBS26).

272 Furthermore, APOBEC-driven tumors were enriched for twin mutations occurring only in the TpC context  
273 (**Figure 4D**). Contrary to the general pattern of ApoHM (enriched in DNA accessible regions), twin  
274 mutations with a high number of alterations were similarly distributed between high and low DNA  
275 accessible regions (**Figure 4E**).

276 Additionally, we found that at higher mutational frequency, at least one of the twin mutations occurs in  
277 the TpC context (**Figure 4F**) and the sequence between the two is 1001 or 101 (0 = A/T, 1 = G/C;  
278 underlined are the positions of the twin mutations) (**Figure 4G**). Because of the unique characteristics  
279 of frequently affected twin mutations, we named them didymi (twins in Greek). In summary, didymi are  
280 two C>T hotspot mutations found in DNA hairpin-loops separated by one or two A/T base-pairs in which  
281 at least one of the twin mutations is located in TpC context and the other in NpC (N = any base-pair;


282 most N bases are A or T). Applying this definition, we identified 231 didymi in the mUC cohort (**Table**  
283 **S3**).

284

#### 285 **Driver hotspot mutations in urothelial carcinoma**

286 After identifying several hotspot mutations that could be attributed to APOBEC activity, we assessed  
287 whether these hotspot mutations had a selective advantage (drivers) or not (passengers). Recent  
288 attempts relying on the stability of hairpin-loops have been proposed to differentiate passengers from  
289 driver ApoHM<sup>8,20</sup>. We confirmed that a more stable loop (Gibbs free energy  $\Delta G$ ; see STAR Methods)  
290 leads to a higher number of alterations (**Figure 5A**). Taking this into account, we developed a statistical  
291 model to identify driver hotspot mutations considering not only the stability of hairpin-loops but also the  
292 tri-nucleotide context, DNA accessibility and the potential for didymi via sequence in the loop (**Figure**  
293 **5A**).





294

295 **Figure 5. Driver hotspot mutations associated with APOBEC mutagenesis in urothelial**  
 296 **carcinoma**

297 (A) Frequency distribution of variables (trinucleotide context, DNA accessibility, DNA hairpin-loop  
 298 stability and sequence in the loop) considered to identify hotspot mutations that were more  
 299 frequently mutated than expected outside (top) and within (bottom) hairpin-loop structures.

300 (B) Driver hotspot mutations were estimated separately for mutations outside and within hairpin  
 301 loops. Per group, p-values were corrected using the Benjamini-Hochberg method.

302 See also Figures S10 and S11, and Table S4.

303

304 Putative ApoHM were divided into those located outside or inside the loop of DNA hairpin-loop  
305 structures. In case of ApoHM outside hairpin-loops, only those in the TpC context were considered. For  
306 ApoHM within loops, all alterations in TpC, ApC, CpC and GpC context were included in the analysis to  
307 account for didymi. In case of TpC, the distribution of hotspot mutations in the trinucleotide context was  
308 considered. A background distribution was modeled as a Poisson process and the significant enrichment  
309 of mutations in a particular genomic site was estimated.

310 We identified 0.40% (n = 27) of ApoHM as drivers (adjusted p < 0.05; **Fig 5B**). Known driver genes  
311 affected by hotspot mutations included coding alterations in *TP53*, *PIK3CA*, *FGFR3*, *RXRA* and the  
312 *TERT* promoter. All other putative driver ApoHM affected non-coding regions including didymi in  
313 *ADGRG6*, *PLEKHS1*, *TBC1D12* and *LEPROTL1* proposed as drivers by other studies<sup>9,22</sup>. Other  
314 potential driver ApoHM include *RNF169* (involved in DNA damage repair)<sup>23</sup>, *BTG3* (angiogenesis)<sup>24</sup>,  
315 *ADM* (adrenomedullin; vasodilator)<sup>25</sup>, *GDF3* (regulation of TGF-beta)<sup>26</sup> and *WDR74* (ribosome  
316 biogenesis)<sup>27</sup>.

317 To validate our method and confirm the driver ApoHM assessment, we used an independent cohort of  
318 non-metastatic urothelial carcinoma of the bladder (n = 23) of the pan-cancer analysis of whole-genomes  
319 (PCAWG) study<sup>28</sup> (**Figure S10**). This analysis confirmed the previously identified ApoHM as potential  
320 cancer drivers of urothelial carcinoma. Moreover, in this cohort, 96% of tumors were APOBEC-driven,  
321 APOBEC enrichment correlated with kataegis and the largest group (35%) had high enrichment for  
322 APOBEC-associated mutations.


323 Furthermore, the performance of the model to identify driver ApoHM in hairpin loops was tested. The Q-  
324 Q plots show that the empirical distribution of ApoHM deviates from the theoretically expected  
325 distribution (Kolmogorov–Smirnov test p < 0.001). However, when outliers that represent highly frequent  
326 ApoHM (>10) are excluded, which according to our analysis are all drivers, we observed a good  
327 agreement between our model and the theoretical distribution (**Figure S11A**; Kolmogorov–Smirnov test  
328 p = 0.19). By simulating a synthetic genome of mUC, we showed that an 80% statistical power is reached  
329 when the cohort size is ~75 samples for highly frequent (>10%) driver ApoHM (**Figure S11B**). However,  
330 the power to detect rare driver ApoHM (≤10%) is considerably reduced, and a larger cohort is needed.  
331 We also evaluated the contribution of different genomic features as covariates (**Table S4**) to identify

332 driver ApoHM. The McFadden's  $R^2$  in the model that only considers the trinucleotide context is low ( $R^2$   
333 = 0.04), but the goodness of fit increases when considering the hairpin loop ( $R^2 = 0.23$ ), hairpin-loop +  
334 sequence in the loop ( $R^2 = 0.27$ ) and when adding DNA accessibility regions ( $R^2 = 0.28$ ) into the model.  
335 DNA accessibility shows a high (anti-)correlation with other genomic features: GC content, RNA  
336 expression levels, mutational load, methylation and replication timing (**Figure S11C**). Therefore, the  
337 addition of the other variables in the model has limited added value (**Figure S11D**).

338

### 339 **Driver hotspot mutations in metastatic breast cancer**

340 To test our statistical framework in other cancer types and to evaluate if our findings were UC-specific,  
341 we analyzed a cohort of 442 mBC (**Figure 6**)<sup>29</sup>. Similar to UC, breast cancer is commonly affected by  
342 APOBEC mutagenic activity<sup>1</sup>. We identified APOBEC-enriched tumors in 76% of patients, and only 19%  
343 of mBC tumors were APOBEC-high. In most patients (39%), tumors were classified as APOBEC-low  
344 and were enriched for HR deficiency (two-sided Fisher's exact  $p < 0.001$ ).



345

346 **Figure 6. Genomic landscape and driver hotspot mutations associated with APOBEC**  
 347 **mutagenesis in metastatic breast cancer**

348 **(A)** Whole-genome sequencing data from 442 metastatic breast cancer were analyzed and patients  
 349 were classified according to the enrichment of APOBEC-associated mutations. The genomic  
 350 features are displayed from top to bottom as follows: APOBEC mutagenesis; genome-wide  
 351 tumor mutational burden (TMB); COSMIC mutational signatures; frequency of kataegis;  
 352 homologous recombination (HR) deficiency and its probable origin; cancer subtype and; the  
 353 most frequent hotspot mutations (known hotspot mutations in *LEPROTL1*, *TBC1D12* and *TERT*  
 354 were also included).

355 **(B)** Putative driver hotspot mutations in APOBEC-enriched breast cancer. P-values were adjusted  
356 using the Benjamini-Hochberg method.

357 See also Figure S12 and Table S5.

358


359 The most frequent coding hotspot mutations affected *PIK3CA*, *ESR1* and *AKT1*. Twin mutations in  
360 hairpin-loops displayed a similar mutational signature as those in mUC, including the APOBEC signature  
361 SBS2 in conjunction with the uncharacterized C>T mutations in ApC context that define didymi (**Figure**  
362 **S12A-B**). Didymi in *PLEKHS1* and *ADGRG6* were the most frequent non-coding hotspot mutations,  
363 while *LEPROTL1* and *TBC1D12*, two other didymi frequently found in mUC, only affected  $\leq 1\%$  of mBC.  
364 A total of 694 didymi were identified in mBC (**Table S5**), but only 19 (2.7%) were shared with mUC  
365 (**Figure S12C**).

366 Our analysis revealed 51 driver ApoHM in APOBEC-enriched mBC (**Figure 6B**), representing only  
367 0.07% of all ApoHM. Drivers included missense hotspot mutations in *PIK3CA*, *AKT1* and *TP53*, and  
368 hotspot mutations outside of the protein-coding region of *MAPKAPK2*, *STAG1* and including didymi in  
369 *PLEKHS1* and *ADGRG6*. In contrast to mUC, and despite being one of the most frequently affected  
370 genes by hotspot mutations in mUC, *LEPROTL1* was not a driver of mBC. This analysis suggests that  
371 driver mutations derived from APOBEC activity are cancer-type-specific.

372

### 373 **Driver hotspot mutations across multiple metastatic cancers**

374 APOBEC mutagenic activity is widespread across multiple cancer types. Here, we analyzed the genome  
375 of 16 additional metastatic cancers, which in total represents 3302 whole genomes (+115 mUC +442  
376 mBC = 3859). Urothelial, breast and uterus cancers have the highest proportion of APOBEC-high tumors  
377 (**Figure 7A**). In mUC and mBC, 95% of hotspot mutations affect non-coding transcripts, introns or  
378 intergenic regions. This proportion varies per cancer type and can represent up to 99% of all hotspot  
379 mutations in esophagus cancer. Missense hotspot mutations are rare, but the highest proportion is found  
380 in liver cancer, reaching 5%.



381

382 **Figure 7. Pan-cancer overview of APOBEC-derived hotspot mutations and drivers.**

383 (A) Proportion of APOBEC-enriched tumors and hotspot mutations across cancer types.

384 (B) Distribution of APOBEC-associated hotspot mutations (ApoHM) across the genome of  
385 APOBEC-enriched tumors.

386 (C) The 10% most frequent genomic positions or genes affected by driver ApoHM. Drivers of  
387 specific cancer type are indicated by stars. The association of driver ApoHM with APOBEC fold  
388 enrichment (continuous values and using all tumor samples) using logistic regression analysis



389 and applying the Wald test ( $p_{adj} < 0.05$ ) shows the “true” APOBEC-derived mutations. All “true”  
390 APOBEC-derived driver ApoHM are included in the figure.

391 See also Tables S6 and S7.

392

393 The frequency of ApoHM increases with the strength of APOBEC mutagenesis and are more recurrent  
394 in hairpin loops (**Figure 7B**). However, skin cancer does not follow this pattern, which has been proven  
395 to be problematic in other studies due to its hypermutated nature, inflating the number of driver  
396 events<sup>30,31</sup>. These studies make special considerations or exclude skin cancer altogether from their  
397 analysis. We found that skin cancer is mostly defined as an APOBEC-low cancer type and we suspected  
398 that a large proportion of mutations that have the APOBEC signature may not have derived from  
399 APOBEC mutagenic activity. This is supported by the relatively low number of driver ApoHM in skin  
400 cancer that correlates with APOBEC enrichment (**Figure 7C**), despite many ApoHM defined as drivers  
401 by our model (**Table S6**). These driver ApoHM considered “true” APOBEC-derived mutations are more  
402 frequent in breast, urothelial, lung and uterus cancers. *TERT*, *PIK3CA*, *PLEKHS1* and *ADGRG6* are the  
403 most affected genes by driver ApoHM. All four genes harbor two driver ApoHM, which are targeted by  
404 APOBEC except for *TERT* that has only one “true” APOBEC-derived hotspot mutation (C250T). *TP53*  
405 is another gene that is frequently affected by driver ApoHM, however, only one out of nine is a “true”  
406 APOBEC-derived mutation. The pan-cancer overview exhibits the distribution of driver ApoHM in  
407 APOBEC-enriched cancers, and the power gained by integrating 3859 samples revealed that of all  
408 potential drivers, only 31 might be “true” APOBEC-derived mutations.

409

410

## 411 DISCUSSION

412 In this study, we describe the genomic landscape of APOBEC-driven tumors, characterize ApoHM and  
413 identify potential cancer drivers in mUC. The in-depth analysis of 115 whole-genomes of mUC identified  
414 chromatin accessibility, hairpin-loop stability and specific sequences within the hairpin-loop as variables  
415 associated with ApoHM. These variables, in combination with the mutational context, were used to  
416 identify ApoHM that were more frequently mutated than expected by chance.

417 The substrate of APOBEC enzymes is single-stranded DNA (ssDNA)<sup>1</sup>, which has led to the following  
418 conflicting hypotheses: 1) APOBEC enzymes are mainly active during replication<sup>32</sup> and 2) APOBEC is  
419 mainly active in open chromatin and transcriptionally active genomic regions<sup>33</sup>. The equal distribution of  
420 all APOBEC-associated mutations across genomic regions supports the hypothesis that these  
421 mutations had been generated during replication when APOBEC enzymes have equal access to ssDNA  
422 across the genome<sup>32</sup>. However, kataegis which has previously been linked to APOBEC activity<sup>34</sup>, and  
423 ApoHM were enriched in high DNA accessible and highly transcribed regions. This observation  
424 reconciliates both views claiming that APOBEC is active during DNA replication (non-clustered and non-  
425 hotspot mutations) and transcriptionally active regions (clustered and hotspot mutations). Additionally,  
426 we observed that APOBEC3A-preferred YTCA mutations are dominant in mUC and are evenly  
427 distributed across genomic regions. This result is in line with experimental observations in human cancer  
428 cell lines<sup>34</sup>, suggesting that APOBEC3A is the main driver of APOBEC mutagenesis. For YTCA  
429 mutations as well as for APOBEC3B-preferred RTCA mutations, the gap between the number of  
430 mutations across genomic regions is smaller at higher APOBEC mutagenesis, which strongly suggests  
431 that APOBEC enzymes do not have a preference for specific genomic regions. However, the enrichment  
432 of highly frequent ApoHM ( $n \geq 4$ ) in open chromatin, of which many are drivers according to our analysis,  
433 may imply a functional effect of these putative driver mutations occurring near gene regulatory  
434 elements<sup>22</sup>.

435 The extensive examination of ApoHM in hairpin-loops revealed twin mutations we termed didymi that  
436 are characterized by a unique mutational pattern. Didymi comprises the APOBEC SBS2 signature and  
437 an unknown signature delineated by C>T mutations in the ApC context. It is remarkable to see only  
438 one of the APOBEC signatures in didymi loci, when they usually appear together in tumor samples  
439 with APOBEC mutagenesis<sup>35,36</sup>. The fact that only C>T mutations that characterize SBS2 are present  
440 in didymi suggests that these mutations may arise predominantly by replication across the uracil

441 bases<sup>37,38</sup> and that the mechanisms to generate C>G mutations that characterize SBS13 are not  
442 operational in this context. Furthermore, the unknown mutational signature of didymi correlates with  
443 spontaneous deamination (SBS1) and defective DNA mismatch repair signatures (SBS6, SBS15,  
444 SBS20 and SBS26), suggesting a potentially different mechanism in ApC from the putative TpC  
445 APOBEC mutations. Although there is a strong correlation with APOBEC mutagenesis<sup>21</sup>, it is unclear if  
446 both mutations in didymi loci are direct targets of APOBEC3A or whether the non-TpC mutations are  
447 just the result of spontaneous deamination followed by DNA mismatch repair.

448 Compared to breast cancer, twice as many bladder cancer tumors were APOBEC-high (19% vs. 41%).  
449 Most driver ApoHM were cancer-specific, possibly reflecting different selective pressures that each  
450 cancer type endured. In both tumor types, APOBEC-low patients had enrichment for HR deficiency,  
451 while APOBEC-high tumors had high tumor mutational burden which may indicate different treatment  
452 options for these two groups of patients with different levels of APOBEC mutagenesis<sup>39-42</sup>.

453 The in-depth analysis performed in this study to characterize the mutational landscape of APOBEC  
454 mutagenesis, revealed the correlation of ApoHM with the stability of DNA hairpin-loops, DNA  
455 accessibility and the potential for didymi associated with these recurrent mutations. These features are  
456 key to modeling the background distribution of hotspot mutations and identifying potential drivers. Most  
457 potential driver ApoHM were in the non-protein coding regions, including didymi. The similar frequency  
458 of these drivers in the metastatic and primary settings of UC indicates a general phenomenon in UC  
459 and the drivers could cause early events of tumorigenesis of UC. APOBEC-associated mutations have  
460 also been identified in normal tissue<sup>43,44</sup>, however, it is unclear if the driver ApoHM we report here are  
461 also present in healthy tissue and to what extent they contribute to cancer development from normal  
462 cells. Nevertheless, experimental validation will be needed to confirm the cancer driver status of these  
463 hotspot mutations.

464 Although several hotspot mutations are defined as drivers by our model, the inclusion of other cancer  
465 types, increased the statistical power revealing that only 31 driver ApoHM have a strong correlation with  
466 APOBEC, and may be considered “true” APOBEC-derived hotspot mutations.

467 In this study, we characterized the genomic landscape of APOBEC-driven mUC and identified novel  
468 mechanisms of genomic alteration patterns associated with APOBEC mutagenesis. The mutational  
469 signatures associated with DNA hairpin-loops targeted by APOBEC in two distinct hotspot positions are

470 unique, demonstrating the exclusive mutational signature of APOBEC-derived hotspot mutations. These  
471 findings were confirmed in non-metastatic UC and in metastatic breast cancer. Additional studies are  
472 needed to clarify the role of APOBEC in these recurrent twin mutations. Also, the enrichment of ApoHM  
473 and kataegis in high DNA accessible regions, suggests a different mechanism compared to the general  
474 APOBEC mutagenesis (non-hotspot mutations) that seems to occur independently of genomic regions,  
475 which may be linked to different mechanisms of APOBEC3A and APOBEC3B<sup>13</sup>. As APOBEC is a major  
476 source of hotspot mutations, it is crucial to identify those in coding and non-coding regions of whole-  
477 genomes that may play an important role in cancer development. The statistical framework we  
478 developed could aid to identify potential driver hotspot mutations derived from APOBEC activity which  
479 may offer novel targeted therapy options for APOBEC-driven cancer patients.

480

#### 481 **Limitations of the study**

482 Despite the thorough analysis we have performed, caution should be exercised when considering these  
483 outliers as true APOBEC-derived driver hotspot mutations as other unknown factors may still explain  
484 the distribution of these highly frequent APOBEC-related hotspot mutations. The sample size for some  
485 tumor types is a limitation when identifying driver hotspot mutations in a cancer-specific manner as we  
486 did.

## 487 **ACKNOWLEDGEMENTS**

488 Hartwig Medical Foundation and the Center of Personalized Cancer Treatment are acknowledged for  
489 making the clinical, genomic and transcriptomic data available. We would also like to thank the Pan-  
490 Cancer Analysis of Whole Genomes study for sharing the genomic data from 23 primary UC patients.  
491 We are particularly grateful to all participating patients and their families. The Stichting Dutch Uro-  
492 Oncology Study (DUOS) group and the Daniel den Hoed Foundation supported this research. J. Alberto  
493 Nakauma-González was collectively supported by the national funding organization the Dutch Cancer  
494 Society (KWF, the Netherlands) under the framework of the ERA-NET TRANSCAN-2 initiative.

495

## 496 **AUTHORS CONTRIBUTIONS**

497 Conceptualization: JAN and HJGvdW; Methodology: JAN and HJGvdW; Software: JAN, HJGvdW and  
498 MTWN; Validation: JAN and HJGvdW; Formal Analysis: JAN and HJGvdW; Investigation: All authors;  
499 Resources: JLB, JWMM; Data Curation: JAN, MR and HJGvdW; Writing – Original Draft: JAN and  
500 HJGvdW; Writing – Review & Editing: All authors; Visualization: JAN; Supervision: JLB, HJGvdW, MPJL  
501 and JWMM; Project Administration: JAN, HJGvdW, MPJL and JLB; Funding Acquisition: JLB, HJGvdW,  
502 and MPJL.

503

## 504 **DECLARATION OF INTERESTS**

505 Joost L. Boormans has received research support from Merck AG / Pfizer, Janssen and Merck Sharp &  
506 Dohme, and consultancy fees from Merck Sharp & Dohme, Bristol-Myers Squibb, Astellas, AstraZeneca,  
507 Ipsen and Janssen (all paid to the Erasmus MC Cancer Institute).

508 Martijn P. J. Lolkema has received research support from JnJ, Sanofi, Astellas and MSD, and  
509 consultancy fees from Incyte, Amgen, JnJ, Bayer, Servier, Roche, INCa, Pfizer, Sanofi, Astellas,  
510 AstraZeneca, Merck Sharp & Dohme, Novartis, Julius Clinical and the Hartwig Medical Foundation (all  
511 paid to the Erasmus MC Cancer Institute).

512 John W. M. Martens has received research support from Pfizer, Sanofi, GSK, Therawis Cergentis, and  
513 Philips (all paid to the Erasmus MC Cancer Institute) and one consultancy fee from Novartis.

514 Astrid A.M. van der Veldt has received consultancy fees from BMS, MSD, Pfizer, Novartis, Eisai, Sanofi,  
515 Pierre Fabre, Ipsen and Roche (all paid to the Erasmus MC Cancer Institute).

516 J. Alberto Nakauma-González, Maud Rijnders, Minouk T.W. Noordsij and Harmen J. G. van de Werken

517 declare no competing interests.

518



519 **STAR METHODS**

520

521 **RESOURCE AVAILABILITY**

522 **Lead contact**

523 Further information and requests for resources and reagents should be directed to and will be fulfilled  
524 by the lead contact, Harmen J. G. van de Werken ([h.vandewerken@erasmusmc.nl](mailto:h.vandewerken@erasmusmc.nl)).

525 **Materials availability**

526 This study did not generate new unique reagents.

527 **Data and code availability**

528 WGS, RNA-seq and clinical data from mUC, mBC and from other metastatic cancers are available  
529 through the Hartwig Medical Foundation at <https://www.hartwigmedicalfoundation.nl>, under request  
530 numbers DR-314, DR-026 and DR-041 respectively. For mUC, samples that were previously analyzed  
531 (DR-031) by Nakauma et al.<sup>10</sup> were retrieved from DR-314. WGS data from primary UC was requested  
532 to the NCBI dbGAP and granted access through request #33427.

533 ChIPseq, replication timing and methylation data experiments, are freely available through The  
534 ENCODE Project Consortium<sup>46</sup> and the Roadmap Epigenomics Consortium<sup>47</sup> on the ENCODE portal  
535 (<https://www.encodeproject.org>)<sup>45</sup>.

536 The scripts, including the algorithm to find hairpin-loops and estimate the thermodynamic stability have  
537 been deposited in a public repository available at [https://github.com/ANakauma/ApobecHM\\_drivers](https://github.com/ANakauma/ApobecHM_drivers).  
538 Additionally, the version v1.0.0 of the code used for this study (ApobecHM\_drivers) is available at  
539 Zenodo (<https://doi.org/10.5281/zenodo.10362579>)<sup>48</sup>. Pre-processed WGS data was provided by the  
540 Hartwig Medical Foundation and scripts are available at <https://github.com/hartwigmedical/hmftools>.  
541 R2CPCT v0.4 was used for additional processing of the WGS (<https://github.com/J0bbie/R2CPCT>),

542

543

544 **EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS**

545 **Patient cohorts**

546 The mUC cohort of this study has been previously described (NCT01855477 and NCT02925234)<sup>10</sup>. In  
547 short, patients with advanced or mUC were prospectively enrolled in these multicenter clinical trials and

548 were scheduled for 1st or 2nd line palliative systemic treatment. Following protocols of the Hartwig  
549 Medical Foundation (HMF)<sup>49</sup>, WGS, with a depth close to 100X<sup>10</sup>, was successfully performed on DNA  
550 from freshly obtained biopsies from metastatic sites, and matched RNA-sequencing (RNA-seq) was  
551 available for 90 patients (97 samples). Sequential biopsies of a metastatic lesion taken at the time of  
552 clinical or radiological disease progression from eight patients were additionally sequenced. Similarly,  
553 the cohorts of other cancer types have been previously described<sup>29</sup>, and the DNA extraction and  
554 sequencing were performed according to the HMF protocols<sup>49</sup>. Only cancer types with >50 samples  
555 were included in the analysis.

556

## 557 **METHOD DETAILS**

### 558 **Whole-genome sequencing and analysis**

559 Alignment and pre-processing of WGS data, detection of genomic subtypes, HR deficiency, MSI,  
560 structural variants, chromothripsis events, APOBEC mutagenesis, and pathway activity have been  
561 previously described<sup>4,10,49,50</sup>. Mutational signatures and kataegis were detected with MutationalPatterns  
562 v3.10.0<sup>51</sup> and Katdetectr v1.2.0<sup>52</sup>. APOBEC enriched tumors (adj.  $p < 0.01$ , otherwise non-APOBEC  
563 tumors) were classified as high when the fold enrichment ( $E$ ) for C>T and C>G mutations in TCW ( $W =$   
564 A or T) context was  $E \geq 3$ , medium when  $2 \leq E < 3$  and low when  $E < 2$ . Similarly, the fold enrichment  
565 for C>T and C>G mutations in the tetra YTCA ( $Y = T$  or C) and RTCA ( $R = G$  or A) context was  
566 calculated.

567

### 568 **Clonal fraction and cancer cell fraction**

569 The clonal fraction of mutations was estimated as previously described<sup>53</sup>. Correcting for tumor purity  
570 and copy number, the variant copy number  $n_{SNV}$  of each SNV was calculated as follows

$$571 \quad n_{SNV} = \frac{f_m}{p} [pC_t + (1 - p)C_h], \quad (1)$$

572 where  $f_m$  is the relative frequency of the mutant variant reads,  $p$  is the tumor purity,  $C_t$  is the copy number  
573 affecting the region where a particular SNV was located and  $C_h$  is the healthy copy number (2 for  
574 autosomes and 1 for allosomes). In this study, mutations were considered clonal when the variant copy  
575 number was >0.75.

576 To identify the proportion of cancer cells carrying a specific mutation, the cancer cell fraction (CCF) was  
577 estimated as previously described<sup>54,55</sup>. Given the number of reference and mutant reads and assuming  
578 binomial distribution, we estimated the expected number of allelic copies ( $n_{chr}$ ) carrying the observed  
579 SNV resulting from  $f_m$  values when the mutation is present in 1, 2, 3, ...,  $N_{chr}$  allelic copies. The resulting  
580 estimated  $n_{chr}$  with the maximum likelihood is used to calculate the CCF as  $n_{SNV}/n_{chr}$ .

581

## 582 **Mutational load across genomic regions**

583 The genome was divided into regions (bins) of one mega base-pair (Mbp). The number of SNVs was  
584 counted in each bin, and the mean number of SNVs was estimated from the entire cohort. These values  
585 represented the average SNVs/Mbp reflecting the mutational load in each genomic region. The average  
586 SNVs/Mbp was smoothed by applying a moving average with a window of  $k = 3$ . For visualization  
587 reasons, in Figure 2 a  $k = 9$  was used.

588

## 589 **DNA accessibility estimation (ChIPseq)**

590 ChIPseq data for healthy urinary bladder, breast and other tissues of adult humans (H3K4me1,  
591 H3K4me3, H3K36me3 and H3K27ac) were downloaded from the ENCODE portal  
592 (<https://www.encodeproject.org>) to our local server. The bed.gz files were imported with narrowPeak  
593 format for analysis. Only peaks with  $q < 0.05$  were kept for analysis. The signal of each experiment was  
594 divided into regions of one Mbp, and a moving average with  $k = 3$  bins was applied. The signals were  
595 normalized using the mean and standard deviation. This procedure was applied to each chromosome.  
596 The sum of all four ChIPseq experiments was considered an approximation of DNA accessibility. High  
597 DNA accessible regions (open chromatin) had values above the median considering the whole genome.  
598 All other regions were considered to be of low DNA accessibility (condensed chromatin). DNA  
599 accessibility for all healthy tissues is available in **Table S7** (for urinary bladder see **Table S4** along with  
600 other covariates). In case that matched normal ChIPseq with tumor type was not available, an average  
601 of all ChIPseq experiments was used.

602

603 **Detection of hairpin loops**

604 All SNVs were assessed to determine whether they occur in the loop of hairpin-loop structures and their  
605 thermodynamic stability. A total of 50 bases upstream and downstream of the mutation site were  
606 considered. The minimum length of the stem was 2 base-pairs and the minimum and maximum loop  
607 size was 3 and 10 bases, respectively (not considering the closing base-pair). Since multiple  
608 configurations are possible, only the structure with the highest stability was considered. One mismatch  
609 was allowed which could be either a non-matching base-pair or a single nucleotide bulge loop.

610

611 **Stability of hairpin loops**

612 We implemented the nearest neighbor stability algorithm (NNSA) to estimate the thermodynamic  
613 stability of hairpin-loops . This algorithm calculates the Gibbs free energy ( $\Delta G$ ) of the DNA hairpin-loop  
614 structure based on known biophysics properties of the base pairs and their interactions<sup>56,57</sup>.

615 The NNSA was applied to the target DNA sequence allowing mismatches in the stem. Each base-pair  
616 contributes to the stability of the stem considering the immediate neighboring base-pairs. Adding the  
617 local  $\Delta G$ 's calculated per base and accounting for the size and sequence of the loop results in a final  $\Delta G$   
618 for the whole hairpin-loop structure. All parameters are available in the literature and the UNAFold web  
619 server was used to infer parameters for mismatches<sup>56-58</sup>. For loops larger than 4, no data was available  
620 for specific sequences and only the loop size was considered.

621

622 **Driver hotspot mutations**

623 To identify driver hotspot mutations, all genomic positions with 2 or more mutations were considered for  
624 analysis. Hotspot mutations were divided into two groups either located within loops of hairpin-loops or  
625 outside of these DNA structures. For hotspot mutations outside of loops, only those in the TpC context  
626 were considered, as these are likely initiated by APOBEC enzymes. For hotspot mutations falling within  
627 loops, all hotspot mutations in NpC (N = any base) context were considered, as APOBEC3A may also  
628 mutate these bases that are not in the TpC context<sup>21</sup>.

629 In cancer, only a few somatic mutations are drivers, while the vast majority are passenger mutations<sup>30</sup>.

630 Under this consideration, we modeled the distribution of the remaining hotspot mutations as a Poisson  
631 process per focal hotspot mutation. For more accurate modeling, we considered the tri-nucleotide

632 context TCW (TCA, TCC, TCG, TCT). In the case of hotspot mutations that do not occur in hairpin loops,  
633 DNA accessibility was used as a predictor variable that can influence the distribution of hotspot  
634 mutations. We modelled this in R as `model_TCW_noloop=glm(n_mut_genpos+DNA_access)`. Where,  
635 `n_muts_genpos` is a vector with the number of mutations per genomic position linked to the DNA  
636 accessibility region (`DNA_access`). Accessibility regions were divided into 10 regions based on  
637 percentiles. Since DNA-accessibility varies per tumor-tissue of origin<sup>59,60</sup>, this was estimated for mUC,  
638 mBC and other tumor types using ChIPseq experiments from normal tissue as described above.

639 In the case of hotspot mutations within hairpin-loops, the model was extended to include mutations in  
640 non-TCW context (grouped as ApC, CpC or GpC context), the hairpin loop stability (`hairpin_stab`) and  
641 the DNA sequence in the loop (`loopSeq`):  
642 `model_TCW_loop=n_mut_genpos+DNA_access+hairpin_stab+loopSeq`. The loop sequence was a  
643 binary variable to indicate whether the mutation occurred in the following sequence: 1001 or 101 (0 =  
644 A/T, 1 = G/C; underlined is the position of the hotspot mutation).

645 Using these models, we estimated the expected number of mutations of the specified hotspot mutation  
646 for which the model was built. Then, the exact Poisson test was applied to estimate the significance of  
647 observing the same or more mutations than expected in a specific genomic position. P-values were  
648 Benjamini-Hochberg adjusted. In rare occasions, only a few mutations (<2) were available to represent  
649 the background distribution of a particular tri-nucleotide. To include these ApoHM in the analysis, a  
650 model that did not consider the tri-nucleotide was used instead.

651

## 652 **RNA-sequencing**

653 Alignment, pre-processing of RNA-seq data and transcript normalization have been previously  
654 described<sup>10,29</sup>. The transcriptomic subtype of each mUC sample was identified when the mean  
655 (normalized) expression of all genes associated with a specific subtype was the highest across all  
656 subtypes.

657

## 658 **mRNA editing**

659 Jalili, et al. identified hotspot mutations in the mRNA of *DDOST* that is targeted by APOBEC3A<sup>17</sup>. The  
660 genomic position of this hotspot mutation reveals a hairpin-loop structure that is an ideal substrate for

661 APOBEC3A. Due to the short life-time of mRNA molecules, the presence of this hotspot mutation  
662 reflects ongoing APOBEC mutagenesis. The proportion of C>U mutations in chr1:20981977 was  
663 estimated to identify the RNA-editing activity of APOEBC3A.

664

#### 665 **Transcriptome expression data mapped to genomic regions**

666 MultiBamSummary from deepTools v1.30.0<sup>61</sup> was used to read BAM files and estimate the number of  
667 reads in genomic regions with a size of one Mbp. The average raw read count per Mbp was calculated,  
668 and a moving average with  $k = 3$  bins was applied. The scale of the read counts was normalized per  
669 chromosome using the mean and standard deviation. High transcriptional regions were defined as such  
670 when the expression value of one region was above the median of the whole genome.

671

#### 672 **Simulations and power calculation**

673 A synthetic genome with 1,000,000 hotspot mutations was reconstructed from the original cohort of  
674 mUC. To reach the number of hotspot mutations, non-hotspot mutations were randomly selected and  
675 the number of mutations per genomic position was drawn from a Poisson distribution using the empirical  
676 lambda from the mUC cohort. The same number of driver ApoHM identified in mUC were simulated as  
677 hypothetical drivers to replicate a 3%-15% prevalence. Hypothetical cohorts with 10-500 samples were  
678 simulated 100 times, using a random number of ApoHM derived from the empirical distribution of the  
679 mUC cohort. The statistical power was estimated as the proportion of driver ApoHM that were correctly  
680 identified. The performance of the model on simulated cohorts of 100 samples, was also tested with  
681 other genomic covariates. These covariates were replication timing and methylation from HeLa cell lines,  
682 and the proportion of GC content from the reference build hg19.

683

## 684 **QUANTIFICATION AND STATISTICAL ANALYSIS**

### 685 **Statistical analysis**

686 Analyses were performed using the statistical analysis platform R v4.1.0<sup>62</sup>. Fisher's exact, Wilcoxon-  
687 rank sum and Wilcoxon signed-rank tests were used for comparison between groups. The correlation  
688 coefficients of continuous values with categorical values were estimated with logistic regression analysis  
689 applying the Wald test. Residuals for QQ plots and the Kolmogorov–Smirnov test were estimated using

690 DHARMa v.0.4.6<sup>63</sup>. The exact Poisson test was applied to identify potential driver hotspot mutations.  
691 The Poisson-binomial method was applied for mutually exclusive mutation events using Rediscover  
692 v0.3.2<sup>64</sup> and the Fisher's exact test was applied for the significance of co-occurred mutations. In all  
693 cases, p-values were adjusted using the Benjamini-Hochberg method.

## 694 **SUPPLEMENTAL INFORMATION TITLES AND LEGENDS**

695

696 Supplementary Figures. Figures S1-S12

697 Supplementary Tables. Tables S1-S7, Related to Figures 3-7

698



## 699 References

- 700 1. Roberts, S.A., Lawrence, M.S., Klimczak, L.J., Grimm, S.A., Fargo, D., Stojanov, P., Kiezun, A., Kryukov,  
701 G.V., Carter, S.L., Saksena, G., et al. (2013). An APOBEC cytidine deaminase mutagenesis pattern is  
702 widespread in human cancers. *Nat. Genet.* *45*, 970-976. 10.1038/ng.2702.
- 703 2. Tate, J.G., Bamford, S., Jubb, H.C., Sondka, Z., Beare, D.M., Bindal, N., Boutselakis, H., Cole, C.G.,  
704 Creatore, C., Dawson, E., et al. (2019). COSMIC: The Catalogue Of Somatic Mutations In Cancer. *Nucleic*  
705 *Acids. Res.* *47*, D941-D947. 10.1093/nar/gky1015.
- 706 3. Boichard, A., Pham, T.V., Yeerna, H., Goodman, A., Tamayo, P., Lippman, S., Frampton, G.M., Tsigelny,  
707 I.F., and Kurzrock, R. (2019). APOBEC-related mutagenesis and neo-peptide hydrophobicity: implications  
708 for response to immunotherapy. *Oncolmunology* *8*, 1550341. 10.1080/2162402X.2018.1550341.
- 709 4. Rijnders, M., Nakauma-González, J.A., Robbrecht, D.G.J., Gil-Jimenez, A., Balcioglu, H.E., Oostvogels,  
710 A.A.M., Aarts, M.J.B., Boormans, J.L., Hamberg, P., van der Heijden, M.S., et al. (2024). Gene-expression-  
711 based T-Cell-to-Stroma Enrichment (TSE) score predicts response to immune checkpoint inhibitors in  
712 urothelial cancer. *Nat. Commun.* *15*, 1349. 10.1038/s41467-024-45714-0.
- 713 5. Law, E.K., Levin-Klein, R., Jarvis, M.C., Kim, H., Argyris, P.P., Carpenter, M.A., Starrett, G.J., Temiz, N.A.,  
714 Larson, L.K., Durfee, C., et al. (2020). APOBEC3A catalyzes mutation and drives carcinogenesis in vivo.  
715 *J. Exp. Med.* *217*. 10.1084/jem.20200261.
- 716 6. Sharma, S., and Baysal, B.E. (2017). Stem-loop structure preference for site-specific RNA editing by  
717 APOBEC3A and APOBEC3G. *PeerJ* *5*, e4136. 10.7717/peerj.4136.
- 718 7. Kumar, S., Warrell, J., Li, S., McGillivray, P.D., Meyerson, W., Salichos, L., Harmanci, A., Martinez-  
719 Fundichely, A., Chan, C.W.Y., Nielsen, M.M., et al. (2020). Passenger Mutations in More Than 2,500  
720 Cancer Genomes: Overall Molecular Functional Impact and Consequences. *Cell* *180*, 915-927.e916.  
721 <https://doi.org/10.1016/j.cell.2020.01.032>.
- 722 8. Shi, M.J., Meng, X.Y., Fontugne, J., Chen, C.L., Radvanyi, F., and Bernard-Pierrot, I. (2020). Identification  
723 of new driver and passenger mutations within APOBEC-induced hotspot mutations in bladder cancer.  
724 *Genome Med.* *12*, 85-85. 10.1186/s13073-020-00781-y.
- 725 9. Wong, J.K.L., Aichmüller, C., Schulze, M., Hlevnjak, M., Elgaafary, S., Lichter, P., and Zapatka, M. (2022).  
726 Association of mutation signature effectuating processes with mutation hotspots in driver genes and non-  
727 coding regions. *Nat. Commun.* *13*, 178. 10.1038/s41467-021-27792-6.
- 728 10. Nakauma-González, J.A., Rijnders, M., van Riet, J., van der Heijden, M.S., Voortman, J., Cuppen, E.,  
729 Mehra, N., van Wilpe, S., Oosting, S.F., Rijstenberg, L.L., et al. (2022). Comprehensive Molecular  
730 Characterization Reveals Genomic and Transcriptomic Subtypes of Metastatic Urothelial Carcinoma. *Eur.*  
731 *Urol.* *81*, 331-336. <https://doi.org/10.1016/j.eururo.2022.01.026>.
- 732 11. Chan, K., Roberts, S.A., Klimczak, L.J., Sterling, J.F., Saini, N., Malc, E.P., Kim, J., Kwiatkowski, D.J.,  
733 Fargo, D.C., Mieczkowski, P.A., et al. (2015). An APOBEC3A hypermutation signature is distinguishable  
734 from the signature of background mutagenesis by APOBEC3B in human cancers. *Nat. Genet.* *47*, 1067-  
735 1072. 10.1038/ng.3378.
- 736 12. Roelofs, P.A., Timmermans, M.A.M., Stefanovska, B., den Boestert, M.A., van den Borne, A.W.M.,  
737 Balcioglu, H.E., Trapman, A.M., Harris, R.S., Martens, J.W.M., and Span, P.N. (2023). Aberrant  
738 APOBEC3B Expression in Breast Cancer Is Linked to Proliferation and Cell Cycle Phase. *Cells* *12*, 1185.
- 739 13. Hirabayashi, S., Shirakawa, K., Horisawa, Y., Matsumoto, T., Matsui, H., Yamazaki, H., Sarca, A.D.,  
740 Kazuma, Y., Nomura, R., Konishi, Y., et al. (2021). APOBEC3B is preferentially expressed at the G2/M  
741 phase of cell cycle. *Biochem. Biophys. Res. Commun.* *546*, 178-184.  
742 <https://doi.org/10.1016/j.bbrc.2021.02.008>.
- 743 14. Cortez, L.M., Brown, A.L., Dennis, M.A., Collins, C.D., Brown, A.J., Mitchell, D., Mertz, T.M., and Roberts,  
744 S.A. (2019). APOBEC3A is a prominent cytidine deaminase in breast cancer. *PLoS Genet.* *15*, e1008545-  
745 e1008545. 10.1371/journal.pgen.1008545.
- 746 15. Glaser, A.P., Fantini, D., Wang, Y., Yu, Y., Rimar, K.J., Podojil, J.R., Miller, S.D., and Meeks, J.J. (2018).  
747 APOBEC-mediated mutagenesis in urothelial carcinoma is associated with improved survival, mutations  
748 in DNA damage response genes, and immune response. *Oncotarget* *9*, 4537-4548.  
749 10.18632/oncotarget.23344.
- 750 16. Robertson, A.G., Kim, J., Al-Ahmadie, H., Bellmunt, J., Guo, G., Cherniack, A.D., Hinoue, T., Laird, P.W.,  
751 Hoadley, K.A., Akbani, R., et al. (2017). Comprehensive Molecular Characterization of Muscle-Invasive  
752 Bladder Cancer. *Cell* *171*, 540-556.e525. 10.1016/J.CELL.2017.09.007.
- 753 17. Jalili, P., Bowen, D., Langenbacher, A., Park, S., Aguirre, K., Corcoran, R.B., Fleischman, A.G., Lawrence,  
754 M.S., Zou, L., and Buisson, R. (2020). Quantification of ongoing APOBEC3A activity in tumor cells by  
755 monitoring RNA editing at hotspots. *Nat. Commun.* *11*, 2971. 10.1038/s41467-020-16802-8.
- 756 18. Allory, Y., Beukers, W., Sagrera, A., Flández, M., Marqués, M., Márquez, M., Van Der Keur, K.A., Dyrskjot,  
757 L., Lurkin, I., Vermeij, M., et al. (2014). Telomerase reverse transcriptase promoter mutations in bladder  
758 cancer: High frequency across stages, detection in urine, and lack of association with outcome. *Eur. Urol.*  
759 *65*, 360-366. 10.1016/j.eururo.2013.08.052.
- 760 19. Siraj, A.K., Bu, R., Iqbal, K., Parvathareddy, S.K., Siraj, N., Siraj, S., Diaz, M.R.F., Rala, D.R., Benito, A.D.,  
761 Sabido, M.A., et al. (2020). Telomerase reverse transcriptase promoter mutations in cancers derived from  
762 multiple organ sites among middle eastern population. *Genomics* *112*, 1746-1753.  
763 <https://doi.org/10.1016/j.ygeno.2019.09.017>.

- 764 20. Buisson, R., Langenbucher, A., Bowen, D., Kwan, E.E., Benes, C.H., Zou, L., and Lawrence, M.S. (2019).  
765 Passenger hotspot mutations in cancer driven by APOBEC3A and mesoscale genomic features. *Science*  
766 364, eaaw2872-eaaw2872. [10.1126/science.aaw2872](https://doi.org/10.1126/science.aaw2872).
- 767 21. Langenbucher, A., Bowen, D., Sakhtemani, R., Bournique, E., Wise, J.F., Zou, L., Bhagwat, A.S., Buisson,  
768 R., and Lawrence, M.S. (2021). An extended APOBEC3A mutation signature in cancer. *Nat. Commun.* 12,  
769 1602. [10.1038/s41467-021-21891-0](https://doi.org/10.1038/s41467-021-21891-0).
- 770 22. Wu, S., Ou, T., Xing, N., Lu, J., Wan, S., Wang, C., Zhang, X., Yang, F., Huang, Y., and Cai, Z. (2019).  
771 Whole-genome sequencing identifies ADGRG6 enhancer mutations and FRS2 duplications as  
772 angiogenesis-related drivers in bladder cancer. *Nat. Commun.* 10, 1-12. [10.1038/s41467-019-08576-5](https://doi.org/10.1038/s41467-019-08576-5).
- 773 23. An, L., Dong, C., Li, J., Chen, J., Yuan, J., Huang, J., Chan, K.M., Yu, C.-h., and Huen, M.S.Y. (2018).  
774 RNF169 limits 53BP1 deposition at DSBs to stimulate single-strand annealing repair. *Proc. Natl. Acad.*  
775 *Sci. U.S.A.* 115, E8286-E8295. doi:10.1073/pnas.1804823115.
- 776 24. Cheng, Y.-C., Chiang, H.-Y., Cheng, S.-J., Chang, H.-W., Li, Y.-J., and Shieh, S.-Y. (2020). Loss of the  
777 tumor suppressor BTG3 drives a pro-angiogenic tumor microenvironment through HIF-1 activation. *Cell*  
778 *Death Dis.* 11, 1046. [10.1038/s41419-020-03248-5](https://doi.org/10.1038/s41419-020-03248-5).
- 779 25. Fischer, J.-P., Els-Heindl, S., and Beck-Sickinger, A.G. (2020). Adrenomedullin – Current perspective on  
780 a peptide hormone with significant therapeutic potential. *Peptides* 131, 170347.  
781 <https://doi.org/10.1016/j.peptides.2020.170347>.
- 782 26. Levine, A.J., and Brivanlou, A.H. (2006). GDF3 at the crossroads of TGF-beta signaling. *Cell Cycle* 5,  
783 1069-1073. [10.4161/cc.5.10.2771](https://doi.org/10.4161/cc.5.10.2771).
- 784 27. Lo, Y.-H., Romes, E.M., Pillon, M.C., Sobhany, M., and Stanley, R.E. (2017). Structural Analysis Reveals  
785 Features of Ribosome Assembly Factor Nsa1/WDR74 Important for Localization and Interaction with  
786 Rix7/NVL2. *Structure* 25, 762-772.e764. <https://doi.org/10.1016/j.str.2017.03.008>.
- 787 28. Campbell, P.J., Getz, G., Korbel, J.O., Stuart, J.M., Jennings, J.L., Stein, L.D., Perry, M.D., Nahal-Bose,  
788 H.K., Ouellette, B.F.F., Li, C.H., et al. (2020). Pan-cancer analysis of whole genomes. *Nature* 578, 82-93.  
789 [10.1038/s41586-020-1969-6](https://doi.org/10.1038/s41586-020-1969-6).
- 790 29. Angus, L., Smid, M., Wiltng, S.M., van Riet, J., Van Hoeck, A., Nguyen, L., Nik-Zainal, S., Steenbruggen,  
791 T.G., Tjan-Heijnen, V.C.G., Labots, M., et al. (2019). The genomic landscape of metastatic breast cancer  
792 highlights changes in mutation and signature frequencies. *Nat. Genet.* 51, 1450-1458. [10.1038/s41588-](https://doi.org/10.1038/s41588-019-0507-7)  
793 [019-0507-7](https://doi.org/10.1038/s41588-019-0507-7).
- 794 30. Martincorena, I., Raine, K.M., Gerstung, M., Dawson, K.J., Haase, K., Van Loo, P., Davies, H., Stratton,  
795 M.R., and Campbell, P.J. (2017). Universal Patterns of Selection in Cancer and Somatic Tissues. *Cell* 171,  
796 1029-1041.e1021. [10.1016/j.cell.2017.09.042](https://doi.org/10.1016/j.cell.2017.09.042).
- 797 31. Juul, R.I., Nielsen, M.M., Juul, M., Feuerbach, L., and Pedersen, J.S. (2021). The landscape and driver  
798 potential of site-specific hotspots across cancer genomes. *NPJ Genom. Med.* 6, 33. [10.1038/s41525-021-](https://doi.org/10.1038/s41525-021-00197-6)  
799 [00197-6](https://doi.org/10.1038/s41525-021-00197-6).
- 800 32. Hoopes, James I.I., Cortez, Luis M.M., Mertz, Tony M.M., Malc, Ewa P.P., Mieczkowski, Piotr A.A., and  
801 Roberts, Steven A.A. (2016). APOBEC3A and APOBEC3B Preferentially Deaminate the Lagging Strand  
802 Template during DNA Replication. *Cell Rep.* 14, 1273-1282.
- 803 33. Kazanov, M.D., Roberts, S.A., Polak, P., Stamatoyannopoulos, J., Klimczak, L.J., Gordenin, D.A., and  
804 Sunyaev, S.R. (2015). APOBEC-Induced Cancer Mutations Are Uniquely Enriched in Early-Replicating,  
805 Gene-Dense, and Active Chromatin Regions. *Cell Rep.* 13, 1103-1109.
- 806 34. Petljak, M., Dananberg, A., Chu, K., Bergstrom, E.N., Striepen, J., von Morgen, P., Chen, Y., Shah, H.,  
807 Sale, J.E., Alexandrov, L.B., et al. (2022). Mechanisms of APOBEC3 mutagenesis in human cancer cells.  
808 *Nature* 607, 799-807. [10.1038/s41586-022-04972-y](https://doi.org/10.1038/s41586-022-04972-y).
- 809 35. Petljak, M., Alexandrov, L.B., Brammied, J.S., Price, S., Wedge, D.C., Grossmann, S., Dawson, K.J., Ju,  
810 Y.S., Iorio, F., Tubio, J.M.C., et al. (2019). Characterizing Mutational Signatures in Human Cancer Cell  
811 Lines Reveals Episodic APOBEC Mutagenesis. *Cell* 176, 1282-1294.e1220. [10.1016/j.cell.2019.02.012](https://doi.org/10.1016/j.cell.2019.02.012).
- 812 36. Alexandrov, L.B., Kim, J., Haradhvala, N.J., Huang, M.N., Tian Ng, A.W., Wu, Y., Boot, A., Covington,  
813 K.R., Gordenin, D.A., Bergstrom, E.N., et al. (2020). The repertoire of mutational signatures in human  
814 cancer. *Nature* 578, 94-101. [10.1038/s41586-020-1943-3](https://doi.org/10.1038/s41586-020-1943-3).
- 815 37. Helleday, T., Eshlad, S., and Nik-Zainal, S. (2014). Mechanisms underlying mutational signatures in  
816 human cancers. *Nat. Rev. Genet.* 15, 585-598. [10.1038/nrg3729](https://doi.org/10.1038/nrg3729).
- 817 38. Petljak, M., and Maciejowski, J. (2020). Molecular origins of APOBEC-associated mutations in cancer.  
818 *DNA Repair* 94, 102905. <https://doi.org/10.1016/j.dnarep.2020.102905>.
- 819 39. Patel, M., Nowsheen, S., Maraboyina, S., and Xia, F. (2020). The role of poly(ADP-ribose) polymerase  
820 inhibitors in the treatment of cancer and methods to overcome resistance: a review. *Cell Biosci.* 10, 35.  
821 [10.1186/s13578-020-00390-7](https://doi.org/10.1186/s13578-020-00390-7).
- 822 40. Jekimovs, C., Bolderson, E., Suraweera, A., Adams, M., O'Byrne, K.J., and Richard, D.J. (2014).  
823 Chemotherapeutic Compounds Targeting the DNA Double-Strand Break Repair Pathways: The Good, the  
824 Bad, and the Promising. *Front. Oncol.* 4, 10.3389/fonc.2014.00086.
- 825 41. Gillyard, T., and Davis, J. (2021). Chapter Two - DNA double-strand break repair in cancer: A path to  
826 achieving precision medicine. In *International Review of Cell and Molecular Biology*, U. Weyemi, and L.  
827 Galluzzi, eds. (Academic Press), pp. 111-137. <https://doi.org/10.1016/bs.ircmb.2021.06.003>.
- 828 42. Marabelle, A., Fakih, M., Lopez, J., Shah, M., Shapira-Frommer, R., Nakagawa, K., Chung, H.C., Kindler,  
829 H.L., Lopez-Martin, J.A., Miller, W.H., Jr., et al. (2020). Association of tumour mutational burden with  
830 outcomes in patients with advanced solid tumours treated with pembrolizumab: prospective biomarker

- 831 analysis of the multicohort, open-label, phase 2 KEYNOTE-158 study. *Lancet Oncol.* 21, 1353-1365.  
832 10.1016/s1470-2045(20)30445-9.
- 833 43. Lawson, A.R.J., Abascal, F., Coorens, T.H.H., Hooks, Y., O'Neill, L., Latimer, C., Raine, K., Sanders, M.A.,  
834 Warren, A.Y., Mahubani, K.T.A., et al. (2020). Extensive heterogeneity in somatic mutation and selection  
835 in the human bladder. *Science* 370, 75-82. 10.1126/science.aba8347.
- 836 44. Wang, Y., Robinson, P.S., Coorens, T.H.H., Moore, L., Lee-Six, H., Noorani, A., Sanders, M.A., Jung, H.,  
837 Katainen, R., Heuschkel, R., et al. (2023). APOBEC mutagenesis is a common process in normal human  
838 small intestine. *Nat. Genet.* 55, 246-254. 10.1038/s41588-022-01296-5.
- 839 45. Davis, C.A., Hitz, B.C., Sloan, C.A., Chan, E.T., Davidson, J.M., Gabdank, I., Hilton, J.A., Jain, K.,  
840 Baymuradov, U.K., Narayanan, A.K., et al. (2018). The Encyclopedia of DNA elements (ENCODE): data  
841 portal update. *Nucleic. Acids. Res.* 46, D794-D801. 10.1093/nar/gkx1081.
- 842 46. Dunham, I., Kundaje, A., Aldred, S.F., Collins, P.J., Davis, C.A., Doyle, F., Epstein, C.B., Frietze, S.,  
843 Harrow, J., Kaul, R., et al. (2012). An integrated encyclopedia of DNA elements in the human genome.  
844 *Nature* 489, 57-74. 10.1038/nature11247.
- 845 47. Roadmap Epigenomics, C., Kundaje, A., Meuleman, W., Ernst, J., Bilenky, M., Yen, A., Heravi-Moussavi,  
846 A., Kheradpour, P., Zhang, Z., Wang, J., et al. (2015). Integrative analysis of 111 reference human  
847 epigenomes. *Nature* 518, 317-329. 10.1038/nature14248.
- 848 48. Nakauma-González, J.A. (2023). Whole-genome mapping of APOBEC mutagenesis in metastatic  
849 urothelial carcinoma identifies driver hotspot mutations and a novel mutational signature. Zenodo.  
850 <https://doi.org/10.5281/zenodo.10362579>.
- 851 49. Priestley, P., Baber, J., Lolkema, M.P., Steeghs, N., de Bruijn, E., Shale, C., Duyvesteyn, K., Haidari, S.,  
852 van Hoeck, A., Onstenk, W., et al. (2019). Pan-cancer whole-genome analyses of metastatic solid tumours.  
853 *Nature* 575, 210-216. 10.1038/s41586-019-1689-y.
- 854 50. van Dessel, L.F., van Riet, J., Smits, M., Zhu, Y., Hamberg, P., van der Heijden, M.S., Bergman, A.M., van  
855 Oort, I.M., de Wit, R., Voest, E.E., et al. (2019). The genomic landscape of metastatic castration-resistant  
856 prostate cancers reveals multiple distinct genotypes with potential clinical impact. *Nat. Commun.* 10, 1-13.  
857 10.1038/s41467-019-13084-7.
- 858 51. Blokzijl, F., Janssen, R., van Boxtel, R., and Cuppen, E. (2018). MutationalPatterns: Comprehensive  
859 genome-wide analysis of mutational processes. *Genome Med.* 10, 33-33. 10.1186/s13073-018-0539-0.
- 860 52. Hazelaar, D.M., van Riet, J., Hoogstrate, Y., and van de Werken, H.J.G. (2023). Katdetectr: an  
861 R/bioconductor package utilizing unsupervised changepoint analysis for robust kataegis detection.  
862 *GigaScience* 12, giad081. 10.1093/gigascience/giad081.
- 863 53. Stephens, P.J., Tarpey, P.S., Davies, H., Van Loo, P., Greenman, C., Wedge, D.C., Nik-Zainal, S., Martin,  
864 S., Varela, I., Bignell, G.R., et al. (2012). The landscape of cancer genes and mutational processes in  
865 breast cancer. *Nature* 486, 400-404. 10.1038/nature11017.
- 866 54. Bolli, N., Avet-Loiseau, H., Wedge, D.C., Van Loo, P., Alexandrov, L.B., Martincorena, I., Dawson, K.J.,  
867 Iorio, F., Nik-Zainal, S., Bignell, G.R., et al. (2014). Heterogeneity of genomic evolution and mutational  
868 profiles in multiple myeloma. *Nat. Commun.* 5, 2997-2997. 10.1038/ncomms3997.
- 869 55. Gundem, G., Van Loo, P., Kremeyer, B., Alexandrov, L.B., Tubio, J.M.C., Papaemmanuil, E., Brewer, D.S.,  
870 Kallio, H.M.L., Högnäs, G., Annala, M., et al. (2015). The evolutionary history of lethal metastatic prostate  
871 cancer. *Nature* 520, 353-357. 10.1038/nature14347.
- 872 56. SantaLucia, J. (1998). A unified view of polymer, dumbbell, and oligonucleotide DNA nearest-neighbor  
873 thermodynamics. *Proc. Natl. Acad. Sci. U.S.A.* 95, 1460-1465. doi:10.1073/pnas.95.4.1460.
- 874 57. John SantaLucia, J., and Hicks, D. (2004). The Thermodynamics of DNA Structural Motifs. *Annu. Rev.*  
875 *Biophys. Biomol. Struct.* 33, 415-440. 10.1146/annurev.biophys.32.110601.141800.
- 876 58. Zuker, M. (2003). Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic. Acids.*  
877 *Res.* 31, 3406-3415. 10.1093/nar/gkg595.
- 878 59. Polak, P., Karlič, R., Koren, A., Thurman, R., Sandstrom, R., Lawrence, M.S., Reynolds, A., Rynes, E.,  
879 Vlahoviček, K., Stamatoyannopoulos, J.A., and Sunyaev, S.R. (2015). Cell-of-origin chromatin  
880 organization shapes the mutational landscape of cancer. *Nature* 518, 360-364. 10.1038/nature14221.
- 881 60. Kübler, K., Karlič, R., Haradhvala, N.J., Ha, K., Kim, J., Kuzman, M., Jiao, W., Gakkhar, S., Mouw, K.W.,  
882 Braunstein, L.Z., et al. (2019). Tumor mutational landscape is a record of the pre-malignant state. *bioRxiv*,  
883 517565. 10.1101/517565.
- 884 61. Ramírez, F., Ryan, D.P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne, S., Dündar, F., and  
885 Manke, T. (2016). deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic.*  
886 *Acids. Res.* 44, W160-W165. 10.1093/nar/gkw257.
- 887 62. Team, R.C. (2017). R Core Team (2017). R: A language and environment for statistical computing. R  
888 Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>. R Foundation for  
889 Statistical Computing-R Foundation for Statistical Computing.
- 890 63. Hartig, F. (2022). DHARMA: residual diagnostics for hierarchical (multi-level/mixed) regression models.  
891 <https://florianhartig.github.io/DHARMA/>.
- 892 64. Ferrer-Bonsoms, J.A., Jareno, L., and Rubio, A. (2021). Rediscover: an R package to identify mutually  
893 exclusive mutations. *Bioinformatics* 38, 844-845. 10.1093/bioinformatics/btab709.

894