# 1 Lessons learned from operationalizing the integration of nutrition-specific and nutrition-

# 2 sensitive interventions in rural Ethiopia

- 3 Girmay Ayana Mersha<sup>1,2\*</sup>, Eshetu Zerihun Tariku <sup>2,3</sup>, Wanzahun Godana Boynito<sup>2,3</sup>, Meseret
- 4 Woldeyohaness<sup>1</sup>, Birhanu Wodajo<sup>1</sup>, Tadese Kebebe<sup>1</sup>, Stefaan De Henauw<sup>2</sup>, Souheila
- 5 Abbeddou<sup>2</sup>
- <sup>6</sup> <sup>1</sup>Food Science and Nutrition Research Directorate, Ethiopian Public Health Institute, Ethiopia.
- <sup>7</sup> <sup>2</sup>Public Health Nutrition, Department of Public Health and Primary Care, Ghent University,
- 8 Belgium.
- 9 <sup>3</sup>School of Public Health, Arba Minch University, Arba Minch, Ethiopia
- 10 \*Corresponding author: <u>GirmayAyana.Mersha@UGent.be</u>

11

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

# 12 Abstract

## 13 Background

14 Undernutrition reduction requires coordinated efforts across sectors to address its causes. A 15 multisectoral approach is important in diagnosing the problem and identifying solutions that 16 would be implemented across different sectors.

The study aims to explore the experience of health and agriculture extension workers in integrating nutrition services provided to households with children under two years of age at community level.

# 20 Methods

A qualitative study has been conducted in agrarian areas of Ethiopia in 2021 at the end of a large intervention program to reduce stunting. In total, 28 key informant interviews were conducted with health- and agriculture-extension workers and mothers. A framework analysis approach was applied to manage and analyze data using NVivo version 12 software.

### 25 Results

Community-level joint intervention is a feasible approach to integrating nutrition services. 26 Farm gardening and cooking demonstrations were practiced jointly with extension workers and 27 households. Because of service integration, extension workers perceived an improved father's 28 role in supporting mothers in childcaring and feeding nutritious diets to children, and decreased 29 severe cases of undernutrition. Integration of health and agriculture sectors for nutrition 30 intervention was challenged by the high workload on extension workers, poor supervision and 31 leadership commitment, lack of appropriate agricultural inputs, and absence of clarity on sector-32 specific roles. In some areas nutrition services are not owned by the health and agriculture 33 sectors, and it was overlooked. 34

# 35 Conclusion

- Integrating existing community-level platforms are key entry points to address undernutrition and promote key agriculture and health interventions in Ethiopia. Empowering extension workers through training, monitoring, and supervision is part of ensuring the sustainability of integration between sectors.
- 40 Keywords: Community health, agriculture, multisectoral, nutrition, Ethiopia

# 42 Introduction

Undernutrition including stunting, wasting, underweight and micronutrient deficiencies 43 plagued populations in low- and middle-income countries for decades and remains one of the 44 most pressing health problems. Young children and women of reproductive age are particularly 45 vulnerable because of their increased nutrient requirements, and, in case of women, their lower 46 social status, which limits their access to nutrient-rich foods [1]. The causes of undernutrition 47 are complex and affected by factors acting at different causal hierarchical levels [2]. The 48 49 political, economy, health, education, agriculture and food systems, water and sanitation, and the environment are all determinants of maternal and child nutrition and health [3]. 50 51 Undernutrition reduction requires coordinated efforts across sectors to address its underlying 52 causes [4, 5]. Addressing the problem of undernutrition needs the collaboration of multiple sectors and a variety of stakeholders in governments, non-government organizations, and the 53 private sector [6]. Undernutrition reduction requires a departure from a health system-centered 54 approach of provision of nutritional services and nutrition-specific interventions only, to an 55 integrated approach that encompasses nutrition-sensitive, food systems-based approaches[7]. 56

The framework of actions proposed by the UNICEF and the Lancet nutrition series recognizes 57 the needs for multiple actions that are at different levels from immediate, underlying to distal 58 [2, 8]. Interventions targeting the improvement of maternal and child nutrition and health act 59 on interlinked factors which influence each other. Nutrition-specific interventions address the 60 61 immediate causes of undernutrition and nutrition-sensitive interventions focus on the underlying causes of good nutrition [9]. Both nutrition-sensitive and specific interventions 62 require an enabling environment, that encompasses actions addressing basic determinants of 63 64 undernutrition. Multiple actions to improve nutrition were undertaken over the recent years, the most prominent is Scaling Up Nutrition which clearly underscores the necessity of actions 65

among all the stakeholders [10]. This approach puts nutrition at the core of each sector plan andinstitutional effort.

68 However, systematic coordination between different sectors to achieve the objectives of improved maternal and child nutrition is problematic, given the bureaucratic barriers that 69 70 characterize the administrative division of responsibilities among sectors. All sectors are expected to be clear about what they can contribute to resolving malnutrition in a particular 71 context specific to their sector role. Agriculture, as an example of nutrition-sensitive 72 intervention, can potentially increase food availability, which would directly improve the 73 dietary accessibly leading to improved nutritional status of targeted populations. However, the 74 multiple factors that influence nutrition, the social dynamics within households that can result 75 in unexpected allocations of resources, and the many indirect ways that agriculture may impact 76 77 nutrition, require purposeful planning and careful consideration of how exactly program inputs will lead to nutritional improvements. This complex relationships has consequently lead to 78 79 limited impact of nutrition intervention, because other determinants are not put in place at different levels. 80

Ethiopia is one of the countries with the highest rate of undernutrition in sub-Saharan Africa. 81 The proportions of stunting, underweight and wasting among children under age of five years 82 are respectively 38%, 24% and 10% in Ethiopia [11]. Multiple efforts have been made to 83 implement the government commitment towards ending undernutrition. To reach this goal, 84 Ethiopia has demonstrated a strong policy commitment to nutrition through the development of 85 a multisectoral food and nutrition policy, which highlights the need to accelerate multi-sectoral 86 approaches [12]. Political willingness is an important factor that helps to implement effective 87 nutrition interventions [10]. However, the Ethiopian situation remains in its debuts with poorly 88 synchronized sectors of food, nutrition, agriculture, and health to prevent the problem of under-89 nutrition. In Ethiopia, the presence of health and agriculture extension systems and frontline 90

workers at the community level is a good framework that facilitates the synergetic linkage of 91 92 nutrition within health and agriculture services [13] Additionally, existing multisectoral nutrition coordination structure at all levels, and the willingness of international and national 93 development partners to support multisectoral nutrition interventions are favorable conditions 94 on the ground [14]. Ethiopia has implemented multiple nutrition-specific and nutrition-sensitive 95 programs such as the Ethiopian national nutrition program, and the Segota declaration program, 96 97 that have required the development of structures and processes that facilitate multisectoral coordination [15]. But, collaborations between the health and agriculture sectors are suboptimal 98 at community levels [16]. The multi-sectoral nutrition programs and interventions lack strong 99 100 coordination that brings on board all the relevant stakeholders.

Based on these facts, Ethiopia designed different intervention modalities whose objective is to 101 102 strengthen community-level multisectoral nutrition interventions. In this context, the Sustainable Undernutrition Reduction program in Ethiopia (SURE) was developed and 103 104 implemented between 2016 and 2020 in rural regions of Ethiopia. The program utilizes the potentials of the Agricultural Development Agents (DAs) and the Health Extension Workers 105 (HEWs) for the promotion of improved agricultural practices and utilization of high nutrient 106 products to promote dietary diversification among rural mothers and children. At present, there 107 is no study that has analyzed the experience and perception of the health (HEWs) and 108 agriculture extension workers (AEWs) in integrating nutrition services provision at the 109 community level in rural Ethiopia. Understanding frontline workers' practices and experience 110 is crucial for identifying appropriate measures to sustain nutrition improvements [9]. This 111 qualitative study aims at appraising the understanding, the knowledge and perception of those 112 women and men who are the most in contact with the communities in rural Ethiopia and who 113 were responsible to implement the SURE program. 114

# 115 Methods

## 116 Study Setting

The Ethiopian Ministry of Health and the Ministry of Agriculture led the implementation of the 117 SURE program whose aim was to reduce stunting by improving complementary feeding 118 practices and dietary diversity of young children through a process of improving local 119 agricultural productivity, food accessibility, food consumption diversity and nutritional status 120 of young children and women [27]. SURE, was implemented between 2016 and 2020 in four 121 agrarian regions (Oromia, Amhara, Southern Nations, Nationalities, and Peoples' Region 122 [SNNPR], and Tigray). The program package included promoting diversified agriculture, 123 promoting infant and young child feeding (IYCF) practices, empowering women's decision-124 making related to agriculture, diversified food intake, and health, and enhancing food security 125 126 and water, sanitation, and hygiene practices (WASH). The SURE intervention districts were selected because of the high food insecurity and stunting prevalence in four regions out of the 127 nine total regions in Ethiopia. Details of the SURE intervention program have been published 128 elsewhere [17]. 129

130 Briefly, a total of 36 districts (Woreda as locally called) were selected and assigned to the SURE intervention. Comparison control districts were selected in equal number (36 districts) in the 131 same four agrarian regional strata as for intervention districts. Districts were roughly matched 132 based on stunting prevalence tertiles (low, medium or high) and food insecurity scores. Eligible 133 households are households with at least one child under the age of 2 years. The intervention 134 135 started in 2016 and included 4299 households, 2146 households from 144 Kebeles (smallest administrative unit) were included in the intervention group and 2153 from 144 Kebeles in the 136 137 control group. After baseline data collection, and all households received counselling at household and community levels. Additionally, 10 % of households with children aged under 138

two years in the intervention regions (they can be enrolled or not in the larger SURE
intervention) were 1) provided improved seed variety and chicken through AEWs and 2) trained
through demonstration gardens at farmer training centers.

The SURE program was delivered through: 1) interpersonal contacts to provide counselling on IYCF and nutrition-sensitive agriculture advice to mothers and fathers of children under 24 months, inclusive of pregnant women and fathers-to-be, jointly delivered by the local HEWs and AEWs during routine household visits; 2) men's and women's group dialogues targeting all men and all women in a given community network, facilitated also by the local HEWs and AEWs, and 3) media campaign to reinforce IYCF and dietary diversity messages.

# 148 Study design

149 A phenomenological qualitative study design was conducted among beneficiaries (mothers)

and implementers (HEWs and AEWs) of the SURE program in the intervention communities.

151 One Kebele per district from the intervention group was randomly selected to contribute in the 152 qualitative study.

#### 153 Study period

This qualitative study was conducted after four years of intervention. The data were collected at the community level in selected districts of Amhara, Oromia, and SNNPR between January and February 2021. Data collection was discontinued in Tigray because of the war.

# 157 Recruitment of participants

All SURE intervention districts and Kebeles in the three agrarian regions of Ethiopia were eligible for inclusion. The recruitment of participants was done purposely to obtain maximum variation of information. The HEWs and AEWs as service providers of the intervention packages, and mothers who were enrolled in the SURE intervention as direct beneficiaries were

invited to participate in this study. The HEWs and AEWs were eligible to participate in this study if they received the SURE program training in 2016 and have provided the service as per their training in their respective Kebele for a minimum of two years. Mothers should be residents of the Kebele, would have had a child 0-24 months old at the time the intervention started, and received joint health and agriculture counseling as described in the study setting.

#### 167 Data collection and tools

Data were collected using key informant interviews (KIIs) at the community level from a total of 28 eligible respondents including HEWs, AEWs and mothers. Participants were asked about their understanding of nutrition, quality of service provided and quality of nutrition education specific to the SURE program, the challenges of implementing nutrition programs jointly, and how the program could be sustained for a better nutrition service provision (Supplementary material 1).

Participants took part in a face-to-face interview. A topic guide was developed by the research team and training on how to conduct KIIs was provided. The interview began with demographic questions about schooling, experience, and professional background. All the interviews were conducted in a private space and were audio recorded. Data was collected in the local languages (Amharic and Oromifa). Audio recordings were transcribed and translated into English. All transcribed data was anonymized.

#### 180 Data Analysis:

A framework analysis approach was used to manage and analyze the data. Framework approach provides clear steps to follow and produces structured outputs of summarized data that can be used with a variety of data sources, including interviews, focus groups, and documents [18]. The steps include familiarization (becoming familiar with the content of the data), Identification of thematic areas (identifying a thematic framework, which is a set of predetermined themes or

categories that are relevant to the research question), Indexing (systematically coding the data 186 187 to identify content related to each theme or category), and Interpretation (identifying patterns and relationships across the data, and interpreting what these patterns and relationships mean 188 in relation to the research question). Qualitative data analysis software NVIVO version 12 189 software was used in coding and constructing thematic areas from the collected data. The 190 transcripts were coded into specific themes. The themes were identified and coded based on the 191 192 topic guide as nutrition understanding, achievements, opportunities, and way forwards for a successful implementation of nutrition-specific and nutrition-sensitive programs and 193 sustainability. 194

# **195** Ethical considerations:

Ethical approval was obtained from the research ethic committees both at the University 196 Hospital of Ghent, Belgium (protocol number BC-08862), and the Ethiopian Public Health 197 Institute, Ethiopia (protocol number EPHI-IRB-306-2020). The qualitative study included 198 199 recipients from the same cohort (participating mothers from the intervention group), in addition 200 to health extension workers from the intervention Kebeles. Written informed consents, which included permission to be recorded using a digital audio recorder, were obtained from 201 participants before interviews. Further administrative permissions were granted by local 202 government bureaus in all regions. 203

# 204 Results

# 205 Demographic characteristics of study participants

From the total of twenty-eight key informants who participated in the study, 43%, 39% and 18% were from Oromia, Amhara and SNNPR regions, respectively. The proportion of HEWs, AEWs and mothers were almost similar. Totally 58 % of the extension workers were female

- and 73% had at least five years of work experience. Sixty-seven percent of the study participants
- 210 have completed college and above.
- Table 1: Demographic characteristics of the qualitative study participants, Ethiopia

Demographic Characteristic(N=28)	Frequency (%)
Age of participants (in years)	
20-30	16 (57.1)
31-40	10 (35.7)
41-50	2 (7.1)
Region	
Amhara	11 (39.3)
Oromia	12 (42.9)
SNNPR	5 (17.9)
Role	
Agriculture extension worker	9 (32.1)
Health extension worker	10 (35.1)
Mother	9 (32.1)
Sex of participants	
Female	20 (71.4)
Male	8 (28.6)
Health and agriculture extension workers experience	
2-4 years	5 (26.3)
5-10 years	7 (36.8)
>10 years	7 (36.8)
Education status of participants	

No formal education	3 (10.7)
Primary school	6 (21.4)
Technical/vocational college	15 (53.6)
Diploma and above	4 (14.3)

212

# 213 Major causes of undernutrition

The causes of undernutrition are seasonal and context-specific. The key informants perceived that, in their community, undernutrition was due to a shortage of land to produce adequate food to sell or to consume for a rapidly growing population. The severity of the problem increases when it is coupled with poor distribution of agricultural supplies, and pest infestation that affect productivity.

Absence of enough farmland is the cause of undernutrition. Since the community is large in
number, farmland is not enough. on average the farmland owned by the farmers is about 0.375
up to 0.5 hectare. AEW,SNNPR.

The problem of land and water scarcity were identified as the main cause of undernutrition in the community in Oromia region. The small plot of land is challenged by a shortage of water and agricultural inputs.

225 The lack of adequate farming place also another problem because it is difficult to produce

enough food for family consumption using one farming place and leads to poverty. Moreover,

227 lack of adequate rain, lack of irrigation service, and the fertilizer supply is not as required in

228 *our kebele, which may lead to poverty. HEW, Oromia.* 

Access to fruits and vegetables is important to improve food security and undernutrition reduction. Frequent drought and absence of irrigation facilities to farm during the dry season

are among the causes of undernutrition that undermine the efforts of AEWs and households toimprove food and nutrition interventions.

233 Malnutrition is mainly due to insufficient production of fruits and vegetables due to drought in 234 this area. When the husbands and wives agree to produce in their home garden, it will vanish, 235 because there is no irrigation. Due to this reason our community failed to produce fruits and 236 vegetables and give less emphasis. HEW, Oromia.

War and civil unrest affect access to- and availability of foods through destructing market linkage. Instability compromises the support of HEWS and AEWs to the community. An extension worker indicated that the causes of undernutrition are related to lack of hygiene and sanitation, diseases and infections, inflation and war that displaced people from their residential areas and hindered them from producing food.

The cause of under nutrition are lack of hygiene, disease, war, and such things are the cause of
under nutrition in addition to the above, there are food security issues. For instance, there is a
lack of milk for children, they can't access enough milk. There is also limited access to eggs.
HEW, SNNPR.

Now in some places, based on market inflation, some people become incapable of buying something, as a result, the child may develop malnutrition. For example, when children need milk, they may fail to give it. As a child fails to get balanced food when they need and passed that time, they may go into malnutrition. Some households are economically incapable to care for their child, there is a household that relies on the market business for leading the family, such a woman gave other foods to their child before six months and leave the child at home because they can't stay home and breastfeed their child regularly. HEW, Oromia.

253

### 254 Challenges of integrating health and agriculture interventions in Ethiopia

255 Many challenges have been reported by extension workers in the implementation of 256 intersectoral interventions. High workload, conflict, shortage of agricultural inputs, limited 257 training and poor supportive supervision and COVID-19 pandemic were the most frequently 258 reported challenges (Figure 1).

259 Intervention in the midst of the war in Ethiopia

Ethiopia is facing civil unrest in the last four years and the project intervention has been affected. Conflict was a challenge of service delivery at community level, which resulted in destruction of facilities, damages of means, and loss of income by restricting people's movement and targeted attacks on workers.

264The current situation is overly sensitive. It is now three years since the conflict began. Every265day there are riots, there is something. So, people think it is turning to something else. At the266beginning of the SURE program, the expert was there to support them every day, and the people267were getting education and had been given training. It was good to take care of that and268practice. But now people are abandoning everything and entering politics. "Who will die today?269Who will live today?' They are living in fear. In this regard, political stability is a bit difficult.270AEW,SNNPR.

Conflict and the COVID pandemic have also compromised the supportive supervision and
monitoring activities by restricting people's movement from regional and zonal levels to front
line workers.

274 ... So, this team has met frequently, and minutes have been kept on our discussion. This meeting
275 was previously held every week, then turns to every month. As I said earlier, recently we have
276 less engagement due to varied factors and we do not have minutes. We fail to gather the
277 participants to give different training as this contradicts the covid-19 restrictive measures.
278 HEW, Oromia.

14

## 279 Shortage of agricultural inputs and material for workshop training

The availability of agricultural inputs for gardening and cooking demonstrations at community level was a challenge and participants were not interested in bringing their own food and material for demonstration. The project gives agricultural inputs to only 10% of the total population, but the demand for agriculture inputs was high and the community perceived that the project does not treat people equally.

- 285 We are very much interested in the work (for example a demonstration of food preparation).
- 286 For this demonstration, if you asked the farmers to bring the ingredients/inputs (like egg, kale,
- 287 *cereals, ...) from their home they will lose their interest. It is difficult to bring by yourself too.*
- 288 The issue is the problem of access to input for the demonstration. AEW, Oromia.

#### 289 Workload and ownership of the program

Nutrition activities are not a priority agenda in all sectors equally. The ownership of nutrition is not clearly defined, and extension experts assume an additional assignment is beyond their regular tasks. There is a lack of awareness and commitment by AEWs to implement nutrition activities at a community level. AEWs assume that tackling issues related to women's and children's nutrition to be the role of HEWs.

Emm... there is often a seasonal job. For example, in the health sector, they will have activities such as the vaccination campaign, as well as in the agriculture sector, such as spring, harvest, watershed and irrigation. We do not keep our appointment because we give priority to our main work. Especially in health, there are increased immunization programs, increased follow up programs. AEW, SNNPR.

Integrating agriculture and health services is challenged by shortage of workforce. Food and nutrition-related interventions are not a priority for the agriculture sector. The leadership does not prioritize nutrition, and they always focus on campaigns that aim to achieve a sector-specific activity like vaccination in the health sector and water shedding and plantation in the agriculture

sector. Extension workers assume that nutrition activities require additional experts assigned to
 the job. The service was also compromised by the absence of continuous monitoring and
 evaluation mechanisms.

307 The main reason was being overburdened or having overlapping duties on our side. The 308 agricultural extension workers push these SURE program activities on us, by mentioning that it 309 is not their duty. We also mentioned that these SURE program duties are their responsibility. Both sectors push the duties to each other because of work overload. Therefore, we emphasize 310 311 only a single duty expected from our major career and letting others perform the SURE duties. 312 We met and discussed the issues only when stakeholders from the zone came to our kebele for supervision, otherwise, it was overlooked by both of us. No one continuously monitors the 313 314 progress as result, it will be forgotten. If this is not the case, we would perform it successfully in collaboration with AEW. HEW, Oromia. 315

316 Absence of adequate training, supervision, and technical support

Supportive supervision and technical support during the implementation of new program is a key factor for the success of a project. The SURE program was designed in a way that supervisors contribute continuously during the whole implementation period. The support provided during the program implementation was limited and many extension workers perceived the support and supervision provided were not adequate.

Let alone frequent training, if at least the concerned bodies at zone level visit our activities and provide the necessary feedback, it will motivate us for further improvement. It is pretty good if those in the zone were actively involved. As said earlier there is no problem at kebele level but at zone level, it seems an overlooked opportunity. HEW, Oromia.

When the support is provided in an interrupted way, the community perceives the activities/supports are useless. If the support comes in an organized manner and reaches the kebele (for the farmers) hierarchically, the farmers did not oppose it. They will curiously take

329 *the supports and adapt to their home/household. But, if the support seems loose from its origin,* 

330 *the community considers it useless. AEW, Oromia.* 

# 331 Lesson learned and sustainability of health and agriculture intersectoral intervention

The SURE intervention packages depended on multi sectoral coordination at the local and district levels (Figure 1). However most agricultural interventions were perceived as the distribution of poultry and other agricultural materials. Although most extension workers captured the intention of the target population, as being mothers, young children and those who are most vulnerable households.

SURE, program covered mainly hen distribution for targeted population segments. For
example, the program distributed up to 6 hens per household. Especially for the lowest quantiles
of fathers and mothers [socioeconomic], lactating mothers, and children less than 2 years of
age. SURE, also supplies seed for vegetables. Training is also given for both the professional
and community. AEW, Amhara.

In the beginning, we provided training. After we took 15 days of training at the start of the program, we mainstreamed this to the level of community. Then, in all kebeles, chicken (for egg production) was purchased and provided for 20 individuals per kebele. Different productive agricultural materials like water pumps and others were given to support them in the production of fruits and vegetables in their backyard. AEW, Oromia.

Health and agriculture extension workers jointly visited households with under two years of age children and provided age-appropriate IYCF counselling and advice on nutrition-sensitive agriculture to improve dietary diversity. Job aids have been developed with food calendars and recipe books for each region. Extension workers describe their experience of joint visit as follows:

we are working together on SURE program. The kebele has a man's and women's development team
[community-level volunteers who provide support to health and agriculture extension workers in

mobilizing communities to deliver service]. We and the health extension staff are going from house 354 to house and teaching them how to feed their families, especially pregnant mothers, and children 355 356 under two years of age, how to grow vegetables and animals on their limited land and use them at 357 home without taking them to market. When we teach together, we or the agriculture extension workers through the men's development team, teach them how to produce, what to sow, how to care 358 for and feed their children together, how to feed nutritious food and how to feed at least six different 359 types of food. The health extension worker shows them cooking demonstration. For example, they 360 361 teach them what to mix, how to prepare and how often to feed. AEW, SNNPR. 362 We gather women twice a month in their group to provide training. We have materials and different

food items on our demonstration site, we show them how to prepare and feed soft foods like porridge
for children. We advise women to prepare food available in the home based on their capacity. For
instance, they sell eggs and buy cabbage, we advise them to avoid this habit and spend time feeding
the child. HEW, Oromia.

Mothers witnessed the service received by extension workers in their community. Although, asking mothers to come with food ingredients could be one factor that limits mothers from attending cooking demonstration sessions.

From HEW I have heard about the preparation of porridge from food items available at home like
cereals (bean, chickpea and pea), seeds (teff, wheat, oats, millet an barley) vegetables and animal
products (meat, egg, milk etc). I have also learnt about feeding the children only breast milk until
the age of six months. .... From the AEW, I have learnt about chicken for baby and garden for baby.
They taught us, as we must have vegetables in our garden and animal products at home. If these are
not available at home, we are expected to buy from the market and prepare porridge from these
variety of food items to feed children. Mother, Oromia.

Farm demonstration gardens were established at the farmer's training centers to facilitate
coordination between the HEW managing the health posts and the AEWs working on the farms.
Farm demonstration gardens were developed and managed by AEWs. Appropriate varieties of

vegetables were selected according to agroecology. Harvests from the farmer's training centers
were used for cooking demonstrations. Cooking demonstrations were managed by HEWs at
health posts near the farmer's training centers. Women also donate ingredients from their homes
to contribute to the exercise.

We give agricultural demonstrations on every occasion. Whenever we get the best performance, we give training and demonstrations on farmer's land or on farmer's training centers to act as the best performed farmers. We also give demonstrations as per the farmers' needs and questions. SURE, the program recommends demonstration every month. We give porridge demonstration for both women and men developmental army every month. Before a year this was done every month

389 *consistently. Currently we give demonstrations every 2 months or longer. AEW, Amhara.* 

When they advised us, they said in recent days, even not the rural, the urban communities are practicing home gardening with the limited land they have in the backyard. Different types of vegetables (tomatoes, kale, potatoes...) are on your hand; you may not need to buy them from the market, plant them. Mother, Oromia.

# 394 Knowledge gained through the SURE program

The SURE program interventions have impact on the knowledge of community to improve child caring practice, agricultural and behavioral practices in different parameters like severe undernutrition workload, women empowerment, and father's support. The extension workers have observed the changes in the community, mainly fathers' role in supporting mothers in childcaring and the importance given to feeding nutritious diet to children instead of selling the food produced. The extension workers felt that severe cases of undernourished children visiting heath centers have decreased.

402 ...after the implementation of the SURE program, we noticed a significant decline in the
403 prevalence of malnutrition. The number of children brought to our health post with malnutrition
404 has significantly declined. Before the implementation of this program, we found a lot of cases

405 and referred them. If in case we found a child with malnutrition, it is the moderate type of
406 malnutrition. The severe form of malnutrition is almost rare after the SURE program.

- 407 *Concerning women empowerment, one of the most important changes SURE bring is, engaging*
- 408 husbands in food demonstration, letting husbands start to think as they are responsible for
- 409 *childcare, for their women and they also believe that women have an equal right to decide on*
- 410 *their asset. The surprising thing was we as our area discuss many issues related to women,*
- 411 *including their right, ability, responsibility and how they can be independent economically and*
- 412 manage their home. For instance, if she generated income by breeding chicken, she could obtain
- 413 money on her own and incorporated it into the household after counselling and significant
- 414 *improvement have been observed. HEW, Oromia.*
- In the past, men thought that caring for children was a women's job. But now it has completely
  changed; men help their wives by caring for their children. AEW, SNNPR.
- 417 Importance of monitoring and regular supervision

Awareness creation activities should be organized frequently for proper implementation of nutrition interventions, and ideally by another person responsible for coordinating nutrition activities at community level. Additionally, the importance of regular meetings and evaluation for integrating services at community level was highlighted.

The first is, giving awareness creation education. For this HEWs and AEWs are responsible.
But HEWS and AEWs are busy with other tasks. Therefore, there should be other responsible
body at kebele level for this task to reduce the work overload and to avoid task overlapping
among HEWs and AEWs. This may increase the effectiveness of the SURE program.

The other is having regular meetings to monitor and evaluate the progress of the program.
During this meeting, all responsible bodies should be present. The program should have a
budget for both monitoring and evaluation as well as for supplies. The porridge and the like

- 429 *demonstration need raw materials. These materials should be funded by the SURE program.*
- 430 *AEW, Amhara.*

At the start of the program there was an evaluation session every six months and sometimes
every year. Now it has totally stopped. Especially after COVID 19. There is a plan, but the
achievement is not checked. The reason is usually lack of budget. Therefore, there should be a
budget for the monitoring and evaluation. HEW, Oromia.

# 435 **Discussion**

This study explored the experience of integrating health and agriculture sectors for improved 436 food security and nutritional status in Ethiopia. This study reported that the major cause of 437 438 undernutrition in the SURE program intervention areas are drought, conflict, and population pressure. Furthermore, the high burden of infectious diseases and WASH problems were the 439 cause of the undernutrition. The finding is aligned with the study of Mohamed (2017) which 440 found drought, population pressure, and conflict as the major causes of food security problems 441 in Ethiopia, resulting in child undernutrition and household food insecurity [19]. Similarly 442 finding on child growth in the time of drought by Hoddinott and Kinsey (2001) has shown that 443 conflict and drought increase stunting prevalence among young children in poor households 444 [20]. 445

Nutrition interventions require intensive effort from multiple sectors. The expansion of 446 governmental services from nutrition-specific to nutrition-sensitive strategies is the preferred 447 approach, as it addresses all determinants of maternal and child nutrition and health including 448 social and economic determinants [15]. Ethiopia has a national structure for multi sectoral 449 nutrition coordination anchored at the Ministry of Health and a working model at district level 450 also exists [14]. Nevertheless, according to this finding nutrition has been perceived as the work 451 of the Ministry of Health and has been framed as a treatment/preventive problem, which mainly 452 involves health sector interventions. One of the reasons to have a multisectoral nutrition policy 453 is to lift the responsibility of improving nutrition into a national concern. SURE is one of the 454 government-led programs that try to bring the health and agriculture sectors to work jointly for 455

improved dietary diversity and improved nutritional status [17, 21]. Integrating nutrition into 456 457 the national development agenda, ensuring all sectors have a mutual understanding of the importance of undernutrition prevention, and having a mechanism that helps to ensure local 458 delivery of nutrition services, through decentralization are key components of multisectoral 459 response for undernutrition reduction. SURE, is a common umbrella that brings HEWs, AEWs 460 and women to promote dietary diversity and better nutrition in Ethiopia. But we found extension 461 462 workers are overloaded by different sectoral activities which can limit their joint efforts and compromise the integration of nutrition-related activities. Similar to this study, frontline 463 workers in Uganda confront various pressures from the institutional and socio-political 464 465 environment, that limit their decisions about their daily activities [9]. Pelletier and collaborators found that in sub Saharan African countries, institutional integration of nutrition in different 466 sectors was challenged by the job description of staff, the lack of sectoral alignment on 467 468 multisectoral nutrition objectives and integration into the planning and reporting of sector activities [14]. A systematic review exploring the integration of nutrition-sensitive and specific 469 470 interventions in fragile environments shows a lack of interest in collaboration, workload and diverting to only sectoral activities as the barriers to integrating nutrition services [22]. 471

Delivering nutrition services to the local level tends to work better in countries that have 472 adequate decentralized structures. In Ethiopia, the vertical structure of the Ministry of Health 473 and Ministry of Agriculture from central to Kebele level offers an opportunity to integrate 474 nutrition services at community level. Front-line HEWs and AEWs are a critical bridge between 475 communities and service provision systems in the health and agriculture sectors. Empowering 476 extension workers, supporting them and responding to the challenges they face will be an 477 important part of ensuring the sustainability of joint activities to promote proper nutrition. 478 Therefore, appropriate training, supportive supervision and performance monitoring should be 479

in place to ensure the successful implementation of joint programs in the health and agriculturesectors.

# 482 Strengths and limitations

Our study collected data from actual implementers and beneficiaries of joint intervention program which can be used to scale up their experience. However, our study focused only on the health and agriculture sectors and did not explore the experiences of other sectors included in the Ethiopian national nutrition program.

# 487 Conclusion

Integrating existing community-level platforms are key entry points to address undernutrition and promote key agriculture and health interventions in Ethiopia. Community-based joint interventions are perceived as appropriate approaches to prevent severe cases of undernutrition and improve the role of family members in supporting mothers. Empowering extension workers through training, monitoring, and supervision, and addressing the challenges they face are part of ensuring the sustainability of integration between sectors.

494

- 495 Author contributions
- 496 Conceptualization GAM, SDH, SA
- 497 Data curation GAM, EZT, MW
- 498 Formal analysis GAM, SA
- 499 Investigation GAM, MW, EZT, MW, TK, BW
- 500 Methodology GAM, SDH, SA, TK, BW, WGB
- 501 Project administration GAM, SDH, SA, EZT, MW

- 502 Resources SDH, SA
- 503 Supervision GAM, SA, EZT, MW, TK, BW
- 504 Validation GAM, SA
- 505 Visualization GAM, SA
- 506 Writing original draft preparation GAM
- 507 Writing review and editing GAM, SDH, SA, EZT, MW, WGB, BW, TK
- 508 Funding
- 509 This work was funded by Ethiopian public health institute and the Global Minds Fund of Ghent
- 510 University<u>https://www.ugent.be/en/research/funding/devcoop/globalmindsfund.htm</u> (GRANT
- 511 GMF.CAB.2020.0025.01). The funder had no role in study design, data collection and analysis,
- decision to publish, or manuscript preparation. The funders had no role in the study design, data
- 513 collection and analysis, decision to publish or preparation of the manuscript.
- 514 Acknowledgment:
- 515 We are grateful to the study participants. In addition, our appreciation goes to the health officials 516 from central to community level, health, and agriculture extension workers for facilitating the
- 517 implementation of the study. Finally, we would like to thank our data collectors and supervisors
- 518 for their support.

# 519 Conflict of interest

520 The authors declare that there are no conflicts of interest regarding the publication of this paper.

521

# 522 References

Girard AW, Self JL, McAuliffe C, Olude O. The effects of household food production strategies
 on the health and nutrition outcomes of women and young children: a systematic review. Paediatric
 and perinatal epidemiology. 2012;26 Suppl 1:205-22. Epub 2012/07/07. doi: 10.1111/j.1365 3016.2012.01282.x. PubMed PMID: 22742612.

United Nations Children's Fund (UNICEF), World Health Organization, International Bank for
 Reconstruction and Development/The World Bank. Levels and trends in child malnutrition: key findings
 of the 2021 edition of the joint child malnutrition estimates. Geneva: World Health Organization, 2021.

530 3. Stewart CP, Iannotti L, Dewey KG, Michaelsen KF, Onyango AW. Contextualising
531 complementary feeding in a broader framework for stunting prevention. Maternal & child nutrition.
532 2013;9 Suppl 2:27-45. doi: 10.1111/mcn.12088. PubMed PMID: 24074316.

King SE, Sawadogo-Lewis T, Black RE, Roberton T. Making the health system work for the
 delivery of nutrition interventions. Maternal & child nutrition. 2021;17(1):e13056. Epub 2020/07/22.
 doi: 10.1111/mcn.13056. PubMed PMID: 32691489; PubMed Central PMCID: PMCPMC7729521.

536 5. Ouedraogo O, Doudou MH, Drabo KM, Kiburente M, Cisse D, Mesenge C, et al. Facilitating 537 factors and challenges of the implementation of multisectoral nutrition programmes at the community 538 level to improve optimal infant and young child feeding practices: a qualitative study in Burkina Faso. 539 Public Health Nutr. 2020:1-12. Epub 2020/10/01. doi: 10.1017/S136898002000347X. PubMed PMID: 540 32993837.

Reinhardt K, Fanzo J. Addressing Chronic Malnutrition through Multi-Sectoral, Sustainable
 Approaches: A Review of the Causes and Consequences. Front Nutr. 2014;1:13. Epub 2014/01/01. doi:
 10.3389/fnut.2014.00013. PubMed PMID: 25988116; PubMed Central PMCID: PMCPMC4428483.

544 7. Isabel Madzorera LJ, Robert Paarlberg, Anna Herforth, Sabri Bromage, Shibani Ghosh, Samuel
545 S Myers, William A Masters, and Wafaie W Fawzi,. Food Systems as Drivers of Optimal Nutrition and
546 Health: Complexities and Opportunities for Research and Implementation. CURRENT DEVELOPMENTS
547 IN NUTRITION. 2021.

5488.Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, de Onis M, et al. Maternal and child549undernutrition and overweight in low-income and middle-income countries. The Lancet.5502013;382(9890):427-51. doi: 10.1016/s0140-6736(13)60937-x.

Namugumya BS, Candel JJL, Talsma EF, Termeer C, Harris J. Integrating Nutrition Actions in
 Service Delivery: The Practices of Frontline Workers in Uganda. International journal of health policy
 and management. 2022. Epub 2022/05/02. doi: 10.34172/ijhpm.2022.5898. PubMed PMID:
 35490257.

555 10. Gillespie S, Haddad L, Mannar V, Menon P, Nisbett N. The politics of reducing malnutrition:
556 building commitment and accelerating progress. The Lancet. 2013;382(9891):552-69. doi:
557 10.1016/s0140-6736(13)60842-9.

558 11. Central Statistical Agency and ICF. Ethiopia Demographic and Health Survey 2016. Addis Ababa,
559 Ethiopia, and Rockville, Maryland, USA: CSA and ICF, 2016.

Kennedy E, Tessema M, Hailu T, Zerfu D, Belay A, Ayana G, et al. Multisector Nutrition Program
Governance and Implementation in Ethiopia: Opportunities and Challenges. Food Nutr Bull.
2015;36(4):534-48. Epub 2015/11/05. doi: 10.1177/0379572115611768. PubMed PMID: 26531747.

Kadiyala S, Morgan EH, Cyriac S, Margolies A, Roopnaraine T. Adapting Agriculture Platforms
for Nutrition: A Case Study of a Participatory, Video-Based Agricultural Extension Platform in India.
PLoS One. 2016;11(10):e0164002. Epub 2016/10/14. doi: 10.1371/journal.pone.0164002. PubMed
PMID: 27736897; PubMed Central PMCID: PMCPMC5063370.

Pelletier D, Gervais S, Hafeez-ur-Rehman H, Sanou D, Tumwine J. Boundary-spanning actors in
 complex adaptive governance systems: The case of multisectoral nutrition. 2017.

Bach A, Gregor E, Sridhar S, Fekadu H, Fawzi W. Multisectoral Integration of Nutrition, Health,
and Agriculture: Implementation Lessons From Ethiopia. Food and nutrition bulletin. 2020;41(2):27592. Epub 2020/03/14. doi: 10.1177/0379572119895097. PubMed PMID: 32166964.

572 16. Mersha GA, Hailu T, Kuche D, Abera A, Eshetu S, Petros A, et al. Linkages between health and 573 agriculture sectors in Ethiopia: a formative research study exploring barriers, facilitators and 574 opportunities for local level coordination to deliver nutritional programmes and services. BMC 575 Nutrition. 2017;3(1). doi: 10.1186/s40795-017-0189-4.

576 17. Moss C, Bekele TH, Salasibew MM, Sturgess J, Ayana G, Kuche D, et al. Sustainable 577 Undernutrition Reduction in Ethiopia (SURE) evaluation study: a protocol to evaluate impact, process and context of a large-scale integrated health and agriculture programme to improve complementary 578 579 feeding in Ethiopia. BMJ open. 2018;8(7):e022028. Epub 2018/07/22. doi: 10.1136/bmjopen-2018-580 022028. PubMed PMID: 30030320; PubMed Central PMCID: PMCPMC6059290.

- 581 Nicola K Gale, Gemma Heath, Elaine Cameron, Rashid S, Redwood S. Using the framework 18. 582 method for the analysis of qualitative data in multi-disciplinary health research. BMC medical research 583 methodology. 2013.
- Mohamed AA. Food Security Situation in Ethiopia: A Review Study International Journal of 584 19. 585 Health Economics and Policy. 2017;2(3):86-96. doi: 10.11648/j.hep.20170203.11.
- 586 20. Hoddinott J, Kinsey B. Child Growth in the Time of Drought. Oxford Bulletin of Economics and Statistics. 2001;63(4):409-36. doi: 10.1111/1468-0084.t01-1-00227. 587
- 588 21. Salasibew MM, Moss C, Ayana G, Kuche D, Eshetu S, Dangour AD. The fidelity and dose of 589 message delivery on infant and young child feeding practice and nutrition sensitive agriculture in 590 Ethiopia: a qualitative study from the Sustainable Undernutrition Reduction in Ethiopia (SURE) programme. J Health Popul Nutr. 2019;38(1):29. Epub 2019/10/23. doi: 10.1186/s41043-019-0187-z. 591 592 PubMed PMID: 31639070; PubMed Central PMCID: PMCPMC6805331.
- 593 22. Abdullahi LH, Rithaa GK, Muthomi B, Kyallo F, Ngina C, Hassan MA, et al. Best practices and 594 opportunities for integrating nutrition specific into nutrition sensitive interventions in fragile contexts: 595 a systematic review. BMC nutrition. 2021;7(1):46. Epub 2021/07/30. doi: 10.1186/s40795-021-00443-
- 596 1. PubMed PMID: 34321101; PubMed Central PMCID: PMCPMC8320180.

# 598 List of figures

- 599 Figure 1: Sustaining the integration of agriculture and health sectors for reduction of
- 600 undernutrition in Ethiopia
- 601
- 602 Supporting information
- 603 S1 File. Interview guide
- 604 S2 File. COREQ (Consolidated criteria for REporting Qualitative research) Checklist



Figure