- 1 2 3
- 4

5 Scanning electron microscopy observations on the

- ⁶ surface morphology of placenta villi from HIV-
- 7 positive women having preterm birth while

8 receiving antiretroviral therapy

- 9
- 10 Moses M. Obimbo^{1,2,3*}, Shem Ochieng^{'4,5}, Samwel R. Gesaka³, Julius Ogeng'o^{2,3}
- 11
- ¹Department of Obstetrics and Gynecology, University of Nairobi, Nairobi, Kenya
- 13 ² Department of Human Anatomy and Physiology, University of Nairobi, Nairobi, Kenya
- ¹⁴ ³ Basic Clinical and Translational (BCT) Research Laboratory, University of Nairobi,
- 15 Nairobi, Kenya
- ⁴ Electron Microscopy Laboratory, International Centre for Insect Physiology and
- 17 Ecology (ICIPE), Nairobi, Kenya
- 18 ⁵ School of Medicine, Kenyatta University, Nairobi, Kenya
- 19
- 20 * Corresponding author
- 21 E-mail: moses.obimbo@uonbi.ac.ke (MMO)

22 Abstract

23 **Background**: Preterm birth is a significant global issue. Antiretroviral therapy (ART) use 24 and human immunodeficiency virus (HIV) infection have both been linked in recent 25 research as independent risk factors for preterm birth. Although there has been 26 evidence linking preterm delivery to significant pathological alterations in the placenta, it 27 is still unclear how exactly HIV and ART harm the placenta and raise the risk of 28 prematurity. To explain the increased risk of preterm birth (PTB), we set out to describe 29 the surface morphological alterations in placenta villi associated with HIV and ART. 30 Methods and materials: We collected and processed 160 placentas from 40 HIV-31 positive women on ART and 40 HIV-negative women who had preterm deliveries, 40 32 HIV- positive women and 40 HIV - negative women with term delivery in Nairobi, Kenya. 33 The placenta biopsies were harvested, washed in phosphate buffer solution, and 34 processed for scanning electron microscopy. The dried tissue was mounted onto 35 specimen stubs, sputter coated with gold and visualized using Zeiss Merlin FESEM in-36 lens. Forty representative samples, 10 from each group, were randomly selected and 37 examined by investigators who were blinded to maternal HIV serostatus. 38 **Results**: The average gestational age for preterm and term births was 34 and 39 39 weeks, respectively. The average age of the mothers of preterm and term babies was 40 26.8 + 4.6 years and 24.3 + 4.3 years, respectively. The villous core of the placenta 41 from HIV-negative patients was covered with microvilli that varied in size and 42 appearance, and there were hardly any residual red blood cells. Placenta from HIVpositive women with preterm birth had widespread damage with shrunken and wrinkled 43 44 villi, predominant blunting of the microvilli, with attendant syncytiotrophoblast disruption.

- 45 and significant erythrocyte adhesion within extensive fibrillar meshwork and on surface
- 46 of the syncytium.
- 47 **Conclusion**: Our results show distinctive alterations in the placenta of HIV-positive
- 48 mothers who gave birth prematurely, which may impair the syncytium's ability to
- 49 function normally. Microvilli blunting, syncytial disruptions, and syncytial erythrocyte
- adhesion might be the symptoms of a deeper biological process. Further work to
- 51 understand the effect of HIV/ART on the syncytiotrophoblast in relation to prematurity is
- 52 recommended.

53 Introduction

54 The surface interface between fetus and mother is formed by fetal syncytiotrophoblasts. These are fused multinucleated cells generated from trophectoderm and are constantly 55 56 bathed in maternal blood. These cells are involved in exchange of nutrients, 57 electrolytes, and gases between the fetus and the mother, hormone production, and are 58 also important in fetomaternal immunotolerance [1–3]. Efficient transfer of substances across the placenta is essential in maintaining fetal growth and development and to 59 60 ensure maternal health [4–6]. The surface of the placenta syncytium exposed to the maternal blood has a brush border on their free surface: This border facilitates diverse 61 functions including absorption, secretion, and mechanotransduction [7–9]. Equally, this 62 63 surface is also exposed to toxic substances and pathogens circulating in maternal blood, putting it at risk of impaired morphology [10-12]. 64

65

Preterm birth (PTB) is a leading cause of perinatal morbidity and mortality in both 66 developing and developed countries. The mechanism by which it occurs remains to be 67 68 elucidated. Human immunodeficiency virus (HIV) and antiretroviral therapy (ART) have 69 recently been identified as independent risk factors for preterm birth [13–17]. Our recent work demonstrated unfavorable histological changes in placenta from HIV positive 70 71 women with preterm birth [18], similar to findings from other studies [19–21]. These 72 findings give credence to the belief that there is a strong association between HIV and 73 ART with adverse pregnancy outcome via diminishing placenta functional capacity. 74 Nevertheless, the exact mechanisms by which HIV and ART cause placental damage 75 and increase the risk of prematurity is far from being fully understood. We set out to

determine the surface morphological changes on placenta villi associated with HIV and
ART that could explain an increased risk of preterm labor (PTB) in this population.
Scanning electron microscopy (SEM) has been used as a standard method of exploring
features on the cell surface and therefore a suitable tool to use to characterize placental
syncytium.

81 Methods and materials

82 Study setting and population.

83 The clinical information and placental samples were collected at the Kenyatta National 84 Hospital (KNH) and Pumwani Maternity Hospital. The two are the busiest maternity hospitals in Nairobi, Kenya each with a delivery rate of over 10,000 annually. 85 86 Forty placentas were collected in the numbers given from each of the following groups: 87 (1) HIV-positive women who received ART and had a PTB, (2) HIV- negative women who delivered preterm, (3) HIV-positive women who received ART and delivered at 88 89 term, and (4) HIV- negative women who delivered at term. All the placentas were from 90 women of African descent aged 18 to 40 years with a singleton live birth. Only women with an early pregnancy obstetric scan before 16 weeks and sure of the last normal 91 92 menstrual period were included in the study. Preterm births were defined as delivery at 93 <37 weeks of gestation and term births were defined as delivery >37 weeks of gestation. We excluded women with medical and obstetric complications during 94 95 pregnancy and at the time of labor.

96

97 Ethical considerations

- 98 Approval to conduct the study and permission to transport specimens was sought from
- 99 the Kenyatta National Hospital/University of Nairobi Ethics and Research Committee
- 100 (registration number P5/01/2017).
- 101

Recruitment and informed consent

A trained research assistant cum midwife identified women who were admitted to labor and delivery meeting our inclusion criteria. He gave a full explanation of the study and obtained a written consent to enroll. Study participants were recruited from February 14, 2017, to June 22, 2017.

107

108 Sample collection and preparation

109 Placentas were collected after obtaining a written consent from the recruited mothers. 110 Six sampling sites were aseptically biopsied systematically from each placenta, trimmed 111 and labeled a-f [22], rinsed in phosphate buffer solution and then immediately fixed in 112 3% glutaraldehyde solution at pH 7.2 (at ambient temperature) and stored at 4°C for 24 113 hours at the Kenya Aids Vaccine Initiative Institute of Clinical Research (KAVI-ICR), 114 University of Nairobi, Kenya. The specimens were transported to the International Centre for Insect Physiology and Ecology, Kenya (ICIPE) and were post fixed in 1% 115 osmium tetroxide in 0.1 M cacodylate buffer for 60 minutes. The blocks were then 116 117 washed in distilled water three times, dehydrated in an ascending series of ethanol starting at 30%, 50%, 70%, 90% to absolute concentration for 20 minutes each and 118

119	stored in epoxy/propylenoxide (1:1) overnight. The blocks were then embedded in
120	epoxy resin polymerized at 60°C for 36 hours and stored. Final drying was done using
121	100% hexamethyldisilazane (HMDS, 30mins each) and left to dry for 24hrs in a fume
122	hood at the Central Analytical Facilities, Stellenbosch University, South Africa. Analysis
123	of tissue ultrastructure with conventional scanning electron microscopes is labor
124	intensive procedure and requires significant resources. To circumvent this challenge ten
125	out of forty representative placenta samples were randomly selected from each of the
126	above groups and examined.

127

128 Scanning electron microscopy imaging

129 After mounting onto standard 12mm aluminum SEM stubs, specimens were sputter-

130 coated with gold and visualized using Zeiss Merlin FESEM in-lens (secondary electron

- 131 (SE) and SE2) detector at 1.8 mm working distance, 60 µm aperture and operated at 3-
- 132 5kV keV acceleration voltage, 100 pA probe current.

133 **Results**

134

A total of 160 placentas were collected and processed, 80 preterm and 80 term. All HIV-

positive women were on treatment with ART. The average gestational ages of the

- preterm placenta was 34 (range 30-37) weeks and 39 (range 38-41) weeks for the term
- 138 placenta, respectively. The mean maternal age for mothers with preterm delivery was
- 139 26.8+4.6 years and 24.3 +4.3 for the mothers with term delivery. Thirty-six (36) and 28
- 140 patients with preterm and term deliveries had cesarean deliveries, respectively.
- 141

142 Structure of the placenta from HIV-negative women with

143 preterm and term birth

144 The chorionic villous structures had a tree-like appearance with gradual branching. 145 Branching seemed to increase gradually with advancing gestation. The terminal 146 ramifications appeared as buds and some extended to the neighboring terminal villous 147 structure via intervillous bridges (Fig 1A, B). The villous core structure was covered with 148 microvilli of variable sizes and appearance, closely apposed against each other (Fig 149 1C). There were syncytial knots which appeared as dome-shaped like elevations on the 150 surface of the syncytium and seemed to increase with advancing gestation. A few red 151 blood cells were observed on the surface, pitting the structure of the syncytium (Fig 1D). 152 In a few areas, the syncytial continuity was interrupted and the gap to a lesser extent 153 filled with fibrinoid material (Fig 1E, F). There were occurrences of tip-to-tip fusion of the 154 terminal villi in some placental sections. The preterm placenta demonstrated 155 intermediate villi transitioning into terminal villi. In some instances, the terminal villi were 156 cross-sectionally cut and revealed capillaries, many of which were dilated and occupied 157 most of the cross-sectional area of the villous core (Fig 1F).

158

159 Fig 1. Electron microscopy structure of term and preterm placentas from HIV-

160 negative women. (A, B) The chorionic structures with a tree-like appearance with 161 gradual branching. The terminal ramifications and joining via intervillous bridges (white 162 arrows). (C) The villous core structure covered with microvilli (MV) of variable sizes and 163 appearance. (D) Red blood cells (asterisks) observed on the syncytial surface, pitting 164 the structure of the syncytium. (E) The syncytial continuity (white arrow) disrupted and

filled with fibrinoid material (black arrow). (F) Cross-sectionally cut terminal villi revealedcapillaries with fetal red blood cells (FRBC).

167 Structure of the placenta from HIV-positive women with

168 preterm birth

169 The general structure of preterm placenta consisted of villous tree structure with 170 intermediate and terminal villi. Most sections of these placentas showed widespread damage with shrunken and wrinkled villi. The terminal villi showed thinning with loss of 171 172 congruently shaped terminal buds. Microvilli were either blunted or absent in significant 173 areas of both the intermediate and the terminal villi with attendant syncytiotrophoblast disruption (Fig 2A, C). Massive fibrinoid material covered the cross-sectional cuts (Fig 174 175 2B). At the bases and sometimes along the surface of the terminal villi, red blood cells 176 appeared to be adherent on the surfaces of these villi. The red blood cells were in 177 different shapes and sizes and seemed to be trapped within extensive fibrillar meshwork 178 (Fig 2D, E).

179

Fig 2. Electron microscopy structure of preterm placentas from HIV-positive women. (A) The villi appeared shrunken (SV) and wrinkled with thinning terminal villi and loss of congruently shaped terminal buds. (B) Microvilli (MV) were either blunted or absent with attendant syncytiotrophoblast disruption (white arrows). Red blood cells (RBC) were seen and massive fibrinoid material (black arrow) covered the villous structures in C, D, and E, red blood cells (asterisks) on the surface of the syncytium at advancing magnification

187 Structure of the placenta from HIV-positive women with term

188 birth

- 189 There was a progressive expansion of the villous tree with predominance of terminal villi
- in the term placenta. Focal villous syncytiotrophoblast thinning as a key feature was
- also apparent (Fig 3A). Club-shaped microvilli were present in most parts of the surface
- 192 of the syncytium (Fig 3B). The red blood cells were observed on the surface of
- 193 syncytium but not as significant as was seen in HIV-positive preterm placenta. The
- 194 syncytium is obliterated (Fig 3C, D).
- 195

Fig 3. Electron microscopy structure of term placentas from HIV-positive women.
(A) Focal thinning and loss of congruence of the terminal villi. (B) Club-shaped syncytial
knots on the villi (black arrow). (C, D) Red blood cells trapped in the fibrillar meshwork
(asterisks) on the surface of the syncytium with diminished microvilli (black arrow).

200

201 **Discussion**

The maturation spectrum of the placenta from the HIV-negative women was in keeping with what has previously been observed in normal placenta. The gradual ramifications of the chorionic villi has been described and forms the functional core of the human placenta [23]. The branches are covered by the syncytialized villous trophoblast that is in direct contact with maternal blood in the intervillous space. The presence of the syncytial knots and intervillous bridges in these cases mostly signify a normal placental ageing process [24,25]. In other cases, they may be associated with pathological

processes such as preeclampsia, obesity, malaria, oxidative stress and intrauterine growth restriction [26–29]. The intervillous bridges are involved in support role within placenta though the formation of an internal strut mechanism [30] and act to protect the villous capillaries from collapse during labor process. Dilatation of the capillary villi occurs at the tip of the terminal villi forming sinusoids that facilitate exchange of substances between the mother and the fetus via a thinned vasculosyncytial membranes [31].

216

The shrinking, wrinkling ant thinning of the placenta villi observed in placenta of HIV-217 218 positive women with preterm birth is in keeping with the decreased villi surface area and 219 perimeter observed in our previous work [18], but has also been noted in some 220 infections involving the placenta [32,33], intrauterine growth restriction [34], and 221 preeclampsia [35]. By the virtue of exposure to maternal blood, the syncytium is always 222 under constant danger of immunological or reactive oxygen species attack that could 223 result in apoptosis, degeneration and sometimes attempt to regeneration. As stated 224 above, the shriveling of the syncytium may be due to reduced trophoblast regeneration 225 capacity occasioned either independently or in combination by HIV infection or the 226 antiretrovirals. The result is that many functions of the villous syncytiotrophoblast like 227 gaseous exchange, nutrient transfer and hormone production may be impaired. 228 229 Microvilli, found on the apical aspect of some epithelial cells are surface specialization 230 that provide an increased surface area for absorption, secretion, mechanosensation,

and also provide cellular polarity enhancing certain cellular functions such as

232 membrane transportation, cytoskeletal distribution, and enzymatic function [8,36]. These 233 cells, in addition, must maintain the barrier function of the compartments being 234 separated [37]. The loss of microvilli on the surface of syncytium has been correlated 235 with diminished placental function and has been reported in eclampsia, gestational 236 diabetes, intrauterine growth restriction, and small for gestational age fetuses [38-40]. In 237 the current study, the loss of microvilli may signify a possible diminished placental 238 function which may have played a role in instigating preterm labor. Syncytiotrophoblast 239 disruption, red cell adhesion, and an extensive fibrillar meshwork that were frequently 240 found on the surface of the syncytium in these patients might be indicative of apoptotic 241 process of the syncytiotrophoblast, attempt to repair, and attendant inflammatory 242 process [41]. Fibrinoid deposition, in small amounts may be physiological as a response 243 to minimize maternal bleeding to improve pregnancy outcome. In extensive amounts, as 244 observed in this study, it may result in obliteration of the intervillous space, with atrophy 245 of the chorionic villi. The basis for this observation is related to syncytiotrophoblastic 246 damage, immunological derangement with initiation of coagulation events that results 247 into adherence of the red blood cells on the surface of the syncytium and within the 248 fibrinoid material [42–44]. This phenomenon heralds adverse pregnancy outcome and 249 has been identified as a signature of maternal anti-fetal rejection [45].

250

The remarkable similarities illustrated in this study between the placenta from term HIVpositive patients and placenta from HIV-negative patients strongly suggest that these pregnancies were minimally affected by HIV/ART. We however, observed to a lesser extent, syncytiotrophoblast thinning, microvilli missing in some parts and the red blood

cells adherence on the surface of syncytium. The difference between this group and

- 256 placenta from HIV-positive women with preterm birth may be explained by
- 257 interindividual biologic variability in response to drug metabolism, immune reaction or
- duration of HIV/ART exposure [46–48]. These results suggest HIV and ART may play a
- role in inducing syncytial morphological changes in placentas.

260 Conclusion

- 261 Our results show distinctive alterations in the placenta of HIV-positive mothers who
- gave birth prematurely, which may impair the syncytium's ability to function normally.
- 263 Microvilli blunting, syncytial disruptions, and syncytial erythrocyte adhesion might be the
- symptoms of a deeper biological process. Further work to understand the effect of
- 265 HIV/ART on the syncytiotrophoblast in relation to prematurity is recommended.

266 Acknowledgments

All our patients who graciously donated the placenta for our research. Our research and

laboratory assistants who made this work possible we say thank you.

270 **References**

- 271 1. Ellery PM, Cindrova-Davies T, Jauniaux E, Ferguson-Smith AC, Burton GJ. Evidence for
- 272 Transcriptional Activity in the Syncytiotrophoblast of the Human Placenta. Placenta.
- 273 2009.
- 274 2. Burton GJ, Jauniaux E, Charnock-Jones DS. Human Early Placental Development:
- 275 Potential Roles of the Endometrial Glands. Placenta. 2007.
- 276 3. Abumaree MH, Chamley LW, Badri M, El-Muzaini MF. Trophoblast debris modulates the
- 277 expression of immune proteins in macrophages: A key to maternal tolerance of the fetal
- allograft? Journal of Reproductive Immunology. 2012.
- 279 4. Cross JC, Werb Z, Fisher SJ. Implantation and the placenta: Key pieces of the
- 280 development puzzle. Science. 1994.
- 5. Fisher SJ. Why is placentation abnormal in preeclampsia? Vol. 213, American Journal of
- 282 Obstetrics and Gynecology. 2015. p. S115–22.
- 283 6. Salafia CM, Charles AK, Maas EM. Placenta and fetal growth restriction. Clinical
- 284 Obstetrics and Gynecology. 2006.
- 285 7. Kenny AJ, Maroux S. Topology of microvillar membrance hydrolases of kidney and
 286 intestine. Physiological Reviews. 2017.
- 287 8. Lange K. Fundamental role of microvilli in the main functions of differentiated cells:
- 288 Outline of an universal regulating and signaling system at the cell periphery. Journal of 289 Cellular Physiology. 2011.
- 290 9. Al-Zuhair AGH, Ibrahim MEA, Mughal S, Abdulla MA. Loss and regeneration of the
- 291 microvilli of human placental syncytiotrophoblast. Archives of Gynecology. 1987.

- 292 10. Obimbo M, Qureshi Z, Ogeng'o J. Infection in pregnancy; understanding impact on
- 293 placental microenvironment and preterm birth: a review. Journal of Obstetrics and
- 294 Gynecology of East and Central Africa. 2018;29(1):17–24.
- 295 11. Rudge C V., Röllin HB, Nogueira CM, Thomassen Y, Rudge MC, Odland JO. The placenta as
- a barrier for toxic and essential elements in paired maternal and cord blood samples of
- 297 South African delivering women. Journal of Environmental Monitoring. 2009.
- 298 12. Delorme-Axford E, Sadovsky Y, Coyne CB. The Placenta as a Barrier to Viral Infections.
- 299 Annual Review of Virology. 2014.
- 300 13. Ategeka J, Wasswa R, Olwoch P, Kakuru A, Natureeba P, Muehlenbachs A, et al. The
- 301 prevalence of histologic acute chorioamnionitis among HIV infected pregnant women in
- 302 Uganda and its association with adverse birth outcomes. Plos One. 2019;14(4):e0215058.
- Mandelbrot L, Sibiude J. A link between antiretrovirals and perinatal outcomes? The
 Lancet HIV. 2017.
- 305 15. Gagnon LH, MacGillivray J, Urquia ML, Caprara D, Murphy KE, Yudin MH. Antiretroviral
- 306 therapy during pregnancy and risk of preterm birth. European Journal of Obstetrics
- 307 Gynecology and Reproductive Biology. 2016.
- 308 16. Cotter AM, Garcia AG, Duthely ML, Luke B, O'Sullivan MJ. Is Antiretroviral Therapy during
- 309 Pregnancy Associated with an Increased Risk of Preterm Delivery, Low Birth Weight, or
- 310 Stillbirth? The Journal of Infectious Diseases. 2006.
- 17. Nlend AEN, Motazé AN, Tetang SM, Zeudja C, Ngantcha M, Tejiokem M. Preterm birth
- 312 and low birth weight after in Utero exposure to antiretrovirals initiated during pregnancy
- in Yaoundé, Cameroon. PLoS ONE. 2016.

- 18. Obimbo MM, Zhou Y, McMaster MT, Cohen CR, Qureshi Z, Ong'ech J, et al. Placental
- 315 Structure in Preterm Birth Among HIV-Positive Versus HIV-Negative Women in Kenya.
- 316 Journal of acquired immune deficiency syndromes (1999). 2019.
- 317 19. Castejón OC, López AJ, Pérez Ybarra LM, Castejón OC. [Placental villous lesions in HIV-1
- 318 infection treated with zidovudine]. Ginecología y obstetricia de México. 2011.
- 319 20. Shapiro RL, Souda S, Parekh N, Binda K, Kayembe M, Lockman S, et al. High prevalence of
- 320 hypertension and placental insufficiency, but no in utero HIV transmission, among
- 321 women on HAART with stillbirths in botswana. PLoS ONE. 2012.
- 322 21. Mohammadi H, Papp E, Cahill L, Rennie M, Banko N, Pinnaduwage L, et al. HIV
- 323 antiretroviral exposure in pregnancy induces detrimental placenta vascular changes that
- 324 are rescued by progesterone supplementation. Scientific Reports. 2018;8(1).
- Burton GJ, Sebire NJ, Myatt L, Tannetta D, Wang YL, Sadovsky Y, et al. Optimising sample
 collection for placental research. Placenta. 2014.
- 327 23. Kingdom J, Huppertz B, Seaward G, Kaufmann P. Development of the placental villous
- 328 tree and its consequences for fetal growth. European Journal of Obstetrics Gynecology
- 329 and Reproductive Biology. 2000.
- 330 24. Loukeris K, Sela R, Baergen RN. Syncytial Knots as a Reflection of Placental Maturity:
- 331 Reference Values for 20 to 40 Weeks' Gestational Age. Pediatric and Developmental
- 332 Pathology. 2009.
- 333 25. Fox H. THE SIGNIFICANCE OF VILLOUS SYNCYTIAL KNOTS IN THE HUMAN PLACENTA.
- BJOG: An International Journal of Obstetrics & Gynaecology. 1965.

- 26. Heazell AEP, Moll SJ, Jones CJP, Baker PN, Crocker IP. Formation of Syncytial Knots is
- 336 Increased by Hyperoxia, Hypoxia and Reactive Oxygen Species. Placenta. 2007.
- 337 27. Narasimha A, Vasudeva D. Spectrum of changes in placenta in toxemia of pregnancy.
- 338 Indian Journal of Pathology and Microbiology. 2011.
- 28. Crocker IP, Tanner OM, Myers JE, Bulmer JN, Walraven G, Baker PN. Syncytiotrophoblast
- 340 degradation and the pathophysiology of the malaria-infected placenta. Placenta. 2004;
- 29. Roberts KA, Riley SC, Reynolds RM, Barr S, Evans M, Statham A, et al. Placental structure
- and inflammation in pregnancies associated with obesity. Placenta. 2011.
- 343 30. Hormann G. No Title. Archiv fur Gynakologie. 1954;184(1):109–23.
- 344 31. Jirkovská M, Janáček J, Kaláb J, Kubínová L. Three-dimensional Arrangement of the
- 345 Capillary Bed and Its Relationship to Microrheology in the Terminal Villi of Normal Term
 346 Placenta. Placenta. 2008.
- 347 32. Chaikitgosiyakul S, Rijken MJ, Muehlenbachs A, Lee SJ, Chaisri U, Viriyavejakul P, et al. A
- 348 morphometric and histological study of placental malaria shows significant changes to
- 349 villous architecture in both Plasmodium falciparum and Plasmodium vivax infection.
- 350 Malaria Journal. 2014.
- 351 33. Walter PR, Garin Y, Blot P. Placental pathologic changes in malaria: A histologic and
- 352 ultrastructural study. Obstetrical and Gynecological Survey. 1983.
- 353 34. Almasry SM, Elfayomy AK. Morphometric analysis of terminal villi and gross
- 354 morphological changes in the placentae of term idiopathic intrauterine growth
- 355 restriction. Tissue and Cell. 2012.

356	35.	Saleh RA, Dkhil MAM. Structural changes of placenta in preeclamptic patients: Light and
357		electron microscopic study. Turkish Journal of Medical Sciences. 2008.
358	36.	Lange K. Microvillar ion channels: Cytoskeletal modulation of ion fluxes. Journal of
359		Theoretical Biology. 2000.
360	37.	Kraehenbuhl JP, Neutra MR. Molecular and cellular basis of immune protection of
361		mucosal surfaces. Physiological Reviews. 2017.
362	38.	Meng Q, Shao L, Luo X, Mu Y, Xu W, Gao C, et al. Ultrastructure of Placenta of Gravidas
363		with Gestational Diabetes Mellitus. Obstetrics and Gynecology International. 2015.
364	39.	Salgado SS, Salgado MKR. Structural changes in pre-eclamptic and eclamptic placentas -
365		an ultrastructural study. Journal of the College of Physicians and Surgeons Pakistan.
366		2011.
367	40.	Ansari TI, Fenlon S, Pasha S, O'Neill B, Gillan JE, Green CJ, et al. Morphometric
368		assessment of the oxygen diffusion conductance in placentae from pregnancies
369		complicated by intra-uterine growth restriction. Placenta. 2003.
370	41.	Roberts DJ. Placental Pathology. In: Reproductive and Developmental Toxicology. 2011.
371	42.	Fitzgerald J, Bonnke C, Brückmann A, Schleußner E, Sossdorf M, Lösche W. Pro-coagulant
372		capacity of syncytiotrophoblastic microparticles (STBMs). Zeitschrift für Geburtshilfe und
373		Neonatologie. 2011.
374	43.	Moe N, Jørgensen L. FIBRIN DEPOSITS ON THE SYNCYTIUM OF THE NORMAL HUMAN
375		PLACENTA: EVIDENCE OF THEIR THROMBOGENIC ORIGIN. Acta Pathologica
376		Microbiologica Scandinavica. 1968.

- 377 44. Guller S. Role of the syncytium in placenta-mediated complications of preeclampsia.
- 378 Thrombosis Research. 2009.
- 379 45. Romero R, Whitten A, Korzeniewski SJ, Than NG, Chaemsaithong P, Miranda J, et al.
- 380 Maternal Floor Infarction/Massive Perivillous Fibrin Deposition: A Manifestation of
- 381 Maternal Antifetal Rejection? American Journal of Reproductive Immunology. 2013.
- 382 46. Yasuda SU, Zhang L, Huang SM. The role of ethnicity in variability in response to drugs:
- 383 Focus on clinical pharmacology studies. Clinical Pharmacology and Therapeutics. 2008.
- 384 47. Niepel M, Spencer SL, Sorger PK. Non-genetic cell-to-cell variability and the
- 385 consequences for pharmacology. Current Opinion in Chemical Biology. 2009.
- 386 48. Sweeney GD. Variability in the human drug response. Thrombosis Research. 1983.

























Fig3