1 Dietary amino acids and risk of stroke subtypes: a prospective analysis of

2 **356,000** participants in seven European countries.

3

4	Tammy Y.N. Tong ¹ , Robert Clarke ² , Julie A. Schmidt ^{1,3} , Inge Huybrechts ⁴ , Urwah Noor ¹ ,
5	Nita G. Forouhi ⁵ , Fumiaki Imamura ⁵ , Ruth C. Travis ¹ , Elisabete Weiderpass ⁶ , Krasimira
6	Aleksandrova ^{7,8} , Christina C. Dahm ⁹ , Yvonne T. van der Schouw ¹⁰ , Kim Overvad ⁹ , Cecilie
7	Kyrø ¹¹ , Anne Tjønneland ¹¹ , Rudolf Kaaks ¹² , Verena Katzke ¹² , Catarina Schiborn ^{13,14} ,
8	Matthias B. Schulze ^{13,15} , Ana-Lucia Mayen-Chacon ⁴ , Giovanna Masala ¹⁶ , Sabina Sieri ¹⁷ ,
9	Maria Santucci de Magistris ¹⁸ , Rosario Tumino ¹⁹ , Carlotta Sacerdote ²⁰ , Jolanda M.A. Boer ²¹ ,
10	W.M. Monique Verschuren ^{10,21} , Magritt Brustad ^{22,23} , Therese Haugdahl Nøst ^{22,24} , Marta
11	Crous-Bou ^{25,26} , Dafina Petrova ²⁷⁻²⁹ , Pilar Amiano ²⁹⁻³¹ , José María Huerta ^{29,32} , Conchi
12	Moreno-Iribas ^{29,33} , Gunnar Engström ³⁴ , Olle Melander ^{34,35} , Kristina Johansson ³⁶ , Kristina
13	Lindvall ³⁷ , Elom K. Aglago ³⁸ , Alicia K. Heath ³⁸ , Adam S. Butterworth ³⁹⁻⁴² , John Danesh ³⁹⁻⁴³ ,
14	Timothy J. Key ¹

15

16 Affiliations:

17	1.	Cancer Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, Oxford,	
18		United Kingdom	
19	2.	Clinical Trial Service Unit and Epidemiological Studies Unit, Nuffield Department of Population Health,	
20		Oxford, United Kingdom	
21	3.	Department of Clinical Epidemiology, Department of Clinical Medicine, Aarhus University Hospital and	
22		Aarhus University, Aarhus, Denmark.	

- 23 4. Nutrition and Metabolism Branch, International Agency for Research on Cancer (IARC), World Health
- 24 Organization (WHO), Lyon, France.

- 25 5. MRC Epidemiology Unit, University of Cambridge School of Clinical Medicine, Cambridge, United
- 26 Kingdom
- 27 6. International Agency for Research on Cancer (IARC), World Health Organization (WHO), Lyon, France.
- 28 7. Department Epidemiological Methods and Etiological Research, Leibniz Institute for Prevention Research
- 29 and Epidemiology BIPS, Bremen, Germany.
- 30 8. Faculty of Human and Health Sciences, University of Bremen, Grazer Straße 2, 28359, Bremen, Germany.
- 31 9. Department of Public Health, Aarhus University, Aarhus, Denmark.
- 32 10. Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht University,
- 33 Utrecht, The Netherlands.
- 34 11. Danish Cancer Society Research Center, Copenhagen, Denmark.
- 12. Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany.
- 36 13. Department of Molecular Epidemiology, German Institute of Human Nutrition Potsdam-Rehbruecke,
- 37 Nuthetal, Germany.
- 38 14. German Center for Diabetes Research (DZD), Neuherberg, Germany
- 39 15. Institute for Nutritional Science, University of Potsdam, Nuthetal, Germany.
- 40 16. Institute for cancer research, prevention and clinical network (ISPRO), Florence, Italy.
- 41 17. Epidemiology and Prevention Unit, Fondazione IRCCS Istituto Nazionale dei Tumori di Milano, Milan,

42 Italy

- 43 18. Azienda Ospedaliera Universitaria Federico II, Napoli, Italy.
- 44 19. Hyblean Association for Epidemiological Research AIRE-ONLUS, Ragusa, Italy.
- 45 20. Unit of Cancer Epidemiology, Città della Salute e della Scienza University-Hospital, Turin, Italy.
- 46 21. National Institute for Public Health and the Environment, Bilthoven, the Netherlands.
- 47 22. Department of Community Medicine, Faculty of Health Sciences, UiT The Arctic University of Norway,
- 48 Tromsø, Norway.
- 49 23. The Public Dental Service Competence Centre of Northern Norway (TkNN), Tromsø, Norway.
- 50 24. K.G. Jebsen Centre for Genetic Epidemiology, Department of Public Health and Nursing, Norwegian
- 51 University of Science and Technology, Trondheim, Norway,
- 52 25. Unit of Nutrition and Cancer, Cancer Epidemiology Research Program, Catalan Institute of Oncology
- 53 (ICO) Bellvitge Biomedical Research Institute (IDIBELL). L'Hospitalet de Llobregat, Barcelona, Spain.
- 54 26. Department of Epidemiology, Harvard T.H. Chan School of Public Health. Boston, MA, USA.

- 55 27. Escuela Andaluza de Salud Pública (EASP), Granada, Spain
- 56 28. Instituto de Investigación Biosanitaria ibs.GRANADA, Granada, Spain
- 57 29. Centro de Investigación Biomédica en Red de Epidemiología y Salud Pública (CIBERESP), Instituto de
- 58 Salud Carlos III, C. de Melchor Fernández Almagro, Madrid, Spain.
- 59 30. Ministry of Health of the Basque Government, Sub Directorate for Public Health and Addictions of
- 60 Gipuzkoa, San Sebastian, Spain.
- 61 31. Biodonostia Health Research Institute, Epidemiology of Chronic and Communicable Diseases Group, , San
- 62 Sebastián, Spain.
- 63 32. Department of Epidemiology, Murcia Regional Health Council, IMIB-Arrixaca, Murcia, Spain.
- 64 33. Navarra Institute for Health Research (IdiSNA), Pamplona, SpainInstituto de Salud Pu blica de Navarra,
- 65 IdiSNA, Navarre Institute for Health Research, Pamplona, Spain
- 66 34. Department of Clinical Science in Malmö, Lund University, Clinical Research Center, Malmö, Sweden.
- 67 35. Department of Emergency and Internal Medicine, Skåne University Hospital, Malmö, Sweden.
- 68 36. Department of Public Health and Clinical medicine, Umeå University, Umeå, Sweden
- 69 37. Department of Epidemiology and Global Health, Umeå University, Umeå, Sweden
- 38. Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London,
- 71 United Kingdom.
- 72 39. British Heart Foundation Cardiovascular Epidemiology Unit, Department of Public Health and Primary
- 73 Care, University of Cambridge, Cambridge, United Kingdom.
- 40. National Institute for Health Research Blood and Transplant Research Unit in Donor Health and Genomics,
- 75 University of Cambridge, Cambridge, United Kingdom.
- 76 41. British Heart Foundation Centre of Research Excellence, University of Cambridge, Hills Road, Cambridge,
 77 United Kingdom.
- 42. Health Data Research UK Cambridge, Wellcome Genome Campus and University of Cambridge, United
- 79 Kingdom.
- 43. Department of Human Genetics, Wellcome Sanger Institute, Hinxton, United Kingdom.
- 81
- 82 Corresponding author: Dr Tammy Y.N. Tong (<u>tammy.tong@ndph.ox.ac.uk</u>)
- 83 Nuffield Department of Population Health, University of Oxford, Richard Doll Building, Old
- 84 Road Campus, Oxford OX3 7LF, UK

85 Abstract

108

epidemiology, prospective cohort

86	Background: Previously reported associations of protein-rich foods with stroke subtypes have
87	prompted interest in assessment of individual amino acids. We examined the associations of
88	dietary amino acids with risks of ischaemic and haemorrhagic stroke in the EPIC study.
89	Methods: We analysed data on 356,142 participants from seven European countries. Dietary
90	intakes of 19 individual amino acids were assessed using validated country-specific dietary
91	questionnaires, calibrated using additional 24-hour dietary recalls. Multivariable-adjusted
92	Cox regression models were used to estimate hazard ratios (HRs) and 95% confidence
93	intervals (CIs) of ischaemic and haemorrhagic stroke in relation to intake of each amino acid.
94	The role of blood pressure as a potential mechanism was assessed in 267,642 (75%)
95	participants.
96	Results: After a median follow-up of 12.9 years, 4,295 participants had an ischaemic stroke
97	and 1,375 participants had a haemorrhagic stroke. After correction for multiple testing, higher
98	intake of proline (as percent of total protein) was associated with 12% lower risk of
99	ischaemic stroke (HR per 1 SD higher intake 0.88; 95% CI 0.82, 0.94). The association
100	persisted after mutual adjustment for all other amino acids, systolic and diastolic blood
101	pressure. The inverse associations of isoleucine, leucine, valine, phenylalanine, threonine,
102	tryptophan, glutamic acid, serine and tyrosine with ischaemic stroke were each attenuated
103	with adjustment for proline intake. For haemorrhagic stroke, no statistically significant
104	associations were observed in the continuous analyses after correcting for multiple testing.
105	Conclusion: Higher proline intake was associated with a lower risk of ischaemic stroke,
106	independent of other dietary amino acids and blood pressure.
107	Keywords: amino acids, dietary protein, ischaemic stroke, haemorrhagic stroke, nutritional

109 Background

110 While stroke is the second leading cause of death worldwide, the incidence of stroke varies 111 substantially between countries reflecting differences in risk factors including variations in 112 dietary intakes [1]. Differences in dietary protein has been suggested to be important, but 113 previous studies of the associations of dietary protein intake with risk of stroke have reported 114 conflicting results, possibly reflecting heterogeneity by pathological stroke types and dietary 115 sources of protein [2, 3]. The different sources of dietary protein reflect a different profile of 116 individual dietary amino acids. For example, meat and meat products are a main source of the 117 essential amino acids and glycine, while grain products are a main source of dietary cysteine 118 [4]. A previous prospective study reported an inverse association of dietary cysteine with risk 119 of total stroke [5], and another reported an inverse association of glutamic acid but positive 120 association of glycine with stroke mortality [6], but overall there is very limited evidence on 121 the topic. Moreover, it is unclear whether the absolute quantity or the proportions of each 122 individual dietary amino acid (i.e. protein quality) that make up total dietary protein would be 123 more relevant for health. The two main stroke subtypes (ischaemic and haemorrhagic) also 124 have different dietary [7] and non-dietary (including genetic) risk factors [8, 9], which 125 highlights the importance of examining stroke subtypes separately. For example, previous 126 analyses in EPIC (the European Prospective Investigation into Cancer and Nutrition) reported 127 inverse associations of ischaemic stroke with consumption of fruit and vegetables, dietary 128 fibre and dairy products, while the risk of haemorrhagic stroke was positively associated with 129 egg consumption [7]. Therefore, any associations with dietary amino acids are likely to vary 130 by stroke type.

The aim of the present study was to assess the associations of both absolute and relative quantities of individual dietary amino acids with risk of stroke subtypes in a prospective study of 356,000 participants from seven European countries. Additionally, we aimed to

- assess if these associations were independent of blood pressure, the major risk factor for both
- 135 subtypes of stroke.

136

- 137 Methods
- 138 Study population
- 139 The present analyses involved 356,142 participants recruited from 20 centres between 1992
- and 2000 in seven European countries (Denmark, Germany, Italy, the Netherlands, Spain,
- 141 Sweden, and the UK), who participated in the EPIC study. Details of the study design have
- been described previously [10, 11]. The inclusion and exclusion criteria for the present study
- are shown in Supplementary figure 1. All participants provided written informed consent, and
- the study protocol was approved by the ethical review board of IARC and the institutions
- 145 where the participants were recruited [10].

146

147 Data collection

148 At recruitment all participants completed questionnaires on medical history and socio-

149 demographic factors, as well as validated country-specific dietary questionnaires (mostly

150 food frequency questionnaires or diet histories) which asked about diet in the previous year

- 151 [10]. In addition, a stratified random sample of 8% of participants across all centres also
- 152 completed a standardised and computerised 24-hour recall, on average ~1.4 years after
- recruitment, which was used to calibrate the dietary exposures to reduce between-centre
- heterogeneity and to correct for measurement error [12, 13]. The calibration method and
- rationale have been described previously [7, 12, 13].

156 Based on reported dietary intakes in both the baseline dietary questionnaires and 24-hour 157 recalls, estimates of individual dietary amino acids were derived by matching to the National 158 Nutrient Database for Standard Reference of the United States (developed at the United 159 States Department of Agriculture, or USDA) food composition tables, and appended to the 160 EPIC Nutrient Database. The matching and validation process has been reported in detail 161 [14]. USDA estimates of dietary amino acids were used because the equivalent estimates 162 were not available from all ten national Food Composition Databases that were originally 163 used in EPIC. Altogether, both calibrated and observed estimates were available for the 164 intakes of 19 individual dietary amino acids. Of the 20 standard amino acids, glutamine and 165 asparagine were not available as these two amino acids were affected by deamination 166 reactions during the acid hydrolysis process used to compile the USDA database [15, 16]. 167 Cystine (two cysteine molecules joined by a disulfide bond) [17] was estimated instead of 168 cysteine, while hydroxyproline, which is not a standard amino acid, was also estimated. The 169 sums of branched-chain amino acids (isoleucine, leucine, valine), other essential amino acids 170 (histidine, lysine, methionine, phenylalanine, threonine and tryptophan) and non-essential 171 amino acids (alanine, arginine, aspartic acid, cystine, glutamic acid, glycine, hydroxyproline, 172 proline, serine and tyrosine) were also calculated. 173 The primary outcomes were ischaemic stroke (ICD 9 433-434 or ICD 10 I63) and 174 haemorrhagic stroke (ICD 9 430-431 or ICD 10 I60-I61). In addition, we also considered 175 intracerebral haemorrhage (ICD 9 431 or ICD 10 I61) and subarachnoid haemorrhage (ICD 9 176 430 or ICD 10 I60) as separate outcomes. Both non-fatal and fatal incident events were 177 considered, and details of the ascertainment process and validation methods have been 178 previously reported [7, 11]. Details on the data collection for blood pressure (available for 179 most of the cohort) and other covariates have also been reported previously [7, 11].

180

181 Statistical analyses

182 Cohort characteristics and dietary intakes of amino acids at baseline were summarised as means (SD), median (25th, 75th percentile) or absolute numbers (%). Spearman correlation 183 184 coefficients were estimated between the individual dietary amino acids, and also for dietary 185 amino acids with dietary protein and food sources of protein. Intakes of dietary amino acids 186 were expressed in three different ways, primarily as individual amino acids modelled as a 187 percentage of total protein intake (% of total protein), and secondarily as individual amino 188 acids modelled as grams per day (g/day) and grams per 1000 kcal (g/1000 kcal). The three 189 approaches tested the hypotheses that the relative protein quality or protein makeup, the 190 absolute quantity of amino acids, and the relative quantity of amino acids to overall diet were 191 relevant to risk of stroke. The primary approach of expressing amino acids as a percentage of 192 total protein was selected since this approach was less susceptible to collinearity with other 193 dietary variables [18]. Because the sum of the individual amino acids did not add up to 100% 194 of total protein (mean=86.2%, partly because glutamine and asparagine were missing), we 195 included post hoc analyses expressing the amino acids as a percentage of the sum of all 196 available amino acids. The associations of total dietary protein with stroke risk were also 197 assessed, by expressing protein in g/day and g/1000 kcal.

Using multivariable-adjusted Cox regression, we assessed the hazard ratios (HRs) and 95% confidence intervals (CIs) for sex-specific SD differences in calibrated intakes of each dietary amino acid, and sex-specific SD differences and sex-specific fifths of observed intakes. For each association, a test of trend was estimated by fitting the calibrated or observed intake of each amino acid as a continuous variable when modelling per SD differences, or by fitting the median values of each fifth as a pseudo-continuous variable when modelling differences by fifths of intake. Tests of non-linearity were performed by using a likelihood-ratio test to

205 compare models with the exposure fitted as a continuous variable versus a categorical206 variable (fifths of intake), based on the observed intake.

207	The underlying time variable for the Cox regression was age at recruitment to age at
208	diagnosis, death or administrative censoring. All analyses were stratified by sex and EPIC
209	centre, and adjusted for age at recruitment (continuous, Model 1). Model 2 additionally
210	adjusted for total energy intake (continuous, calibrated or observed intake based on USDA
211	estimates). Model 3 additionally adjusted for smoking status and intensity (never, former,
212	current<10, 10-19, 20+ cigarettes/day, unknown), current calibrated or observed alcohol
213	consumption (non-drinker (<0.1), 0.1-4.9, 5.0-14.9, 15-29.9, 30-59.9, 60+ g/day based on
214	USDA estimates), physical activity (inactive, moderately inactive, moderately active, active,
215	unknown), employment status (employed or student, neither employed nor student, unknown)
216	and level of education completed (none or primary, secondary, vocational or university,
217	unknown). Model 4 additionally adjusted for self-reported history of diabetes (yes, no,
218	unknown), prior hypertension (yes, no, unknown) and prior hyperlipidaemia (yes, no,
219	unknown). Model 5 additionally adjusted for body mass index (<22.5, 22.5-24.9, 25.0-27.4,
220	27.5-29.9, ≥30.0 kg/m2, unknown), and was considered the main model in these analyses. As
221	total energy intake was held constant (i.e. adjusted for) in the regression model, a higher
222	intake of any individual amino acid, which contributes to total energy, would result in a
223	simultaneously lower intake of some other energy-providing foods or nutrients [19].
224	Therefore, in model 6, we removed the adjustment for total energy intake thus allowing the
225	comparison of the results with (Model 5) and without (Model 6) a theoretical substitution
226	effect of the amino acid of interest for an unspecified energy-providing source. To further
227	assess the influence of major macronutrients and to allow the interpretation of the findings
228	when holding total protein intake constant, we fitted an additional model based on Model 6
229	(without energy intake), but with the inclusion of intakes of total protein (continuous), total

230	carbohydrates (continuous), saturated fats (continuous) and unsaturated fats (continuous).
231	Likelihood-ratio χ^2 statistics were estimated at each adjustment level (for models 1-6) by
232	comparing regression models with and without inclusion of the amino acids of interest using
233	a likelihood ratio test [20]. The proportional hazards assumption was assessed using
234	Schoenfeld residuals. The assumption was not met for two covariates in the models for
235	ischaemic stroke (history of diabetes and prior hypertension) and for haemorrhagic stroke
236	(prior hypertension and BMI). However, fitting these covariates as stratifying variables
237	fulfilled the assumption and yielded close to identical estimates as adjusting, so they were
238	retained as adjustment variables.
239	To investigate whether the observed associations for the amino acids were independent from
240	the other amino acids, we included additional models mutually adjusting for each of the other
241	dietary amino acids, one at a time. We also included post-hoc analyses adjusting the amino
242	acids that remain significant across different models for its major food sources. To assess
243	whether the associations were independent of blood pressure, we estimated the adjusted mean
244	systolic and diastolic blood pressure by fifths of each dietary amino acid in the subset of
245	participants with blood pressure measurements (n=267,642), using the same covariate
246	adjustments as above. Additionally, we repeated the Cox regression analyses in this subset
247	(based on Model 5), with and without further adjustment for systolic or diastolic blood
248	pressure.
249	As secondary analyses, we repeated the main analyses for the two subtypes of haemorrhagic
250	stroke (intracerebral haemorrhage and subarachnoid haemorrhage). To assess potential
251	heterogeneity of the associations by subgroups of the major confounders and to assess
252	residual confounding, we also examined the results stratified by age at recruitment (<55, 55-

253 64, \geq 65 years), sex, BMI (<25, 25-29.9, \geq 30 kg/m²), smoking status (never, former, current

smokers), alcohol drinking (non-drinkers, moderate drinkers <15g/day, heavy drinkers

255	\geq 15g/day) and history of diseases (as a dichotomy of no disease history vs history of diabetes,		
256	hypertension or hyperlipidaemia). We also examined the associations based on observed		
257	intakes stratified by EPIC country and pooled using a meta-analysis approach. Tests of		
258	heterogeneity of trend between subgroups were conducted by comparing the risk coefficients		
259	in each subgroup using inverse variance weighting, testing for statistical significance with a		
260	χ^2 test on k-1 degrees of freedom, where k is the number of subgroups. To assess whether the		
261	overall results might be influenced by reverse causation, we repeated the analyses for		
262	ischaemic and haemorrhagic stroke after excluding the first four years of follow-up. A		
263	complete case analysis was also performed to assess the potential influence of missing		
264	covariates, which were coded as a missing category in the main analyses.		
265	All tests for statistical significance were two sided. Conventional p-values are reported, but		
265 266	All tests for statistical significance were two sided. Conventional p-values are reported, but results were interpreted with consideration of multiple testing. To account for multiple testing		
265 266 267	All tests for statistical significance were two sided. Conventional p-values are reported, but results were interpreted with consideration of multiple testing. To account for multiple testing while allowing for correlation between the exposures, we conducted a principal component		
265 266 267 268	All tests for statistical significance were two sided. Conventional p-values are reported, but results were interpreted with consideration of multiple testing. To account for multiple testing while allowing for correlation between the exposures, we conducted a principal component analysis of the exposure variables, and determined that the first four principal components		
265 266 267 268 269	All tests for statistical significance were two sided. Conventional p-values are reported, but results were interpreted with consideration of multiple testing. To account for multiple testing while allowing for correlation between the exposures, we conducted a principal component analysis of the exposure variables, and determined that the first four principal components explained 99% of the total variation in the exposure data [21, 22]. Consequently, the effective		
265 266 267 268 269 270	All tests for statistical significance were two sided. Conventional p-values are reported, but results were interpreted with consideration of multiple testing. To account for multiple testing while allowing for correlation between the exposures, we conducted a principal component analysis of the exposure variables, and determined that the first four principal components explained 99% of the total variation in the exposure data [21, 22]. Consequently, the effective number of independent tests was determined to be four, and the statistical significance level		
265 266 267 268 269 270 271	All tests for statistical significance were two sided. Conventional p-values are reported, but results were interpreted with consideration of multiple testing. To account for multiple testing while allowing for correlation between the exposures, we conducted a principal component analysis of the exposure variables, and determined that the first four principal components explained 99% of the total variation in the exposure data [21, 22]. Consequently, the effective number of independent tests was determined to be four, and the statistical significance level after Bonferroni correction for multiple testing based on this number was defined as		
265 266 267 268 269 270 271 271	All tests for statistical significance were two sided. Conventional p-values are reported, but results were interpreted with consideration of multiple testing. To account for multiple testing while allowing for correlation between the exposures, we conducted a principal component analysis of the exposure variables, and determined that the first four principal components explained 99% of the total variation in the exposure data [21, 22]. Consequently, the effective number of independent tests was determined to be four, and the statistical significance level after Bonferroni correction for multiple testing based on this number was defined as 0.05/4=0.0125. All analyses were performed using Stata version 17.0 (StataCorp, TX, USA).		
265 267 268 269 270 271 272 273	All tests for statistical significance were two sided. Conventional p-values are reported, but results were interpreted with consideration of multiple testing. To account for multiple testing while allowing for correlation between the exposures, we conducted a principal component analysis of the exposure variables, and determined that the first four principal components explained 99% of the total variation in the exposure data [21, 22]. Consequently, the effective number of independent tests was determined to be four, and the statistical significance level after Bonferroni correction for multiple testing based on this number was defined as 0.05/4=0.0125. All analyses were performed using Stata version 17.0 (StataCorp, TX, USA). All figures were generated using "Jasper makes plots" package [23] version 2-266 in R		

275

276 **Results**

277 Baseline and dietary characteristics

278	The baseline characteristics of the study participants are shown in Table 1 and additionally in		
279	detail in Supplementary table 1. Intakes of dietary amino acids, total protein and total energy		
280	are shown by sex in Supplementary table 2, and by EPIC country in Supplementary table 3.		
281	The Spearman correlation coefficients between the individual dietary amino acids are shown		
282	in Supplementary figure 2, and for amino acids with dietary protein and food sources of		
283	protein in Supplementary figures 3 and 4 respectively.		
284			
285	Associations of dietary amino acids with stroke risk		
286	After a median follow-up of 12.9 years, there were 4,295 cases of ischaemic stroke and 1,375		

cases of haemorrhagic stroke among 356,142 participants (4,529,626 person-years in total).

In the fully adjusted models, associations between dietary amino acids and risks of ischaemic

and haemorrhagic stroke were numerically similar irrespective of whether the amino acids

were expressed as a percentage of total protein (Figure 1, Supplementary tables 4-5), in g/day

291 (Supplementary figure 5, Supplementary tables 4-5), g/1000 kcal (Supplementary figure 6,

Supplementary tables 4-5), or as a percentage of total amino acids (Supplementary figure 7,

Supplementary tables 4-5). Unless otherwise stated, the primary results reported on the

associations that remained statistically significant after correcting for multiple testing, or

results from categorical analyses where we observed non-linear associations.

296 When expressing the amino acids as a percentage of total protein, in the fully adjusted models

297 (Model 5) we found inverse associations of ischaemic stroke for each 1 SD higher calibrated

intake of total and individual branched-chain amino acids (HR [95% CI] per SD higher of

total: 0.93 [0.88, 0.98], isoleucine: 0.93 [0.88, 0.98], leucine: 0.93 [0.87, 0.98], valine: 0.92

300 [0.88, 0.97]), phenylalanine (0.92 [0.87, 0.98]), threonine (0.93 [0.88, 0.99]), tryptophan

301 (0.94 [0.89, 0.99]), total non-essential amino acids (0.92 [0.87, 0.98]), glutamic acid (0.90

302	[0.84, 0.96]), proline (0.88 [0.82, 0.94]), serine (0.92 [0.88, 0.97]) and tyrosine (0.92 [0.88,
303	0.98]) (Figure 1). Glycine showed a non-linear association with ischaemic stroke in the
304	categorical analysis (HR [95% CI] 1.28 [1.14, 1.43] for top versus bottom fifth, p non-
305	linearity<0.001), but only when expressing glycine as a percentage of estimated intakes of
306	total protein or total amino acids (Supplementary table 4). For haemorrhagic stroke, no
307	statistically significant associations were observed in the continuous analyses after correcting
308	for multiple testing (Figure 1). In categorical analyses, lower risks were observed from the
309	second to top fifth of proline intake compared to bottom fifth (top versus bottom fifth: 0.72
310	[0.58, 0.90], p-trend across categories=0.006, Supplementary table 5). No statistically
311	significant associations were observed for total protein with either ischaemic or haemorrhagic
312	stroke (Supplementary figures 4-5, Supplementary tables 4-5), or in any analyses examining
313	the two haemorrhagic stroke subtypes as outcomes (Supplementary table 6). Results from the
314	different covariate adjustment models are shown in Supplementary tables 7-9.

315

316 Mutual adjustment of dietary amino acids

317 Results with mutual adjustment for each of the other amino acids for the ten amino acids that 318 were inversely associated with ischaemic stroke risk, as reported above, are shown in Figure 319 2 (for the five amino acids that were significantly associated in both the calibrated and 320 observed analyses) and in Supplementary figure 8 (for the five amino acids that were 321 significantly associated in the calibrated analyses only). The inverse associations for proline 322 persisted after adjustment for all other amino acids, but the associations for other amino acids 323 were attenuated upon adjustment for proline. However, the higher risk of ischaemic stroke 324 comparing top to bottom fifth of glycine intake remained with adjustment for proline (1.27 325 [1.12, 1.43], results not shown in tables). In post-hoc analyses, the inverse associations for

proline with ischaemic stroke remained upon mutual adjustment for its different food sources (e.g. 0.89; 0.83, 0.96 after mutual adjustment for cheese, with no significant association for cheese in the same model, results not shown in tables). Based on these findings, the changes in the HRs and likelihood ratio χ^2 statistics for proline are also depicted graphically in Figure 3, by each level of covariate adjustment.

331

332 Assessment of role of blood pressure

- 333 The adjusted mean levels of systolic and diastolic blood pressure across fifths of dietary
- amino acids are shown in Supplementary tables 10-11. Across all amino acids, higher intakes

were associated with higher blood pressure, though the differences were very small (up to 1

336 mmHg difference in most cases). As expected from these small differences, the results for

both ischaemic and haemorrhagic stroke were almost unchanged after further adjustment for

either systolic or diastolic blood pressure (Supplementary table 12).

339

340 *Sensitivity analyses*

- 341 In stratified analyses, the main findings were consistent between subgroups of age, sex, BMI,
- 342 smoking status, alcohol drinking and disease status for both ischaemic and haemorrhagic
- 343 stroke (Supplementary tables 13-24). The pooled estimates by country using a meta-analysis
- approach were similar (Supplementary tables 25-26), and results were consistent after
- excluding the first four years of follow-up (Supplementary table 27) or based on a complete
- case analysis (Supplementary table 28).

347

348 Discussion

349	In this large prospective study, we observed inverse associations of dietary intakes of proline
350	with risk of ischaemic stroke after adjustment of potential confounders, independent of the
351	other amino acids and of total protein intake. The associations were consistent regardless of
352	how the amino acids were modelled (i.e. percentage of protein, g/day and g/1000 kcal), and
353	across all subgroup analyses. Although inverse associations with ischaemic stroke were
354	initially observed for some other amino acids, these might be explained by the high
355	correlations between these amino acids and proline, since the associations were attenuated
356	after adjustment for proline. We further observed evidence of a higher risk of ischaemic
357	stroke in individuals with the highest intakes of dietary glycine, and a lower risk of
358	haemorrhagic stroke in people with higher intakes of dietary proline, though these
359	associations differed depending on the way dietary amino acids were modelled in the
360	analyses.
361	The present analyses provide important novel evidence on the associations between dietary
361 362	The present analyses provide important novel evidence on the associations between dietary amino acids and stroke risk, as only two prospective studies have previously published on the
361 362 363	The present analyses provide important novel evidence on the associations between dietary amino acids and stroke risk, as only two prospective studies have previously published on the topic [5, 6]. The Swedish Mammography Cohort, with 1751 cases of total stroke (1311
361 362 363 364	The present analyses provide important novel evidence on the associations between dietary amino acids and stroke risk, as only two prospective studies have previously published on the topic [5, 6]. The Swedish Mammography Cohort, with 1751 cases of total stroke (1311 ischaemic and 264 haemorrhagic after 10 years), reported an inverse association between
361 362 363 364 365	The present analyses provide important novel evidence on the associations between dietary amino acids and stroke risk, as only two prospective studies have previously published on the topic [5, 6]. The Swedish Mammography Cohort, with 1751 cases of total stroke (1311 ischaemic and 264 haemorrhagic after 10 years), reported an inverse association between dietary cysteine and the risk of total stroke, but no other associations including for proline
361 362 363 364 365 366	The present analyses provide important novel evidence on the associations between dietary amino acids and stroke risk, as only two prospective studies have previously published on the topic [5, 6]. The Swedish Mammography Cohort, with 1751 cases of total stroke (1311 ischaemic and 264 haemorrhagic after 10 years), reported an inverse association between dietary cysteine and the risk of total stroke, but no other associations including for proline were observed after mutual adjustment of the amino acids [5]. In contrast to that study, we
361 362 363 364 365 366 367	The present analyses provide important novel evidence on the associations between dietary amino acids and stroke risk, as only two prospective studies have previously published on the topic [5, 6]. The Swedish Mammography Cohort, with 1751 cases of total stroke (1311 ischaemic and 264 haemorrhagic after 10 years), reported an inverse association between dietary cysteine and the risk of total stroke, but no other associations including for proline were observed after mutual adjustment of the amino acids [5]. In contrast to that study, we did not find an association of cystine with either stroke subtype; however, although cystine is
361 362 363 364 365 366 367 368	The present analyses provide important novel evidence on the associations between dietary amino acids and stroke risk, as only two prospective studies have previously published on the topic [5, 6]. The Swedish Mammography Cohort, with 1751 cases of total stroke (1311 ischaemic and 264 haemorrhagic after 10 years), reported an inverse association between dietary cysteine and the risk of total stroke, but no other associations including for proline were observed after mutual adjustment of the amino acids [5]. In contrast to that study, we did not find an association of cystine with either stroke subtype; however, although cystine is expected to be highly correlated with cysteine, their molecular structures are different, so the
361 362 363 364 365 366 367 368 369	The present analyses provide important novel evidence on the associations between dietary amino acids and stroke risk, as only two prospective studies have previously published on the topic [5, 6]. The Swedish Mammography Cohort, with 1751 cases of total stroke (1311 ischaemic and 264 haemorrhagic after 10 years), reported an inverse association between dietary cysteine and the risk of total stroke, but no other associations including for proline were observed after mutual adjustment of the amino acids [5]. In contrast to that study, we did not find an association of cystine with either stroke subtype; however, although cystine is expected to be highly correlated with cysteine, their molecular structures are different, so the two sets of results are not directly comparable. The Takayama Study in Japan, with 677
361 362 363 364 365 366 367 368 369 370	The present analyses provide important novel evidence on the associations between dietary amino acids and stroke risk, as only two prospective studies have previously published on the topic [5, 6]. The Swedish Mammography Cohort, with 1751 cases of total stroke (1311 ischaemic and 264 haemorrhagic after 10 years), reported an inverse association between dietary cysteine and the risk of total stroke, but no other associations including for proline were observed after mutual adjustment of the amino acids [5]. In contrast to that study, we did not find an association of cystine with either stroke subtype; however, although cystine is expected to be highly correlated with cysteine, their molecular structures are different, so the two sets of results are not directly comparable. The Takayama Study in Japan, with 677 deaths from stroke (393 ischaemic and 153 intracerebral haemorrhage after 16 years), showed
361 362 363 364 365 366 367 368 369 370 371	The present analyses provide important novel evidence on the associations between dietary amino acids and stroke risk, as only two prospective studies have previously published on the topic [5, 6]. The Swedish Mammography Cohort, with 1751 cases of total stroke (1311 ischaemic and 264 haemorrhagic after 10 years), reported an inverse association between dietary cysteine and the risk of total stroke, but no other associations including for proline were observed after mutual adjustment of the amino acids [5]. In contrast to that study, we did not find an association of cystine with either stroke subtype; however, although cystine is expected to be highly correlated with cysteine, their molecular structures are different, so the two sets of results are not directly comparable. The Takayama Study in Japan, with 677 deaths from stroke (393 ischaemic and 153 intracerebral haemorrhage after 16 years), showed that glutamic acid intake was inversely associated with total stroke mortality among women,

373 no prior history of hypertension; the study did not report on proline [6]. In our study, we also

374 observed an inverse association of glutamic acid intake with ischaemic stroke, but this 375 association was attenuated when we further adjusted for dietary proline. In accordance with 376 the Takayama Study, we also found a higher risk of ischaemic stroke for people in the highest 377 fifth of glycine intake. The differences in intakes of the amino acids across populations 378 should be considered when comparing the findings. For example, the average intakes of both 379 glutamic acid and glycine when expressed as a percentage of total protein in the EPIC study 380 is closer to the lower end of the intakes reported in the Takayama study. Overall, compared to 381 the two previous studies, the current study is over three- and ten-fold larger in case numbers. 382 We also focused on examining ischaemic and haemorrhagic stroke separately, based on prior 383 evidence of heterogeneity in risk factors by stroke subtypes. Overall, the associations of 384 dietary amino acids with haemorrhagic stroke were less clear. As the number of 385 haemorrhagic stroke cases in the study was comparatively low, further investigations in other 386 populations with higher numbers of haemorrhagic stroke are warranted. 387 Previous studies have largely attributed the observed associations of dietary amino acids with 388 stroke risk to the influences of the amino acids on blood pressure. The INTERMAP study 389 reported inverse associations of glutamic acid and to a lesser degree proline [18], but positive 390 associations of glycine intake [24], with blood pressure. Our results did not suggest an 391 important role of blood pressure in explaining any of the findings, given that we did not 392 observe any meaningful differences in blood pressure by amino acid intake, nor did we 393 observe any differences in the HRs after further adjusting for blood pressure. However, blood 394 pressure and diet in EPIC were both measured at recruitment, which does not allow the 395 evaluation of a temporal association. Additionally, the use of blood pressure measurements 396 assessed at a single time point as an indicator of long-term blood pressure is also prone to 397 random measurement error, which could result in an underestimation of its role in explaining 398 any associations between amino acid intakes and stroke risk. Furthermore, blood pressure

399 measurements in EPIC were not taken using a standardised protocol across centres [25], but 400 our analyses were stratified by centre, so any influence on the results should be small. 401 The possible biological mechanisms for the association between proline and stroke risk are 402 not yet well defined. Despite being non-essential amino acids, proline, glycine and 403 hydroxyproline are the major constituents of collagen, which accounts for about one third of 404 proteins in the human body [26]. Collagen has an important role in maintaining the structure 405 and strength of connective tissues, cartilage and blood vessels [26, 27], and studies from 406 animal models have suggested that endogenous synthesis of these amino acids may not be 407 adequate for growth or collagen production [26, 28]. Because proline is the only precursor for 408 hydroxyproline, which in turn is a precursor for glycine [26, 29], dietary intakes of proline 409 may be of particular importance for collagen synthesis. 410 In contrast to other amino acids, proline contains an imino group in its structure instead of a 411 primary amino group, which results in its distinctive cyclic structure and exclusion from 412 standard amino acid metabolism [30, 31]. For this reason, the family of proline-containing

413 peptides, collectively called glycoprolines, cannot be degraded by most peptidases owing to

the presence of proline in their structures, and thus are more likely to enter the bloodstream

and exert various biological effects, including possible atheroprotective or neuroprotective

416 roles, regulation of insulin-like growth factor-I (IGF-I) homeostasis and lipid metabolism

417 including lowering of total cholesterol [32]. Glycoprolines may be derived from collagen or

418 IGF-Ia study in rats has shown that oral administration of cyclic glycine-proline, a

419 glycoproline derived from IGF-I, promoted neural plasticity and remodelling in animals with

420 focal ischaemic lesions, suggesting a neuroprotective effect of cyclic glycine-proline against

421 ischaemic stroke or its recovery [33]. In human studies, both cyclic glycine-proline

422 concentrations [34] and proline concentrations [35–37] have been proposed as prognostic

423 markers of stroke recovery or stroke events. However, the directions of the changes observed

424 have been inconsistent. Other studies have also reported low correlations between dietary 425 intake and circulating levels of proline [38]. Therefore, further research into the link between 426 proline intake and the physiological effects of circulating proline, as well as the role of 427 proline in both stroke incidence and prognosis (recovery or mortality) is needed. Among dietary sources, the results of the present study showed that dairy products, especially 428 429 cheese, was most highly correlated with proline intake. Therefore, the findings of lower risks 430 of ischaemic stroke in people with higher proline intake in the present study are consistent 431 with previous observations of lower ischaemic stroke risk in people with higher intake of 432 cheese and yogurt in the same cohort [7]. Despite being a major source of saturated fat, 433 higher intakes of dairy products have been inversely associated with cardiovascular disease 434 outcomes in a meta-analysis of prospective studies [39], suggesting that some components in 435 cheese or dairy products might be protective. For example, studies had suggested that the 436 specific saturated fatty acid isomer composition of different foods [40], or the calcium [41] or 437 probiotics content [42] of dairy foods may account for their associations with stroke risk. As 438 the observed association for proline remained upon adjustment for cheese and other food 439 sources, the findings of the present study suggest that the amino acid profiles of dairy 440 products and their associations with CVD warrant further research. Meanwhile, dietary intake 441 of glycine was strongly correlated with red meat intake, which was previously found to be 442 modestly but positively associated with ischaemic stroke risk in EPIC, though this association 443 was attenuated after adjustment for fibre intake [7]. Further population-based and mechanistic 444 studies are needed to replicate the observed associations between dietary amino acids and 445 risks of stroke subtypes, prior to investigating any implications for potential dietary 446 interventions.

The chief strengths of this study include the large sample size with participants recruited fromseven European countries, the prospective design and prolonged duration of follow-up. EPIC

449 is also a well-characterised cohort with extensive dietary and lifestyle/health behavioural 450 data, allowing us to examine estimated intakes of 19 dietary amino acids with stroke risk, 451 while adjusting for multiple confounders. The ascertainment of stroke cases in this cohort 452 was either completely or partially validated depending on EPIC country, and previous dietary 453 analyses have found no heterogeneity in the associations by the extent of stroke validation 454 [7]. To adjust for potential bias due to differences between the country-specific dietary 455 questionnaires, we also applied calibration to the dietary data using additional standardised 456 24-hour recalls to reduce between-centre heterogeneity and to reduce measurement error. The 457 study also had several limitations. Diet was ascertained using a single dietary questionnaire 458 collected at baseline, which meant that we were not able to account for dietary changes 459 during follow-up. We were unable to assess the associations of dietary glutamine, asparagine 460 or cysteine with stroke risk, as estimates of these amino acids were not available from the 461 USDA database. The matching of amino acids data based on a single US database in different 462 countries across Europe may also be a limitation, and the absolute validity of the amino acid 463 estimates assessed using FFQs and 24-hour recalls is not known, although good agreement 464 and thus relative validity has been shown for estimates of total energy and protein intakes 465 between the USDA and country-specific databases in the previous validation study [14]. 466 Because of the large number of tests conducted, false positive findings due to chance are 467 possible, but we focused on the findings that survived correction for multiple testing. We also 468 conducted analyses further adjusting for each one of the other amino acids, which given their 469 high correlations might constitute over-adjustment, but this approach also helped highlight 470 the clear associations with proline. As with any observational study, the present study could 471 not establish causality, and cannot fully exclude residual confounding or reverse causation. However, we assessed the former by comparing the changes in χ^2 statistics across models, 472 473 and limited the latter possibility by our sensitivity analysis excluding the first four years of

follow-up. Information on other potential mediators such as cholesterol concentrations was
only available in a small subset of participants at study baseline, so we were unable to access
their potential effects. Lastly, the present study was largely restricted to white European
individuals, therefore the findings may not be generalisable to other populations which have
different sources of protein intake and dietary amino acid profiles, as well as genetic
differences.

481 **Conclusions**

482 This large study demonstrated an inverse association of both absolute intake and relative

483 proportions of dietary proline with risk of ischaemic stroke, independent of the other amino

484 acids. The suggestive associations of lower risk of haemorrhagic stroke in people with higher

485 proline intake, and higher risk of ischaemic stroke in people with higher proportions of

486 glycine intake, also warrant further investigation. These observations were largely

unexplained by differences in blood pressure. Further research is needed from both large-

488 scale population-based studies and mechanistic studies to replicate these findings, in addition

to genetic studies and intervention trials to assess the causal relevance of these associations

490 and possible implications for prevention.

491 *Acknowledgments*: The authors thank all participants in the EPIC cohort for their invaluable

492 contribution to the study.

Authors' contributions: The study was conceived and designed by TYNT and TJK with input
from all other authors. TYNT analysed the data and wrote the first draft of the manuscript,
with input from TJK, RC, JAS, IH, UN, NGF, FI, RCT, EW, KA, CCD and YTVS. All other
authors provided the data and revised the manuscript critically for important intellectual
content. TYNT had primary responsibility for final content. All authors read and approved the
final manuscript. *Disclaimer*: Where authors are identified as personnel of the International Agency for

500 Research on Cancer / World Health Organization, the authors alone are responsible for the

views expressed in this article and they do not necessarily represent the decisions, policy or

views of the International Agency for Research on Cancer / World Health Organization.

503 *Conflicts of interest*: The authors have no conflicts of interest to declare.

504 Funding: This work was supported by a Nuffield Department of Population Health 505 Fellowship (TYNT), the UK Medical Research Council (MR/M012190/1), Cancer Research 506 UK (C8221/A19170 and 570/A16491), and the Wellcome Trust (Our Planet Our Health, 507 Livestock Environment and People 205212/Z/16/Z). EPIC-CVD has been supported by the 508 European Union Framework 7 (HEALTH-F2-2012-279233), the European Research Council (268834), the UK Medical Research Council (G0800270 and MR/L003120/1), the British 509 510 Heart Foundation (SP/09/002 and RG/08/014 and RG13/13/30194), and the UK National 511 Institute of Health Research. The establishment of the study subcohort was supported by the 512 EU Sixth Framework Programme (FP6) (grant LSHM CT 2006 037197 to the InterAct 513 project) and the Medical Research Council Epidemiology Unit (grants MC_UU_12015/1 and 514 MC UU 12015/5). The coordination of the European Prospective Investigation into Cancer

515 and Nutrition (EPIC) is financially supported by International Agency for Research on 516 Cancer (IARC) and also by the Department of Epidemiology and Biostatistics, School of 517 Public Health, Imperial College London which has additional infrastructure support provided 518 by the NIHR Imperial Biomedical Research Centre (BRC). The national cohorts are 519 supported by: Danish Cancer Society (Denmark); Ligue Contre le Cancer, Institut Gustave 520 Roussy, Mutuelle Générale de l'Education Nationale, Institut National de la Santé et de la 521 Recherche Médicale (INSERM) (France); German Cancer Aid, German Cancer Research 522 Center (DKFZ), German Institute of Human Nutrition Potsdam-Rehbruecke (DIfE), Federal 523 Ministry of Education and Research (BMBF) (Germany); Associazione Italiana per la 524 Ricerca sul Cancro-AIRC-Italy, Compagnia di SanPaolo and National Research Council 525 (Italy); Dutch Ministry of Public Health, Welfare and Sports (VWS), LK Research Funds, 526 Dutch Prevention Funds, Dutch ZON (Zorg Onderzoek Nederland), World Cancer Research 527 Fund (WCRF) (The Netherlands), and Statistics Netherlands is acknowledged for providing 528 the causes of death; Health Research Fund (FIS) - Instituto de Salud Carlos III (ISCIII), 529 Regional Governments of Andalucía, Asturias, Basque Country, Murcia and Navarra, and the 530 Catalan Institute of Oncology - ICO (Spain); Swedish Cancer Society, Swedish Research 531 Council and County Councils of Skåne and Västerbotten and the funds supporting the 532 Northern Sweden Diet Database (Sweden); Cancer Research UK (14136 to EPIC-Norfolk; 533 C8221/A29017 to EPIC-Oxford), Medical Research Council (1000143 to EPIC-Norfolk; 534 MR/M012190/1 to EPIC-Oxford) (United Kingdom). The funders had no role in the design 535 of the study; in the collection, analyses, or interpretation of data; in the writing of the 536 manuscript; or in the decision to publish the results.

537 *Data sharing*: For information on how to submit an application for gaining access to EPIC
538 data and/or biospecimens, please follow the instructions at http://epic.iarc.fr/access/index.php

539 **References**

- Feigin VL, Stark BA, Johnson CO, et al (2021) Global, regional, and national burden of stroke and its risk factors, 1990-2019: A systematic analysis for the Global Burden of Disease Study 2019. Lancet Neurol 20:1–26. https://doi.org/10.1016/S1474-4422(21)00252-0
- Zhang Z, Xu G, Yang F, et al (2014) Quantitative analysis of dietary protein intake and stroke risk. Neurology 83:19–25. https://doi.org/10.1212/WNL.00000000000551
- Stang XW, Yang Z, Li M, et al (2016) Association between dietary protein intake and risk of stroke: A meta-analysis of prospective studies. Int J Cardiol 223:548–551.
 https://doi.org/10.1016/j.ijcard.2016.08.106
- Górska-Warsewicz H, Laskowski W, Kulykovets O, et al (2018) Food Products as
 Sources of Protein and Amino Acids-The Case of Poland. Nutrients 10:.
 https://doi.org/10.3390/nu10121977
- 5. Larsson SC, Håkansson N, Wolk A (2015) Dietary Cysteine and Other Amino Acids
 and Stroke Incidence in Women. Stroke 46:922–6.
 https://doi.org/10.1161/STROKEAHA.114.008022
- Nagata C, Wada K, Tamura T, et al (2015) Dietary intakes of glutamic acid and
 glycine are associated with stroke mortality in Japanese adults. J Nutr 145:720–728.
 https://doi.org/10.3945/jn.114.201293
- Tong TYN, Appleby PN, Key TJ, et al (2020) The associations of major foods and
 fibre with risks of ischaemic and haemorrhagic stroke: a prospective study of 418 329
 participants in the EPIC cohort across nine European countries. Eur Heart J 44:1–11.
 https://doi.org/10.1093/eurheartj/ehaa007
- Sun L, Clarke R, Bennett D, et al (2019) Causal associations of blood lipids with risk
 of ischemic stroke and intracerebral hemorrhage in Chinese adults. Nat Med 25:569–
 574. https://doi.org/10.1038/s41591-019-0366-x
- 9. Price AJ, Wright FL, Green J, et al (2018) Differences in risk factors for 3 types of
 stroke: UK prospective study and meta-analyses. Neurology 90:e298–e306.
 https://doi.org/10.1212/WNL.00000000004856
- Riboli E, Hunt K, Slimani N, et al (2002) European Prospective Investigation into
 Cancer and Nutrition (EPIC): study populations and data collection. Public Health
 Nutr 5:1113. https://doi.org/10.1079/PHN2002394
- 571 11. Danesh J, Saracci R, Berglund G, et al (2007) EPIC-Heart: The cardiovascular
 572 component of a prospective study of nutritional, lifestyle and biological factors in
 573 520,000 middle-aged participants from 10 European countries. Eur J Epidemiol
 574 22:129–141. https://doi.org/10.1007/s10654-006-9096-8
- Slimani N, Ferrari P, Ocké M, et al (2000) Standardization of the 24-hour diet recall
 calibration method used in the European Prospective Investigation into Cancer and
 Nutrition (EPIC): general concepts and preliminary results. Eur J Clin Nutr 54:900–17
- Ferrari P, Day NE, Boshuizen HC, et al (2008) The evaluation of the diet/disease
 relation in the EPIC study: considerations for the calibration and the disease models.
 Int J Epidemiol 37:368–378. https://doi.org/10.1093/ije/dym242

581 582 583 584	14.	uacel I, Perez-Cornago A, Schmidt JA, et al (2022) Evaluation of protein and amino id intake estimates from the EPIC dietary questionnaires and 24-h dietary recalls ing different food composition databases. Nutr Metab Cardiovasc Dis 32:80–89. tps://doi.org/10.1016/j.numecd.2021.09.012	
585 586 587	15.	Lenders CM, Liu S, Wilmore DW, et al (2009) Evaluation of a novel food composition database that includes glutamine and other amino acids derived from gene sequencing data. Eur J Clin Nutr 63:1433–1439. https://doi.org/10.1038/ejcn.2009.110	
588 589	16.	Mustă 🗆 ea G, Ungureanu EL, Iorga E (2019) Protein acidic hydrolysis for amino acids analysis in food - progress over time: A short review. J Hyg Eng Des 26:81–87	
590 591 592	17.	National Center for Biotechnology Information (2022) PubChem Compound Summary for CID 595, Cystine. https://pubchem.ncbi.nlm.nih.gov/compound/Cystine. Accessed 11 Feb 2022	
593 594 595 596	18.	Stamler J, Brown IJ, Daviglus ML, et al (2009) Glutamic acid, the main dietary amino acid, and blood pressure: The intermap study (international collaborative study of macronutrients, micronutrients and blood pressure). Circulation 120:221–228. https://doi.org/10.1161/CIRCULATIONAHA.108.839241	
597 598 599	19.	Ibsen DB, Laursen ASD, Würtz AML, et al (2021) Food substitution models for nutritional epidemiology. Am J Clin Nutr 113:294–303. https://doi.org/10.1093/ajcn/nqaa315	
600 601 602 603	20. Floud S, Balkwill A, Moser K, et al (2016) The role of health-related behavioural factors in accounting for inequalities in coronary heart disease risk by education and area deprivation: prospective study of 1.2 million UK women. BMC Med 14:145. https://doi.org/10.1186/s12916-016-0687-2		
604 605 606	21.	Gao X, Starmer J, Martin ER (2008) A multiple testing correction method for genetic association studies using correlated single nucleotide polymorphisms. Genet Epidemiol 32:361–369. https://doi.org/10.1002/gepi.20310	
607 608 609	22.	dams CD, Richmond R, Santos Ferreira DL, et al (2019) Circulating metabolic iomarkers of screen-detected prostate cancer in the ProtecT study. Cancer Epidemiol Giomarkers Prev 28:208–216. https://doi.org/10.1158/1055-9965.EPI-18-0079	
610 611	23.	Arnold M (2020) Jasper: Jasper makes plots. 2020. https://github.com/arnhew99/Jasper	
612 613 614	24.	Stamler J, Brown IJ, Daviglus ML, et al (2013) Dietary glycine and blood pressure: The international study on macro/micronutrients and blood pressure. Am J Clin Nutr 98:136–145. https://doi.org/10.3945/ajcn.112.043000	
615 616 617 618	25.	Schulze MB, Kroke A, Saracci R, Boeing H (2002) The effect of differences in measurement procedure on the comparability of blood pressure estimates in multi-centre studies. Blood Press Monit 7:95–104. https://doi.org/10.1097/00126097-200204000-00002	
619 620 621	26.	Li P, Wu G (2018) Roles of dietary glycine, proline, and hydroxyproline in collagen synthesis and animal growth. Amino Acids 50:29–38. https://doi.org/10.1007/s00726-017-2490-6	
622 623	27.	Xu J, Shi G-P (2014) Vascular wall extracellular matrix proteins and vascular diseases. Biochim Biophys Acta 1842:2106–2119. https://doi.org/10.1016/j.bbadis.2014.07.008	

624 625 626	28.	Hou Y, Yao K, Yin Y, Wu G (2016) Endogenous synthesis of amino acids limits growth, lactation, and reproduction in animals. Adv Nutr 7:331–342. https://doi.org/10.3945/an.115.010850	
627 628 629	29.	Wu G, Bazer FW, Burghardt RC, et al (2011) Proline and hydroxyproline metabolism Implications for animal and human nutrition. Amino Acids 40:1053–1063. https://doi.org/10.1007/s00726-010-0715-z	
630 631 632	30.	Phang JM, Donald SP, Pandhare J, Liu Y (2008) The metabolism of proline, a stress substrate, modulates carcinogenic pathways. Amino Acids 35:681–690. https://doi.org/10.1007/s00726-008-0063-4	
633 634	31.	Wyse ATS, Netto CA (2011) Behavioral and neurochemical effects of proline. Metab Brain Dis 26:159–172. https://doi.org/10.1007/s11011-011-9246-x	
635 636	32.	Misiura M, Miltyk W (2019) Proline-containing peptides-New insight and implications: A Review. Biofactors 45:857–866. https://doi.org/10.1002/biof.1554	
637 638 639	33.	Kaneko H, Namihira M, Yamamoto S, et al (2022) Oral administration of cyclic glycyl-proline facilitates task learning in a rat stroke model. Behav Brain Res 417:113561. https://doi.org/10.1016/j.bbr.2021.113561	
640 641 642	34.	Fan D, Krishnamurthi R, Harris P, et al (2019) Plasma cyclic glycine proline/IGF-1 ratio predicts clinical outcome and recovery in stroke patients. Ann Clin Transl Neurol 6:669–677. https://doi.org/10.1002/acn3.743	
643 644 645	35.	Goulart VAM, Sena MM, Mendes TO, et al (2019) Amino acid biosignature in plasma among ischemic stroke subtypes. Biomed Res Int 2019:. https://doi.org/10.1155/2019/8480468	
646 647 648	36.	Szpetnar M, Hordyjewska A, Malinowska I, et al (2016) The fluctuation of free amino acids in serum during acute ischemic stroke. Curr Issues Pharm Med Sci 29:151–154. https://doi.org/10.1515/cipms-2016-0031	
649 650 651	37.	Wang X, Zhang L, Sun W, et al (2021) Changes of Metabolites in Acute Ischemic Stroke and Its Subtypes. Front Neurosci 14:1–8. https://doi.org/10.3389/fnins.2020.580929	
652 653 654	38.	Iguacel I, Schmidt JA, Perez-Cornago A, et al (2021) Associations between dietary amino acid intakes and blood concentration levels. Clin Nutr 40:3772–3779. https://doi.org/10.1016/j.clnu.2021.04.036	
655 656 657	39.	Alexander DD, Bylsma LC, Vargas AJ, et al (2016) Dairy consumption and CVD: a systematic review and meta-analysis. Br J Nutr 115:737–750. https://doi.org/10.1017/S0007114515005000	
658 659 660 661	40.	Steur M, Johnson L, Sharp SJ, et al (2021) Dietary fatty acids, macronutrient substitutions, food sources and incidence of coronary heart disease: Findings from the EPIC-CVD case-cohort study across nine european countries. J Am Heart Assoc 10:. https://doi.org/10.1161/JAHA.120.019814	
662 663 664	41.	Lorenzen JK, Astrup A (2011) Dairy calcium intake modifies responsiveness of fat metabolism and blood lipids to a high-fat diet. Br J Nutr 105:1823–1831. https://doi.org/10.1017/S0007114510005581	
665	42.	Mozaffarian D, Wu JHY (2018) Flavonoids, dairy foods, and cardiovascular and	

metabolic health: A review of emerging biologic pathways. Circ Res 122:369–384.
https://doi.org/10.1161/CIRCRESAHA.117.309008

Table 1: Participant characteristics at recruitment by sex in the EPIC study

Participant characteristics	Men	Women
Number of participants	129,005	227,137
Age, years, mean (SD)	52.1 (9.9)	50.6 (10.8)
Current smokers, n (%)	36,561 (28.3%)	48,287 (21.3%)
Moderately active or active, n (%)	63,284 (49.1%)	91,104 (40.1%)
Employed or student, n (%)	85,841 (66.5%)	124,753 (54.9%)
Vocational or university degree, n (%)	67,344 (52.2%)	109,160 (48.1%)
Alcohol consumption ≥ 15.0 g/day, n (%)	60,989 (47.2%)	42,452 (18.7%)
Alcohol consumption, g/day, median (25 th , 75 th percentile)	13.8 (4.5, 31.4)	4.0 (0.6, 12.4)
Overweight or obese (BMI $\ge 25.0 \text{ kg/m}^2$), n (%)	81,921 (63.5%)	107,011 (47.1%)
BMI, kg/m ² , mean (SD)	26.5 (3.6)	25.5 (4.5)
History of diabetes, n (%)	3,981 (3.1%)	4,668 (2.1%)
History of hypertension, n (%)	24,828 (19.2%)	41,442 (18.2%)
History of hyperlipidaemia, n (%)	21,093 (16.4%)	24,149 (10.6%)
Total protein intake, g/day, mean (SD)	90.0 (26.6)	74.9 (21.4)
Total energy intake, kcal/day, mean (SD)	2444.5 (634.8)	1965.9 (517.4)

			Ischaemic stroke			Haemorrhagic stroke		
Dietary amino acids (% of total protein)	SD (men)	SD (women)		Hazard Ratios (95% CI)	p-trend		Hazard Ratios (95% CI)	p-trend
Branched-chain amino acids	1.7	1.7	— — —	0.93 (0.88, 0.98)	0.006*	e .	0.97 (0.88, 1.07)	0.5
Isoleucine	0.4	0.4	e	0.93 (0.88, 0.98)	0.008*	e	0.97 (0.88, 1.07)	0.5
Leucine	0.8	0.8	e	0.93 (0.87, 0.98)	0.007*	e	0.97 (0.88, 1.07)	0.6
Valine	0.5	0.5	e	0.92 (0.88, 0.97)	0.003*	-	0.97 (0.88, 1.07)	0.5
Other essential amino acids	2.2	2.2	e	0.93 (0.88, 0.99)	0.02		0.97 (0.88, 1.08)	0.6
Histidine	0.3	0.3		0.94 (0.88, 1.00)	0.04		0.96 (0.86, 1.07)	0.5
Lysine	0.8	0.9	_	0.94 (0.88, 1.00)	0.04	e	0.98 (0.88, 1.10)	0.8
Methionine	0.3	0.3		0.93 (0.88, 0.99)	0.03		0.98 (0.87, 1.09)	0.7
Phenylalanine	0.4	0.4	_ 	0.92 (0.87, 0.98)	0.005*		0.97 (0.88, 1.07)	0.6
Threonine	0.4	0.4	e	0.93 (0.88, 0.99)	0.01*	e	0.97 (0.88, 1.07)	0.5
Tryptophan	0.1	0.1	_ _	0.94 (0.89, 0.99)	0.01*	e	0.98 (0.89, 1.08)	0.7
Non-essential amino acids	5.6	5.2	e	0.92 (0.87, 0.98)	0.007*	-	0.97 (0.87, 1.08)	0.6
Alanine	0.5	0.5	_ _	0.96 (0.90, 1.03)	0.2	e	0.95 (0.84, 1.06)	0.4
Arginine	0.4	0.4	_ _ _	0.97 (0.92, 1.02)	0.2	_	0.95 (0.86, 1.04)	0.2
Aspartic acid	0.6	0.6		0.96 (0.92, 1.00)	0.05	+	1.00 (0.93, 1.08)	0.9
Cystine	0.2	0.1	_	0.97 (0.90, 1.04)	0.4	B	0.92 (0.80, 1.05)	0.2
Glutamic acid	2.4	2.1	_ _	0.90 (0.84, 0.96)	0.002*		0.98 (0.86, 1.11)	0.7
Glycine	0.4	0.4	_	0.99 (0.93, 1.06)	0.8	e	0.94 (0.84, 1.06)	0.3
Hydroxyproline	0.04	0.04	_	1.01 (0.94, 1.08)	0.9	_ →	1.14 (1.02, 1.27)	0.02
Proline	1.0	0.9	• ·	0.88 (0.82, 0.94)	<0.001*	_	0.96 (0.86, 1.08)	0.5
Serine	0.5	0.4	_ _	0.92 (0.88, 0.97)	0.003*	_	1.00 (0.90, 1.10)	0.9
Tyrosine	0.4	0.4	- e	0.92 (0.88, 0.98)	0.004*	e	0.99 (0.90, 1.09)	0.9
		0.8	0.9 1 1.1 1.2 Hazard ratios and 95% CI		0.8	0.9 1 1.1 1.2 Hazard ratios and 95% Cl		

Figure 1 Hazard ratios (95% confidence intervals) for ischaemic (4295 cases) and haemorrhagic stroke (1375 cases) by increments of calibrated intakes of dietary amino acids (as percent of total protein).

Hazard ratios modelled per 1 sex-specific SD increment in dietary amino acids, expressed as percent of total protein. The model was stratified by sex and centre, and adjusted for age, calibrated energy intake, smoking, calibrated alcohol consumption, physical activity, employment status, highest level of education completed, history of diabetes, prior hypertension, prio hyperlipidaemia, body mass index. Estimates with an asterisk denote ones which were statistically significant after correcting for multiple testing.





Results are shown for amino acids which were statistically significantly associated with ischemic stroke in both calibrated and uncalibrated analyses. Hazard ratios modelled per 1 sex-spe SD (as shown in Figure 1) increment in dietary amino acids in column heading, expressed as percent of total protein, with multivariable adjustment plus amino acid on the left panel. If the amino acid in column and in left panel are the same, the result is interpretable as the hazard ratio for the amino acid based on the multivariable adjusted model.

HR (95% CI) for proline across adjustment levels	Proline and ischaemic stroke	Hazard Ratios (95% CI)	p-trend	Likelihood-ratio statistic for proline
Minimally adjusted		0.88 (0.84, 0.92)	<0.001	30.7
Plus energy intake		0.88 (0.82, 0.94)	<0.001	15.6
Plus smoking		0.91 (0.85, 0.97)	0.003	9
Plus physical activity		0.91 (0.85, 0.97)	0.003	8.6
Plus employment status		0.91 (0.85, 0.97)	0.004	8.6
Plus education level		0.92 (0.86, 0.98)	0.007	7.3
Plus alcohol consumption		0.94 (0.88, 1.00)	0.06	3.6
Plus history of diabetes		0.91 (0.85, 0.97)	0.003	8.7
Plus history of hypertension	_ _	0.89 (0.84, 0.95)	0.001	11.7
Plus history of hyperlipidaemia		0.89 (0.84, 0.95)	0.001	11.8
Plus BMI		0.88 (0.82, 0.94)	<0.001	15.1
	0.8 0.9 1 1.1 Hazard ratios and 95% Cl			

Figure 3 Hazard ratios (95% confidence intervals) for the association between dietary proline and ischaemic stroke (4295 cases), with additional adjustment each covariate.

Hazard ratios modelled per 1 SD increment (1 SD=1.0 in men, 0.9 in women) in calibrated proline intake, expressed as percent of total protein. The minimally adjusted model was stratific sex and centre, and adjusted for age (continuous). The likelihood-ratio test statistic for proline is the χ^2 statistic based on a likelihood ratio test comparing models with and without proline the varying levels of covariate adjustment. The changes in the χ^2 statistic across models could be interpreted as a measure of the extent to which the covariates could account for any associations between proline and ischaemic stroke risk.