

Kinesiophobia and Physical Activity: A Systematic Review and Meta-Analysis

Miriam Goubran^{1,2}, Ata Farajzadeh^{1,2}, Ian M. Lahart³, Martin Bilodeau^{1,2}, Matthieu P. Boisgontier^{1,2,*}

Cite this article:

Goubran M, Farajzadeh A, Lahart IM, Bilodeau M, Boisgontier MP. (2023). Kinesiophobia and physical activity: a systematic review and meta-analysis. *MedRxiv*.

Preregistration:

https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42022364063

Data, code, and materials:

<https://doi.org/10.5281/zenodo.8216112>

*Author for correspondence:

Matthieu P. Boisgontier
E-mail: matthieu.boisgontier@uottawa.ca

Key words

Chronic Pain, Exercise, Fear of Movement, Health, Humans, Physical Fitness

¹ School of Rehabilitation Sciences, Faculty of Health Sciences, University of Ottawa, Canada

² Bruyère Research Institute, Ottawa, Canada

³ Faculty of Education, Health and Wellbeing, Institute of Human Sciences, University of Wolverhampton, Wolverhampton, UK

Abstract

Objective. Physical activity is known to reduce the risk of disability, disease, and mortality. However, in some patients, an excessive, irrational, and debilitating fear of movement (i.e., kinesiophobia) is thought to induce avoidance behaviors, contributing to decreased engagement in physical activity. The aim of this study was to examine whether kinesiophobia is negatively associated with physical activity and what factors influence this relationship.

Methods. Three databases were searched for articles including both a measure of kinesiophobia and physical activity. Two reviewers screened articles for inclusion, assessed risk of bias, and extracted data from each study. Pearson product-moment correlations were pooled from eligible studies using the generic inverse pooling and random effects method to examine the relationship between kinesiophobia and physical activity.

Results. Forty-nine articles were included in the systematic review and 41 studies ($n = 4,848$) in the meta-analysis. Results showed a moderate negative correlation between kinesiophobia and physical activity ($r = -0.31$; 95% CI: -0.42 to -0.20 ; $I^2 = 1.8\%$; $p < 0.0001$). Subgroup meta-analyses revealed that the correlation was statistically significant only in patients with a cardiovascular or arthritis condition and in studies using a self-reported measure of physical activity. There was no evidence of an effect of age, gender, or pain.

Conclusions. Higher levels of kinesiophobia were moderately associated with lower levels of physical activity. However, between-study heterogeneity was considerable, and results showed no evidence of this association when physical activity was assessed with accelerometers or pedometers. Additional studies using device-based measures of physical activity are required to confirm these results and to understand the factors and mechanisms influencing this potential relationship.

Impact. Our results suggest that kinesiophobia could be considered as a limiting factor when developing physical activity promotion strategies for inactive patients.

Several decades ago, the seminal work of Morris et al. (1953)¹ showed that conductors on London double-decker buses, who were responsible for checking tickets, assisting passengers with luggage, and supervising the loading and unloading of passengers, had a lower incidence and less severe coronary heart disease than bus drivers. Since then, the scientific literature demonstrating the health benefits of physical activity has grown exponentially and expanded to include multiple health conditions². These benefits include reduced risk of disability, disease, and mortality^{2,3}. Specifically, higher levels of physical activity have been shown to contribute to a reduced risk of cardiovascular disease⁴, obesity⁵, depression⁶, hypertension⁷, cancer⁸, and dementia⁹. Yet, one in four adults worldwide does not meet the recommendations for physical activity¹⁰.

Several factors may explain physical inactivity¹¹. Environmental, interpersonal, and intrapersonal factors¹². Environmental factors include lack of access, weather conditions, and safety concerns¹³. Interpersonal factors include family responsibilities, lack of support, and lack of a gym partner¹⁴. Intrapersonal factors include gender¹⁵, age¹⁶, cognitive function^{17,18}, and socioeconomic circumstances¹⁹. One intrapersonal factor of interest in patients is kinesiophobia²⁰, which can be defined as an excessive, irrational, and debilitating fear of movement and activity resulting from a sense of vulnerability to pain, injury, or a medical condition. Kinesiophobia is typically measured using self-administered questionnaires, such as the Tampa Scale of Kinesiophobia (TSK)^{21,22}, which assesses an individual's belief that physical activity can lead to injury or pain and that the severity of their medical condition is underestimated.

The relationship between kinesiophobia and physical activity can be explained within the framework of dual models, which propose that physical activity behavior is governed by two main processes: Controlled and automatic processes²³⁻²⁶. Controlled processes rely on higher brain functions, are deliberative, require cognitive resources, and are developed through conscious thought. Automatic processes, on the other hand, rely on learned associations, are unintentional, require fewer cognitive

resources, and do not require conscious thought. Regarding automatic processes, these models propose that the perception of a cue related to physical activity automatically activates the concept of physical activity as well as the unpleasant (or pleasant) affective memories associated with this concept^{25,26}. This activation leads to an impulse that favors the tendency to avoid (or approach) physical activity²⁷. Thus, negative affective associations are likely to hinder physical activity. Accordingly, an aversive fear of pain, injury, or aggravation of a medical condition that has been associated with the concept of movement may result in the development of automatic avoidance behaviors that contribute to the maintenance and exacerbation of this fear, and ultimately lead to a phobic state (i.e., kinesiophobia) that diminishes the ability to engage in regular physical activity.

The main objective of this study was to systematically review and meta-analyze the direct relationship between kinesiophobia and physical activity. We hypothesized that levels of kinesiophobia would be negatively associated with levels of physical activity. In addition, we examined whether this association was influenced by kinesiophobia measures, physical activity measures (i.e., accelerometers, pedometers, questionnaires), physical activity outcome (e.g., total physical activity, moderate or vigorous physical activity, steps per day), health status, age, gender, or pain.

Methods

Search Strategy

This review was reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines²⁸. Potential studies were identified by searching the MEDLINE (via PubMed), PsychInfo, and CINAHL databases. In October and November 2022, two reviewers (MG and AF) searched for all available records using the following combination of keywords in the title or abstract of the article: “kinesiophobia” and “physical activity”. Specifically, in PubMed, the search was “physical activity” (MeSH terms) AND “kinesiophobia” (All Fields)”. In PsychInfo and CINAHL, the search was “kinesiophobia”

AND “physical activity” in all fields. To reduce literature bias^{29,30}, this systematic review was pre-registered in PROSPERO³¹.

Eligibility Criteria and Study Selection

Inclusion Criteria

To be included in this systematic review, articles had to be published in a peer-reviewed journal, be written in English, report original data collected from human participants, include at least one self-reported measure of kinesiophobia and one measure of physical activity, and formally test the association between these two variables. The physical activity measure could be a self-reported measure of the level of physical activity or the use of an accelerometer while participants are engaged in their normal daily activities.

Exclusion Criteria

Studies were excluded if they were published as a book chapter, study protocol, conference abstract, or were based on laboratory-based measures of physical fitness (e.g., maximal muscle force, $\dot{V}O_2$ max) and not on a measure of physical activity.

Study Selection

Article screening was performed in Covidence systematic review software (Veritas Health Innovation, Melbourne, Australia; www.covidence.org), a web-based collaborative software platform that streamlines the production of systematic reviews. After removing duplicates, titles and abstracts were independently reviewed by two reviewers (MG, AF) according to the inclusion and exclusion criteria using a systematic 5-step process. If there was any doubt at any step, the full text was further reviewed. Step 1: Articles not written in English were excluded. Step 2: Articles that did not report original empirical data were excluded (e.g., reviews, meta-analyses, commentaries, technical reports, case studies). Step 3: Articles that did not involve human participants were excluded. Step 4: Articles that did not assess both kinesiophobia and physical activity were excluded. Step 5: Articles that did not formally test the association between

kinesiophobia and physical activity were excluded. In addition, we performed reference screening and forward citation tracking on the articles remaining after step 5. Any disagreements between the two reviewers were resolved by consensus among four reviewers (MG, AF, MB, MPB).

Data Extraction

Data extracted from selected articles included first author’s name, article title, publication year, digital object identifier (DOI), number of participants, number of men and women, age range, mean age, mean weight, mean height, mean body mass index, health status, mean pain intensity, type of kinesiophobia measure, level of kinesiophobia, type of physical activity measure, type of physical activity outcome, level of physical activity, as well as statistical estimates and significance of the association between kinesiophobia and physical activity.

Bias Assessment

The risk of bias of the studies included in the systematic review was estimated using the critical appraisal tool for assessing the quality of cross-sectional studies (AXIS)³². AXIS is a 20-item checklist designed to assess the introduction, methods, results, discussion, conflict of interest, and ethical approval or consent of the included studies.

Meta-Analysis

In our meta-analysis, we pooled Pearson product-moment correlations from eligible studies to examine the relationship between kinesiophobia and physical activity. When a study measured physical activity with both a questionnaire and accelerometers, the correlation included in the analysis was that of the most reliable outcome, i.e., the accelerometer-based outcome. Correlations were pooled using the generic inverse pooling method via the ‘metacor’ function in the R ‘meta’ package³³. This function automatically performs a necessary Fisher’s z-transformation on the original, untransformed correlations prior to pooling. The ‘metacor’ function also reconverts the pooled association back to its original form for ease of interpretation.




Figure 1. PRISMA 2020 flow diagram

We anticipated considerable between-study heterogeneity, and therefore used a random-effects model to pool correlations. The restricted maximum likelihood (RML) estimator³⁴ was used to calculate the heterogeneity variance τ^2 . In addition to τ^2 , to quantify between-study heterogeneity, we report the I^2 statistic, which provides the percentage of variability in the correlations that is not caused by sampling error³⁵. The I^2 statistic was interpreted as follows: 0-40%, may not be important; 30-60%, may represent moderate heterogeneity; 50-90%, may represent substantial heterogeneity; and 75-100%, may represent considerable heterogeneity. To reduce the risk of false positives, we used a Knapp-Hartung adjustment³⁶ to calculate the confidence interval around the pooled association. We also report the prediction interval, which provides a range within which we can expect the associations of future studies to fall based on the current evidence. The pooled correlation was interpreted using Cohen's conventions³⁷: $r \approx -0.10$, small negative correlation; $r \approx -0.30$, moderate negative correlation; $r \approx -0.50$, large negative correlation. Egger's regression test of funnel plot

asymmetry³⁸ and a p-curve analysis³⁹ were conducted to assess potential publication bias in our meta-analysis. The Rücker's limit meta-analysis method⁴⁰, which explicitly includes the heterogeneity variance in the model, was used to compute bias-corrected estimate of the true effect size.

We conducted subgroup analyses to examine the differences in correlations between studies including participants with different health conditions and using different types of physical activity measures (i.e., device-based vs. self-reported), physical activity measurement instruments (i.e., type of questionnaires, type of devices), physical activity outcomes, and kinesiophobia measures. In addition, we used meta-regressions to examine if the average age of participants or the proportion of women in a study predicted the reported correlation between kinesiophobia and physical activity.

A secondary meta-analysis was conducted using the same approach, but based on Spearman's rho values, to further test the relationship between kinesiophobia and device-based physical activity.

A sensitivity analysis was conducted to examine whether the quality of the studies (AXIS score) affected the results.

All analyses were performed in RStudio integrated development environment (IDE) (2023.06.1+524, “Mountain Hydrangea” release) for R software environment⁴¹ using the ‘meta’, ‘metafor’, and ‘metasens’ R packages^{33,40,42,43}.

Results

Literature Search

The primary search identified 395 potentially relevant articles from the three databases (Figure 1), including 80 duplicates. Of the 315 articles screened, disagreement occurred in 25 cases (7.9%), all of which were resolved by consensus. All articles remained after step 1 as they were all written in English. Ninety-six articles were excluded in step 2 because they did not report original data. No articles were excluded in step 3 because they all involved human participants. One hundred and thirty-six articles were excluded in step 4 because they did not assess kinesiophobia ($k = 3$) or physical activity ($k = 133$). Forty-five articles were excluded at step 5 because they did not formally test the association between the kinesiophobia and physical activity. The corresponding authors of these latter articles were contacted by email to request the Pearson correlation value of this association and the sample size used to calculate it or the raw data for the physical activity and kinesiophobia measures. Fifteen authors replied to our email: 3 authors provided their datasets⁴⁴⁻⁴⁶, 5 authors provided the Pearson’s correlation value⁴⁷⁻⁵¹, 4 authors replied that they would contact their co-authors but did not get back to us. Based on reference screening and forward citation tracking, the authors identified 13 studies that assessed both physical activity and kinesiophobia, but none provided an estimate of their relationship. The corresponding authors of these 13 studies were asked by email to share this estimate or their data. Three of them responded and provided the Pearson’s correlation value⁵²⁻⁵⁴. The remaining 10 authors did not respond to our email. In total, data from 49 articles were included in the

systematic review. AXIS scores ranged from 14 to 18 out of 20, with a mean of 16.9 ± 0.6 (Table 1).

Descriptive Results

Participants

The 49 articles identified by the systematic review included a total of 6,084 participants aged 11 to 85 years, including 2,655 women, 1,967 men, and 1,462 participants whose gender and sex was not reported. The studies investigated populations with pain ($k = 20$ ^{44,46,49,55-71}), cardiovascular ($k = 6$ ^{52,53,72-75}), surgery ($k = 6$ ^{45,75-79}), arthritis ($k = 6$ ^{51,80-84}), neurological ($k = 2$ ^{85,86}), pulmonary ($k = 2$ ^{87,88}), and cancer ($k = 1$ ⁵⁰) conditions, as well as healthy adults ($k = 6$ ^{47,48,54,89-91}) (Table 1).

Kinesiophobia

In 44 of the 49 studies, kinesiophobia was assessed using the 17-item TSK ($k = 30$)^{44,47-49,51,54-58,61-65,68,70,77,79-81,83-88,90,92}, shorter versions of the TSK [TSK-11⁹³, $k = 8$ ^{45,60,66,67,71,75,78,91}; TSK-14, $k = 1$ ⁵⁰; TSK-13⁹⁴, $k = 2$ ^{59,76}], or its adaptation for patients with coronary artery disease [TSK-Heart⁹⁵, $k = 2$ ^{52,72}]. The TSK is a questionnaire that assesses the belief that movements can lead to (re)injury, pain, or aggravation of an underlying and serious medical condition²². Each item is rated on a Likert scale ranging from 1 (strongly disagree) to 4 (strongly agree). On the TSK-17, a score of 37 is used to distinguish between low (≤ 37) and high (> 37) levels of kinesiophobia²¹. On the TSK-13, scores inferior to 23 are considered sub-clinical⁹⁶. The other measures that were used are the Kinesiophobia Causes Scale⁹⁷ (KCS; $k = 2$ ^{54,89}), the Fear of Activities in Situations scale⁷⁴ (FActS; $k = 1$ ⁷⁴), the Brief Fear of Movement Scale for Osteoarthritis⁹⁸ (BFMSO; $k = 1$ ⁵¹), and the Fear-Avoidance Belief Questionnaire⁹⁹ (FABQ; $k = 1$ ⁵³).

Forty-three studies reported mean levels of kinesiophobia (Table 1). The studies based on the TSK-17 or TSK-Heart (mean range: 17 to 68) reporting the highest levels of kinesiophobia were those involving participants with a cardiovascular condition (41.4 to 49.7), followed by studies testing participants with arthritis (31.8 to 45.0), chronic pain (30.5 to 44.4), or a pulmonary condition (39.6 to 42). Levels of kinesiophobia were lower in parti-

Study (Year)	N (# women)	Mean age (SD or range)	Health status	Mean kinesiophobia (SD or range; measure)	Mean physical activity (SD or range; measure)	Level of pain (measure)	Corr.	p-value	AXIS score
Alschuler (2011)	20 (9)	46.1 (9.35)	Chronic low back pain	30.55 (TSK-17)	228 counts/min (wrist accelero.)	4.87 (NRS)	n.a.	0.03	18
Altuğ (2016)	112 (73)	45.0 (14.6)	Chronic low back pain	44.30 (TSK-17)	5495 MET-min/week (IPAQ)	3.45 (VAS)	r = -0.096	0.313	16
Assadourian (2020)	147 (88)	49 (12)	Chronic low back pain	n.a. (TSK-17)	n.a. (diary; < vs. ≥ 1h/week)	6.7 (NRS)	r = -0.022†	0.813†	18
Atici (2022)	254 (171)	n.a. (>65)	Older adults	54.55 (KCS)	182.8 (PASE)	n.a.	ρ = -0.345	<0.001	16
Aydemir (2022)	37 (25)	58.8 (8.6)	Knee osteoarthritis	40.3 (TSK-17)	4.8 (UCLA)	52.0 (KOOS-P)	r = -0.773	<0.05	17
Aykut Selcuk (2020)	67 (67)	60.6 (8.0)	Knee osteoarthritis	44.8 (TSK-17)	n.a. (IPAQ; low vs. moderate vs. high)	4.6 (VAS)	r = -0.247	0.019	17
	29 (0)	61.6 (8.1)	Knee osteoarthritis	42.0 (TSK-17)	n.a. (IPAQ; low vs. moderate vs. high)	2.9 (VAS)	r = -0.309	0.116	
Baday-Keskin (2022)	88 (67)	52 (n.a.)	Rheumatoid arthritis	45 (TSK-17)	594 MET-min/week (IPAQ)	4.8 (VAS)	r = -0.12	>0.05	17
	93 (67)	45 (n.a.)	Healthy adults	39 (TSK-17)	971 MET-min/week (IPAQ)	n.a.	n.a.		
Baez (2020)	40 (24)	24.3 (4.1)	Surgery (ACLR)	18.2 (TSK-11)	8 657 steps/day (hip pedo.) 7.7 (TAS)	81.5 (KOOS-P)	r = 0.181†	0.265†	18
							n.a.	n.a.	
Bahar Özdemir (2021)	101 (59)	33.9 (6.0)	Healthy	36.4 (5.8; TSK-17)	756 MET-min/week (1090; IPAQ)	3.1 (3.3; NRS)	r = -0.007†	0.944†	18
Baykal Sahin (2021)	98 (35)	58.1 (10.4)	Coronary artery disease	41.4 (6.2; TSK-17)	839 MET-min/week (1212; IPAQ)	60.1 (27.2; SF-36)	r = -0.315	0.002	17
Bernard (2015)	121 (121)	65.5 (57-75)	Post-menopausal women	36 (TSK-17)	n.a. (PAQE)	n.a.	r = -0.05†	0.55†	18
Bossenbroek (2009)	15 (10)	53 (6.3)	Pulmonary (COPD)	42 (TSK-17)	1407 steps/day (pedo.) 1349 MET-min/week (SQUASH)	n.a.	n.a.	n.a.	17
	47 (29)	55 (5.5)	Lung transplant	30 (TSK-17)	6642 steps/day (pedo.) 5434 MET-min/week (SQUASH)				
Carvalho (2017)	119 (82)	39.1 (11.2)	Chronic low back pain	41 (TSK-17)	6844 steps/day (hip accelero.) 296 counts/min (hip accelero.) 22 min MVPA/day (hip accelero.) 333 min LPA/day (hip accelero.) 6.7 (BHPAQ)	6.7 (NRS)	ρ = -0.15 r = -0.02 ρ = -0.13 r = 0.09 r = -0.18	>0.05 >0.05 >0.05 >0.05 <0.05	17
Coronado (2021)	248 (126)	62.2 (11.9)	Surgery (laminectomy)	28.4 (TSK-13)	427 counts/min (hip accelerometer)	3.1 (NRS)	r = -0.05	>0.05	16
Corrigan (2018)	53 (18)	54.8 (34-65)	Achilles tendinopathy	35.4 (TSK-17)	n.a. (SGPALS)	n.a.	d = 0.027	0.969	16
Dabek (2020)	130 (n.a.)	n.a.	Coronary disease	44.3 (TSK-Heart)	1545 MET-min/week (IPAQ)	n.a.	r = -0.523	<0.001	17
	119 (n.a.)		Hypertension	44.4 (TSK-Heart)	1509 MET-min/week (IPAQ)		r = -0.410	<0.001	
	27 (n.a.)		Heart valve defect	44.4 (TSK-Heart)	1308 MET-min/week (IPAQ)		r = -0.201	>0.05	
	72 (n.a.)		Myocardial infarction	46.7 (TSK-Heart)	1369 MET-min/week (IPAQ)		r = -0.428	<0.001	
	86 (n.a.)		Rhythm disorder	43.3 (TSK-Heart)	1660 MET-min/week (IPAQ)		r = -0.563	<0.001	
	18 (n.a.)		Stroke	49.7 (TSK-Heart)	1135 MET-min/week (IPAQ)		r = -0.868	<0.001	
	15 (n.a.)		Other CVD	44.2 (TSK-Heart)	2207 MET-min/week (IPAQ)		r = -0.663	0.01	
Demirbüken (2016)	99 (65)	43.5 (12.8)	Chronic neck pain	41.82 (TSK-17)	3749 MET-min/week (IPAQ)	6.47 (VAS)	r = -0.153	n.a.	17
Donnarumma (2017)	51 (12)	61.9 (13.9)	Surgery (laminectomy)	n.a. (TSK-17)	n.a. (IPAQ)	3.5 (GRS)	n.a.	0.01	16
Elfving (2007)	64 (39)	47 (19-64)	Chronic low back pain	n.a. (TSK-13)	n.a. (SGPALS)	n.a.	n.a.	0.010	17
González de La Flor (2022)	42 (32)	36.7 (13.2)	Chronic headache	9 (TSK-11)	n.a. (IPAQ)	7.14 (NRS)	ρ = 0.204	n.a.	17
Helmus (2012)	53 (37)	39.9 (11.3)	Chronic MSK pain	35.4 (TSK-17)	138 counts/min (hip accelero.)	5.8 (VAS)	r = -0.05	0.75	17
Huijnen (2010)	111 (n.a.)	44.1 (10.3)	Subacute low back pain	36.0 (TSK-17)	n.a. (trunk accelero.)	n.a. (NRS)	n.a.	>0.05	17
Kiliç (2019)	200 (120)	53.2 (6.0)	Knee osteoarthritis	31.8. (TSK-17)	1947 MET-min/week (IPAQ)	24.1 (OKS)	r = -0.693	<0.001	17
Knapik (2019)	135 (59)	71.9 (4.8)	Coronary artery disease	43.02 (TSK-Heart)	2.60 (ad-hoc questionnaire)	n.a.	r = -0.8†	n.a.	17
Koho (2011)	93 (60)	44.0 (17-68)	Chronic pain	n.a. (TSK-17)	n.a. (LTPAQ)	6.4 (VAS)	r = 0.10	>0.05	16
Lotzke (2018)	118 (63)	46 (8)	Chronic low back pain	38.1 (TSK-17)	198 min MVPA/week (accelero.)	6.1 (VAS)	n.a.	0.034	17
Luthi (2018)	433 (n.a.)	n.a.	Chronic MSK pain	44.6 (TSK_17)	4.45 (BHPAQ)	4.45 (BPI-S)	r = 0.067†	0.759†	18
Marques-Sule (2022)	117 (51)	56 (12.1)	Heart transplantation	32.5 (TSK-11)	219 MET-min/week (IPAQ)	n.a.	r = -0.32	0.001	16
Miller (2018)	52 (32)	67.4 (5.1)	Healthy Older Adults	18.9 (TSK-17)	6743 steps/day (hip accelero.)	1.4 (QWBS-P)	r = -0.54	<0.001	16
Minetama (2022)	71 (36)	71.6 (5.6)	Lumbar spinal stenosis	24.8 (TSK-11)	3601 steps/day (pedo.)	6.2 (NRS)	r = -0.229	0.055	17
Navarro-Ledesma (2022)	41 (41)	52.6 (8.0)	Fibromyalgia	27.5 (6.9; TSK-11)	n.a.	n.a.	r = -0.059	>0.05	17
Norte (2019)	77 (35)	21.6 (7.8)	Surgery (ACLR)	32.9 (6.0; TSK-17)	72.7 (34.9; GLTEQ)	91.4 (9.2; KOOS-P)	r = -0.312	<0.05	17
Ohlman (2018)	52 (33)	67.4 (5.1)	Older adults	18.8 (4.5; TSK-11)	n.a. (hip accelero.)	n.a.	ρ = -0.29	<0.05	17
Olsson (2014)	81 (12)	40.0 (9.6)	Achilles tendon rupture	35.9 (7.5; TSK-17)	2.9 (1.0; SGPALS)	n.a.	ρ = -0.275	0.013	18
Ozer (2022)	62 (30)	36.8 (6.1)	Asthma	39.6 (5.8; TSK-17)	2249 MET-min/week (1333; IPAQ)	n.a.	r = -0.889	0.001	17
Pazzinatto (2022)	92 (92)	n.a. (18-35)	Patellofemoral pain	35.3 (6.8; TSK-17)	7.8 (1.5; BHPAQ)	5.1 (2.1; VAS)	ρ = -0.14	n.a.	17
Pedler (2018)	103 (74)	39.7 (13.9)	Whiplash injury	26 (TSK-11)	9.9% of active time (8.2; trunk accelero.)	4.0 (2.4; VAS)	r = 0.140	>0.05	16
Priore (2020)	50 (37)	22.4 (3.9)	Patellofemoral pain	36.7 (TSK-17)	3088 MET-min/week (IPAQ)	n.a. (VAS)	r = -0.251†	0.072†	17
Roaldsen (2009)	98 (62)	76 (60-86)	Leg ulcer	12 (FABQ)	2.6 (SGPALS)	1.3 (VRS)	r = -0.39†	n.a.	17

Saulicz (2016)	105 (105)	n.a.	Older adults	45.2 (15.6; KCS)	n.a. (BHPAQ)	n.a.	r = -0.577 [†]	<0.001 [†]	16
Sertel (2021)	163 (76)	71.4 (6.0)	Chronic pain	44.4 (7.7; TSK-17)	171.3 (76.2; PASE)	n.a. (VAS)	r = -0.021	>0.05	17
Spaderna (2020)	61 (13)	67.5 (10.7)	Heart failure	1.5 (FactS)	2332 kcal/day (361; hip accelero.)	n.a.	r = -0.28	<0.05	17
Strandberg (2022)	451 ^{††} (n.a.)	n.a.	Cancer	n.a. (TSK-14)	1.3 h MVPA/week (0.8; arm accelero.)	n.a.	r = -0.084 [†]	0.074 [†]	16
Sütçü (2021)	20 (10)	69.8 (9.4)	Parkinson's disease	39.8 (7.4; TSK-17)	3078 steps/day (arm accelero.) 2055 kcal/day (475; arm accelero.)	n.a.	n.a.	0.157 0.013	16
Uritani (2020)	167 (105)	62.2 (7.5)	Knee osteoarthritis	12.5 (BFMSO)	7998 steps/day (thigh accelero.)	5.7 (NRS)	r = -0.163 [†]	0.035 [†]	16
Verbunt (2005)	123 (57)	44.1 (10.3)	Subacute low back pain	36.0 (TSK-17)	n.a. (hip accelero.)	4.2 (VAS)	ρ = 0.06	>0.05	17
Wasiuk-Zowada (2022)	80 (60)	45.5 (8.6)	Multiple sclerosis	36.6 (TSK-17)	5.1 (BHPAQ)	3.5 (VAS)	r = -0.363	0.001	16
Yüksel Karsli (2021)	34 (12)	41 (n.a.)	Radiographic SpA	42 (TSK-17)	2203 min LPA/day (1377; hip accelero.) 210 min MPA/day (109; hip accelero.) 0 min VPA/day (3; hip accelero.)	n.a.	ρ = -0.16 ρ = -0.158 ρ = -0.394	0.929 0.373 0.021	17
	33 (10)	33 (n.a.)	Non-radiographic SpA	36 (TSK-17)	2576 min LPA/day (1858; hip accelero.) 265 min MPA/day (216; hip accelero.) 2 min VPA/day (12; hip accelero.)		ρ = -0.001 ρ = 0.013 ρ = -0.240	0.997 0.947 0.209	
Zelle et al. (2016)	487 (209)	51.6 (12.5)	Renal transplantation	n.a. (TSK-11)	165 METs-min/day (MLTPAQ & TOAQ)	n.a.	r = -0.22	<0.001	17

Table 1. Sample characteristics of the studies included in the systematic review.

Notes. Accelero. = Accelerometer, ACLR = anterior cruciate ligament reconstruction, BHPAQ = Baecke Habitual Physical Activity Questionnaire (3 – 15), BFMSO = Brief Fear of Movement Scale for Osteoarthritis (6 – 24), BPI-S = Brief Pain Inventory-Severity (1 – 10), COPD = Chronic Obstructive Pulmonary Disease, Corr. = Correlation, CVD = cardiovascular disease, FABQ = Fear-Avoidance Belief Questionnaire (0 – 24), FactS = Fear of Activity in Situations (0 – 30), GLTEQ = Godin Leisure Time Exercise Questionnaire (0 – 119), GRS = Graphic Rating Scale (0 – 10), IPAQ = short form of the International Physical Activity Questionnaire, KCS = Kinesiophobia Causes Scale (0 – 100), KOOS-P = Knee Injury and Osteoarthritis Outcome Score – Pain (0 – 100), LPA = light physical activity, n.a. = not available, MPA = moderate physical activity, MSK = Musculoskeletal, MVPA = moderate-to-vigorous physical activity, NRS = Numeric Rating Scale (maximum score = 0 – 10), OKS = Oxford Knee Score – Pain, PA = Physical Activity, PASE = Physical Activity Scale for the Elderly, Pedom. = Pedometer, r = Pearson's correlation coefficient, ρ = Spearman's correlation coefficient, SF-36 = 36-Item Short Form Survey (0 – 100), QWBS-P = Quality of Well-Being Scale-Pain, SGPALS = Saltin-Grimby Physical Activity Level Scale (1 – 4), SpA = axial spondyloarthritis, TAS = Tegner Activity Scale (0 – 10), TOAQ = Tecumseh Occupational Activity Questionnaire, TSK = Tampa Scale of Kinesiophobia [TSK-17: 17 – 68, TSK-Heart = 17 – 68, TSK-14 = 14 – 56, TSK-13 = 13 – 52, TSK-11 = 11 – 44], UCLA = University of California Los Angeles activity score (1 – 10), VAS = Visual Analog Scale (0 – 10), VPA = vigorous physical activity, VRS = Verbal Rating Scale for pain assessment (0 – 5), † was obtained by email from authors, ††Number of participants used to calculate the correlation, according to the email sent by the authors.

cipants with a neurological (36.0 to 39.8) or surgical condition (32.9) and in healthy adults (18.9 to 39.0).

Forty-three studies reported mean levels of kinesiophobia (Table 1). The studies based on the TSK-17 or TSK-Heart (mean range: 17 to 68) reporting the highest levels of kinesiophobia were those involving participants with a cardiovascular condition (41.4 to 49.7), followed by studies testing participants with arthritis (31.8 to 45.0), chronic pain (30.5 to 44.4), or a pulmonary condition (39.6 to 42). Levels of kinesiophobia were lower in participants with a neurological (36.0 to 39.8) or surgical condition (32.9) and in healthy adults (18.9 to 39.0).

Physical Activity

Thirty-four studies assessed physical activity using a self-reported measure (Table 1). Most of these questionnaire-based studies used the short form of the International Physical Activity Questionnaire (IPAQ-SF) ($k = 13^{48,49,56,58,60,72,73,75,77,81-83,87}$), which consists of 6 items assessing time spent in light (i.e., walking), moderate (e.g., carrying light loads, cycling at moderate speed, doubles tennis), and vigorous physical activity (e.g., digging, fast cycling, heavy lifting, aerobics) over the last 7 days¹⁰⁰. Other questionnaires were used to assess physical activity, such as the Baecke Habitual Physical Activity Questionnaire¹⁰¹ (BHPAQ; $k = 5^{46,54,57,65,85}$), the Saltin-Grimby Physical Activity Level Scale¹⁰² (SGPALS; $k = 4^{53,59,70,92}$), the Godin-Shephard Leisure-Time Exercise Questionnaire¹⁰³ (GLTEQ; $k = 2^{66,79}$), the Minnesota Leisure Time Physical Activity Questionnaire¹⁰⁴ (MLTPAQ; $k = 1^{78}$), the Physical Activity Scale for the Elderly¹⁰⁵ (PASE; $k = 2^{68,89}$), the Physical Activity Questionnaire for the Elderly¹⁰⁶ (PAQE; $k = 1^{47}$), the Short Questionnaire to Assess Health Enhancing Questionnaires¹⁰⁷ (SQUASH; $k = 1^{88}$), the Tegner Assessment Scale¹⁰⁸ (TAS; $k = 1^{45}$), and the University of California Los Angeles (UCLA) activity score¹⁰⁹ ($k = 1^{80}$).

Physical activity was also assessed with devices such as accelerometers measuring accelerations in 3 dimensions ($k = 15^{50,51,55,57,61,62,64,67,69,74,76,84,86,90,91}$) and pedometers measuring the number of steps ($k = 3^{45,71,88}$) (Table 1). In most studies, the device was worn at the hip ($k = 9^{45,57,61,69,74,76,84,90,91}$).

Other positions included trunk ($k = 2^{62,67}$), arm ($k = 2^{50,86}$), wrist ($k = 1^{55}$), and thigh ($k = 1^{51}$), with three studies not reporting where the device was worn^{64,71,88}. Most studies that employed accelerometer-based measures used the ActiGraph (ActiGraph, LLC, Pensacola, FL, USA) GT3X+ ($k = 4^{64,76,90,91}$) or wGT3X-BT ($k = 1^{84}$). The other accelerometers were the RT3 (Stayhealthy Inc., Monrovia, CA, USA; $k = 3^{61,62,69}$), the SenseWear Pro3 Armband (BodyMedia, Pittsburgh, PA, USA; $k = 2^{50,86}$), the Activity Sensory Move II (movisens GmbH, Karlsruhe, Germany; $k = 1^{74}$), the LifeShirt (Vivometrics, Inc., Ventura, CA, USA; $k = 1^{67}$), and the ActiWatch (Mini Mitter Co., Inc., Bend, OR, USA; $k = 1^{55}$). The type of accelerometer was not reported in one study⁵⁷. The pedometers were the Digi-Walker SW-200 (New Lifestyles Inc., Lees Summit, MO, USA; $k = 2^{45,88}$) and the Active Style Pro HJA-350IT (Omron Healthcare, Kyoto, Japan; $k = 1^{71}$).

To assess physical activity, the studies used the following outcomes: Score from a questionnaire (e.g., TAS, PAQE, BHPAQ, SGPALS, MLTPAQ; $k = 18^{45-47,52-54,57,59,63,65,66,70,78-80,85,89,92}$), MET-min/week ($k = 13^{48,49,56,58,72,73,75,78,81-83,87,88}$), steps per day ($k = 7^{45,51,57,71,86,88,90}$), hours per day or week ($k = 6^{44,50,57,64,83,84}$), counts per minute ($k = 4^{55,57,61,76}$), kilocalories per day ($k = 2^{74,86}$), or percentage of active time ($k = 1^{67}$). Five studies used multiple physical activity outcomes^{45,57,84,86,88}.

Association Between Physical Activity and Kinesiophobia

Thirty-two of the 49 articles reported a correlation estimate of the association between physical activity and kinesiophobia. Twenty-three articles reported a correlation estimate of the association between physical activity and kinesiophobia. Twenty-three articles reported at least one Pearson's r estimate, eight reported at least one Spearman's ρ estimate, and one reported a Cohen's d . Eleven additional Pearson's r values were obtained by email from the authors⁴⁴⁻⁵⁴, for a total of 34 articles including 41 Pearson's r estimates that were used in the meta-analysis (Table 1).




Figure 2. Main meta-analysis: Correlation between kinesiophobia and physical activity (k = 41, n = 4,848). Heterogeneity: $\tau^2 = 0.1204$; $\chi^2 = 490.59$, df = 40 ($P < 0.01$); $I^2 = 92\%$

Notes: 95% CI = 95% confidence interval, IV = Inverse-variance method, Random = Random-effects method.

Pain

Mean pain intensity at rest was reported in 28 out of the 49 articles included in the systematic review. Most studies used the Visual Analog Scale¹¹⁰ (VAS; $k = 11^{56,61,63-65,67,69,81,83,85}$) or the Numeric Rating Scale¹¹¹ (NRS; $k = 8^{44,48,51,55,57,60,71,76}$). Other studies used the Knee Injury Osteoarthritis Outcome Score pain subscale¹¹² (KOOS-P; $k = 3^{45,79,80}$), Brief Pain Inventory¹¹³ ($k = 1^{46}$), Oxford Knee Score¹¹⁴ (OKS; $k = 1^{82}$), the Quality of Well-Being Scale – Self-administered Pain Scale¹¹⁵ (QWBS-P; $k = 1^{90}$), the Short Form 36 bodily pain¹¹⁶ (SF-36; $k = 1^{73}$), the Graphic Rating Scale¹¹⁷ (GRS; $k = 1^{77}$), and the Verbal Rating Scale¹¹⁸ (VRS; $k = 1^{53}$). In the meta-analysis, scores that were not on a 0-100 scale in the initial measure were scaled to that range.

Meta-Analysis

Main Meta-Analysis

Our meta-analysis of 41 studies ($n = 4,848$) revealed a statistically significant moderate negative correlation between kinesiophobia and physical activity ($r = -0.31$; 95% confidence interval [95% CI]: -0.42 to -0.20 ; $p < 0.0001$; Table 2; Figure 2). However, we observed considerable between-study statistical heterogeneity ($Tau^2 = 0.12$, 95% CI: 0.08 to 0.22 ; $I^2 = 91.8\%$, 95% CI: 89.9 to 93.5%), and the prediction interval ranged from $r = -0.78$ to 0.37 , indicating that a moderate positive correlation cannot be ruled out for future studies.




Figure 3. Contour-enhanced funnel plot of the main meta-analysis




Figure 4. P-curve analysis

Publication Bias Assessment

Egger's regression test showed that the data in the funnel plot was asymmetric ($b = -2.85$, 95% CI: -5.55 to -0.14 , $p = 0.046$; Figure 3), which may be explained by publication bias, but also by other potential causes, such as different study procedures and between-study heterogeneity¹¹⁹, which was considerable here. The bias-corrected estimate of the true effect size, calculated using Rücker's limit meta-analysis method, showed that the correlation would remain significant if there was publication bias ($r = -0.18$; 95% CI: -0.34 to -0.01 ; $p = 0.0378$). A total of 41 studies were provided to the p-curve analysis, including 21 (51.2%) studies with $p < 0.05$ and 18 studies (43.9%) with $p < 0.025$ (Figure 4). The p-value of the right-skewness test was < 0.0001 for both the half and full curve, suggesting that evidential value was present¹²⁰.

	k	n	cor.	95% CI	I² (%)	p
Main						<.0001
Kinesiophobia and physical activity	41	4848	-.31	[-.42; -.20]	92	
Subgroup: Health status						<.0001
Chronic pain	11	1410	-.03	[-.10; .04]	18	
Cardiovascular	11	958	-.47	[-.59; -.32]	84	
Arthritis	6	588	-.42	[-.69; -.04]	93	
Surgery	4	852	-.12	[-.42; .21]	73	
Older adults	3	278	-.40	[-.86; .40]	91	
Neurological	2	98	-.62	[-1.0; 1.0]	91	
Cancer	1	451	-.08	[-.42; -.20]		
Young adults	1	101	-.01	[-.20; .19]		
Pulmonary	1	62	-.89	[-.93; -.82]		
Acute pain	1	50	-.25	[-.49; .03]		
Subgroup: Physical activity measure						.011
Self-reported	32	3934	-.36	[-.48; -.24]	93	
Device-based	9	914	-.12	[-.28; -.06]	70	
Secondary: rho values						.217
Kinesiophobia and device-based physical activity	5	361	-.10	[-.27; .09]	30	
Subgroup: Physical activity measurement instrument						.014
IPAQ	18	1490	-.43	[-.57; -.26]	90	
Accelerometer	8	1254	-.13	[-.30; .05]	69	
BHPAQ	3	618	-.30	[-.84; .54]	96	
Pedometer	2	111	-.04	[-.99; .99]	76	
MLTPAQ	2	580	-.07	[-.97; .96]	87	
GLTEQ	2	118	-.20	[-.95; .90]	43	
PASE	1	163	-.31	[-.42; -.20]		
PAQE	1	135	-.80	[-.85; -.73]		
Diary	1	123	-.02	[-.20; .16]		
Ad-hoc questionnaire	1	121	-.05	[-.23; .13]		
SGPALS	1	98	-.39	[-.55; -.21]		
UCLA	1	37	-.77	[-.88; -.60]		
Subgroup: Physical activity outcome						<.0001
MET-min/week	19	1977	-.42	[-.56; -.25]	90	
Score	11	1383	-.33	[-.56; -.06]	95	
Steps/day	4	330	-.21	[-.61; .28]	78	
Active time	3	677	.01	[-.27; .29]	52	
Counts/min	3	420	-.04	[-.08; .00]	0	
Kcal/day	1	61	-.28	[-.50; -.03]		
Subgroup: Kinesiophobia measure						.032
TSK-17	21	2257	-.28	[-.43; -.11]	93	
TSK-Heart	8	602	-.59	[-.75; -.36]	85	
TSK-11	6	859	-.09	[-.31; .13]	74	
TSK-14	1	451	-.08	[-.17; .01]		
TSK-13	1	248	-.05	[-.17; .08]		
KCS	1	105	-.58	[-.69; -.43]		
FABQ	1	98	-.39	[-.55; -.21]		
FActS	1	61	-.28	[-.50; -.03]		

Table 2. Results of the main, subgroup, and secondary meta-analyses

Notes. 95% CI = 95% confidence interval, BHPAQ = Baecke Habitual Physical Activity Questionnaire, Cor. = Correlation estimate, FABQ = Fear-Avoidance Belief Questionnaire, FActS = Fear of Activity in Situations, GLTEQ = Godin Leisure Time Exercise Questionnaire, IPAQ = short form of the International Physical Activity Questionnaire, k = number of studies, KCS = Kinesiophobia Causes Scale, MLTPAQ = Minnesota Leisure Time Physical Activity Questionnaire, n = number of participants, p = p-value for between-group difference, PAQE = Physical Activity Questionnaire for the Elderly, PASE = Physical Activity Scale for the Elderly, SGPALS = Saltin-Grimby Physical Activity Level Scale, TSK = Tampa Scale of Kinesiophobia, UCLA = University of California Los Angeles activity score.

Subgroup Meta-Analyses

The test of subgroup differences between health status measures was possible between studies comprising people with chronic pain (k = 11), car-




Figure 5. Subgroup meta-analysis: Differences according to health status.

diovascular condition (k = 11), arthritis (k = 6), and neurological conditions (k = 2), people who had received surgery (k = 4), and older adults (k = 3) (Table 4; Figure 5). We found statistical differences between these studies (p < 0.0001). The relationship between kinesiophobia and physical activity was statistically significant only in studies that included participants with cardiovascular disease (r = -0.47; 95% CI: -0.59 to -0.32) and arthritis (r = -0.42; 95% CI: -0.69 to -0.05). The effect of arthritis remained significant when focusing on osteoarthritis (k = 5; r = -0.48; 95% CI: -0.76 to -0.35). Statistical heterogeneity was higher in the studies comprising people with arthritis (I² = 92.7%) than the studies comprising people with cardiovascular disease (I² = 83.9%).




Figure 6. Subgroup meta-analysis: Differences according to physical activity measure (self-report vs. device).

The test of subgroup differences between self-reported ($k = 32$) and device-based ($k = 9$) measures of physical activity showed a statistically significant difference ($p < 0.0001$; Table 4), with only the self-reported measures showing a correlation ($r = -0.36$; 95% CI: -0.48 to -0.24 ; Figure 6). However, we observed considerable between-study statistical heterogeneity ($I^2 = 92.9\%$). Results of the secondary meta-analysis based on Spearman rho values ($k = 5$, $n = 361$) were consistent with this subgroup analysis as they showed no statistical evidence of an association between kinesiophobia and accelerometer-based measures of physical activity ($r = -0.10$; 95% CI: -0.27 to 0.09 ; $I^2 = 30.2\%$; $p = 0.217$; Table 4; Figure 7).




Figure 7. Secondary meta-analysis based on Spearman's rho value of studies using accelerometer-based physical activity.

The test of subgroup differences according to physical activity measurement instruments was possible between studies using the IPAQ ($k = 18$), BHPAQ ($k = 3$), MLTPAQ ($k = 2$), GLTEQ ($k = 2$), as well as accelerometers ($k = 8$) and pedometers ($k = 2$). We found statistical differences between these studies ($p = 0.014$). The relationship between kinesiophobia and physical activity was statistically significant only in studies that used the IPAQ ($r = -0.43$; 95% CI: -0.57 to -0.26 ; Table 4; Figure 8). However, we observed considerable between-study statistical heterogeneity ($I^2 = 89.7\%$).

The test of subgroup differences between physical activity outcomes was possible between studies using MET-min/week, typically from the IPAQ ($k = 19$), a score from a questionnaire ($k = 11$), steps/day ($k = 4$), counts/min ($k = 3$), and active time in hours per day or week ($k = 3$) (Table 4; Figure 9). We observed statistical differences between these studies ($p < 0.0001$), with only the studies using the MET-min/week ($r = -0.42$; 95% CI: -0.56 to -0.26)




Figure 8. Subgroup meta-analysis: Differences according to physical activity measurement instruments.

Notes: 95% CI = 95% confidence interval, BFMSO = Brief Fear of Movement Scale for Osteoarthritis, FABQ = Fear-Avoidance Belief Questionnaire, FAcIS = Fear of Activity in Situations, IV = Inverse-variance method, KCS = Kinesiophobia Causes Scale, PA = physical activity, Random = Random-effects method, TSK = Tampa Scale of Kinesiophobia

and score outcome ($r = -0.33$; 95% CI: -0.56 to -0.06) showing a statistical correlation. Heterogeneity was considerable in studies relying on these outcomes ($I^2 = 90.1$ and 95.1%).

The test of subgroup differences between kinesiophobia measures was possible only for TSK-17 ($k = 21$), TSK-Heart ($k = 8$), and TSK-11 ($k = 6$) (Table 4; Figure 10), with only the former two

showing a statistical correlation. The correlation was stronger in studies using the TSK-Heart ($r = -0.59$; 95% CI: -0.75 to -0.36) than with the TSK-17 ($r = -0.28$; 95% CI: -0.43 to -0.11). Heterogeneity was lower, albeit substantial to considerable in the studies using TSK-Heart ($I^2 = 85.0\%$) than TSK-17 ($I^2 = 92.6\%$).




Figure 9. Subgroup meta-analysis: Differences according to physical activity outcome.




Figure 10. Subgroup meta-analysis: Differences according to kinesiophobia measure.




Figure 11. Meta-Regressions testing the influence of age ($k = 31$) (A), the proportion of women ($k = 33$) (B), and pain intensity ($k = 23$) (C) on the correlation estimates of the meta-analysis studies.

Meta-Regressions

Age did not statistically influence the correlation values of the meta-analysis studies ($k = 31$; $p = 0.263$) (Figure 11A). Due to the similar I^2 between this meta-regression and the main meta-analysis (92.6% vs. 91.8%, respectively), the addition of age as a predictor did not explain the considerable heterogeneity observed between study correlations. The R^2 revealed that only 1.0% of the differences between study correlations could be explained by age. Similarly, the proportion of women ($k = 33$; Figure 11B) and the mean level of pain in the studies ($k = 23$; Figure 11C) did not influence correlation values. The R^2 revealed that these variables explained less than 0.01% of the differences between study correlations.

Sensitivity Analysis

The meta-regression by AXIS score ($k = 41$) showed that a study's quality did not influence correlation values. The R^2 revealed that this variable explained less than 0.01% of the differences between study correlations. However, it should be noted that there was very little variation in axis scores with all scores ranging between 16 and 18.

Discussion

The main objective of this study was to systematically review and meta-analyze the direct relationship between kinesiophobia and physical

activity. In addition, we examined the influence of potential moderators. To our knowledge, this is the first review of its kind on this emerging research topic.

Patients with Arthritis, Cardiovascular, and Pulmonary conditions

The meta-analysis ($k = 41$, $n = 4,848$) showed a moderate negative correlation between kinesiophobia and physical activity. This result is consistent with our hypothesis and the dual models of physical activity^{23,24,26}. According to the theoretical models, this main result suggests that the fear of movement characteristic of kinesiophobia triggers an impulse to avoid physical activity behaviors, which contributes to the maintenance or exacerbation of the initial fear. Accordingly, kinesiophobia and physical inactivity would be self-perpetuating or even self-reinforcing.

Our results suggest that patients with cardiovascular or arthritic conditions may be at greater risk for this negative relationship between kinesiophobia and physical activity than those with other conditions, such as chronic pain. This latter finding was surprising because fear of pain is a key component of kinesiophobia, appearing in 10 of the 17 items on the TSK-17 and TSK-Heart scales, and reinforces the importance of considering the multidimensional nature of kinesiophobia, which also reflects fear of injury and fear of worsening a health condition. Although our results showed no evidence of an association between kinesiophobia and physical activity in other health conditions such as

cancer, acute pain, post-surgery, neurological and pulmonary conditions, these effects cannot be fully ruled out, as the lack of statistical significance could be explained by a lack of statistical power in these subgroup meta-analyses including fewer studies ($k = 1$ to 5). Of note, only one study examined the relationship between kinesiophobia and physical activity in people with a pulmonary condition (i.e., chronic obstructive pulmonary disease, COPD), but this study was the one that reported the highest correlation ($r = -0.89$; 95% CI: -0.93 to -0.82), suggesting that this population should also be closely monitored for kinesiophobia.

No Evidence Based on Objective Measures of Physical Activity

Importantly, the results showed that the negative association between kinesiophobia and physical activity was statistically significant in studies using self-report measures (e.g., IPAQ), but not in studies using device-based measures (i.e., accelerometers or pedometers). This difference persisted in analyses that further disaggregated the physical activity measures (i.e., specifying the types of questionnaires and devices) and tested the effect of the different outcomes. The secondary analysis based on Spearman's rho values further supported these results as it provided no evidence suggesting an association between kinesiophobia and device-based measures of physical activity, albeit in a small number of studies. Because device-based measures of physical activity are more valid and reliable than self-reported measures, these findings call into question the robustness of the relationship between kinesiophobia and physical activity. Therefore, additional studies relying on device-based measures are required to confirm the negative associations we observed between kinesiophobia and physical activity we observed in people with a cardiovascular, arthritis, or pulmonary condition, which were mainly based on self-reported measures.

Pain vs. Fear

Contrary to our expectations, we found no statistical evidence showing that the level of actual pain intensity at rest influenced the effect of

kinesiophobia on physical activity, despite the substantial number of studies included in this analysis ($k = 23$). This result is consistent with the weak relationship that has been shown between kinesiophobia and pain¹²¹, further suggesting that it is not the actual pain that prevents physical activity, but the fear of triggering pain, injury, or aggravating an underlying condition. However, our results also suggest that the effect of pain may be better assessed by pain history (e.g., pain duration in months) or pain intensity during exercise. The absence of significant effects in the group of studies based on shorter versions of the TSK suggests that the items removed from these versions may be important to accurately assess kinesiophobia.

Kinesiophobia Measure

Studies using the TSK-17 or the TSK-Heart to assess kinesiophobia both showed a statistically significant relationship with physical activity. This consistency is not surprising as these two measures are very similar. They differ only in specific vocabulary that makes the TSK-Heart specific to individuals with a cardiac condition (e.g., “heart problem” instead of “pain” and “injury”) and the TSK-17 more broadly applicable to different populations. In addition, the stronger correlation observed in studies using the TSK-Heart compared with the TSK-17 is consistent with our results showing stronger correlations in studies of people with a cardiovascular condition.

Limitations and Strengths

The results of this systematic review and meta-analysis should be considered in light of several limitations. Some of our results showed considerable heterogeneity, which may be explained by the diversity of the methods used to assess physical activity (questionnaires vs. accelerometers vs. pedometers), the instruments used in these methods (10 different questionnaires, 8 different accelerometers and pedometers), and the physical activity outcomes ($n = 9$), but also by the type of questionnaires used to assess kinesiophobia ($n = 9$). This heterogeneity suggests that the measures of kinesiophobia and physical activity used in the literature reflect different dimensions of these two constructs. For example, self-reported

measures of physical activity do not accurately reflect actual levels of physical activity¹²². Only articles published in English were included. Inclusion of articles published in other languages may have influenced the results. Only 11 of the 58 authors we contacted (19 %) shared their estimates or data with us, which is more than reported in previous literature¹²³. Including these missing data may have had an impact on the results. However, these limitations are counterbalanced by several strengths including the preregistration of our work, a substantial number of studies in the meta-analysis, as well as the assessment and correction for publication bias.

Conclusions

Higher levels of kinesiophobia were moderately associated with lower levels of physical activity, especially in people with a cardiovascular, arthritis, and pulmonary condition, and when self-reported physical activity was used. According to theoretical models, this relationship between kinesiophobia and physical activity results from automatic processes that may be self-reinforcing and should therefore not be overlooked. However, there was considerable heterogeneity between studies, and the lack of evidence based on objective measures of physical activity calls for cautious conclusions about this potential relationship. In sum, our results suggest that kinesiophobic patients have only a moderate, if any, risk of being more physically inactive than other patients. Therefore, more evidence is required to determine the impact kinesiophobia should have on therapeutic decisions when aiming to reduce physical inactivity¹¹. Further studies using device-based measures of physical activity are needed to confirm our findings and to understand the factors and mechanisms that influence the potential relationship between kinesiophobia and physical activity.

Article Information

Data and Code Availability

According to good research practices³⁰, the dataset, R Markdown script, and supplemental

material are freely available in Zenodo: <https://doi.org/10.5281/zenodo.8216112>¹²⁴

Authorship Contribution Statement

Based on the Contributor Roles Taxonomy (CRediT)^{125,126} individual author contributions to this work are as follows:

- **Miriam Goubran:** Conceptualization; Methodology (Systematic Review); Investigation; Writing – Original Draft; Writing – Review and Editing.
- **Ata Farajzadeh:** Investigation; Writing – Review and Editing.
- **Ian M. Lahart:** Methodology (Meta-Analysis); Formal Analysis; Visualization; Data Curation; Writing – Original Draft; Writing – Review and Editing.
- **Martin Bilodeau:** Conceptualization; Methodology (Systematic Review); Writing – Original Draft; Writing – Review and Editing; Supervision; Project Administration.
- **Matthieu P. Boisgontier:** Conceptualization (Lead); Methodology; Investigation; Formal Analysis; Data Curation; Visualization; Writing – Original Draft (Lead); Writing – Review and Editing; Supervision; Project Administration; Funding Acquisition.

Funding

Matthieu P. Boisgontier is supported by Natural Sciences and Engineering Research Council of Canada (NSERC; RGPIN-2021-03153), the Canada Foundation for Innovation (CFI), Mitacs, and the Banting Research Foundation. Ata Farajzadeh is supported by an Admission Scholarship, a Doctoral International Scholarship, and a Special Merit Scholarship from the University of Ottawa, and by a BMO Graduate Scholarship.

Conflict of Interest Disclosure

The other authors declare that they have no financial conflict of interest related to the content of this article. Matthieu P. Boisgontier is the founder, representative, and manager of Peer Community In (PCI) Health & Movement Sciences (<https://healthmovsci.peercommunityin.org/about>),

a free and transparent peer review service provided by a community of researchers who review and recommend preprints. He is a former co-chair and current member of the Society for Transparency, Openness, and Replication in Kinesiology (STORK; <https://storkinesiology.org>), current editor-in-chief for Communications in Kinesiology (<https://storkjournals.org/index.php/cik>), and associate editor for the European Rehabilitation Journal (<https://rehab-journal.com>), both of which are Diamond Open Access journals publishing articles in the field of health and rehabilitation sciences.

References

1. Morris JN, Heady JA, Raffle PA, Roberts CG, Parks JW. Coronary heart-disease and physical activity of work. *Lancet*. 1953;262(6795):1053-1057. [https://doi.org/10.1016/s0140-6736\(53\)90665-5](https://doi.org/10.1016/s0140-6736(53)90665-5)
2. Lee IM, Shiroma EJ, Lobelo F, et al. Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *Lancet*. 2012;380(9838):219-229. [https://doi.org/10.1016/S0140-6736\(12\)61031-9](https://doi.org/10.1016/S0140-6736(12)61031-9)
3. Salinas-Rodríguez A, Manrique-Espinoza B, Palazuelos-González R, Rivera-Almaraz A, Jáuregui A. Physical activity and sedentary behavior trajectories and their associations with quality of life, disability, and all-cause mortality. *Eur Rev Aging Phys Act*. 2022;19(1):13. <https://doi.org/10.1186/s11556-022-00291-3>
4. Wahid A, Manek N, Nichols M, et al. Quantifying the association between physical activity and cardiovascular disease and diabetes: a systematic review and meta-analysis. *J Am Heart Assoc*. 2016;5(9):e002495. <https://doi.org/10.1161/JAHA.115.002495>
5. Bleich SN, Vercammen KA, Zatz LY. Interventions to prevent global childhood overweight and obesity: a systematic review. *Lancet Diabetes Endocrinol*. 2018;6:332-346. [https://doi.org/10.1016/S2213-8587\(17\)30358-3](https://doi.org/10.1016/S2213-8587(17)30358-3)
6. Schuch F, Vancampfort D, Firth J, et al. Physical activity and sedentary behavior in people with major depressive disorder: a systematic review and meta-analysis. *J Affect Disord*. 2017;210:139-150. <https://doi.org/10.1016/j.jad.2016.10.050>
7. Liu X, Zhang D, Liu Y, et al. Dose-response association between physical activity and incident hypertension: a systematic review and meta-analysis of cohort studies. *Hypertension*. 2017;69(5):813-820. <https://doi.org/10.1161/HYPERTENSIONA.116.08994>
8. Moore SC, Lee IM, Weiderpass E, et al. Association of leisure-time physical activity with risk of 26 types of cancer in 1.44 million adults. *JAMA Intern Med*. 2016;176(6):816-825. <https://doi.org/10.1001/jamainternmed.2016.1548>
9. Najjar J, Östling S, Gudmundsson P, et al. Cognitive and physical activity and dementia: a 44-year longitudinal population study of women. *Neurology*. 2019;92(12):E1322-E1330. <https://doi.org/10.1212/WNL.00000000000007021>
10. World Health Organization. Physical activity - Key facts. October 5, 2022. Accessed August 3, 2023. <https://www.who.int/news-room/fact-sheets/detail/physical-activity>
11. Boisgontier MP, Iversen MD. Physical inactivity: a behavioral disorder in the physical therapist's scope of practice. *Phys Ther*. 2020;100(5):743-746. <http://doi.org/10.1093/PTJ/PZAA011>
12. O'Donoghue G, Perchoux C, Mensah K, et al. A systematic review of correlates of sedentary behaviour in adults aged 18-65 years: a socio-ecological approach. *BMC Public Health*. 2016;16(1):163. <https://doi.org/10.1186/s12889-016-2841-3>
13. Garcia L, Mendonça G, Benedetti TRB, et al. Barriers and facilitators of domain-specific physical activity: a systematic review of

- reviews. *BMC Public Health*. 2022;22(1):1964.
<https://doi.org/10.1186/s12889-022-14385-1>
14. Biddle SJH, Gorely T, Faulkner G, Mutrie N. Psychology of physical activity: a 30-year reflection on correlates, barriers, and theory. *Int J Sport Exerc Psychol*. 2023;21(1):1-14.
<https://doi.org/10.1080/1612197X.2022.2147261>
 15. Azevedo MR, Araújo CLP, Reichert FF, Siqueira FV, da Silva MC, Hallal PC. Gender differences in leisure-time physical activity. *Int J Public Health*. 2007;52(1):8-15.
<https://doi.org/10.1007/s00038-006-5062-1>
 16. Caspersen CJ, Pereira MA, Curran KM. Changes in physical activity patterns in the United States, by sex and cross-sectional age. *Med Sci Sports Exerc*. 2000;32(9):1601-1609.
<https://doi.org/10.1097/00005768-200009000-00013>
 17. Cheval B, Rebar AL, Miller MW, et al. Cognitive resources moderate the adverse impact of poor perceived neighborhood conditions on self-reported physical activity of older adults. *Prev Med*. 2019;126:105741.
<https://doi.org/10.1016/j.ypmed.2019.05.029>
 18. Cheval B, Orsholits D, Sieber S, Courvoisier D, Cullati S, Boisgontier MP. Relationship between decline in cognitive resources and physical activity. *Health Psychol*. 2020;39(6):519-528.
<https://doi.org/10.1037/hea0000857>
 19. Cheval B, Sieber S, Guessous I, et al. Effect of early- and adult-life socioeconomic circumstances on physical inactivity. *Med Sci Sports Exerc*. 2018;50(3):476-485.
<https://doi.org/10.1249/MSS.0000000000000472>
 20. Kori SH, Miller RP, Todd DD. Kinesiophobia: a new view of chronic pain behavior. *Pain Manag*. 1990;3:35-43.
 21. Vlaeyen JWS, Kole-Snijders AMJ, Boeren RGB, van Eek H. Fear of movement/(re)injury in chronic low back pain and its relation to behavioral performance. *Pain*. 1995;62(3):363-372.
[https://doi.org/10.1016/0304-3959\(94\)00279-N](https://doi.org/10.1016/0304-3959(94)00279-N)
 22. Miller R, Kori S, Todd D. The Tampa Scale: a measure of kinesiophobia. *Clin J Pain*. 1991;7(1):51-52.
 23. Brand R, Ekkekakis P. Affective-Reflective Theory of physical inactivity and exercise. *Ger J Exerc Sport Res*. 2018;48:48-58.
<https://doi.org/10.1007/s12662-017-0477-9>
 24. Conroy DE, Berry TR. Automatic affective evaluations of physical activity. *Exerc Sport Sci Rev*. 2017;45(4):230-237.
<http://doi.org/10.1249/JES.0000000000000120>
 25. Cheval B, Boisgontier MP. Promouvoir une activité physique régulière chez les patients : l'importance de la perception de l'effort. *Staps*. 2023.
 26. Cheval B, Boisgontier MP. The theory of effort minimization in physical activity. *Exerc Sport Sci Rev*. 2021;49(3):168-178.
<https://doi.org/10.1249/JES.0000000000000052>
 27. Farajzadeh A, Goubran M, Beehler A, et al. Relationship between age and physical and sedentary stimuli. *Peer Community J*. 2023;3:e21.
<https://doi.org/10.24072/pcjournal.246>
 28. Page MJ, Moher D, Bossuyt PM, et al. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ*. 2021;372:n160.
<https://doi.org/10.1136/bmj.n160>
 29. Caldwell AR, Vigotsky AD, Tenan MS, et al. Moving sport and exercise science forward: a call for the adoption of more transparent research practices. *Sports Med*. 2020;50(3):449-459.
<https://doi.org/10.1007/s40279-019-01227-1>
 30. Boisgontier MP. Research integrity requires to be aware of good and questionable research practices. *Eur Rehabil J*. 2021;2(1):1-3.
<https://doi.org/10.52057/erj.v2i1.24>
 31. Goubran M, Boisgontier M, Bilodeau M. Kinesiophobia and physical activity: a systematic review. PROSPERO (CRD42022364063) [preregistration]; October 17, 2022.
https://www.crd.york.ac.uk/prospéro/display_record.php?ID=CRD42022364063

32. Downes MJ, Brennan ML, Williams HC, Dean RS. Development of a critical appraisal tool to assess the quality of cross-sectional studies (AXIS). *BMJ Open*. 2016;6:1-7. <https://doi.org/10.1136/bmjopen-2016>
33. Schwarzer G. *meta: general package for meta-analysis* [R package]. Version 6.5-0; 2023. <https://cran.r-project.org/web/packages/meta/meta.pdf>
34. Viechtbauer W. Bias and efficiency of meta-analytic variance estimators in the random-effects model. *J Educ Behav Stat*. 2005;30(3):261-293. <https://doi.org/10.3102/10769986030003261>
35. Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med*. 2002;21(11):1539-1558. <https://doi.org/10.1002/sim.1186>
36. Knapp G, Hartung J. Improved tests for a random effects meta-regression with a single covariate. *Stat Med*. 2003;22(17):2693-2710. <https://doi.org/10.1002/sim.1482>
37. Cohen J. *Statistical power analysis for the behavioral sciences*. 2nd ed.; 1988.
38. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997;315(7109):629-634. <https://doi.org/10.1136/bmj.315.7109.629>
39. Simonsohn U, Nelson LD, Simmons JP. P-curve: A key to the file-drawer. *J Exp Psychol Gen*. 2014;143(2):534-547. <http://doi.org/10.1037/a0033242>
40. Schwarzer G, Carpenter RJ, Rücker G. *metasens: advanced statistical methods to model and adjust for bias in meta-analysis* [R package]. Version 1.5-2; 2023. <https://cran.r-project.org/web/packages/metasens/metasens.pdf>
41. R Core Team. *R: a language and environment for statistical computing* [Computer software]. Version 4.3.1. Vienna, Austria: Foundation for Statistical Computing; 2023. <https://www.r-project.org>
42. Viechtbauer W. *metafor: meta-analysis package for R* [R package]. Version 4.2-0; 2023. <https://cran.r-project.org/web/packages/metafor/metafor.pdf>
43. Viechtbauer W. Conducting meta-analyses in R with the metafor package. *J Stat Softw*. 2010;36(3):1-48. <https://doi.org/10.18637/jss.v036.i03>
44. Assadourian M, Bailly F, Letellier P, et al. Criteria for inclusion in programs of functional restoration for chronic low back pain: pragmatic study. *Ann Phys Rehabil Med*. 2020;63(3):189-194. <https://doi.org/10.1016/j.rehab.2019.06.019>
45. Baez SE, Hoch MC, Hoch JM. Psychological factors are associated with return to pre-injury levels of sport and physical activity after ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc*. 2020;28(2):495-501. <https://doi.org/10.1007/s00167-019-05696-9>
46. Luthi F, Vuistiner P, Favre C, Hilfiker R, Léger B. Avoidance, pacing, or persistence in multidisciplinary functional rehabilitation for chronic musculoskeletal pain: an observational study with cross-sectional and longitudinal analyses. *PLoS One*. 2018;13(9):e0203329. <https://doi.org/10.1371/journal.pone.0203329>
47. Bernard P, Ninot G, Bernard PL, et al. Effects of a six-month walking intervention on depression in inactive post-menopausal women: a randomized controlled trial. *Aging Ment Health*. 2015;19(6):485-492. <https://doi.org/10.1080/13607863.2014.948806>
48. Bahar Özdemir Y. Investigation of low back pain in the white-collar population working from home due to the COVID-19 pandemic. *J Phys Med Rehabil Sci*. 2021;24(2):135-142. <https://doi.org/10.31609/jpmrs.2021-81527>
49. Priore LB, Lack S, Garcia C, Azevedo FM, de Oliveira Silva D. Two weeks of wearing a knee Brace compared with minimal intervention on kinesiophobia at 2 and 6 weeks in people with patellofemoral pain: a randomized controlled trial. *Arch Phys Med Rehabil*. 2020;101(4):613-623. <https://doi.org/10.1016/j.apmr.2019.10.190>
50. Strandberg E, Bean C, Vassbakk-Svindland K, et al. Who makes it all the way?

- Participants vs. decliners, and completers vs. drop-outs, in a 6-month exercise trial during cancer treatment. Results from the Phys-Can RCT. *Support Care Cancer*. 2022;30(2):1739-1748. <https://doi.org/10.1007/s00520-021-06576-0>
51. Uritani D, Kasza J, Campbell PK, Metcalf B, Egerton T. The association between psychological characteristics and physical activity levels in people with knee osteoarthritis: a cross-sectional analysis. *BMC Musculoskelet Disord*. 2020;21(1):269. <https://doi.org/10.1186/s12891-020-03305-2>
52. Knapik A, Dąbek J, Brzęk A. Kinesiophobia as a problem in adherence to physical activity recommendations in elderly Polish patients with coronary artery disease. *Patient Prefer Adherence*. 2019;13:2129-2135. <http://doi.org/10.2147/PPA.S216196>
53. Roaldsen KS, Elfving B, Stanghelle JK, Talme T, Mattsson E. Fear-avoidance beliefs and pain as predictors for low physical activity in patients with leg ulcer. *Physiother Res Int*. 2009;14(3):167-180. <https://doi.org/10.1002/pri.433>
54. Saulicz M, Saulicz E, Knapik A, et al. Impact of physical activity and fitness on the level of kinesiophobia in women of perimenopausal age. *Prz Menopauzalny*. 2016;15(2):104-111. <https://doi.org/10.5114/pm.2016.61193>
55. Alschuler KN, Hoodin F, Murphy SL, Rice J, Geisser ME. Factors contributing to physical activity in a chronic low back pain clinical sample: a comprehensive analysis using continuous ambulatory monitoring. *Pain*. 2011;152(11):2521-2527. <https://doi.org/10.1016/j.pain.2011.07.017>
56. Altuğ F, Ünal A, Kilavuz G, Kavlak E, Çitişli V, Cavlak U. Investigation of the relationship between kinesiophobia, physical activity level and quality of life in patients with chronic low back pain. *J Back Musculoskelet Rehabil*. 2016;29(3):527-531. <https://doi.org/10.3233/BMR-150653>
57. Carvalho FA, Maher CG, Franco MR, et al. Fear of movement is not associated with objective and subjective physical activity levels in chronic nonspecific low back pain. *Arch Phys Med Rehabil*. 2017;98(1):96-104. <http://doi.org/10.1016/j.apmr.2016.09.115>
58. Demirbüken İ, Özgül B, Kuru Çolak T, Aydoğdu O, Sarı Z, Yurdalan SU. Kinesiophobia in relation to physical activity in chronic neck pain. *J Back Musculoskelet Rehabil*. 2016;29(1):41-47. <http://doi.org/10.3233/BMR-150594>
59. Elfving B, Andersson T, Grooten WJ. Low levels of physical activity in back pain patients are associated with high levels of fear-avoidance beliefs and pain catastrophizing. *Physiother Res Int*. 2007;12(1):14-24. <http://doi.org/10.1002/pri.355>
60. González de la Flor Á, García Pérez de Sevilla G, Domínguez Balmaseda D, Martín Vera D, Montero Martínez M, Del Blanco Muñoz JA. Relationship between self-efficacy and headache impact, anxiety, and physical activity levels in patients with chronic tension-type headache: an observational study. *Behav Neurol*. 2022;2022:8387249. <http://doi.org/10.1155/2022/8387249>
61. Helmus M, Schiphorst Preuper HR, Hof AL, Geertzen JH, Reneman MF. Psychological factors unrelated to activity level in patients with chronic musculoskeletal pain. *Eur J Pain*. 2012;16(8):1158-1165. <http://doi.org/10.1002/j.1532-2149.2011.00109.x>
62. Huijnen IP, Verbunt JA, Peters ML, Seelen HA. Is physical functioning influenced by activity-related pain prediction and fear of movement in patients with subacute low back pain? *Eur J Pain*. 2010;14(6):661-666. <http://doi.org/10.1016/j.ejpain.2009.10.014>
63. Koho P, Orenius T, Kautiainen H, Haanpää M, Pohjolainen T, Hurri H. Association of fear of movement and leisure-time physical activity among patients with chronic pain. *J Rehabil Med*. 2011;43(9):794-799. <http://doi.org/10.2340/16501977-0850>
64. Lotzke H, Jakobsson M, Gutke A, et al. Patients with severe low back pain exhibit a low level of physical activity before lumbar fusion surgery: a cross-sectional study. *BMC*

- Musculoskelet Disord.* 2018;19(1):365.
<http://doi.org/10.1186/s12891-018-2274-5>
65. Pazzinatto MF, Silva DO, Willy RW, Azevedo FM, Barton CJ. Fear of movement and (re)injury is associated with condition specific outcomes and health-related quality of life in women with patellofemoral pain. *Physiother Theory Pract.* 2022;38(9):1254-1263.
<http://doi.org/10.1080/09593985.2020.1830323>
66. Navarro-Ledesma S, Pruijboom L, Lluch E, Dueñas L, Mena-Del Horno S, Gonzalez-Muñoz A. The relationship between daily physical activity, psychological factors, and vegetative symptoms in women with fibromyalgia: a cross-sectional observational study. *Int J Environ Res Public Health.* 2022;19(18):11610.
<http://doi.org/10.3390/ijerph191811610>
67. Pedler A, Kamper SJ, Maujean A, Sterling M. Investigating the fear avoidance model in people with whiplash: the association between fear of movement and in vivo activity. *Clin J Pain.* 2018;34(2):130-137.
<http://doi.org/10.1097/AJP.0000000000000524>
68. Sertel M, Aydoğan Arslan S, Tütün Yümin E, Demirci CS, Tarsuslu Şimşek T. Investigation of the relationship between physical activity, kinesiophobia and fear of falling in older adults with chronic pain. *Somatosens Mot Res.* 2021;38(3):241-247.
<http://doi.org/10.1080/08990220.2021.1958774>
69. Verbunt JA, Sieben JM, Seelen HA, et al. Decline in physical activity, disability and pain-related fear in sub-acute low back pain. *Eur J Pain.* 2005;9(4):417-425.
<http://doi.org/10.1016/j.ejpain.2004.09.011>
70. Corrigan P, Cortes DH, Pontiggia L, Silbernagel KG. The degree of tendonitis is related to symptom severity and physical activity levels in patients with midportion Achilles tendinopathy. *Int J Sports Phys Ther.* 2018;13(2):196-207.
<http://doi.org/10.26603/ijsp20180196>
71. Minetama M, Kawakami M, Teraguchi M, et al. Associations between psychological factors and daily step count in patients with lumbar spinal stenosis. *Physiother Theory Pract.* 2022;38(10):1519-1527.
<http://doi.org/10.1080/09593985.2020.1855685>
72. Dąbek J, Knapik A, Gallert-Kopyto W, Brzęk AM, Piotrkowicz J, Gąsior Z. Fear of movement (kinesiophobia) - an underestimated problem in Polish patients at various stages of coronary artery disease. *Ann Agric Environ Med.* 2020;27(1):56-60.
<http://doi.org/10.26444/aaem/106143>
73. Baykal Şahin H, Kalaycıoğlu E, Şahin M. The effect of cardiac rehabilitation on kinesiophobia in patients with coronary artery disease. *Turk J Phys Med Rehabil.* 2021;67(2):203-210.
<http://doi.org/10.5606/TFTRD.2021.5164>
74. Spaderna H, Hoffman JM, Hellwig S, Brandenburg VM. Fear of physical activity, anxiety, and depression: barriers to physical activity in outpatients with heart failure? *Eur J Health Psychol.* 2020;27(1):3-13.
<http://doi.org/10.1027/2512-8442/a000042>
75. Marques-Sule E, Söderlund A, Almenar L, Espí-López GV, López-Vilella R, Bäck M. Influence on kinesiophobia by disability, physical, and behavioural variables after a heart transplantation. *Eur J Cardiovasc Nurs.* 2022;21(6):537-543.
<http://doi.org/10.1093/eurjcn/zvab134>
76. Coronado RA, Robinette PE, Henry AL, et al. Bouncing back after lumbar spine surgery: early postoperative resilience is associated with 12-month physical function, pain interference, social participation, and disability. *Spine J.* 2021;21(1):55-63.
<http://doi.org/10.1016/j.spinee.2020.07.013>
77. Donnarumma P, Presaghi F, Tarantino R, Fragale M, Rullo M, Delfini R. The impact of pelvic balance, physical activity, and fear-avoidance on the outcome after decompression and instrumented fusion for degenerative lumbar stenosis. *Eur Spine J.* 2017;26(2):428-433.
<http://doi.org/10.1007/s00586-016-4644-8>

78. Zelle DM, Corpeleijn E, Klaassen G, Schutte E, Navis G, Bakker SJL. Fear of movement and low self-efficacy are important barriers in physical activity after renal transplantation. *PLoS One*. 2016;11(2):e0147609. <http://doi.org/10.1371/journal.pone.0147609>
79. Norte GE, Solaas H, Saliba SA, Goetschius J, Slater LV, Hart JM. The relationships between kinesiophobia and clinical outcomes after ACL reconstruction differ by self-reported physical activity engagement. *Phys Ther Sport*. 2019;40:1-9. <http://doi.org/10.1016/j.pts.2019.08.002>
80. Aydemir B, Huang CH, Foucher KC. Strength and physical activity in osteoarthritis: The mediating role of kinesiophobia. *J Orthop Res*. 2022;40(5):1135-1142. <http://doi.org/10.1002/jor.25151>
81. Baday-Keskin D, Ekinçi B. The relationship between kinesiophobia and health-related quality of life in patients with rheumatoid arthritis: a controlled cross-sectional study. *Joint Bone Spine*. 2022;89(2):105275. <http://doi.org/10.1016/j.jbspin.2021.105275>
82. Kiliç H, Karahan S, Atilla B, Kinikli Gİ. Can fear of movement, depression and functional performance be a predictor of physical activity level in patients with knee osteoarthritis? *Arch Rheumatol*. 2019;34(3):274-280. <http://doi.org/10.5606/ArchRheumatol.2019.7160>
83. Aykut Selçuk M, Karakoyun A. Is there a relationship between kinesiophobia and physical activity level in patients with knee osteoarthritis? *Pain Med*. 2020;21(12):3458-3469. <http://doi.org/10.1093/pm/pnaa180>
84. Yuksel Karsli T, Bayraktar D, Ozer Kaya D, et al. Comparison of physical activity levels among different sub-types of axial spondyloarthritis patients and healthy controls. *Mod Rheumatol*. 2021;31(6):1202-1207. <http://doi.org/10.1080/14397595.2021.1891676>
85. Wasiuk-Zowada D, Brzęk A, Krzystanek E, Knapik A. Kinesiophobia in people with multiple sclerosis and its relationship with physical activity, pain and acceptance of disease. *Medicina (Kaunas)*. 2022;58(3):414. <http://doi.org/10.3390/medicina58030414>
86. Sütçü G, Ayvat E, Kiliç M. Effects of fatigue and kinesiophobia on functional capacity, physical activity and quality of life in Parkinson's disease. *Int J Rehabil Res*. 2021;44(1):65-68. <http://doi.org/10.1097/MRR.0000000000000449>
87. Ozer AY, Karaca S, Senocak E, Oğuz S, Polat MG. Does kinesiophobia limit physical activity and quality of life in asthmatic patients? *Int J Rehabil Res*. 2022;45(3):230-236. <http://doi.org/10.1097/MRR.0000000000000534>
88. Bossenbroek L, ten Hacken NH, van der Bij W, Verschuuren EA, Koëter GH, de Greef MH. Cross-sectional assessment of daily physical activity in chronic obstructive pulmonary disease lung transplant patients. *J Heart Lung Transplant*. 2009;28(2):149-155. <http://doi.org/10.1016/j.healun.2008.11.905>
89. Atıcı E, Girgin N, Çevik Saldıran T. The effects of social isolation due to COVID-19 on the fear of movement, falling, and physical activity in older people. *Australas J Ageing*. 2022;41(3):407-413. <http://doi.org/10.1111/ajag.1306390>
90. Miller L, Ohlman T, Naugle KM. Sensitivity to physical activity predicts daily activity among pain-free older adults. *Pain Med*. 2018;19(8):1683-1692. <http://doi.org/10.1093/pm/pnx251>
91. Ohlman T, Miller L, Naugle KE, Naugle KM. Physical activity levels predict exercise-induced hypoalgesia in older adults. *Med Sci Sports Exerc*. 2018;50(10):2101-2109. <http://doi.org/10.1249/MSS.0000000000001661>
92. Olsson N, Karlsson J, Eriksson BI, Brorsson A, Lundberg M, Silbernagel KG. Ability to perform a single heel-rise is significantly related to patient-reported outcome after Achilles tendon rupture. *Scand J Med Sci Sports*. 2014;24(1):152-158.

- <http://doi.org/10.1111/j.1600-0838.2012.01497.x>
93. Woby SR, Roach NK, Urmston M, Watson PJ. Psychometric properties of the TSK-11: a shortened version of the Tampa Scale for Kinesiophobia. *Pain*. 2005;117(1-2):137-144. <http://doi.org/10.1016/j.pain.2005.05.029>
 94. Jørgensen MB, Damsgård E, Holtermann A, Anke A, Søgaard K, Røe C. Properties of the Tampa Scale for Kinesiophobia across workers with different pain experiences and cultural backgrounds: a Rasch analysis. *J Appl Meas*. 2015;16(2):218-227. <http://jampress.org/abst2015.htm>
 95. Bäck M, Jansson B, Cider A, Herlitz J, Lundberg M. Validation of a questionnaire to detect kinesiophobia (fear of movement) in patients with coronary artery disease. *J Rehabil Med*. 2012;44(4):363-369. <http://doi.org/10.2340/16501977-0942>
 96. Neblett R, Hartzell MM, Mayer TG, Bradford EM, Gatchel RJ. Establishing clinically meaningful severity levels for the Tampa Scale for Kinesiophobia (TSK-13). *Eur J Pain*. 2016;20(5):701-710. <http://doi.org/10.1002/ejp.795>
 97. Knapik A, Sauliez E, Gnat R. Kinesiophobia - introducing a new diagnostic tool. *J Hum Kinet*. 2011;28:25-31. <http://doi.org/10.2478/v10078-011-0019-8>
 98. Shelby RA, Somers TJ, Keefe FJ, et al. Brief Fear of Movement Scale for osteoarthritis. *Arthritis Care Res (Hoboken)*. 2012;64(6):862-871. <http://doi.org/10.1002/acr.21626>
 99. Waddell G, Newton M, Henderson I, Somerville D, Main CJ. A Fear-Avoidance Beliefs Questionnaire (FABQ) and the role of fear-avoidance beliefs in chronic low back pain and disability. *Pain*. 1993;52(2):157-168. [http://doi.org/10.1016/0304-3959\(93\)90127-B](http://doi.org/10.1016/0304-3959(93)90127-B)
 100. Craig CL, Marshall AL, Sjöström M, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc*. 2003;35(8):1381-1395. <http://doi.org/10.1249/01.MSS.0000078924.61453.FB>
 101. Baecke JA, Burema J, Frijters JE. A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *Am J Clin Nutr*. 1982;36(5):936-942. <http://doi.org/10.1093/ajcn/36.5.936>
 102. Grimby G, Börjesson M, Jonsdottir IH, Schnohr P, Thelle DS, Saltin B. The "Saltin-Grimby Physical Activity Level Scale" and its application to health research. *Scand J Med Sci Sports*. 2015;25 Suppl 4:119-125. <http://doi.org/10.1111/sms.12611>
 103. Godin G, Shephard RJ. A simple method to assess exercise behavior in the community. *Can J Appl Sport Sci*. 1985;10(3):141-146.
 104. Taylor HL, Jacobs DR Jr, Schucker B, Knudsen J, Leon AS, Debacker G. A questionnaire for the assessment of leisure time physical activities. *J Chronic Dis*. 1978;31(12):741-755. [http://doi.org/10.1016/0021-9681\(78\)90058-9](http://doi.org/10.1016/0021-9681(78)90058-9)
 106. Voorrips LE, Ravelli AC, Dongelmans PC, Deurenberg P, Van Staveren WA. A physical activity questionnaire for the elderly. *Med Sci Sports Exerc*. 1991;23(8):974-979.
 107. Wendel-Vos GC, Schuit AJ, Saris WH, Kromhout D. Reproducibility and relative validity of the short questionnaire to assess health-enhancing physical activity. *J Clin Epidemiol*. 2003;56(12):1163-1169. [http://doi.org/10.1016/S0895-4356\(03\)00220-8](http://doi.org/10.1016/S0895-4356(03)00220-8)
 108. Tegner Y, Lysholm J. Rating systems in the evaluation of knee ligament injuries. *Clin Orthop Relat Res*. 1985;198:43-49. <http://doi.org/10.1097/00003086-198509000-00007>
 109. Zahiri CA, Schmalzried TP, Szuszczewicz ES, Amstutz HC. Assessing activity in joint replacement patients. *J Arthroplasty*. 1998;13(8):890-895. [http://doi.org/10.1016/s0883-5403\(98\)90195-4](http://doi.org/10.1016/s0883-5403(98)90195-4)
 110. Melzack R, Katz J. The McGill Pain Questionnaire: appraisal and current status. In: Turk DC, Melzack R, eds. *Handbook of Pain Assessment*. 2nd ed. The Guilford Press; 2001:35-52.

111. Ross RLP. Clinical assessment of pain. In: van Dieen JH, ed. *Assessment in Occupational Therapy and Physical Therapy*. WB Saunders; 1997:123-133.
112. Roos EM, Roos HP, Lohmander LS, Ekdahl C, Beynnon BD. Knee Injury and Osteoarthritis Outcome Score (KOOS)—development of a self-administered outcome measure. *J Orthop Sports Phys Ther*. 1998;28(2):88-96. <http://doi.org/10.2519/jospt.1998.28.2.88>
113. Cleeland CS, Ryan KM. Pain assessment: global use of the Brief Pain Inventory. *Ann Acad Med Singap*. 1994;23(2):129-138.
114. Murray DW, Fitzpatrick R, Rogers K, et al. The use of the Oxford hip and knee scores. *J Bone Joint Surg Br*. 2007;89(8):1010-1014. <http://doi.org/10.1302/0301-620X.89B8.19424>
115. Bush JW, Kaplan RM. The quality of well-being scale (formerly the index of well-being). In: McDowell I, ed. *Measuring Health: A Guide to Rating Scales and Questionnaires*. 3rd ed. Oxford University Press; 2006:675-683.
116. Ware JE Jr. SF-36 health survey update. *Spine (Phila Pa 1976)*. 2000;25(24):3130-3139. <http://doi.org/10.1097/00007632-200012150-00008>
117. Freyd M. The Graphic Rating Scale. *J Educ Psychol*. 1923;14:83-102. <https://doi.org/10.1037/h0074329>
118. Melzack R, Torgerson WS. On the language of pain. *Anesthesiology*. 1971;34(1):50-59. <http://doi.org/10.1097/00000542-197101000-00017>
119. Page MJ, Sterne JAC, Higgins JPT, Egger M. Investigating and dealing with publication bias and other reporting biases in meta-analyses of health research: a review. *Res Synth Methods*. 2021;12(2):248-259. <http://doi.org/10.1002/jrsm.1468>
120. Simonsohn U, Simmons JP, Nelson LD. Better P-curves: making P-curve analysis more robust to errors, fraud, and ambitious P-hacking, a reply to Ulrich and Miller (2015). *J Exp Psychol Gen*. 2015;144(6):1146-1152. <http://doi.org/10.1037/xge0000104>
121. Rethman KK, Mansfield C, Moeller J, et al. Kinesiophobia is associated with poor function and modifiable through interventions in people with patellofemoral pain: a systematic review with individual participant data correlation meta-analysis. *Phys Ther*. 2023. <http://doi.org/10.1093/ptj/pzad074>
122. Prince SA, Adamo KB, Hamel ME, Hardt J, Connor Gorber S, Tremblay M. A comparison of direct versus self-report measures for assessing physical activity in adults: a systematic review. *Int J Behav Nutr Phys Act*. 2008;5:56. <http://doi.org/10.1186/1479-5868-5-56>
123. Gabelica M, Bojčić R, Puljak L. Many researchers were not compliant with their published data sharing statement: a mixed-methods study. *J Clin Epidemiol*. 2022;150:33-41. <http://doi.org/10.1016/j.jclinepi.2022.05.019>
124. Goubran M, Farajzadeh A, Lahart IM, Bilodeau M, Boigontier MP. Kinesiophobia and physical activity: a systematic review and meta-analysis [Data set, Rmd script, supplementary material]. Version 1.0; 2023. <https://doi.org/10.5281/zenodo.8216112>
125. Allen L, O'Connell A, Kiermer V. How can we ensure visibility and diversity in research contributions? How the contributor role taxonomy (CRediT) is helping the shift from authorship to contributorship. *Learn Publ*. 2019;32(1):71-74. <https://doi.org/http://doi.org/10.1002/leap.1210>
126. Brand A, Allen L, Altman M, Hlava M, Scott J. Beyond authorship: attribution, contribution, collaboration, and credit. *Learn Publ*. 2015;28(2):151-155. <http://doi.org/10.1087/20150211>