

# **Fractional flow reserve versus angiography for revascularization in patients with coronary artery disease: A systematic review and meta-analysis of randomized controlled trials**

Sunny Goel, MD<sup>\*1,2</sup>, Sharon Slomovich, MD<sup>\*1,3</sup>, Sami Edris, MD<sup>1</sup>, Gal Rubinstein, MD<sup>3,4</sup>, Chirag Agarwal, MD<sup>5</sup>, Won Jun Park, MD<sup>1</sup>, Daniel Zinkovsky, MD<sup>1</sup>, Amit Hooda MD<sup>2</sup>, Parasuram Melarcode Krishnamoorthy MD<sup>2</sup>, Umesh K. Gidwani, MD<sup>2</sup>, Samin K. Sharma, MD<sup>2</sup>, Annapoorna Kini, MD<sup>2</sup>

<sup>1</sup>Division of Cardiology, Department of Medicine, Mount Sinai South Nassau, Oceanside, NY

<sup>2</sup>Division of Cardiology, Department of Medicine, Mount Sinai Hospital, New York, NY

<sup>3</sup>Division of Cardiology, Columbia University Irving Medical Center, New York, NY

<sup>4</sup>Department of Internal Medicine, Jacobi Medical Center, Albert Einstein College of Medicine, Bronx, NY

<sup>5</sup>Maimonides Medical Center, Department of Cardiology, Brooklyn, NY

\*Both authors contributed equally to the manuscript

**Short title:** FFR versus angiography in coronary artery disease

## **Corresponding author's contact information:**

Sharon Slomovich, MD  
Mount Sinai South Nassau  
One Healthy Way  
Oceanside, NY 11572  
Sharon.Slomovich@snch.org

**Manuscript word count:** 4,798

## ABSTRACT

**Background:** Fractional flow reserve (FFR) is routinely used to assess the ischemic potential of a coronary artery lesion. However, recently published randomized control trials have questioned the advantage of FFR over angiography to guide revascularization. Whether FFR guided revascularization provides clinical benefit over angiography remains unclear.

**Methods:** We performed a meta-analysis in patients with stable coronary artery disease (CAD), acute coronary syndrome (ACS), multivessel or single vessel CAD undergoing revascularization comparing FFR versus angiography to guide revascularization. Randomized control trials comparing FFR versus angiography guided revascularization were searched through PubMed, Cumulative Index to Nursing and Allied Health Literature, Cochrane Central, Scopus, Google Scholar, and Web of Science databases. The primary endpoints included cardiovascular mortality, repeat revascularization, myocardial infarction, major adverse cardiac events, stroke or transient ischemic attack and target lesion revascularization. We also evaluated the procedural outcomes including the average number of stents used between the two groups, procedure time and contrast volume used. Event rates were compared using a forest plot of odds ratios using a random-effects model assuming interstudy heterogeneity.

**Results:** The meta-analysis included 13 trials in which 7415 patients met the eligibility criteria. There was no significant difference between the FFR versus angiography guided revascularization groups across all clinical measures including all-cause mortality (OR = 1.06, 95% CI = 0.74-1.53, P = 0.74, I<sup>2</sup> = 27%), cardiovascular mortality (OR = 0.81, 95% CI = 0.43-1.52, P = 0.51, I<sup>2</sup> = 44%), repeat revascularization (OR =

1.02, 95% CI = 0.83-1.26,  $P = 0.83$ ,  $I^2 = 17\%$ ), myocardial infarction (OR = 0.92, 95% CI = 0.69-1.21,  $P = 0.54$ ,  $I^2 = 36\%$ ), major adverse cardiac event (OR = 0.82, 95% CI = 0.62-1.08,  $P = 0.15$ ,  $I^2 = 41\%$ ), stroke or transient ischemic attack (OR = 1.49, 95% CI = 0.87-2.55,  $P = 0.15$ ,  $I^2 = 0\%$ ) and target lesion revascularization (OR = 0.86, 95% CI = 0.44-1.69,  $P = 0.67$ ,  $I^2 = 0\%$ ). A sensitivity analysis was performed for studies that included patients exclusively with an ACS and studies that used FFR coronary artery bypass grafting (CABG) as a revascularization strategy. There was no difference in any of the clinical outcomes between the two groups in the sensitivity analysis. In terms of procedural outcomes, the average number of stents used was lower in the FFR group as compared to the angiography group, mean difference (MD) of -0.79 (95% CI = - 1.10, - 0.48),  $P < 0.00001$ ) with no difference in procedure time or contrast volume used.

**Conclusion:** This meta-analysis suggests that FFR when used in conjunction with angiography prevents unnecessary PCI without any difference in clinical outcomes between the two groups.

## INTRODUCTION

In patients with coronary artery disease (CAD), localizing lesions responsible for reversible ischemia can be challenging. Identifying these physiologically significant lesions by coronary angiography, particularly those of indeterminate significance (40-70% diameter narrowing) is subject to limitations due to inter-operator variability and inaccuracies in visual interpretation<sup>1-4</sup>. Fractional flow reserve (FFR) is routinely used to estimate the functional impact of these intermediate coronary stenosis and to guide percutaneous intervention in everyday practice<sup>3,5</sup>.

Fractional flow reserve is a diagnostic technique that measures pressure differences proximal and distal to a stenotic coronary lesion at maximal flow, indicating the degree of stenosis. FFR is utilized during coronary angiography to guide revascularization by assessing the ischemic potential of a coronary artery lesion<sup>4,6,7</sup>. An FFR index of  $\leq 0.80$  identifies a coronary lesion causing ischemia with >90% accuracy. Clinical trials have demonstrated that FFR-guided angiography can prevent unnecessary percutaneous intervention and improve clinical outcomes<sup>7-11</sup>.

The advantages of FFR over angiography to guide revascularization have been well established and FFR has remained the clinical standard in guiding clinical decision-making for lesions with questionable hemodynamic impact<sup>3,10,12</sup>. However, recently published randomized controlled trials have questioned the clinical benefits of using FFR over angiography to guide coronary artery revascularization in patients with multi-vessel CAD, especially in patients with acute coronary syndromes<sup>13-16</sup>. We, therefore, conducted a meta-analysis of all randomized control trials to compare the clinical outcomes in patients with stable CAD, acute coronary syndrome, single vessel CAD

and multivessel disease who underwent FFR versus angiography-guided revascularization either with PCI or CABG.

# **MATERIALS AND METHODS**

## **Study Design**

A systematic review of the literature was performed according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement<sup>17</sup>.

## **Data sources and search strategy**

We conducted a systematic search of electronic databases including PubMed, Cumulative Index to Nursing and Allied Health Literature (CINAHL), Cochrane Central, Scopus, Google Scholar, and Web of Science databases for all randomized control trials (RCT) that compared FFR versus angiography for patients with CAD. All relevant combinations of keywords relating to FFR and angiography for CAD were searched: “fractional flow reserve”, “angiography”, “revascularization,” “multi-vessel coronary artery disease” and “randomized control trial.” The search was conducted from the inception of these databases to December 31st, 2022. No language or age restrictions were applied. Pertinent trials were also searched on <http://www.clinicaltrials.gov> and in the proceedings of major international cardiology meetings (American College of Cardiology, American Heart Association, European Society of Cardiology, and Transcatheter Cardiovascular Therapeutics [TCT]). We also manually searched the references of these articles to find additional relevant articles. Two independent reviewers SG and SS conducted the search. This meta-analysis was registered at PROSPERO (CRD42022301144). This study was exempted from Institutional Review Board approval, as it was a study-level meta-analysis.

## Study Selection

We included in our meta-analysis all studies that met the following criteria: (1) randomized control trial (2) study on human subjects with participants of any age (3) study reporting clinical outcomes between FFR versus angiography guided revascularization either with PCI or CABG (4) full-length article in a peer-reviewed journal.

In addition, we also included the FAME 3 trial, one of the largest clinical trials evaluating the role of physiological assessment of coronary artery lesions to provide additional strength to our analysis. It compared FFR guided percutaneous intervention (PCI) versus angiography guided CABG in patients with 3 vessel CAD<sup>18</sup>. Although PCI and CABG are different means of revascularization, we wanted to evaluate the totality of the evidence. CABG is routinely used to revascularize patients with multivessel CAD and FFR is employed during the pre-CABG diagnostic angiography to determine hemodynamic significance of intermediate coronary artery lesions. Separate sensitivity analyses were performed to exclude the FAME 3 trial (Figure S1), studies with ACS patients exclusively (Figure S2) and the studies that used FFR guided CABG as a revascularization strategy (Figure S3). The longest follow-up data available for the included studies was used in the final analysis.

Several trials that randomized patients to culprit versus non-culprit lesions, with subsequent FFR versus angiography guided revascularization, such as the COMPLETE and CVLPRIT trials, were excluded from our analysis as they did not strictly randomize patients to FFR versus angiography<sup>19,20</sup>. FFR was left at the operator's discretion in these trials. The recently published RIPCORDER-2 trial, and FRAME AMI (acute

myocardial infarction) trials are also included in our analysis<sup>16,21</sup>. For our analysis, the events rates were calculated only for the patients with angiographic obstructive coronary disease in accordance with previous trials<sup>16</sup>.

## **Data Extraction**

Reviewers (SG and SS) independently screened the titles and abstracts that met the inclusion criteria. Discrepancies between reviewers were discussed until a consensus was reached. Using the above-mentioned selection criteria, these two reviewers independently determined which articles were to be included and excluded, and the data from the relevant articles were extracted using predefined extraction forms. Any disagreements in data extraction were discussed until a consensus was reached. Bibliographies of relevant publications were hand-searched to attempt complete inclusion of all studies of interest.

## **Study endpoints**

The clinical endpoints included were all-cause mortality, cardiovascular mortality, repeat revascularization including target vessel revascularization, myocardial infarction (MI), major adverse cardiac event (MACE), stroke or transient ischemic attack (TIA), and target lesion revascularization (TLR). The definition of outcomes as described in the trials are defined in Table S1. The procedural outcomes evaluated were the average number of stents used, total contrast volume and procedure time.

## **Data analysis**

To analyze the data, we used Review Manager Software (Rev Man, version 5.3). Analysis was performed on an intention-to-treat basis. Data were summarized across

treatment arms using the Mantel– Haenszel odds ratio (OR) using a random-effects model. Heterogeneity between studies was assessed using the Cochrane Q test and  $I^2$  statistics, which denotes the percentage of total variation across studies that is a result of heterogeneity rather than chance. Heterogeneity was considered significant if the p-value was  $<0.05$ . Sensitivity analyses were performed as described in the study selection.

### **Risk of bias assessment**

The risk of the bias for the included trials was evaluated using the Cochrane risk assessment tool for bias, which comprises 7 criteria: random sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessment, incomplete outcome data, selective reporting, and other sources of bias. Based on the fulfillment of these criteria, studies were classified as low risk, unclear risk, or high risk for bias. Publication bias was evaluated using the Egger test and funnel plots were created (Figure S4).

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

## RESULTS:

Figure 1 shows the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram describing the search strategy. The initial search yielded 2,163 abstracts of which 1,849 were excluded based on the title and reading of the abstract. Six hundred and seventy-five articles were reviewed out of which 361 were excluded as they were either non-randomized studies or had no comparison group.

Table 1 outlines the included studies for the final analysis. A total of thirteen RCTs met the inclusion criteria. The weighted mean follow-up period was 22 months (1.8 years) with a total of 7415 patients (FFR group = 3719, angiography group = 3714).

The baseline characteristics were similar between both groups (Table 1). The mean age of patients in the angiography group was  $64.4 \pm 2.8$  years, whereas it was  $64.8 \pm 2.2$  years in the FFR group. Males comprised 77.4% of patients in the angiography group compared to 77.9% of the patients in the FFR group. A history of diabetes mellitus was present in 23.8% of the patients in the angiography group and 24.4% of the patients in the FFR group. The prevalence of hypertension was 59.7% and 58.7% in the angiography and FFR groups respectively. Of the patients in the angiography group, 58.3 % and 57.3% of patients in the FFR group had a history of hyperlipidemia. In the angiography group, 33.9% of patients were current smokers compared to 33.0% of those in the FFR group. Previous MI was noted in 18.0% of patients in the angiography group and 21.6% of patients in the FFR group (Table 2).

## STUDY OUTCOMES

There was no significant difference between FFR versus angiography across all clinical outcomes; all-cause mortality (OR = 1.06, 95% CI = 0.74-1.53, P = 0.74, I<sup>2</sup>= 27%)

[Figure 2], cardiac mortality (OR = 0.81, 95% CI = 0.43-1.52, P = 0.51, I<sup>2</sup> = 44%) [Figure 2], repeat revascularization (OR = 1.02, 95% CI = 0.83-1.26, P = 0.83, I<sup>2</sup> = 17%) [Figure 2], target lesion revascularization (OR = 0.86, 95% CI = 0.44-1.69, P = 0.67, I<sup>2</sup> = 0%) [Figure 2], major adverse cardiac events [MACE] (OR = 0.82, 95% CI = 0.62-1.08, P = 0.15, I<sup>2</sup> = 41%) [Figure 2], myocardial infarction (OR = 0.92, 95% CI = 0.69-1.21, P = 0.54, I<sup>2</sup> = 36%) [Figure 2], incidence of Stroke or TIA (OR = 1.49, 95% CI = 0.87-2.55, P = 0.15, I<sup>2</sup> = 0%) [Figure 2]. In terms of procedural outcomes, the average number of stents used was lower in the FFR group as compared to the angiography group, MD of -0.79 (95% CI = -1.10, -0.48), P < 0.00001, with no difference in procedure time or contrast volume used [Figure 3].

A sensitivity analysis was performed for studies with ACS patients exclusively and the studies that used FFR guided CABG as a revascularization strategy. There was no statistically significant difference between the two groups (Figure S2 and S3).

## DISCUSSION

The main finding of our analysis is that there is no significant difference in clinical outcomes (all-cause mortality, cardiovascular mortality, repeat revascularization, MI, MACE, Stroke/TIA, TLR) in patients undergoing FFR versus conventional angiography guided revascularization at the mean follow-up period of 1.8 years. FFR does however reduce the average number of stents used, with no difference in the contrast volume used and procedure times.

FFR is currently utilized as the gold standard for determining the hemodynamic influence of an intermediate coronary lesion. FFR-guided PCI has been shown to reduce unnecessary revascularization and improve clinical outcomes in both single and multivessel CAD<sup>3,11,12</sup>. The FAME trial was one of the earliest multicenter, randomized control trials to demonstrate favorable clinical outcomes of FFR as compared to angiography to guide PCI in multivessel CAD<sup>9</sup>. In this cohort, there were a significantly higher number of stents placed in the angiography-guided group as compared to the FFR-guided group with a significant decrease in major adverse cardiac events in the FFR group in the first two years<sup>11</sup>. Several RCTs followed with encouraging results in patients with both acute coronary syndromes (ACS) and stable CAD<sup>10,11,22–24</sup>. However, recent studies have emerged questioning the role of FFR versus angiography to guide revascularization. In the FUTURE trial, there was no evidence to support that the FFR-guided treatment strategy reduced the risk of ischemic cardiovascular events or death at 1-year-follow-up<sup>15</sup>. Of note, the Data Safety and Monitoring Board of the FUTURE trial recommended early discontinuation of the trial for safety concerns as the all-cause mortality was increased in the FFR group as compared to the control group (4.3%

versus 1.8%;  $P = 0.038$ ). However, this difference in death was not statistically significant after the collection of all data and the final analysis ( $P = 0.06$ ). Similarly, the recently published RIPCORDER-2 trial showed no difference in clinical outcomes between the strategy of systematic FFR assessment compared with angiography<sup>16</sup>.

In the FAMOUS-NSTEMI trial, there were no statistically significant differences in clinical outcomes between the FFR guided revascularization versus routine angiography guided revascularization of non-culprit vessels. Patients underwent less revascularization in the FFR-guided group at 12 months [79.0 versus 86.8%, difference 7.8% (20.2%, 15.8%),  $P=0.054$ ]<sup>10</sup>. In the FLOWER-MI trial, patients with ST-elevation myocardial infarction (STEMI) that underwent complete revascularization, the FFR guided group showed no significant benefit over an angiography, regarding the risk of death, MI, or urgent revascularization at 1 year for non-culprit vessels<sup>14</sup>.

In addition, FFR versus angiographic evaluation to guide CABG showed no impact on graft patency rate in two small-sized randomized clinical trials. In the FARGO trial, at 6 months, for 72 patients in the cohort, similar graft failure rates (16% versus 12%;  $p = 0.97$ ) and clinical outcomes (rates of death, MI, and stroke) were seen in both the FFR and angiography guided groups<sup>25</sup>. Similarly in the GRAFFITI trial, no difference was seen in graft patency at one year<sup>13</sup>.

FAME 3 trial was one of the largest clinical trials comparing the noninferiority of FFR guided PCI using the latest-generation stents to conventional angiography guided CABG for patients with three-vessel disease. The results showed no difference in clinical outcomes at 1 year between the two groups<sup>18</sup>.

Prior meta-analyses investigating clinical outcomes in FFR versus angiography-guided PCI are outdated and lack current evidence. Furthermore, these analyses include non-randomized and observational studies which confer more bias than randomized control trials<sup>26</sup>. Our meta-analysis includes a comprehensive overview of the role of FFR versus angiography-guided revascularization in both patients with stable and unstable CAD as studied in randomized control trials. Also, we included patients undergoing FFR guided CABG. A recent meta-analysis investigated the role of FFR versus angiography to guide revascularization in patients with STEMI and multivessel disease for non-culprit lesions. However, patients were not randomized to FFR or angiography to guide PCI (apart from in FLOWER-MI), and thus there is no direct comparison between angiography and FFR-guided revascularization of non-culprit lesions<sup>27</sup>.

Our meta-analysis reflects the real-world population with a mix of patients with stable CAD, ACS, single vessel CAD as well as multivessel disease who are deemed candidates for revascularization with PCI and/or CABG. Our analysis shows that FFR provides no additional clinical benefit over conventional angiography. However, FFR can help reduce the number of stents used, prevent unnecessary revascularization procedures and is therefore more cost-effective when compared with angiography<sup>28</sup>.

## **LIMITATIONS**

There are several limitations in our study. First, there was no patient-level data available for our analysis. Additionally, there is heterogeneity in study designs, some studies compared FFR versus angiography guided PCI whereas other studies compared FFR versus angiography guided revascularization (PCI and CABG). However, sensitivity analysis was similar across all groups. FFR cutoff values also varied between studies

and ranged from less than 0.75 to less than or equal to 0.80. Specifically, the RIPCORD study evaluated lesions <50% which maybe stretching the limit of FFR as there is no expected benefit of its use in these small lesions. The significant angiographic stenosis was based on operator discretion and ranged between 30% and 70%. Definitions of MACE were variable between studies. The duration of follow-up time also differed amongst the studies. Finally, none of the studies looked at patients exclusively with stable CAD.

## **CONCLUSION**

There was no difference in clinical outcomes in patients undergoing FFR-guided versus angiography-guided revascularization for patients with CAD.

## **ACKNOWLEDGEMENT, SOURCES OF FUNDING & DISCLOSURES**

**a) Acknowledgments:** None

**b) Funding/Support:** None

**c) Conflict of Interest Disclosures:** No disclosures for the manuscript

### **Supplemental Materials:**

Tables S1

Figures S1-S4

**Registration information:** This meta-analysis was registered at PROSPERO (CRD42022301144). A review protocol was not prepared.

## REFERENCES

1. Fischer JJ, Samady H, McPherson JA, et al. Comparison between visual assessment and quantitative angiography versus fractional flow reserve for native coronary narrowings of moderate severity. *Am J Cardiol.* 2002;90(3):210-215. doi:10.1016/S0002-9149(02)02456-6
2. Kern MJ, Lerman A, Bech JW, et al. Physiological assessment of coronary artery disease in the cardiac catheterization laboratory: A scientific statement from the American Heart Association Committee on diagnostic and interventional cardiac catheterization, council on clinical cardiology. *Circulation.* 2006;114(12):1321-1341. doi:10.1161/CIRCULATIONAHA.106.177276
3. Tonino P AL, De Bruyne B, Pijls NH, et al. *Fractional Flow Reserve versus Angiography for Guiding Percutaneous Coronary Intervention.* Vol 360.; 2009.
4. Pijls NHJ, Fearon WF, Tonino PAL, et al. Fractional flow reserve versus angiography for guiding percutaneous coronary intervention in patients with multivessel coronary artery disease: 2-Year follow-up of the FAME (fractional flow reserve versus angiography for multivessel evaluation) study. *J Am Coll Cardiol.* 2010;56(3):177-184. doi:10.1016/j.jacc.2010.04.012
5. Chowdhury M, Osborn EA. Physiological Assessment of Coronary Lesions in 2020. *Curr Treat Options Cardiovasc Med.* 2020;22(1):2. doi:10.1007/S11936-020-0803-7
6. Pijls NHJ, Van Gelder B, Van Der Voort P, et al. Fractional Flow Reserve. *Circulation.* 1995;92(11):3183-3193. doi:10.1161/01.CIR.92.11.3183
7. Pijls NHJ, de Bruyne B, Peels K, et al. Measurement of fractional flow reserve to assess the functional severity of coronary-artery stenoses. *N Engl J Med.* 1996;334(26):1703-1708. doi:10.1056/NEJM199606273342604
8. Pijls NHJ, Van Gelder B, Van Der Voort P, et al. Fractional flow reserve. A useful index to evaluate the influence of an epicardial coronary stenosis on myocardial blood flow. *Circulation.* 1995;92(11):3183-3193. doi:10.1161/01.CIR.92.11.3183
9. Tonino PAL, Fearon WF, De Bruyne B, et al. Angiographic versus functional severity of coronary artery stenoses in the FAME study fractional flow reserve versus angiography in multivessel evaluation. *J Am Coll Cardiol.* 2010;55(25):2816-2821. doi:10.1016/J.JACC.2009.11.096
10. Layland J, Oldroyd KG, Curzen N, et al. Fractional flow reserve vs. angiography in guiding management to optimize outcomes in non-ST-segment elevation myocardial infarction: the British Heart Foundation FAMOUS-NSTEMI randomized trial. *Eur Heart J.* 2015;36(2):100-111. doi:10.1093/EURHEARTJ/EHU338
11. Van Nunen LX, Zimmermann FM, Tonino PAL, et al. Fractional flow reserve versus angiography for guidance of PCI in patients with multivessel coronary artery disease (FAME): 5-year follow-up of a randomised controlled trial. *Lancet.*

- 2015;386(10006):1853-1860. doi:10.1016/S0140-6736(15)00057-4
12. Bech GJW, De Bruyne B, Pijls NHJ, et al. Fractional flow reserve to determine the appropriateness of angioplasty in moderate coronary stenosis: a randomized trial. *Circulation*. 2001;103(24):2928-2934. doi:10.1161/01.CIR.103.24.2928
13. Toth GG, de Bruyne B, Kala P, et al. Graft patency after FFR-guided versus angiography-guided coronary artery bypass grafting: The GRAFFITI trial. *EuroIntervention*. 2019;15(11):E999-E1005. doi:10.4244/EIJ-D-19-00463
14. Puymirat E, Cayla G, Simon T, et al. Multivessel PCI Guided by FFR or Angiography for Myocardial Infarction. *N Engl J Med*. 2021;385(4):297-308. doi:10.1056/nejmoa2104650
15. Rioufol G, Dérimey F, Roubille F, et al. Fractional Flow Reserve to Guide Treatment of Patients With Multivessel Coronary Artery Disease. *J Am Coll Cardiol*. 2021;78(19):1875-1885. doi:10.1016/j.jacc.2021.08.061
16. Stables RH, Mullen LJ, Elguindy M, et al. Routine Pressure Wire Assessment Versus Conventional Angiography in the Management of Patients With Coronary Artery Disease: The RIPCARD 2 Trial. *Circulation*. 2022;146(9):687-698. doi:10.1161/CIRCULATIONAHA.121.057793
17. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *Int J Surg*. 2010;8(5):336-341. doi:10.1016/j.ijsu.2010.02.007
18. Fearon WF, Zimmermann FM, De Bruyne B, et al. Fractional Flow Reserve–Guided PCI as Compared with Coronary Bypass Surgery. *N Engl J Med*. Published online November 4, 2021. doi:10.1056/nejmoa2112299
19. Mehta SR, Wood DA, Storey RF, et al. Complete Revascularization with Multivessel PCI for Myocardial Infarction. *N Engl J Med*. 2019;381(15):1411-1421. doi:10.1056/NEJMOA1907775/SUPPL\_FILE/NEJMOA1907775\_DATA-SHARING.PDF
20. Gershlick AH, Khan JN, Kelly DJ, et al. Randomized Trial of Complete Versus Lesion-Only Revascularization in Patients Undergoing Primary Percutaneous Coronary Intervention for STEMI and Multivessel Disease: The CvLPRIT Trial. *J Am Coll Cardiol*. 2015;65(10):963-972. doi:10.1016/J.JACC.2014.12.038
21. Lee JM, Kim HK, Park KH, et al. Fractional flow reserve versus angiography-guided strategy in acute myocardial infarction with multivessel disease: a randomized trial. *Eur Heart J*. Published online December 20, 2022. doi:10.1093/EURHEARTJ/EHAC763
22. Zhang Z, Li K, Tian J. Efficacy and safety outcomes of fractional flow reserve in guiding clinical therapy of non-ST-segment elevation myocardial infarction compared with angiography alone in elderly Chinese patients. *Clin Interv Aging*. 2016;11:1751-1754. doi:10.2147/CIA.S123735
23. Quintella EF, Ferreira E, Azevedo VMP, et al. Clinical outcomes and cost-

- effectiveness analysis of FFR compared with angiography in multivessel disease patient. *Arq Bras Cardiol.* 2019;112(1):40-47. doi:10.5935/abc.20180262
24. Park SH, Jeon KH, Lee JM, et al. Long-Term Clinical Outcomes of Fractional Flow Reserve-Guided Versus Routine Drug-Eluting Stent Implantation in Patients with Intermediate Coronary Stenosis: Five-Year Clinical Outcomes of DEFER-DES Trial. *Circ Cardiovasc Interv.* 2015;8(12). doi:10.1161/CIRCINTERVENTIONS.115.002442
  25. Thuesen AL, Riber LP, Veien KT, et al. Fractional Flow Reserve Versus Angiographically-Guided Coronary Artery Bypass Grafting. *J Am Coll Cardiol.* 2018;72(22):2732-2743. doi:10.1016/J.JACC.2018.09.043
  26. Enezate T, Omran J, Al-Dadah AS, et al. Fractional flow reserve versus angiography guided percutaneous coronary intervention: An updated systematic review. *Catheter Cardiovasc Interv.* 2018;92(1):18-27. doi:10.1002/CCD.27302
  27. Baine KR, Engström T, Smits PC, et al. Complete vs Culprit-Lesion-Only Revascularization for ST-Segment Elevation Myocardial Infarction: A Systematic Review and Meta-analysis. *JAMA Cardiol.* 2020;5(8):1. doi:10.1001/JAMACARDIO.2020.1251
  28. Hong D, Kim H, Lee H, et al. Long-Term Cost-Effectiveness of Fractional Flow Reserve–Based Percutaneous Coronary Intervention in Stable and Unstable Angina. *JACC Adv.* 2022;1(5):100145. doi:10.1016/j.jacadv.2022.100145

**Table 1.** Baseline Characteristics

Study	Trial Name	Year	Analysis	Population	Follow-up time, y	FFR Cutoff	No.	Group	Mean age, y	Male, No. (%)
Chen et al. <sup>28</sup>	DKCRUSH-VI	2015	FFR vs. ANGIO-guided PCI	ACS, Stable CAD	1	<0.80	160	ANGIO	65.4±9.2	116 (72.5)
							160	FFR	65.2±9.6	121 (75.6)
Park et al. <sup>24</sup>	DEFER-DES	2015	FFR vs. ANGIO-guided PCI	ACS, Stable CAD	1	<0.75	115	ANGIO	63±10	87 (75)
							114	FFR	62±10	83 (73)
Nunen et al. <sup>11</sup>	FAME: 5-year	2015	FFR vs. ANGIO-guided PCI	ACS, Stable CAD	5	<0.80	429	ANGIO	63.9	318 (74)
							436	FFR	64.5	328 (75)
Layland et al. <sup>10</sup>	FAMOUS-NSTEMI	2015	FFR vs. ANGIO-guided PCI	ACS	1	≤0.80	174	ANGIO	61.6	127 (73.0)
							176	FFR	62.3	133 (75.6)
Zhang et al. <sup>22</sup>	N/A	2016	FFR vs. ANGIO-guided PCI	ACS	1	≤0.80	110	ANGIO	70±3.4	78 (70.9)
							110	FFR	70±3.7	75 (68.2)
Thuesen et al. <sup>25</sup>	FARGO	2018	FFR vs. ANGIO-guided CABG	ACS, Stable CAD	0.5	≤0.80	49	ANGIO	65.3±8.8	44 (89)
							48	FFR	66.4±6.4	44 (88)
Toth et al. <sup>13</sup>	GRAFFITI	2019	FFR vs. ANGIO-guided CABG	ACS, Stable CAD	1	≤0.80	84	ANGIO	67	66 (79)
							88	FFR	67	73 (83%)
Quintella et al. <sup>23</sup>	N/A	2019	FFR vs. ANGIO-guided PCI	ACS, Stable CAD	1	<0.75	100	ANGIO	59.5±9.4	22 (46.8)
							100	FFR	62.7±8.4	25 (53.2)
Rioufol et al. <sup>15</sup>	FUTURE	2021	FFR vs. ANGIO-guided PCI	ACS, Stable CAD	1	≤0.80	467	ANGIO	66±11	385 (82.0)
							460	FFR	65±10	393 (85.0)
Puymirat et al. <sup>14</sup>	FLOWER-MI	2021	FFR vs. ANGIO-guided PCI	ACS	1	≤0.80	577	ANGIO	61.9±11.4	468 (81.1)
							586	FFR	62.5±11.0	498 (85.0)
Fearon et al. <sup>18</sup>	FAME:3-year	2021	FFR guided PCI vs. ANGIO-guided CABG	ACS, Stable CAD	1	≤0.80	743	ANGIO	65.1±8.3	619 (83.3)
							757	FFR	65.2±8.6	616 (81.4)
Stables et al. <sup>16</sup>	RIPCORDER 2	2022	FFR vs. ANGIO-guided PCI	ACS, Stable CAD	1	≤0.80	552	ANGIO	64.3 (10.2)	426 (77.2)
							548	ANGIO+FFR	64.3 (10.0)	403 (73.5)
Lee et al. <sup>21</sup>	FRAME-AMI	2022	FFR vs. ANGIO-guided PCI	ACS	3.5	≤0.80	284	FFR	63.9±11.4	240 (84.5)
							278	ANGIO	62.7±11.5	234 (84.2)


Abbreviations: FFR, fractional flow reserve; ANGIO, angiography; PCI, percutaneous intervention; ACS, acute coronary syndromes; CAD, coronary artery disease

**Table 2.** Baseline comorbidities

Study	Group	Diabetes Mellitus, No. (%)	Hypertension, No. (%)	Hyperlipidemia, No. (%)	Current smoker, No. (%)	Previous Myocardial Infarction, No. (%)
Chen et al. <sup>28</sup>	ANGIO	43 (26.9)	106 (68.3)	32 (20.0)	64 (40.0)	19 (11.9)
	FFR	48 (30.0)	116 (72.5)	27 (16.9)	66 (41.3)	12 (7.5)
Park et al. <sup>24</sup>	ANGIO	39 (34)	65 (57)	78 (68)	38 (33)	20 (17)
	FFR	30 (26)	73 (64)	80 (70)	30 (26)	22 (19)
Nunen et al. <sup>11</sup>	ANGIO	107 (25)	277 (65)	316 (74)	130 (30)	155 (36)
	FFR	98 (22)	259 (59)	307 (70)	111 (25)	154 (35)
Layland et al. <sup>10</sup>	ANGIO	26 (14.9)	81 (46.6)	56 (32.2)	71 (40.8)	24 (13.8)
	FFR	26 (14.8)	78 (44.3)	71 (40.3)	72 (40.9)	22 (12.5)
Zhang et al. <sup>22</sup>	ANGIO	36 (32.7)	83 (75.5)	93 (84.5)	31 (28.2)	23 (20.9)
	FFR	40 (36.4)	81 (73.6)	90 (81.8)	29 (26.4)	24 (21.8)
Thuesen et al. <sup>25</sup>	ANGIO	11 (23)	33 (67)	36 (75)	8 (17)	10 (21)
	FFR	11 (22)	33 (67)	42 (86)	13 (27)	14 (29)
Toth et al. <sup>13</sup>	ANGIO	33 (40)	59 (70)	66 (79)	35 (42)	5 (6)
	FFR	31 (35)	68 (77)	70 (80)	47 (53)	15 (17)
Quintella et al. <sup>23</sup>	ANGIO	12 (50.0)	26 (50.9)	26 (52.0)	9 (47.4)	7 (46.7)
	FFR	12 (50.0)	25 (49.0)	24 (42.0)	10 (52.6)	8 (53.3)
Rioufol et al. <sup>15</sup>	ANGIO	147 (32.0)	283 (61.0)	286 (61.0)	118 (26.0)	100 (21.0)
	FFR	143 (31.0)	265 (58.0)	275 (60.0)	109 (24.0)	90 (20.0)
Puymirat et al. <sup>14</sup>	ANGIO	82(14.2)	262 (45.4)	237 (41.1)	210 (36.4)	31 (5.4)
	FFR	107(18.3)	253 (43.2)	232 (39.6)	235 (40.1)	45 (7.7)
Fearon et al. <sup>18</sup>	ANGIO	214 (28.8)	556 (75.0)	531 (71.7)	136 (18.4)	248 (33.5)
	FFR	214 (28.3)	538 (71.2)	521 (68.9)	145 (19.2)	252 (33.3)
Stables et al. <sup>16</sup>	ANGIO	97 (17.6)	294 (53.5)	317 (57.6)	356 (65)	129 (23.4)
	ANGIO+FFR	113 (20.6)	315 (57.6)	315 (57.5)	316 (58.5)	117 (21.4)
Lee et al. <sup>21</sup>	ANGIO	86 (30.9)	152 (54.7)	107 (38.5)	105 (37.8)	7 (2.5)
	FFR	97 (34.2)	151 (53.2)	121 (42.6)	91 (32.0)	7 (2.5)


Abbreviations: FFR, fractional flow reserve; ANGIO, angiography

**Figure 1.** Preferred Reporting Items for Systemic Reviews and Meta-Analysis (PRISMA) flow diagram.




**Figure 2. (A)** Forest plot showing all-cause mortality comparing FFR versus angiography **(B)** Forest plot showing cardiac mortality comparing FFR versus angiography **(C)** Forest plot showing repeat revascularization comparing FFR versus angiography **(D)** Forest plot showing target lesion revascularization (TLR) comparing FFR versus angiography **(E)** Forest plot showing MACE comparing FFR versus angiography **(F)** Forest plot showing myocardial infarction comparing FFR versus angiography **(G)** Forest plot showing stroke or TIA comparing FFR versus angiography.


**(A) All-cause mortality**




**(B) Cardiovascular mortality**




**(C) Repeat revascularization**




**(D) Target lesion revascularization**




**(E) Major adverse cardiac event (MACE)**



**(F) Myocardial infarction**




**(G) Stroke or TIA**




**Figure 3. (A) Forest plot showing the average number of stents (B) Forest plot showing contrast volume (C) Forest plot showing total procedural time**


**(A) Average Number of Stents**



**(B) Contrast Volume (liters)**



**(C) Total Procedure Time (hours)**



## **Figure Titles and Legends:**

### **Forest Plot Risk of bias legend:**

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Binding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias