

The interplay between air pollution, built environment, and physical activity: perceptions of children and youth in rural and urban India

Authors: Jamin Patel¹, Tarun Reddy Katapally¹⁻⁴, Anuradha Khadilkar⁴, Jasmin Bhawra^{4,5*}

¹DEPth Lab, Faculty of Health Sciences, Western University, London, Ontario, Canada N6A 5B9

²Department of Epidemiology and Biostatistics, Schulich School of Medicine and Dentistry, Western University, London, Ontario, Canada N6A 3K7

³Children's Health Research Institute, Lawson Health Research Institute, 750 Base Line Road East, Suite 300, London, Ontario, Canada N6C 2R5

⁴Hirabai Cowasji Jehangir Medical Research Institute, Pune, Maharashtra, India 411 001

⁵School of Occupational and Public Health, Toronto Metropolitan University, Toronto, Ontario, Canada M5B 2K3

* Corresponding author

Dr. Jasmin Bhawra

E-mail: jasmin.bhawra@torontomu.ca

School of Occupational and Public Health

Toronto Metropolitan University

350 Victoria Street

Toronto, ON M5B 2K3

ABSTRACT

The role of physical inactivity as a contributor to non-communicable diseases (NCDs) risk in children and youth is widely recognized. Air pollution and built environment can limit participation in physical activity and exacerbate NCD risk; however, the relationships between perceptions of air pollution, built environment, and health behaviors are not fully understood, particularly among children and youth in low and middle-income countries. Currently, there are no studies capturing how child and youth perceptions of air pollution and built environment influence physical activity in India, thus, this study investigates the association between perceived air pollution and built environmental factors on moderate-to-vigorous physical activity (MVPA) levels of children and youth in both rural and urban India. Online surveys captured MVPA, perception of air pollution and built environment factors, as well as relevant sociodemographic characteristics from parents and children aged 5 to 17 years in partnership with 41 schools across 28 urban and rural locations during the Coronavirus disease lockdowns in 2021. After adjusting for age, gender, and location, a significant association was found between the perception of air pollution and MVPA levels ($\beta = -18.365$, $p < 0.001$). Similarly, the perception of a high crime rate was associated with lower MVPA levels ($\beta = -23.383$, $p = 0.002$). Reporting the presence of zebra crossings and pedestrian signals or attractive natural sightings was associated with higher MVPA levels; however, this association varied across sociodemographic groups. These findings emphasize the importance of addressing air pollution and improving the built environment to facilitate outdoor active living, including active transportation – solutions that are particularly relevant not only for NCD risk mitigation, but also for climate change adaptation.

Keywords: air pollution; built environment; physical activity; children and youth; India

INTRODUCTION

The health benefits of physical activity have been extensively researched, with studies associating physical activity with improved physical, psychological, social, and cognitive health indicators among children and youth (Castelli, 2022; Pascoe et al., 2020; Poitras et al., 2016). However, we are in the midst of a global physical inactivity pandemic (Kohl et al., 2012; Pratt et al., 2020) with recent studies indicating that a large proportion of children and youth do not meet the recommended levels of physical activity necessary to maintain good health (Aubert et al., 2022; Author, 2018; Author et al., 2023; Kuzik et al., 2023). Physical inactivity has not only been linked to the growing prevalence of chronic diseases such as heart disease, type-2 diabetes, and cancer (CDC, 2022), but it is also considered to be the fourth leading risk factor for mortality (WHO, 2012).

This is a pressing concern, as the burden of disability and mortality of non-communicable diseases (NCDs) has risen among children and youth (Akseer et al., 2020), with physical inactivity as a key risk factor (Budreviciute et al., 2020). Low to middle-income countries such as India experience a significant burden of NCDs, which is compounded by the presence of multiple syndemics (Frumkin and Haines, 2019; World Health Organization, 2009). These syndemics result from the coexistence and interaction of infectious diseases and NCDs, which further intensify health challenges in low to middle-income countries (Boutayeb, 2006; Oni and Unwin, 2015). Socioeconomic factors, including poverty, limited healthcare access, and inadequate infrastructure in many low to middle-income countries also exacerbate the burden of

NCDs and infectious diseases (Gouda et al., 2019). In India, NCDs are a major health concern, accounting for nearly 60% of all deaths (Nethan et al., 2017).

Studies have found that behaviours such as physical activity can be effective in preventing NCDs (Budreviciute et al., 2020), thus promoting physical activity can not only help prevent a substantial number of NCD-related premature deaths, but also the overall burden on healthcare systems and society. In particular, meeting the globally recommended 60 minutes of daily moderate-to-vigorous physical activity (MVPA) is an effective strategy for preventing and managing NCDs (Baran et al., 2020; Sprengeler et al., 2021). A recent study indicated that a one-hour-per-day increase in light-intensity physical activity resulted in 10% less mortality, while the same one-hour-per-day increase in MVPA resulted in 40% less mortality among adults over the age of 40 (Saint-Maurice et al., 2018). As a result, identifying the socioeconomic and environmental factors that influence physical activity levels, particularly MVPA, is imperative to help create conducive environments which reduce the NCD burden over the lifespan.

Air pollution is one environmental issue that influences outdoor physical activity, particularly in urban centres of India (Singh et al., 2021). Due to its recognized impact on physical and respiratory health, high levels of air pollution can deter outdoor activity among children and youth (Yu and Zhang, 2023). Multiple studies have found air pollution to be associated with increased odds of physical inactivity among children and adults (An et al., 2018), with one study identifying air pollution to be associated with lower physical activity and higher sedentary behaviour among children and youth (Yu and Zhang, 2023).

In addition to air pollution, the built environment can influence the health behaviours of children and youth. Built environment factors such as poor pedestrian safety, traffic hazards, and prevalence of crime have been associated with lower physical activity levels among children and youth in low-and middle-income countries (Hermosillo-Gallardo et al., 2020; Lv and Wang, 2023; Odunitan-Wayas et al., 2021), as well as in high-income countries such as Canada (Hunter et al., 2022). In contrast, built environment factors such as neighbourhood aesthetics have been associated with higher levels of physical activity among children and youth (Lv and Wang, 2023; Odunitan-Wayas et al., 2021; Qu et al., 2021).

Despite evidence indicating the growing impact of air pollution and the built environment on physical activity, there is a lack of research in India capturing the perceptions of children and youth regarding air pollution and the built environment in their communities. While other studies have focused specifically on the impacts of air quality or built environment features on MVPA (Author et al., 2023; Rees-Punia et al., 2018; Singh et al., 2021), this is the only study to date exploring location-specific experiences of air pollution, built environment, and MVPA from a youth perspective. Irrespective of location, most physical activity studies have focused on the perspectives of parents or caregivers, overlooking the unique viewpoints and concerns of younger populations (Daniel et al., 2020; Fan et al., 2017; Hunter et al., 2022; Wang et al., 2022; Waters et al., 2021). Understanding how children and youth perceive their environment is crucial in designing effective interventions and strategies to promote physical activity. More importantly, in the most populous country in the world, with diverse urban and rural environments of varying scales (Hertog and Gerland, 2023; Sciubba, 2023), it is important to understand the perception of children and youth across jurisdictions. The objective of this study is to understand the


relationship between child and youth perceptions of air pollution, built environment, and MVPA across rural and urban regions in India.

METHODS

Design

A cross-sectional, observational study was conducted during the Coronavirus disease lockdown in India in 2021. Data were collected through online surveys as a part of a multi-center cohort study (Author et al., 2022; Vispute et al., 2023). The study implemented a multi-stage stratified random sampling method involving urban and rural schools in five Indian states across 28 different cities and villages as shown in **Figure 1**. Ethics approval was obtained from the Ethics Committee Jehangir Clinical Development Centre Pvt. Ltd in Pune, Maharashtra (EC registration number –ECR/352/Inst/MH/2013/RR-19). All study methods adhered to the Declaration of Helsinki for biomedical research involving human participants.

Figure 1 – Map of India showing 28 cities and villages sampled in the study



Recruitment and Participants

School principals (n = 50) from urban and rural areas across five Indian states (Maharashtra, Gujarat, Telangana, Madhya Pradesh, and Tamil Nadu) were invited to participate in the study.

Of the invited principals, 41 agreed to participate and electronically shared the study information and consent forms with parents and children from their respective schools. Informed consent was obtained from 1042 children and youth aged 5 to 17 years and their parents. After receiving consent, anonymized questionnaires were administered via Google Forms between March 15, 2021, and May 20, 2021. Children under 13 years of age received parental assistance in completing the questionnaire. Children and youth were asked to self-report information on various factors, including physical activity and sedentary behaviour, perception of air pollution, participation in sports, active transportation, peer support, built environment, and school policies promoting physical activity.

Sociodemographic Characteristics

In order to assess the relationship between various sociodemographic characteristics and behaviours, this study collected information on participants' age, gender, and geographic location. Gender was collected by asking participants to report whether they are male or female. Children and youth provided their date of birth, which was used to calculate the age of each participant at the time of the study. As schools from both urban and rural centres were selected from five Indian states, the urban or rural residence was determined based on the location of the respective school in a city or village.

Physical Activity Measures

To assess physical activity levels, participants were asked to indicate the frequency of their involvement in various activities. Children and youth completed a questionnaire with an extensive list of activities, including sports, weight training, running, and a range of other

activities like jumping rope, playing garden games, and swimming. For each selected activity, participants reported the duration in minutes spent per day. The collected physical activity data were then categorized into three intensity types: inactivity, light activity, and MVPA [28,29]. In this study, MVPA was calculated by summing the daily average time spent in moderate and vigorous activities, including sports, garden games, running, aerobic activity, cycling, swimming, and weight training.

Air Pollution Measures

Participants were asked to indicate their level of agreement using a 5-point Likert scale (strongly disagree, somewhat disagree, neither agree nor disagree, somewhat agree, strongly agree) with four statements related to air pollution in their neighbourhood. These statements included whether there is an air pollution problem, whether air pollution prevents them from being active outside, whether their parents restrict their outdoor activity due to air pollution, and if they take precautions against air pollution when going outside or engaging in outdoor physical activity. Responses obtained in this study using a 5-point Likert scale were analyzed as dichotomous variables. Responses falling within the “agree” category included “strongly agree” and “agree”, while responses falling within the “disagree” category included “strongly disagree” and “disagree.” If responses were “neither agree nor disagree,” they were excluded from the analyses.

Built Environment Measures

Perceptions about the built environment were collected by asking children and youth about their level of agreement using a 5-point Likert scale (strongly disagree, somewhat disagree, neither

agree nor disagree, somewhat agree, strongly agree) with statements such as: “There are attractive natural sightings in my neighbourhood,” “There is a high crime rate in my neighbourhood,” and “There are zebra crossing and pedestrian signals to help walkers cross busy streets in my neighbourhood.” The full list of questions can be found in the **Supplementary File**.

Peer Support Measures

To assess peer support, participants were asked about the physical activity levels of their closest friends: "Your closest friends are the ones you enjoy spending the most time with. How many of your closest friends engage in regular physical activity?" Participants were provided with response options ranging from zero to four or more friends.

Statistical Analysis

All statistical analyses were performed using R Studio version 2022.12.0+353 (RStudio Team, 2015). The primary independent variable was the perception of air pollution as a problem in the city, while the primary dependent variable was the average daily minutes spent on MVPA. Following established procedures for analyzing physical activity data, participants reporting over two standard deviations of average daily MVPA were considered outliers and removed from the analysis. The sample was divided based on gender (male and female), geographic location (urban and rural), and age cohorts (5-10 years old, 11-13 years old, and 14-17 years old), resulting in eight multiple linear regression models, including the overall model that used data from all participants. The estimated coefficients in these models indicated the expected change in daily MVPA associated with the participant’s perception of air pollution as a problem while controlling for age, community and built environment, and peer support. The gender models were also

adjusted for location, while the location models were adjusted for gender. T-tests for unequal variances were conducted to compare average MVPA across various demographic and behavioural factors. A chi-square analysis was also conducted to compare the frequency distribution of responses to the perception of air pollution as a problem. All reported results in this study were considered statistically significant at a threshold of $p < 0.05$.

RESULTS

The study consisted of 1042 participants. A total of 50 participants were excluded as their age was above the study's target range of 5-17 years. The overall sample was further segregated by sociodemographic characteristics to assess differences in perceptions of air pollution across subgroups. As shown in **Table 1**, the study included 499 males (49.7%) and 493 females (50.3%), comprising 592 urban residents (59.7%) and 400 rural residents (40.3%). There were 339 participants aged 5 to 10 years (34.2%), 318 participants aged 11 to 13 years (32.1%), and 335 participants aged 14 to 17 years (33.8%).

Table 1. Children and youth sample summary by gender, age, and location (N = 992)

	Total (N=992)	
Sample characteristic	N	%
Age Group		
5 to 10 years	339	34.2
11 to 13 years	318	32.1
14 to 17 years	335	33.8
Gender		

Male	499	49.7
Female	493	50.3
Location		
Rural	400	40.3
Urban	592	59.7

Of the total participants, 663 were included in the regression models after removing missing data and outliers for MVPA. As shown in **Table 2**, the average daily MVPA for all children and youth was 82.5 minutes. Those who perceived air pollution as a problem had lower MVPA levels compared to those who did not perceive air pollution as a problem ($t(757.22) = -2.306$, $p = 0.033$). Males had significantly higher MVPA levels compared to females ($t(917.92) = 10.571$, $p < 0.001$). There was no significant difference in MVPA levels between urban and rural areas; however, more urban residents reported an air pollution problem than rural residents ($\chi^2 = 87.297$, $p < 0.001$). Moreover, among participants from rural areas, those who perceived an air pollution problem had lower MVPA levels compared to those who disagreed ($t(355.36) = 2.3762$, $p = 0.018$). When examining MVPA levels across different age groups, there was no significant difference observed.

Table 2. Daily minutes of engagement in MVPA among children and youth based on gender, age, and location

		Overall Sample	There is an air pollution problem: Agree	There is an air pollution problem: Disagree

Category	Group	Average minutes (SD)	Average minutes (SD)	Average minutes (SD)
Gender	Male	107.6 (80.7)	101.0 (81.0)	116.1 (77.3)
	Female	57.8 (65.0)	55.1 (63.3)	61.6 (68.1)
Location	Urban	83.4 (79.4)	81.0 (78.1)	91.9 (82.4)
	Rural	81.2 (74.0)	71.4 (71.5)	89.3 (75.4)
Age (years)	5-10	76.8 (68.7)	72.5 (68.2)	88.8 (72.6)
	11-13	87.7 (81.2)	80.5 (77.0)	95.5 (81.7)
	14-17	83.4 (81.5)	82.2 (84.0)	86.7 (79.1)
Overall	NA	82.5 (77.3)	78.0 (76.2)	90.2 (78.0)

Table 3 shows the association between air pollution and MVPA for the full sample of children and youth. Participants who agreed that there is an air pollution problem were associated with lower MVPA levels ($\beta = -18.365$, $p = 0.002$), compared to those who disagreed. Furthermore, the presence of a high crime rate in the neighbourhood was also associated with lower MVPA levels ($\beta = -23.383$, $p = 0.002$) compared to those who disagreed. Having one or more active friends was associated with higher levels of MVPA ($\beta = 33.531$, $p = 0.005$) in comparison to having no active friends. However, other built-environment factors such as the presence of footpaths, attractive natural sightings, zebra crossings, and pedestrian signals did not show significant associations with MVPA levels in the overall model.

Table 3. Relationship between reporting that there is an air pollution problem and moderate to vigorous physical activity (Model 1)

	Beta coefficients (95% Confidence Intervals)^a
There is an air pollution problem: Agree	-18.365* (-30.115, -6.615)
Disagree – (Ref)	
There is a high crime rate in my neighborhood: Agree	-23.383* (-37.856, -8.910)
Disagree – (Ref)	
There are attractive natural sightings in my neighborhood: Agree	-1.532 (-13.623, 10.558)
Disagree – (Ref)	
There are zebra crossings and pedestrian signals to help walkers cross busy streets in my neighborhood: Agree	10.590 (-3.038, 24.219)
Disagree – (Ref)	
Has one or more active friends: Yes	33.531* (10.354, 56.708)
Has no active friends – (Ref)	
Constant	7.980 (-25.378, 41.338)

* Indicates a statistically significant relationship at the $p < 0.05$ level

^aThe model was adjusted for age, gender, and location.

Table 4 demonstrates that across the three age groups, significant associations were found between reporting air pollution and MVPA levels. In the 5 to 10-year age group, the perception of an air pollution problem was associated with lower levels of MVPA ($\beta = -26.725$, $p = 0.012$)

compared to those who disagreed. However, no significant association was found between the perception of an air pollution problem and MVPA levels in the 11 to13-year age group (n = 202) and 14 to 17 age group. Children and youth in the 5 to10 and 11 to13-year age groups, who agreed to a high crime rate in the neighbourhood, were associated with lower MVPA levels than their counterparts who disagreed (Age 5-10: $\beta = -26.833$, $p = 0.025$, Age 11-13: $\beta = -31.251$, $p = 0.038$). Reporting the presence of pedestrian infrastructure, such as zebra crossings and signals, was associated with higher MVPA levels in the 14 to17-year age model ($\beta = 22.860$, $p = 0.0303$) but not in the 5 to 10 and 11 to 13-year age groups. Further, the presence of attractive natural sightings in the neighbourhood was associated with higher MVPA levels in the 5 to 10-year age group ($\beta = 23.557$, $p = 0.026$), in comparison to those who disagreed.

Table 4. Relationship between reporting that there is an air pollution problem and moderate to vigorous physical activity across various sociodemographic groups

	Beta coefficients (95% Confidence Intervals)						
	Age Group^a			Gender^b		Location^c	
	Model 2a-2c			Model 3		Model 4	
	5 – 10 years	11 – 13 years	14 – 17 years	Male	Female	Urban	Rural
There is an air pollution problem: Agree	-26.726* (-47.321, -6.131)	-20.194* (-42.809, 2.421)	-8.575 (- 27.501, 10.352)	-25.090* (-43.372, -6.809)	-9.286 (- 23.686, 5.115)	-14.947 (-33.292, 3.397)	-22.310* (-38.267, -6.354)
Disagree – (Ref)							

There is a high crime rate in my neighborhood: Agree	-26.833* (-50.150, -3.515)	-31.251* (-60.497, -2.005)	-13.643 (-37.049, 9.763)	-19.521 (-43.387, 4.345)	-27.620* (-44.253, -10.986)	-24.311* (-45.918, -2.704)	-22.652* (-42.802, -2.502)
Disagree – (Ref)							
There are attractive natural sightings in my neighborhood: Agree	23.557* (2.973, 44.141)	-14.572 (-36.266, 7.122)	-11.143 (-31.854, 9.568)	-12.749 (-31.347, 5.849)	12.454 (- 2.373, 27.281)	7.061 (- 14.247, 28.368)	-7.243 (- 21.557, 7.071)
Disagree – (Ref)							
There are zebra crossings and pedestrian signals to help walkers cross busy streets in my neighborhood	-17.589 (-44.031, 8.852)	18.053 (- 8.975, 45.080)	22.860* (2.296, 43.425)	15.693 (- 5.452, 36.838)	6.408 (- 10.190, 23.006)	3.881 (- 16.721, 24.483)	17.666 (- 1.223, 36.556)

d: Agree							
Disagree – (Ref)							
Has one or more active friends: Yes	17.728 (-17.370, 52.827)	33.947 (-4.270, 72.163)	45.860 (-3.075, 94.796)	22.554 (-24.566, 69.674)	40.085* (16.432, 63.738)	36.132* (10.028, 62.236)	7.007 (-67.381, 81.396)
Has no active friends – (Ref)							
Constant	-24.944 (-86.262, 36.373)	-114.433 (-262.640, 33.775)	110.565 (-33.903, 255.033)	54.252 (-4.561, 113.065)	29.208 (-8.814, 67.230)	24.872 (-25.045, 74.789)	32.884 (-45.400, 111.169)

Note: Three separate regression models were run for the built environment variables

* Indicates a statistically significant relationship at the $p < 0.05$ level

^aModel 2 was adjusted for age, gender, and location.

^bModel 3 was adjusted for age and location.

^cModel 4 was adjusted for age and gender.

When comparing the models by gender, in the male model, agreeing that air pollution is a problem was associated with higher MVPA levels (Males: $\beta = -25.090$, $p = 0.008$); however, this association was not significant in the female model. Among females, the presence of a high crime rate in their neighbourhood was associated with lower MPVA levels (Females: $\beta = -27.620$, $p = 0.001$), however; this relationship was not significant for males. In contrast, having

one more active friend was associated with higher MVPA levels in females ($\beta = 40.085$, $p = 0.001$).

In the urban model, agreeing with the statement that air pollution is a problem was not significantly associated with MVPA levels. However, in the rural model, the perception of air pollution as a problem was associated with lower MVPA levels ($\beta = -22.310$, $p = 0.006$), compared to those who did not perceive air pollution to be a problem. Contrastingly, having one or more active friends was associated with more MVPA in the urban model ($\beta = 36.132$, $p = 0.007$) but not in the rural model. Further, reporting high crime was associated with lower MVPA levels in both the rural model ($\beta = -22.652$, $p = 0.028$) and the urban model ($\beta = -24.311$, $p = 0.028$).

DISCUSSION

This is the first study to investigate the relationship between child and youth perceptions of air pollution, built environment, and MVPA in urban and rural regions in India. In building on the key findings and gaps identified in the 2022 India Report Card on Physical Activity for Children and Adolescents (Author et al., 2023), this study focused on perceptions of children and youth that have thus far been largely ignored in active living research in India.

Given the wide-ranging impacts of air pollution on vulnerable groups such as children and youth, there is an urgent need to understand how this group's physical activity levels are affected by air pollution, especially considering the worsening air quality in India that is being attributed to climate change (Singh et al., 2023; Van Oldenborgh et al., 2018). By including the perceptions of

children and youth, this study provides a distinct understanding of the impact of air pollution and the built environment on physical activity in younger populations.

The overall findings suggest that perceptions of an air pollution as a problem and negative perceptions of the built environment are both associated with lower levels of MVPA among children, a finding that aligns with active living research in other parts of the world (Goon et al., 2020; Gu et al., 2023; Loh et al., 2019; Taylor et al., 2018). This finding suggests that while air pollution is generally high in Indian cities and towns (Dandotiya et al., 2020; Hussain et al., 2023; Mahendra et al., 2023), perception of air pollution -- irrespective of actual levels of air pollution -- can potentially influence the physical activity of children and youth.

As the evidence indicates that there are differences in not only air pollution, but also MVPA across urban and rural areas in India (Author et al., 2023; Vieira et al., 2023), we developed statistical models segregated by location (urban vs. rural). More children and youth in urban areas perceived air pollution as a problem, a finding that aligns not only with current evidence of air pollution perceptions (Author et al., 2023), but also with actual air pollution levels in India that tend to be higher in urban areas (WHO, 2023). However, contrastingly, rural children and youth who perceived higher air pollution reported significantly lower MVPA in comparison with children and youth who reported lower perception of air pollution – a finding that was not significant in urban areas. This finding also contrasts with previous research which indicates that rural residents lack awareness of the impacts of air pollution on health (Yang, 2020). It is plausible that children and youth in urban areas may get habituated or desensitized to the level of pollution and stop perceiving it as problematic. On the other hand, it is also possible that

pollution levels are increasing in rural areas of India due to rapid industrialization (Karambelas et al., 2018; Ravishankara et al., 2020). These findings require further exploration, including linking citizen perception data with objective weather data.

In addition to location-based differences, by examining perceptions of air pollution and MVPA levels in males and females, this study adds to the existing evidence on gender-based differences in physical activity engagement (Author, 2018; Author et al., 2023). Males reported higher MVPA compared to females, a finding that aligns with existing literature that suggests males in India tend to accumulate higher levels of physical activity compared to females, driven by factors such as social acceptability, cultural norms, and general perception of lack of safety that limit outdoor activities among female children and youth neighbourhoods (Author et al., 2023; Mathur et al., 2021; Tyagi and Raheja, 2020). Among males, the perception of air pollution as a problem was associated with lower levels of MVPA; however, this association was not significant in females, a finding that further suggests that female children and youth might not be obtaining as much MVPA outdoors in comparison to males. These findings indicate the need for targeted interventions to promote physical activity in both male and female populations.

The study findings also reveal significant associations between the perception of air pollution and MVPA levels among children and youth in different age groups. Perceiving air pollution to be a problem was linked to lower MVPA in the 5 to 10-year age group, but not in the 11 to 13 and 14 to 17-year age groups. These findings could be due to younger children being more influenced by their immediate surroundings and parental guidance, and likely having less independent mobility than older children (Pickhardt, 2010; Schoeppe et al., 2013). Moreover,

studies have shown younger children to be more physiologically susceptible to the negative effects of air pollution, which may lead to changes in their physical activity behaviours (Landrigan et al., 2010; Sunyer et al., 2015; UN, 2018).

In terms of the built environment, the presence of attractive natural sightings in the neighbourhood was related to higher MVPA in the 5 to 10-year age group. One potential explanation for this association is young children's fascination with novel sightings in nature, as engaging with nature supports imaginative play and interaction with their environment (Dowdell et al., 2011). Similarly, community-related factors such as perception of high crime rates in the neighbourhood were associated with reduced MVPA in the 5 to 10 and 11 to 13-year age groups, but not among 14 to 17-year-olds. It is possible that older adolescents might develop different coping mechanisms for dealing with perceived environmental challenges and have learned to adapt to their surroundings when making decisions about physical activity engagement (Dwyer et al., 2012). Furthermore, among females, the significant association between a high crime rate in their neighbourhood and lower MVPA levels highlights the potential impact of environmental safety concerns on physical activity. This finding is consistent with research indicating that safety concerns, particularly for women, can act as a barrier to physical activity, limiting opportunities for outdoor exercise and leading to reduced overall activity levels (Author et al., 2023; Mathur et al., 2021; Tyagi and Raheja, 2020). This discrepancy emphasizes the need for gender-specific interventions and strategies to address safety concerns and promote physical activity among females.

In this study, the presence of zebra crossings and pedestrian signals was associated with increased levels of MVPA among children and youth aged 14 to 17; however, this association was not observed in the younger population. These findings could potentially be explained by older adolescents having more independent mobility, including their choice of transportation and routes. As zebra crossings provide safer and more convenient opportunities for pedestrians to cross busy streets (Ismail et al., 2023), they may encourage older adolescents to engage in more outdoor physical activity. In contrast, younger children might rely more on their parents or guardians to dictate their travel routes (Pickhardt, 2010; Schoeppe et al., 2013), and the presence of zebra crossings may not have the same impact on their physical activity as with older adolescents. These findings demonstrate the need for policymakers to prioritize the implementation and maintenance of zebra crossings and pedestrian signals as a means to promote physical activity, particularly in older adolescents.

Furthermore, the impact of peer support on MVPA was found to vary based on gender and location. Previous research has found that by facilitating social connections and promoting enjoyable physical activities, active friends play a vital role in enhancing both physical and mental well-being (Author et al., 2023; Jago et al., 2011; Leggett et al., 2012). In this study, female and urban children and youth who reported having active friends had higher MVPA; however, the same association was not observed in males or rural residents, indicating that other factors might influence their MVPA levels, i.e., the impact of peer support on their physical activity behaviours may be different from those observed in females or urban residents. Further research is required to better understand the specific mechanisms and contextual factors that contribute to these disparities among different demographic groups in India.

The study findings on perceptions of built environmental factors, perceptions of air pollution, and demographic-related variations in physical activity are particularly relevant in light of climate change impacts, with rising temperatures and heat waves exacerbating the effects of air pollution, particularly in the global south (Singh et al., 2023; Van Oldenborgh et al., 2018; Yang and Shao, 2021). In low to middle-income countries such as India where there are increasing levels of particulate matter pollution due to industrial emissions, vehicle exhaust, and the burning of biomass, understanding how people perceive and respond to air pollution is not only crucial for promoting health in children and youth, but is also important for NCD prevention and management (Gargava and Rajagopalan, 2016; India State-Level Disease Burden Initiative Air Pollution Collaborators, 2019; Upadhyay et al., 2018). Nevertheless, with the impact of climate change being felt across the world (Dahl et al., 2023; Hanes et al., 2019), irrespective of location, child and youth perceptions are critical in understanding changing patterns of MVPA globally – an area of research that will require more attention going forward.

Strengths and Limitations

By examining the unique perceptions of air pollution among children and youth across 28 cities and villages from urban and rural populations, this study allowed for cross-jurisdictional comparisons. However, as data were collected at one time point during the COVID-19 pandemic lockdown, reported pollution perceptions and physical activity may differ from a non-pandemic period. Future studies should focus on conducting longitudinal data collection, where both objective air pollution and MVPA data should be linked with the subjective perception of children and youth.

CONCLUSION

This is the first study to investigate the association between child and youth perceptions of air pollution, built environment, and MVPA in both urban and rural regions of India. The study findings emphasize the importance of considering children and youth's unique perceptions of air pollution and their built environment, as they significantly impact their physical activity levels. The study's findings not only highlight the location-, gender-, and age-specific criteria in informing targeted physical activity interventions, but the results indicate the need to focus on child and youth perceptions of air pollution in this age of climate emergency, irrespective of location to understanding changing patterns of MVPA.

REFERENCES

- Akseer, N., Mehta, S., Wigle, J., Chera, R., Brickman, Z.J., Al-Gashm, S., Sorichetti, B., Vandermorris, A., Hipgrave, D.B., Schwalbe, N., Bhutta, Z.A., 2020. Non-communicable diseases among adolescents: current status, determinants, interventions and policies. BMC Public Health 20, 1908. <https://doi.org/10.1186/s12889-020-09988-5>
- An, R., Zhang, S., Ji, M., Guan, C., 2018. Impact of ambient air pollution on physical activity among adults: a systematic review and meta-analysis. Perspect Public Health 138, 111–121. <https://doi.org/10.1177/1757913917726567>
- Aubert, S., Barnes, J.D., Demchenko, I., Hawthorne, M., Abdeta, C., Nader, P.A., Sala, J.C.A., Aguilar-Farias, N., Aznar, S., Bakalár, P., Author, Brazo-Sayavera, J., Bringas, M., Cagas, J.Y., Carlin, A., Chang, C.-K., Chen, B., Christiansen, L.B., Christie, C.J.-A., Roia, G.F.D., Nyström, C.D., Demetriou, Y., Djordjic, V., Emeljanovas, A., Endy, L.F., Gába, A., Galaviz, K.I., González, S.A., Hesketh, K.D., Huang, W.Y., Hubona, O., Jeon, J.Y.,

Jurakić, D., Jürimäe, J., Author, Katewongsa, P., Katzmarzyk, P.T., Kim, Y.-S., Lambert, E.V., Lee, E.-Y., Levi, S., Lobo, P., Löf, M., Loney, T., López-Gil, J.F., López-Taylor, J., Mäestu, E., Mahendra, A., Makaza, D., Mallari, M.F.T., Manyanga, T., Masanovic, B., Morrison, S.A., Mota, J., Müller-Riemenschneider, F., Bermejo, L.M., Murphy, M.H., Naidoo, R., Nguyen, P., Paudel, S., Pedišić, Ž., Pérez-Gómez, J., Reilly, J.J., Reimers, A.K., Richards, A.B., Silva, D.A.S., Saonuam, P., Sarmiento, O.L., Sember, V., Shahril, M.R., Smith, M., Standage, M., Stratton, G., Subedi, N., Tammelin, T.H., Tanaka, C., Tesler, R., Thivel, D., Tladi, D.M., Tlučáková, L., Vanderloo, L.M., Williams, A., Wong, S.H.S., Wu, C.-L., Zembura, P., Tremblay, M.S., 2022. Global Matrix 4.0 Physical Activity Report Card Grades for Children and Adolescents: Results and Analyses From 57 Countries. *Journal of Physical Activity and Health* 19, 700–728.

<https://doi.org/10.1123/jpah.2022-0456>

Author, 2018. Results from India's 2018 Report Card on Physical Activity for Children and Youth. *Journal of Physical Activity and Health* 15, S373–S374.

<https://doi.org/10.1123/jpah.2018-0475>

Author, Author, Krishnaveni, G.V., Kumaran, K., Author, 2023. The 2022 India Report Card on physical activity for children and adolescents. *Journal of Exercise Science & Fitness* 21, 74–82. <https://doi.org/10.1016/j.jesf.2022.10.013>

Author, Kajale, N., Oza, C., Oke, R., Gondhalekar, K., Patwardhan, V., Khadilkar, V., Mughal, Z., Padidela, R., 2022. Vitamin D status and determinants in Indian children and adolescents: a multicentre study. *Sci Rep* 12, 16790. <https://doi.org/10.1038/s41598-022-21279-0>

- Baran, J., Weres, A., Wyszynska, J., Pitucha, G., Czenczek-Lewandowska, E., Rusek, W., Leszczak, J., Mazur, A., 2020. 60 Minutes Per Day in Moderate to Vigorous Physical Activity as a Natural Health Protector in Young Population. *Int J Environ Res Public Health* 17, 8918. <https://doi.org/10.3390/ijerph17238918>
- Boutayeb, A., 2006. The double burden of communicable and non-communicable diseases in developing countries. *Transactions of the Royal society of Tropical Medicine and Hygiene* 100, 191–199.
- Budreviciute, A., Damiati, S., Sabir, D.K., Onder, K., Schuller-Goetzburg, P., Plakys, G., Katileviciute, A., Khoja, S., Kodzius, R., 2020. Management and Prevention Strategies for Non-communicable Diseases (NCDs) and Their Risk Factors. *Frontiers in Public Health* 8.
- Castelli, D.M., 2022. Physical Activity, Fitness, and Cognitive Function in Children and Adolescents, in: Wachira, L.-J. (Ed.), *Sport and Fitness in Children and Adolescents*. IntechOpen, Rijeka. <https://doi.org/10.5772/intechopen.104719>
- CDC, 2022. Chronic Disease Fact Sheet: Physical Inactivity | CDC [WWW Document]. URL <https://www.cdc.gov/chronicdisease/resources/publications/factsheets/physical-activity.htm> (accessed 7.31.23).
- Dahl, K.A., Abatzoglou, J.T., Phillips, C.A., Ortiz-Partida, J.P., Licker, R., Merner, L.D., Ekwurzel, B., 2023. Quantifying the contribution of major carbon producers to increases in vapor pressure deficit and burned area in western US and southwestern Canadian forests. *Environ. Res. Lett.* 18, 064011. <https://doi.org/10.1088/1748-9326/acbce8>

- Dandotiya, B., Sharma, H.K., Jadon, N., 2020. Ambient Air Quality and Meteorological Monitoring of Gaseous Pollutants in Urban Areas of Gwalior City India. *Environmental Claims Journal* 32, 248–263. <https://doi.org/10.1080/10406026.2020.1744854>
- Daniel, L., Michot, M., Esvan, M., Guérin, P., Chauvet, G., Pelé, F., 2020. Perceptions, Knowledge, and Practices Concerning Indoor Environmental Pollution of Parents or Future Parents. *Int J Environ Res Public Health* 17, 7669. <https://doi.org/10.3390/ijerph17207669>
- Dowdell, K., Gray, T., Malone, K., 2011. Nature and its Influence on Children’s Outdoor Play. *Journal of Outdoor and Environmental Education* 15, 24–35. <https://doi.org/10.1007/BF03400925>
- Dwyer, J.J.M., Chulak, T., Maitland, S., Allison, K.R., Lysy, D.C., Faulkner, G.E.J., Sheeshka, J., 2012. Adolescents’ self-efficacy to overcome barriers to Physical Activity Scale. *Res Q Exerc Sport* 83, 513–521. <https://doi.org/10.1080/02701367.2012.10599140>
- Fan, S., Yuan, Z., Liao, X., Tu, H., Lan, G., Maddock, J.E., Lu, Y., 2017. INTERNATIONAL PERSPECTIVES: Study of Parents’ Perception of Air Pollution and Its Effect on Their Children’s Respiratory Health in Nanchang, China. *Journal of Environmental Health* 79, E1–E9.
- Frumkin, H., Haines, A., 2019. Global Environmental Change and Noncommunicable Disease Risks. *Annu Rev Public Health* 40, 261–282. <https://doi.org/10.1146/annurev-publhealth-040218-043706>
- Gargava, P., Rajagopalan, V., 2016. Source apportionment studies in six Indian cities—drawing broad inferences for urban PM10 reductions. *Air Qual Atmos Health* 9, 471–481. <https://doi.org/10.1007/s11869-015-0353-4>

- Goon, S., Kontulainen, S., Muhajarine, N., 2020. Neighborhood Built Environment Measures and Association with Physical Activity and Sedentary Time in 9–14-Year-Old Children in Saskatoon, Canada. *Int J Environ Res Public Health* 17, 3837.
<https://doi.org/10.3390/ijerph17113837>
- Gouda, H.N., Charlson, F., Sorsdahl, K., Ahmadzada, S., Ferrari, A.J., Erskine, H., Leung, J., Santamauro, D., Lund, C., Aminde, L.N., Mayosi, B.M., Kengne, A.P., Harris, M., Achoki, T., Wiysonge, C.S., Stein, D.J., Whiteford, H., 2019. Burden of non-communicable diseases in sub-Saharan Africa, 1990–2017: results from the Global Burden of Disease Study 2017. *The Lancet Global Health* 7, e1375–e1387.
[https://doi.org/10.1016/S2214-109X\(19\)30374-2](https://doi.org/10.1016/S2214-109X(19)30374-2)
- Gu, X., Keller, J., Zhang, T., Dempsey, D.R., Roberts, H., Jeans, K.A., Stevens, W., Borchard, J., VanPelt, J., Tulchin-Francis, K., 2023. Disparity in Built Environment and Its Impacts on Youths’ Physical Activity Behaviors During COVID-19 Pandemic Restrictions. *J. Racial and Ethnic Health Disparities* 10, 1549–1559. <https://doi.org/10.1007/s40615-022-01341-3>
- Hanes, C.C., Wang, X., Jain, P., Parisien, M.-A., Little, J.M., Flannigan, M.D., 2019. Fire-regime changes in Canada over the last half century. *Can. J. For. Res.* 49, 256–269.
<https://doi.org/10.1139/cjfr-2018-0293>
- Hermosillo-Gallardo, M.E., Sebire, S.J., Jago, R., 2020. Perception of Safety and Its Association With Physical Activity in Adolescents in Mexico. *Am J Prev Med* 58, 748–755.
<https://doi.org/10.1016/j.amepre.2019.12.007>
- Hertog, S., Gerland, P., 2023. UN DESA Policy Brief No. 153: India overtakes China as the world’s most populous country | Department of Economic and Social Affairs [WWW

- Document]. UN DESA Policy Brief No. 153: India overtakes China as the world's most populous country. URL <https://www.un.org/development/desa/dpad/publication/un-desa-policy-brief-no-153-india-overtakes-china-as-the-worlds-most-populous-country/> (accessed 4.26.23).
- Hunter, S., Spence, J.C., Leatherdale, S.T., Carson, V., 2022. Associations Between Parent's Perceived Neighborhood Environment and Objectively Measured Walkability With Their Children's Physical Activity. *Journal of Physical Activity and Health* 19, 811–819. <https://doi.org/10.1123/jpah.2022-0310>
- Hussain, A.J., Sankar, T.K., Vithanage, M., Ambade, B., Gautam, S., 2023. Black Carbon Emissions from Traffic Contribute Sustainability to Air Pollution in Urban Cities of India. *Water Air Soil Pollut* 234, 217. <https://doi.org/10.1007/s11270-023-06232-9>
- India State-Level Disease Burden Initiative Air Pollution Collaborators, 2019. The impact of air pollution on deaths, disease burden, and life expectancy across the states of India: the Global Burden of Disease Study 2017. *Lancet Planet Health* 3, e26–e39. [https://doi.org/10.1016/S2542-5196\(18\)30261-4](https://doi.org/10.1016/S2542-5196(18)30261-4)
- Ismail, A., Yusof, L.M., Razlan, I.S.M., 2023. Pedestrian Behaviour at Unsignalized Zebra Crossing: A Case Study in UMP Gambang Campus. *IOP Conf. Ser.: Earth Environ. Sci.* 1140, 012023. <https://doi.org/10.1088/1755-1315/1140/1/012023>
- Jago, R., Macdonald-Wallis, K., Thompson, J.L., Page, A.S., Brockman, R., Fox, K.R., 2011. Better with a Buddy: Influence of Best Friends on Children's Physical Activity. *Medicine & Science in Sports & Exercise* 43, 259. <https://doi.org/10.1249/MSS.0b013e3181edefaa>

- Karambelas, A., Holloway, T., Kinney, P.L., Fiore, A.M., DeFries, R., Kieseewetter, G., Heyes, C.,
2018. Urban versus rural health impacts attributable to PM_{2.5} and O₃ in northern India.
Environ. Res. Lett. 13, 064010. <https://doi.org/10.1088/1748-9326/aac24d>
- Kohl, H.W., Craig, C.L., Lambert, E.V., Inoue, S., Alkandari, J.R., Leetongin, G., Kahlmeier, S.,
Lancet Physical Activity Series Working Group, 2012. The pandemic of physical
inactivity: global action for public health. Lancet 380, 294–305.
[https://doi.org/10.1016/S0140-6736\(12\)60898-8](https://doi.org/10.1016/S0140-6736(12)60898-8)
- Kuzik, N., Cameron, C., Carson, V., Chaput, J.-P., Colley, R., Doiron, J., Faulkner, G., Janssen,
I., Saunders, T., Spence, J.C., Tucker, P., Vanderloo, L.M., Tremblay, M.S., 2023. The
2022 ParticipACTION Report Card on Physical Activity for Children and Youth: Focus
on the COVID-19 pandemic impact and equity-deserving groups. Front Public Health 11,
1172168. <https://doi.org/10.3389/fpubh.2023.1172168>
- Landrigan, P.J., Rauh, V.A., Galvez, M.P., 2010. Environmental Justice and the Health of
Children. Mount Sinai Journal of Medicine: A Journal of Translational and Personalized
Medicine 77, 178–187. <https://doi.org/10.1002/msj.20173>
- Leggett, C., Irwin, M., Griffith, J., Xue, L., Fradette, K., 2012. Factors associated with physical
activity among Canadian high school students. Int J Public Health 57, 315–324.
<https://doi.org/10.1007/s00038-011-0306-0>
- Loh, V.H.Y., Veitch, J., Salmon, J., Cerin, E., Thornton, L., Mavoa, S., Villanueva, K., Timperio,
A., 2019. Built environment and physical activity among adolescents: the moderating
effects of neighborhood safety and social support. Int J Behav Nutr Phys Act 16, 132.
<https://doi.org/10.1186/s12966-019-0898-y>

- Lv, H., Wang, R., 2023. Association between the built environment and moderate to vigorous leisure-time physical activity among suzhou adolescents: a cross-sectional study. *BMC Public Health* 23, 1313. <https://doi.org/10.1186/s12889-023-16243-0>
- Mahendra, H.N., Mallikarjunaswamy, S., Kumar, D.M., Kumari, S., Kashyap, S., Fulwani, S., Chatterjee, A., 2023. Assessment and Prediction of Air Quality Level Using ARIMA Model: A Case Study of Surat City, Gujarat State, India. *Nat. Env. Poll. Tech* 22, 199–210. <https://doi.org/10.46488/NEPT.2023.v22i01.018>
- Mathur, P., Kulothungan, V., Leburu, S., Krishnan, A., Chaturvedi, H.K., Salve, H.R., Amarchand, R., Nongkynrih, B., Ganeshkumar, P., S, V.U.K., Laxmaiah, A., Boruah, M., Kumar, S., Patro, B.K., Raghav, P.R., Rajkumar, P., Sarma, P.S., Sharma, R., Tambe, M., Arlappa, N., Mahanta, T.G., Bhuyan, P.J., Joshi, R.P., Pakhare, A., Galhotra, A., Kumar, D., Behera, B.K., Topno, R.K., Gupta, M.K., Rustagi, N., Trivedi, A.V., Thankappan, K.R., Gupta, S., Garg, S., Shelke, S.C., 2021. Baseline risk factor prevalence among adolescents aged 15–17 years old: findings from National Non-communicable Disease Monitoring Survey (NNMS) of India. *BMJ Open* 11, e044066. <https://doi.org/10.1136/bmjopen-2020-044066>
- Nethan, S., Sinha, D., Mehrotra, R., 2017. Non Communicable Disease Risk Factors and their Trends in India. *Asian Pac J Cancer Prev* 18, 2005–2010. <https://doi.org/10.22034/APJCP.2017.18.7.2005>
- Odunitan-Wayas, F., Musera, D., Africa, A., West, S., Venter, T., Shung-King, M., Oni, T., King, A., Lambert, E., 2021. Association between the perceived neighbourhood environmental walkability and self-reported physical activity in South African adolescents: Mini-Oral

Presentation B1.6. The Health & Fitness Journal of Canada 14.

<https://doi.org/10.14288/hfjc.v14i3.595>

Oni, T., Unwin, N., 2015. Why the communicable/non-communicable disease dichotomy is problematic for public health control strategies: implications of multimorbidity for health systems in an era of health transition. *International health* 7, 390–399.

Pascoe, M., Bailey, A.P., Craike, M., Carter, T., Patten, R., Stepto, N., Parker, A., 2020. Physical activity and exercise in youth mental health promotion: a scoping review. *BMJ Open Sport & Exercise Medicine* 6, e000677. <https://doi.org/10.1136/bmjsem-2019-000677>

Pickhardt, 2010. Adolescence and Parental Influence | Psychology Today Canada [WWW Document]. URL <https://www.psychologytoday.com/ca/blog/surviving-your-childs-adolescence/201009/adolescence-and-parental-influence> (accessed 8.18.23).

Poitras, V.J., Gray, C.E., Borghese, M.M., Carson, V., Chaput, J.-P., Janssen, I., Katzmarzyk, P.T., Pate, R.R., Connor Gorber, S., Kho, M.E., Sampson, M., Tremblay, M.S., 2016. Systematic review of the relationships between objectively measured physical activity and health indicators in school-aged children and youth. *Appl. Physiol. Nutr. Metab.* 41, S197–S239. <https://doi.org/10.1139/apnm-2015-0663>

Pratt, M., Varela, A.R., Salvo, D., Iii, H.W.K., Ding, D., 2020. Attacking the pandemic of physical inactivity: what is holding us back? *Br J Sports Med* 54, 760–762. <https://doi.org/10.1136/bjsports-2019-101392>

Qu, P., Luo, M., Wu, Y., Zhang, F., Vos, H., Gu, X., Mi, Y., Luo, X., Jia, P., 2021. Association between neighborhood aesthetics and childhood obesity. *Obes Rev* 22, e13079. <https://doi.org/10.1111/obr.13079>

- Ravishankara, A.R., David, L.M., Pierce, J.R., Venkataraman, C., 2020. Outdoor air pollution in India is not only an urban problem. *Proc Natl Acad Sci U S A* 117, 28640–28644.
<https://doi.org/10.1073/pnas.2007236117>
- Rees-Punia, E., Hathaway, E.D., Gay, J.L., 2018. Crime, perceived safety, and physical activity: A meta-analysis. *Preventive Medicine* 111, 307–313.
<https://doi.org/10.1016/j.ypmed.2017.11.017>
- RStudio Team, 2015. RStudio: Integrated Development Environment for R.
- Saint-Maurice, P.F., Troiano, R.P., Berrigan, D., Kraus, W.E., Matthews, C.E., 2018. Volume of Light Versus Moderate-to Vigorous Physical Activity: Similar Benefits for All-Cause Mortality? *J Am Heart Assoc* 7, e008815. <https://doi.org/10.1161/JAHA.118.008815>
- Schoeppe, S., Duncan, M.J., Badland, H., Oliver, M., Curtis, C., 2013. Associations of children’s independent mobility and active travel with physical activity, sedentary behaviour and weight status: A systematic review. *Journal of Science and Medicine in Sport* 16, 312–319. <https://doi.org/10.1016/j.jsams.2012.11.001>
- Sciubba, J.D., 2023. What India Becoming the World’s Most Populous Country Means.
- Singh, R., Singh, V., Gautam, A.S., Gautam, S., Sharma, M., Soni, P.S., Singh, K., Gautam, A., 2023. Temporal and Spatial Variations of Satellite-Based Aerosol Optical Depths, Angstrom Exponent, Single Scattering Albedo, and Ultraviolet-Aerosol Index over Five Polluted and Less-Polluted Cities of Northern India: Impact of Urbanization and Climate Change. *Aerosol Sci Eng* 7, 131–149. <https://doi.org/10.1007/s41810-022-00168-z>
- Singh, V., Singh, S., Biswal, A., 2021. Exceedances and trends of particulate matter (PM_{2.5}) in five Indian megacities. *Science of The Total Environment* 750, 141461.
<https://doi.org/10.1016/j.scitotenv.2020.141461>

- Sprengeler, O., Pohlabein, H., Bammann, K., Buck, C., Lauria, F., Verbestel, V., Eiben, G., Konstabel, K., Molnár, D., Moreno, L.A., Pitsiladis, Y., Page, A., Reisch, L., Tornaritis, M., Ahrens, W., 2021. Trajectories of objectively measured physical activity and childhood overweight: longitudinal analysis of the IDEFICS/I.Family cohort. *International Journal of Behavioral Nutrition and Physical Activity* 18, 103. <https://doi.org/10.1186/s12966-021-01171-2>
- Sunyer, J., Esnaola, M., Alvarez-Pedrerol, M., Forn, J., Rivas, I., López-Vicente, M., Suades-González, E., Foraster, M., Garcia-Esteban, R., Basagaña, X., Viana, M., Cirach, M., Moreno, T., Alastuey, A., Sebastian-Galles, N., Nieuwenhuijsen, M., Querol, X., 2015. Association between Traffic-Related Air Pollution in Schools and Cognitive Development in Primary School Children: A Prospective Cohort Study. *PLOS Medicine* 12, e1001792. <https://doi.org/10.1371/journal.pmed.1001792>
- Taylor, L.G., Clark, A.F., Gilliland, J.A., 2018. Context Matters: Examining children's perceived barriers to physical activity across varying Canadian environments. *Health & Place* 54, 221–228. <https://doi.org/10.1016/j.healthplace.2018.10.002>
- Tyagi, M., Raheja, G., 2020. Indian parents' perception of children's independent mobility in urban neighbourhoods: a case study of Delhi. *Children's Geographies* 19, 390–406. <https://doi.org/10.1080/14733285.2020.1797996>
- UN, 2018. Young and old, air pollution affects the most vulnerable [WWW Document]. URL <https://www.unep.org/news-and-stories/blogpost/young-and-old-air-pollution-affects-most-vulnerable> (accessed 8.18.23).

- Upadhyay, A., Dey, S., Chowdhury, S., Goyal, P., 2018. Expected health benefits from mitigation of emissions from major anthropogenic PM2.5 sources in India: Statistics at state level. *Environmental Pollution* 242, 1817–1826. <https://doi.org/10.1016/j.envpol.2018.07.085>
- Van Oldenborgh, G.J., Philip, S., Kew, S., Van Weele, M., Uhe, P., Otto, F., Singh, R., Pai, I., Cullen, H., AchutaRao, K., 2018. Extreme heat in India and anthropogenic climate change. *Nat. Hazards Earth Syst. Sci.* 18, 365–381. <https://doi.org/10.5194/nhess-18-365-2018>
- Vieira, D., Gomes, E.C., Negrão, Â.S., Thuany, M., Gomes, T.N., 2023. Movement Behaviour and Health Outcomes in Rural Children: A Systematic Review. *Int J Environ Res Public Health* 20, 2514. <https://doi.org/10.3390/ijerph20032514>
- Vispute, S., Mandlik, R., Author, Gondhalekar, K., Patwardhan, V., Khadilkar, V., 2023. Influence of ecoregional and lifestyle factors on growth and body composition of Indian children and adolescents aged 9–18 years—A multicenter study. *American Journal of Human Biology* 35, e23850. <https://doi.org/10.1002/ajhb.23850>
- Wang, Y., He, G., Ma, K., Li, D., Wang, C., 2022. Preschool Children’s Physical Activity and Community Environment: A Cross-Sectional Study of Two Cities in China. *Int J Environ Res Public Health* 19, 14797. <https://doi.org/10.3390/ijerph192214797>
- Waters, A.R., Warner, E.L., Vaca Lopez, P.L., Kirchhoff, A.C., Ou, J.Y., 2021. Perceptions and knowledge of air pollution and its health effects among caregivers of childhood cancer survivors: a qualitative study. *BMC Cancer* 21, 1070. <https://doi.org/10.1186/s12885-021-08739-y>

WHO, 2023. WHO Ambient Air Quality Database [WWW Document]. URL

[https://www.who.int/publications/m/item/who-ambient-air-quality-database-\(update-2023\)](https://www.who.int/publications/m/item/who-ambient-air-quality-database-(update-2023)) (accessed 8.18.23).

WHO, 2012. Physical inactivity [WWW Document]. URL

<https://www.who.int/data/gho/indicator-metadata-registry/imr-details/3416#> (accessed 7.31.23).

World Health Organization, 2009. 2008-2013 action plan for the global strategy for the prevention and control of noncommunicable diseases□: prevent and control cardiovascular diseases, cancers, chronic respiratory diseases and diabetes. Plan d'action 2008-2013 pour la stratégie mondiale de lutte contre les maladies non transmissibles 42.

Yang, J., Shao, M., 2021. Impacts of Extreme Air Pollution Meteorology on Air Quality in

China. *Journal of Geophysical Research: Atmospheres* 126, e2020JD033210.

<https://doi.org/10.1029/2020JD033210>

Yang, T., 2020. Association between perceived environmental pollution and health among urban and rural residents-a Chinese national study. *BMC Public Health* 20, 194.

<https://doi.org/10.1186/s12889-020-8204-0>

Yu, H., Zhang, H., 2023. Impact of ambient air pollution on physical activity and sedentary

behavior in children. *BMC Public Health* 23, 357. <https://doi.org/10.1186/s12889-023-15269-8>