#### "Emotional stress is more detrimental than the virus itself": Towards an understanding of

#### HIV testing and pre-exposure prophylaxis (PrEP) use among internal migrant men in **South Africa**

- Maria F. Nardell,<sup>1,2 §</sup> Caroline Govathson,<sup>3,4</sup> Salomé Garnier,<sup>5</sup> Ashley Watts,<sup>6</sup> Dolapo Babalola,<sup>7</sup> Nkosinathi Ngcobo,<sup>3,4</sup> Lawrence Long,<sup>3,4,8</sup> Mark N. Lurie,<sup>9</sup> Jacqui Miot,<sup>3,4</sup> Sophie Pascoe,\*<sup>3,4</sup> Ingrid T.
- Katz\*2,10
- <sup>1</sup>Division of Global Health Equity, Brigham and Women's Hospital, Boston, MA, USA
- <sup>2</sup>Harvard Medical School, Boston, MA, USA
- <sup>3</sup>Health Economics and Epidemiology Research Office (HE<sup>2</sup>RO), Johannesburg, South Africa
- <sup>4</sup>Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa
- <sup>5</sup>EHESP French School of Public Health, Paris, France
- <sup>6</sup>Harvard University, Boston, MA, USA
- <sup>7</sup>College of Medicine, University of Ibadan, Nigeria
- <sup>8</sup>Department of Global Health, Boston University School of Public Health, Boston, MA, USA
- <sup>9</sup>Brown University School of Public Health, Brown University, Providence, Rhode Island, USA
- <sup>10</sup>Division of Women's Health, Brigham and Women's Hospital, Boston, MA, USA
- \*Equal contribution

<sup>§</sup>Corresponding author: Maria Francesca Nardell 

### 41 Abstract

42

# 43 Introduction

44 South Africa has one of the highest rates of internal migration on the continent, largely

45 comprised of men seeking labor in urban centers. South African men who move within the

46 country (internal migrants) are at higher risk than non-migrant men of acquiring HIV yet are less

47 likely to test or use pre-exposure prophylaxis (PrEP). However, little is known about the

48 mechanisms that link internal migration and challenges engaging in HIV services.

49

# 50 Methods

51 We recruited 30 internal migrant men (born outside Gauteng Province) for in-depth qualitative

52 interviews at sites in Johannesburg (Gauteng) where migrants may gather (i.e., factories,

53 homeless shelters). Interviewers used open-ended questions, based in the Theory of Triadic

54 Influence, to explore experiences and challenges with HIV testing and/or PrEP. A mixed

55 deductive inductive content analytic approach was used to review data and explain why

56 participants may or may not use these services.

57

# 58 **Results**

59 Migrant men come to Johannesburg to find work, but their struggle to survive without reliable 60 income causes daily stress. Stress and time constraints limit their availability to seek health 61 services, and many men lack knowledge about the opportunity for PrEP should they test 62 negative. In addition, migrants must also adjust to life in Johannesburg, where they may be 63 unfamiliar with where to access HIV services and lack social support to help them do so. 64 Migrants may also continue to travel intermittently for work or social purposes, which can make 65 it hard to take a daily pill like PrEP. Yet Johannesburg also presents opportunities for HIV 66 services for migrant men, especially those who perceive greater availability and anonymity of 67 HIV information and services in the city as compared to their rural homes of origin.

68

# 69 Conclusions

70 Bringing HIV services to migrant men at community sites may ease the burden of accessing

71 these services. Including PrEP counseling and services alongside HIV testing may further

- real encourage men to test, particularly if integrated into counseling for livelihood and coping
- rategies, as well as support for navigating health services in Johannesburg.

74

# 76 Introduction

77

Despite substantial progress, South Africa continues to have one of the world's largest HIV 78 epidemics, with approximately 7.5 million people living with HIV.<sup>1</sup> While women shoulder a 79 greater burden of HIV than men in South Africa,<sup>2</sup> there are major gaps in reaching men with HIV 80 services.<sup>3–6</sup> In particular, it is estimated that only half of men living with HIV in South Africa 81 know their status,<sup>3,7</sup> and levels of HIV testing for men remain consistently lower than those for 82 women.<sup>8–10</sup> Men with HIV also have lower rates of treatment adherence, viral suppression, and 83 survival.<sup>3,11,12</sup> Alongside efforts to help men living with HIV to start and stay on treatment, it is 84 imperative to focus on testing and prevention for men not already infected.<sup>13</sup> Oral pre-exposure 85 prophylaxis (PrEP) is highly effective at preventing HIV.<sup>14–17</sup> South African men have shown 86 interest in using PrEP,<sup>18,19</sup> but barriers to men's use of HIV services include masculine norms 87 which discourage care seeking, anticipated stigma in health facilities, poverty and 88 unemployment, and lack of knowledge about PrEP.<sup>3,20</sup> In addition, most PrEP services in South 89 Africa have targeted women, men who have sex with men, and sex workers.<sup>21–24</sup> Concerns about 90 the cost-effectiveness of providing PrEP to heterosexual men may limit its implementation.<sup>25</sup> 91 92 Despite this, South Africa's National Strategic Plan for HIV, TB, and STIs 2017–2022 recommends PrEP for anyone who reports being at risk for HIV, including heterosexual men.<sup>26</sup> 93 94 Thus, it is important to understand how to target PrEP delivery to men for whom it may provide 95 greatest benefit.

96

South African men who move within the country, called internal migrants,<sup>27,28</sup> bear as much as 97 two to three times the burden of HIV as compared to non-migrant men in South Africa,<sup>29,30</sup> as 98 well as a higher burden than international migrants.<sup>30</sup> Associations between HIV vulnerability 99 and migration are complex and vary among migrant groups.<sup>31</sup> Some studies have shown that 100 migrant men are at higher risk for HIV acquisition during the disruptive process of relocation<sup>32</sup> 101 and engage in higher sexual risk behaviors.<sup>33</sup> Other research has shown that men living with HIV 102 103 are more likely to become migrants than those who are HIV negative, in part due to marital dissolution, which people living with HIV are more likely to experience.<sup>27</sup> Compared to non-104 migrants, internal migrants are less likely to use health services overall, more likely to use 105 private services or traditional healers, and less likely to test for HIV.<sup>34,35</sup> Migrants are identified 106

as a "key population" for HIV prevention,<sup>31</sup> but only 15% internal migrant men surveyed in 107 community sites in Johannesburg were aware of PrEP.<sup>30</sup> HIV interventions that seek to engage 108 men in testing and prevention, such as workplace programs and partner notification strategies,<sup>36</sup> 109 may exclude men who are unemployed or unpartnered, as is common for migrant men.<sup>30</sup> 110 111 Moreover, large trials to improve HIV testing uptake among men in South Africa have found substantial challenges reaching men in communities with high mobility and in-migration.<sup>37</sup> South 112 Africa has one of the highest rates of internal migration on the continent<sup>38</sup> with an estimated net 113 migration of a million internal migrants settling in urban Gauteng Province from 2016 to 2021 114 alone, dwarfing the number of international immigrants by as many as five to one.<sup>39</sup> 115 116 117 Research to unpack the relationship between migration, masculinity, and barriers to HIV testing

118 and prevention in South Africa has been limited. In particular, there is a lack of research on 119 internal migrants' experience with HIV care, for whom factors such as xenophobia within the 120 healthcare system and lack of knowledge of the South African health care system may not 121 apply.<sup>40–42</sup> In this paper, we seek to fill this gap by exploring the psychosocial and structural factors that influence HIV testing and prevention uptake among internal migrants through 122 123 qualitative interviews with internal migrant men in Johannesburg, South Africa. South Africa 124 must engage this key population if it is to meet its national goal of eliminating HIV as a public 125 health threat with "nobody left behind."<sup>43</sup>

126

#### 127 Methods

128

130

129 Study Setting and Participants

131 This study is based in Johannesburg, the most populous city in South Africa. In our team's prior 132 pilot work among migrant men in Johannesburg, we chose community-based recruitment sites in 133 the Hillbrow, Midrand, Woodmead and Roodeport areas following a period of observation and 134 discussion with community stakeholders. For this study focusing specifically on internal migrant 135 men, we selected two sites where there were high proportions of internal migrant men, a site in 136 Midrand area north of Johannesburg close to factories and the Displaced Persons Unit (DPU) in Hillbrow (Johannesburg central.) Similar to our prior research among this population,<sup>30,44,45</sup> we 137 chose to focus on community sites in order to reach men who may not seek care at clinics. 138

#### 139

Recruitment occurred in August 2022. Team members recruited participants by first asking leadership at the sites if the study team could recruit there on specific days. On site, the research assistants (RA) distributed study flyers. These flyers gave information about the study and invited men to contact the study team if they were interested in participating. Those men who made contact were then given further information and taken through the consent process inperson. The team checked telephone numbers, names, and identifiers to ensure that the same participant did not enroll twice.

147

# 148 Eligibility and Sampling

149 We included men who were 18 years or older, self-reported as an internal migrant (defined as a 150 South African born outside of Gauteng Province), were willing to participate in a 45 to 60-151 minute interview, and able to provide informed consent. Men were excluded from the study if 152 they were believed to be intoxicated at the time of consent (in which case, they were invited to 153 return the following day), unable to understand the study information, or had previously enrolled 154 in the study. In addition to these eligibility criteria, we purposively sampled to seek perspectives 155 both from migrants who reported living in Johannesburg for a year or less, and those who lived 156 in Johannesburg for longer durations.

157

# 158 Study Design Conceptual Framework

159 Interviews followed an in-depth semi-structured interview guide developed to explore internal 160 migrant men's experiences, knowledge, and attitudes towards HIV testing and prevention. The 161 development of our guide was informed by the Theory of Triadic Influence (TTI), a theoretical 162 model of health behavior used in several HIV-related studies, including in South Africa and by our team previously.<sup>46–48</sup> TTI posits that there are three streams of influence on health-seeking 163 164 behavior: individual factors (e.g., personal motivations, knowledge), social factors (e.g., 165 interpersonal relationships, social norms), and *structural factors* (e.g., healthcare system 166 structures and policies). Interview questions examined attitudes and behaviors toward HIV testing and PrEP within each of these streams of influence, such as knowledge of PrEP 167 168 (individual factors), anticipated HIV stigma (social factors), and access to healthcare services (structural factors). 169

# 170 Data Collection and Preparation

171 Qualitative interviews were completed in person by one of three RAs in a private, secure setting 172 in August 2022. (See Interview guides in Appendix.) RAs were trained in qualitative 173 interviewing and fluent in isiZulu, isiXhosa, English and seSotho so as to be able to conduct the 174 interview in the participant's preferred language. Interviews lasted 45 to 60 minutes. After 175 completing the survey, participants were reimbursed for their time with a ZAR150 (~USD10) 176 electronic shopping voucher, sent directly to the participant's cell phone number. The RA 177 conducting the interview recorded it and took notes. All interview recordings were sent to a 178 third-party transcription service who transcribed it, and if in a language other than English, translated it. Each interview was reviewed by the RA who conducted the interview to ensure 179 180 accurate transcription and translation. Transcripts were assigned a unique study identification number to protect participant confidentiality. 181 182

### 183 Data Analysis

184 Analysis of data aimed to characterize participants' reasons for engaging or not engaging with HIV testing and PrEP services. We used a mixed deductive and inductive approach informed by 185 grounded theory.<sup>49</sup> Two team members (MFN, SG) developed an initial codebook based on key 186 187 concepts from TTI and the interview protocol, as well as emergent codes from a review of the 188 initial five interviews. The coding team (MFN, SG, DB, AW) then discussed emerging themes 189 through repeated review of the transcripts and memoing. Through weekly meetings, the coding 190 team further refined the codebook, resolving discrepancies through discussion, maintaining an audit trail of notes and each codebook version. Interviews were coded in Dedoose.<sup>50</sup> Each coding 191 192 team member coded the first 20% of transcripts, and interrater reliability testing resulted in a 193 pooled Cohen's kappa of 0.80. The four coders individually coded the remaining transcripts. We 194 then reviewed excerpts for each code, identifying repeated patterns of content which formed the basis for broader themes.<sup>49</sup> 195

196

# 197 *Ethical Approval*

198 The ethics committees at the University of the Witwatersrand (M191068), Mass General

199 Brigham (Harvard University) (2020P002251), and Boston University (H-40529) approved the

study. All participants provided written informed consent.

# 201 **Results**

- 202
- 203 Participant Characteristics

204 We randomly recruited 30 participants for this study, half at the Midrand factories site and half at

the Hillbrow homeless shelter. The demographic information for these participants is shown in

Table 1. Nearly a third (30%) of participants were born in Limpopo Province, followed by nearly

- 207 a fifth (16.7%) in KwaZulu-Natal. Almost half (46.7%) of participants spoke isiZulu as their
- 208 primary language and came to Johannesburg looking for work. Most reported no regular intimate
- 209 or married partner (66.7%), though half (50%) had children. Nearly half (43.3%) of participants
- 210 reported living in Johannesburg for less than three years, and nearly a fourth (23.3%) for less
- than six months. Half (50.0%) of participants reported traveling to see family. Most participants
- 212 had no regular form of employment. One participant reported living with HIV.

	Median	Kange			
Age	30	19-45		Number	Percentage
			Time in Johannesburg		
	Number	Percentage	<6 months	7	23.3
Site of recruitment			6 months – 3 years	6	20.0
Factories	15	50.0	>3 years	10	33.3
Shelter	15	50.0	No response	7	23.3
South African province of birth			Employment		
Limpopo	9	30.0	Regular employment	3	10.0
Kwazulu-Natal	5	16.7	No regular employment	23	76.7
Eastern Cape	4	13.3	No response	4	13.3
Mpumalanga	4	13.3	Travel since being in Johannesburg		
Western Cape	2	6.7	Travel to see family for holidays	6	20.0
Free State	2	6.7	Unspecified travel to see family	9	30.0
North West	0	0	No reported travel	10	33.3
Northern Cape	0	0	No response	11	36.7
No response	4	13.3	Partner		
Primary language			Yes	9	30.0
Zulu	14	46.7	No	20	66.7
Sisotho	10	33.3	No response	1	3.3
Xhosa	4	13.3	Children		
Setswana	4	13.3	Yes	15	50.0
Sepedi	4	13.3	No	13	43.3
Reason for coming to Johannesburg			No response	2	6.7
Looking for work	14	46.7	Self-reported HIV status		
Unspecified opportunity	7	23.3	Negative or unknown	29	96.7
Family reason	2	6.7	Positive	1	3.3
Problems in home of origin	2	6.7			
No response	5	16.7			

Table 1.	Participant	Characteristics.	Socio-demo	graphic cha	aracteristics of	of 30 parti	cipants.
			-				

- 213
- 214
- \_\_\_\_
- 215

216

# 217 *Qualitative Results*

We organized themes into barriers to and opportunities for care engagement. Within each of these sections, we identified and organized factors at individual, social, and structural levels. Within each of these levels, we identified factors that apply to men more generally as well as factors specific to migrants.

222

# 223 Individual Barriers

224 Participants framed their decisions around HIV testing within the context of the high levels of 225 stress they already experience on a day-to-day basis. As shown in Table 2, migrant men 226 described a "weight" over their lives "that is really heavy from problems and life situations," 227 including "depressing thoughts" about finding work. Adding to their stress was their awareness 228 of HIV risk from unprotected sexual encounters. For most participants, this risk awareness was 229 not a motivator to seek HIV testing. Rather, HIV testing was more often perceived as an added 230 burden because of fear that a positive result "might be too stressful for them." While almost all 231 participants were knowledgeable about the fact that ART can manage HIV, they nonetheless 232 described the possibility of a positive diagnosis in dire terms, including "about to die," "ruin my 233 life," "like suicide," "painful," and "make [me] more sick." As one participant described, "We as 234 men are already carrying a lot. To add another weight on top of that... Emotional stress is more 235 detrimental than the virus itself." This burden was further exacerbated for many participants by 236 virtue of their move to Johannesburg. Migrant men described challenges with navigating a new city, finding health services, and sometimes personal and family challenges that preceded their 237 238 move to Johannesburg, which further taxed their cognitive and emotional bandwidth.

239

Complicating this narrative was limited awareness about PrEP among most (60%) participants.
Even those who had heard of it had little knowledge of anyone who had used it before, where to
find it, or how to use it. Participants also mentioned that awareness of HIV prevention options
was particularly low in rural areas of the country, where many migrants had been raised. Given
most participants did not know PrEP would be an option if they tested negative, HIV testing was
associated with few upsides. Thus, fears of testing positive outweighed the potential for relief if
they tested negative. While all participants reported testing for HIV at least once in their life,

- 247 most tested only when required by a healthcare provider, rather than for prevention. The
- 248 interviewer's mention of PrEP was usually met with interest, with several respondents
- 249 mentioning how they would trust PrEP if it came from healthcare professionals and after they
- 250 "see it works for certain individuals."

### Table 2. Individual-level barriers to HIV testing and PrEP use

Barriers	Examples
Challenges coping with life stressors	"Naturally men have that weight over them that is really heavy from problems and life situations. We know that it is in our nature to always make mistakes. Knowing that I have to test is going to put on more weight on me. I feel it will be adding to the problems to go and test. It is going to give me depression to find out that I am HIV positive. It is not that we do not want to test. It's just that we do not want to know at that certain time." (Migrant from Western Cape)
Fears of testing HIV positive	"People are scared to test. They will rather live in fear not knowing then to find out they are about to die." (Migrant from Limpopo)
Lack awareness or knowledge of PrEP	"I do not know about [PrEP]. I have only heard rumours. I have never come across someone who have taken them." (Migrant from Mpumalanga)
Seek healthcare for symptoms	"When I need health care I go to a public clinic because it is free. I wait and see if the sickness lasts for 48 hours then I go to the clinic. Anything that lasts for less than that I can handle." (Migrant from Mpumalanga)

251 252

# 253 *Social Barriers*

254 Gender norms played a large role in influencing participants' beliefs and behaviors around HIV 255 testing and PrEP. Participants described how HIV testing was felt to be a woman's 256 responsibility, and a female partner's status was assumed to reflect their own status as well. As 257 shown in Table 3, most participants reported that clinics are designed to serve women's needs, 258 and the predominance of female health care providers in clinics could be a disincentive to 259 seeking care. Some migrants described how traditional gender norms were particularly common 260 in their rural homes of origin, and it was hard to shift this perspective in Johannesburg. In 261 addition, social gatherings with other men, particularly involving alcohol, emerged as a barrier to 262 PrEP use. Participants described how men "are most vulnerable" when they drink because it distracts them from taking pills. Drinking also increases their HIV risk because men are less 263 264 likely to use condoms when intoxicated. 265

266 Anticipated stigma was a major concern reported by nearly all men, including stigma from clinic 267 staff and within communities. Although most participants reported that they themselves were 268 comfortable with others having HIV, they nonetheless worried that testing positive would 269 negatively affect their relationship with partners, family, or friends, who "discourage each other 270 from testing." PrEP use posed additional concerns, including the fear that PrEP would be 271 mistaken for ART or signal infidelity. In addition, a few participants who knew of PrEP 272 mentioned that PrEP is associated with men who have sex with men (MSM), which prompted 273 some men "to shy away from using it." Although this study focused on migrants within South 274 Africa, one participant described himself as "a foreigner" within Johannesburg, even though he 275 had been in the city for 12 years, which raised concerns for him about how he would be treated 276 in healthcare spaces. Another participant felt that travel and being a newcomer in Johannesburg 277 for less than a year increased the risk for being "misunderstood" by people who did not know 278 him.

279

Barriers	Examples
Social gender norms	"Women do not have such problems because the clinics are created for them. Women bear children. So already, they are used to the clinic. Women have many illnesses. I think it is women that transfer infections to men. Maybe going to a men's clinic would be better. It is quite a challenge to be seen by women going to the clinic." (Migrant from Western Cape)
Social alcohol use	"It is really difficult to help men. Most men like to drinkBeing drunk can make them miss the pill if it to be taken daily." (Migrant from KwaZulu-Natal)
Anticipated stigma	
Testing HIV positive	"You worry that when you go for an HIV test your results might be different and that will affect the relationship negatively. I think that knowing your status puts a lot of pressure on you." (Migrant from Eastern Cape)
PrEP	"I think the problem might be to take [PrEP] pills in front of other people." (Migrant from Eastern Cape)
Travel and migration	"Different people, different places. You must be prepared to be misunderstood [when traveling]." [Migrant from Free State]

#### Table 3. Social-level barriers to HIV testing and PrEP use

280

281 *Structural Barriers* 

- 282 Structural factors emerged as the largest barrier to HIV testing and PrEP use for migrant men.
- Finding and maintaining work was the biggest priority for all participants. Some managed to find
- intermittent "piece jobs," i.e., unpredictable day jobs in which men are paid by the hour or day

285 for the work that is done. Others described how their jobs were unexpectedly terminated or how 286 their pay was insufficient to afford housing or food. Given pressures to work, time was a major 287 barrier to HIV testing, especially at clinics where participants reported queuing all day. As 288 shown in Table 4, one participant said, "I have to spend the entire day on the street...waiting for 289 an opportunity for piece work. If I go for testing, I might miss that opportunity." Many migrants 290 reported coming to Johannesburg given its reputation as the "City of Gold" where "opportunities 291 are broad." For them, the gap between their dreams and reality was especially stark. Participants 292 reported that work was not just a means of survival but the necessary precursor to "a home, wife, 293 and kids...a dream [which] has not materialized." On the other hand, for those participants with 294 a family, the immediate need to provide for their children emerged as a higher priority than the 295 time or transportation costs involved for HIV testing.

296

297 On top of their livelihood concerns, migrants described challenges adjusting to life in

298 Johannesburg, including concerns about the fast pace of the city and their own safety.

299 Participants also described how migrants may be unfamiliar with where HIV services are

300 located, lack community support to access health services, or not trust services from people they

301 don't know. Migrants reported that travel to other parts of the city or country could increase HIV

risk as they were more likely to have unprotected sexual encounters during these times. None of

the participants used PrEP currently, though some hypothesized that traveling, including

304 overnight travel for piece jobs, might pose a challenge for taking a daily pill. Another participant

also mentioned how "illegal immigrants" have difficulty accessing healthcare in South Africa.

Barriers	Examples
Employment opportunity costs	"I have to spend the entire day on the street by the robot waiting for an opportunity for piece-work. If I go for testing I might miss that opportunity." (Migrant from KwaZulu-Natal)
Traveling for work	"Because I am always on the go. Even with the peace-jobs sometimes I don't sleep at home. I am concerned that I might miss a pill." (Migrant from Limpopo)
Adjustment to Johannesburg	"I am still new in Johannesburg. It is a busy city. It is full of criminals. You cannot even ask people for directions to the clinic. Sometimes health facilities are located down town where we cannot go because it is dangerous. Those are the challenges." (Migrant from Mpumalanga)
Lack of residency documentation	"The issue of illegal immigrants is a serious problem in South Africa. We have many illegal immigrants in the country. You find people sick they cannot access health services because they are here illegally. I feel that clinics should be open even to illegal immigrants. If a person has a passport, they should be able to access health services." (Migrant from Limpopo)

# Table 4. Structural-level barriers to HIV testing and PrEP use

307

## 308 *Opportunities*

309 As shown in Table 5, participants described several opportunities for engagement in HIV testing,

310 prevention, and care, including those that relate to many of the themes above. Despite the

311 challenges of daily life in Johannesburg, participants noted that HIV services in community-

based settings such as pharmacies, pop-up tents, and shops made access easier. Most appreciated

the sense of anonymity they felt seeking healthcare in Johannesburg as compared to rural areas,

314 where they had more concern for unintentional disclosure from clinic staff or other patients who

knew them. In fact, some participants reported that if diagnosed with HIV, they would feel most

316 comfortable seeking support from healthcare staff in Johannesburg. More readily available HIV

317 information in Johannesburg was also perceived as an opportunity to be harnessed to promote

318 greater care engagement. As one participant shared, "I think people need to be informed about

319 PrEP. This will encourage more people to do HIV testing so they can use PrEP if their results are

320 negative."

321

322 Social support from other men surfaced as an important resource for some participants, including

323 through religious organizations, bars, and sporting activities. At the homeless shelter,

324 participants described how ten men in a room together would "share stories every night before

325 we go to bed," while others had support from family members or friends already in

Johannesburg. In contrast to participants who described their sense of masculinity as a deterrent to care seeking (e.g., considering their wife's HIV status as a proxy for their own), for other men, their sense of their role as a man motivated HIV testing and prevention. For example, men described a sense of responsibility for their family: "You don't want to die and leave your family behind." Others drew on their sense of masculinity or felt compelled to test for the sake of their partner: "I am the man and I have to protect my partner." Lastly, some participants described positive coping skills which help them to feel "courage to deal with whatever results" from HIV testing.

# Table 5. Opportunities for HIV testing and PrEP use

Opportunities	Examples		
Differentiated service delivery	"I would be comfortable accessing PrEP even outside the clinic, in pharmacies and shops. That will make it even more accessible to people." (Migrant from Eastern Cape)		
Anonymity	"Yes I am comfortable [testing for HIV] in Johannesburg and free. There are no chances of meeting somebody I know." (Migrant from Limpopo)		
PrEP information	"PrEP need to be advertised more. It needs to get out there." (Migrant from KwaZulu-Natal)		
Social support	"At the shelter there is tons of us. There are ladies and guys. There is six of us in a room. They are my new family." (Migrant from Limpopo)		
Masculinity	"I sit down and talk to myself and tell myself that I am a man and that I can face any challenge." (Migrant from Limpopo)		
Positive coping skills	"In the past I was very scared to test. But I have matured and I always want to go and test for HIV. When I go I come back happy. I always expect anything and I have courage to deal with whatever results I get. I think I am comforted by the fact that HIV can be managed by treatment." (Migrant from KwaZulu-Natal)		
Discussion			
Our findings of barri	iers to and opportunities for HIV testing and PrEP for internal migrant men		
in Johannesburg reve	ealed themes common to men generally as well as themes exacerbated or		
unique to men who i	nigrate internally within South Africa across all levels of influence		
(individual, social, structural), as shown in Figure 1. Similar to prior research, we found that			
substantial fears of a	In HIV-positive diagnosis <sup>47–49</sup> and anticipated stigma <sup>53,54</sup> remain major		

344 barriers to seeking HIV testing for men. Building on this research, our data highlight that men 345 worry that a positive HIV diagnosis on top of pre-existing livelihood stressors would overwhelm 346 their ability to cope and thus their mental wellbeing. Unemployment rates in South Africa are greater than 30%,<sup>55</sup> and studies have shown the stressful toll that unemployment takes on South 347 348 African men, leading not only to financial strain but also lower self-esteem and decreased social status.<sup>56</sup> This suggests the need for interventions for men which address both livelihood stressors 349 350 and coping skills. For example, a livelihood support intervention among persons living with HIV in Kenya showed evidence of mental health improvements,<sup>57</sup> and the MenStar Strategy to engage 351 men in HIV care in South Africa includes a focus on building coping potential.<sup>58</sup> However, 352 353 problematic alcohol use among men – prevalent in resource-constrained settings in South Africa and often the foundation of social gatherings $^{59}$  – further complicates this picture as alcohol use 354 may be associated with HIV risk behaviors.<sup>60</sup> Efforts to address alcohol use and HIV risk have 355 356 had mixed success, with evidence showing that social barriers (e.g., social norms around 357 substance use) and structural level barriers (e.g., unemployment, poverty, violence, limited clinic 358 resources) constrain long-lasting behavior change.<sup>61,62</sup>

359

360 Building on these barriers for men, our data uncovered challenges that are exacerbated for and/or 361 unique to South African migrants. Migrants may have a higher likelihood of life stressors, given 362 the conditions that prompted their move as well as their near-universal need to find employment. 363 Migrants are also faced with numerous additional social and structural stressors by virtue of 364 needing to start afresh in a new city, including securing housing, navigating a new health system, 365 building a new social network, prioritizing physical safety, and adjusting to the fast pace of life 366 in Johannesburg. Each stressor takes up some degree of an individual's "cognitive bandwidth," 367 which encompasses both cognitive capacity (psychological mechanisms which allow for 368 reasoning, problem solving, and retaining information) and executive control (ability to plan, manage attention, and control actions).<sup>63</sup> Each individual has a finite amount of cognitive 369 370 bandwidth. An accumulation of stressors can tax this bandwidth and impede cognitive 371 functioning, making it more difficult to manage immediate stressors, let alone to process new information (e.g., about PrEP) or prioritize HIV testing or preventative care.<sup>63–65</sup> Similarly, 372 373 research among international migrants has found that high levels of unemployment and 374 competing life priorities, especially in the early phase of resettlement, can make it harder for

- 375 migrants to seek preventative care.<sup>66</sup> In addition to interventions focusing on livelihood and
- 376 coping strategies, interventions which help migrants to adjust to life in their new city and
- 377 navigate available health resources may also facilitate care engagement. For example,
- 378 community navigator health workers in North America have been used to offer culturally
- tailored educational and communication support to guide migrants in overcoming barriers to
- 380 healthcare.<sup>67,68</sup>
- 381

**Figure 1**. Barriers to HIV testing and prevention for men and internal migrant men in Johannesburg. In the yellow oval in the figure below, we show themes found to be common to men more generally, including migrants, organized by individual, social and structural levels of influence. In addition, we found themes which were heightened for migrant men, shown in the pink oval.



382

- 383
- 384

Despite these challenges, our data identified opportunities to engage migrants in HIV care, particularly for new arrivals to Johannesburg. As is the case in many countries worldwide, rural communities in South Africa often have less access to healthcare as compared to access in urban centers,<sup>69</sup> and fears of unintentional disclosure may be heightened in rural clinics where seeking care means being seen by known community members.<sup>70</sup> As our data showed, Johannesburg offers migrants greater opportunities to not only seek convenient, anonymous care but also to learn about HIV prevention options, such as PrEP. Similarly, research among international

migrants has shown that many first learn about PrEP in their destination countries.<sup>66</sup> In addition, 392 393 migration may lessen the influence of family and culture norms from one's home of origin, e.g. social constructs of gender which may limit men's care-seeking.<sup>53</sup> While some of our 394 395 participants reported either social isolation or relying on pre-existing family members in 396 Johannesburg, others described information sharing and new bonds that formed between 397 themselves and other migrant men who were new to the city, especially for participants in the 398 shelter. Thus, particularly for newcomers to Johannesburg, there may be an opportunity to 399 provide easily accessible HIV services to migrant men at a time when they may be less constrained by prior social obligations and freer to adopt new social norms and behaviors.<sup>71</sup> 400

401

402 This study must be interpreted within limitations of this work. Many participants in our sample 403 were unaware of PrEP, and none reported using it, similar to other research on barriers and facilitators of PrEP uptake for men in sub-Saharan Africa.<sup>20,72,73</sup> Thus, it is possible that PrEP 404 405 may have theoretical appeal in the context of first hearing about it, and social desirability bias 406 may impact participants' enthusiasm for PrEP as part of a research study. Barriers to PrEP use 407 may emerge more prominently for men with experience using it. For example, accessibility 408 issues were found to impede continuation of PrEP among women who previously used it as part of a study in South Africa.<sup>74</sup> In addition, given the nature of this qualitative study and our focus 409 410 on two recruitment sites in Johannesburg, the results cannot be generalized to all migrant 411 populations within or outside of Johannesburg, nor can we statistically analyze differences 412 between different sub-populations, e.g., newcomers versus longer-term migrants. Rather, we 413 aimed to shed light on mechanisms underlying engagement in HIV testing and prevention to 414 deepen our understanding of experiences for a random sample of internal migrants seeking 415 shelter and work in the city. Many of the themes that we uncovered were similar to those found 416 previously for men more generally, suggesting that strategies to engage migrant men (e.g., 417 workplace HIV testing and PrEP availability) may have benefits for other men as well. Future 418 research focusing on international migrants in South Africa is also warranted to understand the factors influencing their HIV care engagement. 419

- 420
- 421
- 422

# 423 Conclusions

425	Migrant men come to Johannesburg to find work, but their struggle to survive without reliable
426	income causes daily stress. Despite caring about their health, constraints on their time and
427	cognitive bandwidth limit their ability to seek health services, particularly given fears of testing
428	positive for HIV, anticipated stigma, and limited knowledge about the opportunity for PrEP
429	should they test negative. Yet Johannesburg also presents opportunities for HIV testing and PrEP
430	use for migrant men who perceive greater availability and anonymity of HIV information and
431	services in the city as compared to rural homes of origin, and also more willingness to adopt new
432	beliefs and behaviors. Bringing HIV services to migrant men at community sites may ease the
433	burden of accessing these services. Including PrEP counseling and services alongside HIV
434	testing may further encourage men to test, particularly if integrated into counseling on livelihood
435	and coping strategies, as well as support for navigating health services in Johannesburg.
436	
437	
438	
439	
440	
441	
442	
443	
444	
445	
446	
447	
448	
449	
450	
451	
452	
453 454	

455 456	Refe	erences
457 458 459	1.	Country Factsheets: South Africa [Internet]. Joint United Nations Programme on HIV/AIDS (UNAIDS). 2021. Available from: https://www.unaids.org/en/regionscountries/countries/southafrica
460 461 462	2.	Human Sciences Research Council (HSRC). South African National HIV Prevalence, Incidence, Behaviour and Communication Survey, 2017.
463 464 465	3.	Nardell MF, Adeoti O, Peters C, Kakuhikire B, Govathson-Mandimika C, Long L, et al. Men missing from the HIV care continuum in sub-Saharan Africa: a meta-analysis and meta-synthesis. J Int AIDS Soc. 2022 Mar;25(3):e25889.
466 467	4.	Tsai AC, Siedner MJ. The Missing Men: HIV Treatment Scale-Up and Life Expectancy in Sub-Saharan Africa. PLOS Med. 2015 Nov 24;12(11):e1001906.
468 469	5.	Cornell M, Cox V, Wilkinson L. Public health blindness towards men in HIV programmes in Africa. Trop Med Int Health. 2015 Dec;20(12):1634–5.
470 471 472	6.	Makusha T, Rooyen H, Cornell M. Reframing the approach to heterosexual men in the HIV epidemic in sub Saharan Africa. J Int AIDS Soc [Internet]. 2020 Jun [cited 2020 Dec 11];23(S2). Available from: https://onlinelibrary.wiley.com/doi/abs/10.1002/jia2.25510
473 474 475 476	7.	Grobler A, Cawood C, Khanyile D, Puren A, Kharsany ABM. Progress of UNAIDS 90-90-90 targets in a district in KwaZulu-Natal, South Africa, with high HIV burden, in the HIPSS study: a household-based complex multilevel community survey. Lancet HIV. 2017 Nov;4(11):e505–13.
477 478 479	8.	Zuma K, Simbayi L, Zungu N, Moyo S, Marinda E, Jooste S, et al. The HIV Epidemic in South Africa: Key Findings from 2017 National Population-Based Survey. Int J Environ Res Public Health. 2022 Jul 1;19(13):8125.
480	9.	A snapshot of men and HIV in South Africa. UNAIDS; 2017.
481 482 483 484 485	10.	Lippman SA, El Ayadi AM, Grignon JS, Puren A, Liegler T, Venter WDF, et al. Improvements in the South African HIV care cascade: findings on 90 90 90 targets from successive population representative surveys in North West Province. J Int AIDS Soc [Internet]. 2019 Jun [cited 2021 Feb 16];22(6). Available from: https://onlinelibrary.wiley.com/doi/10.1002/jia2.25295
486 487 488 489	11.	Cornell M, Schomaker M, Garone DB, Giddy J, Hoffmann CJ, Lessells R, et al. Gender Differences in Survival among Adult Patients Starting Antiretroviral Therapy in South Africa: A Multicentre Cohort Study. Celentano DD, editor. PLoS Med. 2012 Sep 4;9(9):e1001304.
490 491 492	12.	Bor J, Rosen S, Chimbindi N, Haber N, Herbst K, Mutevedzi T, et al. Mass HIV Treatment and Sex Disparities in Life Expectancy: Demographic Surveillance in Rural South Africa. Tsai AC, editor. PLOS Med. 2015 Nov 24;12(11):e1001905.

493 494 495	13.	De Oliveira T, Kharsany ABM, Gräf T, Cawood C, Khanyile D, Grobler A, et al. Transmission networks and risk of HIV infection in KwaZulu-Natal, South Africa: a community-wide phylogenetic study. Lancet HIV. 2017 Jan;4(1):e41–50.
496 497 498	14.	Van Damme L, Corneli A, Ahmed K, Agot K, Lombaard J, Kapiga S, et al. Preexposure Prophylaxis for HIV Infection among African Women. N Engl J Med. 2012 Aug 2;367(5):411–22.
499 500 501	15.	Marrazzo JM, Ramjee G, Richardson BA, Gomez K, Mgodi N, Nair G, et al. Tenofovir- Based Preexposure Prophylaxis for HIV Infection among African Women. N Engl J Med. 2015 Feb 5;372(6):509–18.
	10	

502 16. Grant RM, Lama JR, Anderson PL, McMahan V, Liu AY, Vargas L, et al. Preexposure
503 Chemoprophylaxis for HIV Prevention in Men Who Have Sex with Men. N Engl J Med.
504 2010 Dec 30;363(27):2587–99.

Baeten JM, Donnell D, Ndase P, Mugo NR, Campbell JD, Wangisi J, et al. Antiretroviral
Prophylaxis for HIV Prevention in Heterosexual Men and Women. N Engl J Med. 2012
Aug 2;367(5):399–410.

18. Hannaford A, Lim J, Moll AP, Khoza B, Shenoi SV. 'PrEP should be for men only': Young heterosexual men's views on PrEP in rural South Africa. Glob Public Health. 2020 Sep 1;15(9):1337–48.

511 19. Cheng CY, Quaife M, Eakle R, Cabrera Escobar MA, Vickerman P, Terris-Prestholt F.
512 Determinants of heterosexual men's demand for long-acting injectable pre-exposure
513 prophylaxis (PrEP) for HIV in urban South Africa. BMC Public Health. 2019
514 Dec;19(1):996.

Shamu S, Shamu P, Khupakonke S, Farirai T, Chidarikire T, Guloba G, et al. Pre-exposure
prophylaxis (PrEP) awareness, attitudes and uptake willingness among young people:
gender differences and associated factors in two South African districts. Glob Health
Action. 2021 Jan 1;14(1):1886455.

S19 21. Rousseau E, Bekker LG, Julies RF, Celum C, Morton J, Johnson R, et al. A communitybased mobile clinic model delivering PrEP for HIV prevention to adolescent girls and
young women in Cape Town, South Africa. BMC Health Serv Res. 2021 Dec;21(1):888.

- 522 22. Bekker LG, Rebe K, Venter F, Maartens G, Moorhouse M, Conradie F, et al. Southern
  523 African guidelines on the safe use of pre-exposure prophylaxis in persons at risk of
  524 acquiring HIV-1 infection. South Afr J HIV Med. 2016;17(1):455.
- Pillay D, Stankevitz K, Lanham M, Ridgeway K, Murire M, Briedenhann E, et al. Factors
  influencing uptake, continuation, and discontinuation of oral PrEP among clients at sex
  worker and MSM facilities in South Africa. Galea JT, editor. PLOS ONE. 2020 Apr
  30;15(4):e0228620.

529 24. Emmanuel G, Folayan M, Undelikwe G, Ochonye B, Jayeoba T, Yusuf A, et al. 530 Community perspectives on barriers and challenges to HIV pre-exposure prophylaxis 531 access by men who have sex with men and female sex workers access in Nigeria. BMC 532 Public Health. 2020 Dec;20(1):69. 533 25. Vogelzang M, Terris-Prestholt F, Vickerman P, Delany-Moretlwe S, Travill D, Quaife M. 534 Cost-Effectiveness of HIV Pre-exposure Prophylaxis Among Heterosexual Men in South 535 Africa: A Cost-Utility Modeling Analysis. JAIDS J Acquir Immune Defic Syndr. 2020 Jun 536 1;84(2):173-81. 26. Bekker LG, Brown B, Joseph-Davey D, Gill K, Moorhouse M, Delany-Moretlwe S, et al. 537 538 Southern African guidelines on the safe, easy and effective use of pre-exposure 539 prophylaxis: 2020. South Afr J HIV Med [Internet]. 2020 Dec 10 [cited 2023 May 540 18];21(1). Available from: https://sajhivmed.org.za/index.php/hivmed/article/view/1152 541 27. Anglewicz P, VanLandingham M, Manda-Taylor L, Kohler HP. Cohort profile: internal 542 migration in sub-Saharan Africa—The Migration and Health in Malawi (MHM) study. 543 BMJ Open. 2017 May;7(5):e014799. 544 28. Vearey J. Learning from HIV: Exploring migration and health in South Africa. Glob Public 545 Health. 2012 Jan;7(1):58-70. 546 29. Lurie MN, Williams BG, Zuma K, Mkaya-Mwamburi D, Garnett GP, Sturm AW, et al. The 547 Impact of Migration on HIV-1 Transmission in South Africa: A Study of Migrant and Nonmigrant Men and Their Partners. Sex Transm Dis. 2003 Feb;30(2):149-56. 548 549 30. Nardell MF, Govathson C, Mngadi Ncube S, Ngcobo N, Letswalo D, Long L, et al. 550 Migratory Men and Care Engagement: A Community-Based Study in Johannesburg, South 551 Africa. Oral Presentation presented at: International AIDS Society; 2021. 552 31. Human Mobility and HIV [Internet]. IOM UN Migration. [cited 2023 May 19]. Available 553 from: https://www.iom.int/human-mobility-and-hiv 554 32. Dobra A, Bärnighausen T, Vandormael A, Tanser F. Space-time migration patterns and risk 555 of HIV acquisition in rural South Africa: AIDS. 2017 Jan;31(1):137-45. 556 33. McGrath N, Eaton JW, Newell ML, Hosegood V. Migration, sexual behaviour, and HIV 557 risk: a general population cohort in rural South Africa. Lancet HIV. 2015 Jun;2(6):e252-9. 558 34. Ginsburg C, Collinson MA, Gómez-Olivé FX, Gross M, Harawa S, Lurie MN, et al. 559 Internal migration and health in South Africa: determinants of healthcare utilisation in a 560 young adult cohort. BMC Public Health. 2021 Dec;21(1):554. 561 35. Kranzer K, van Schaik N, Karmue U, Middelkoop K, Sebastian E, Lawn SD, et al. High 562 Prevalence of Self-Reported Undiagnosed HIV despite High Coverage of HIV Testing: A 563 Cross-Sectional Population Based Sero-Survey in South Africa. Landay A, editor. PLoS 564 ONE. 2011 Sep 28;6(9):e25244.

- 36. Beia T, Kielmann K, Diaconu K. Changing men or changing health systems? A scoping
  review of interventions, services and programmes targeting men's health in sub-Saharan
  Africa. Int J Equity Health. 2021 Dec;20(1):87.
- Floyd S, Shanaube K, Yang B, Schaap A, Griffith S, Phiri M, et al. HIV testing and
  treatment coverage achieved after 4 years across 14 urban and peri-urban communities in
  Zambia and South Africa: An analysis of findings from the HPTN 071 (PopART) trial.
  Geng EH, editor. PLOS Med. 2020 Apr 2;17(4):e1003067.
- 38. Bell M, Charles-Edwards E, Ueffing P, Stillwell J, Kupiszewski M, Kupiszewska D.
  Internal Migration and Development: Comparing Migration Intensities Around the World.
  Popul Dev Rev. 2015 Mar;41(1):33–58.
- 39. Mass migration to Johannesburg and Pretoria here's how many South Africans are moving
  [Internet]. Business Tech. 2020. Available from:
- https://businesstech.co.za/news/property/379407/mass-migration-to-johannesburg-and pretoria-heres-how-many-south-africans-are-moving/
- 40. Crush J, Tawodzera G. Medical Xenophobia and Zimbabwean Migrant Access to Public
  Health Services in South Africa. J Ethn Migr Stud. 2014 Apr 3;40(4):655–70.
- 41. Mukumbang FC, Ambe AN, Adebiyi BO. Unspoken inequality: how COVID-19 has
  exacerbated existing vulnerabilities of asylum-seekers, refugees, and undocumented
  migrants in South Africa. Int J Equity Health. 2020 Dec;19(1):141.
- Falcao J, Ahoua L, Zerbe A, di Mattei P, Baggaley R, Chivurre V, et al. Willingness to use
  short-term oral pre-exposure prophylaxis (PrEP) by migrant miners and female partners of
  migrant miners in Mozambique. Cult Health Sex. 2017 Dec 2;19(12):1389–403.
- 587 43. Let Our Actions Count: South Africa's National Stategic Plan for HIV, TB, and STIs 2017588 2022 [Internet]. 2023 May. Available from:
  589 https://www.gov.za/sites/default/files/gcis\_document/201705/nsp-hiv-tb-stia.pdf
- 44. Pascoe S, Govathson C, Mngadi Ncube S, Ngcobo N, Nardell MF, Katz IT, et al. Fast and
  friendly is key to keeping men on HIV treatment! Results from a discrete choice
  experiment to understand men's preferences in Johannesburg, South Africa for HIV
  treatment services. Oral Presentation presented at: International AIDS Society; 2021.
- 45. Pascoe S, Mngadi-Ncube S, Ngcobo N, Govathson C, Letswalo D, Nardell M, et al.
  Understanding HIV and health care utilisation among men in Gauteng, South Africa. In
  2021.
- Flay B, Snyder F, Petraitis J. The Theory of Triadic Influence. In: DiClemente RJ, Crosby
  RA, Kegler MC, editors. Emerging theories in health promotion practice and research. 2nd
  ed. New York: Jossey-Bass; 2009.
- Katz IT, Ngure K, Kamolloh K, Ogello V, Okombo M, Thuo NB, et al. Multi-level Factors
  Driving Pre-exposure Prophylaxis Non-initiation Among Young Women at High Risk for

602 HIV in Kenya. AIDS Behav [Internet]. 2022 Aug 5 [cited 2022 Nov 9]; Available from: 603 https://link.springer.com/10.1007/s10461-022-03748-9 604 48. Earnshaw VA, Bogart LM, Laurenceau JP, Chan BT, Maughan-Brown BG, Dietrich JJ, et 605 al. Internalized HIV stigma, ART initiation and HIV 1 RNA suppression in South Africa: 606 exploring avoidant coping as a longitudinal mediator. 2018;7. 607 49. Draucker CB, Martsolf DS, Ross R, Rusk TB. Theoretical Sampling and Category 608 Development in Grounded Theory. Qual Health Res. 2007 Oct;17(8):1137–48. 609 50. Dedoose [Internet]. Los Angeles, CA: SocioCultural Research Consultants, LLC; 2021 610 [cited 2022 Dec 9]. Available from: www.dedoose.com 611 51. Bell J, Sharma S, Malone S, Levy M, Reast J, Ciecielag J, et al. Targeting interventions for 612 HIV testing and treatment uptake: An attitudinal and behavioural segmentation of men aged 613 20–34 in KwaZulu-Natal and Mpumalanga, South Africa. Wechsberg W, editor. PLOS 614 ONE. 2021 Mar 10:16(3):e0247483. 615 52. Naugle DA, Tibbels NJ, Hendrickson ZM, Dosso A, Van Lith L, Mallalieu EC, et al. 616 Bringing fear into focus: The intersections of HIV and masculine gender norms in Côte 617 d'Ivoire. Galea JT, editor. PLOS ONE. 2019 Oct 23;14(10):e0223414. 618 53. Treves-Kagan S, El Ayadi AM, Pettifor A, MacPhail C, Twine R, Maman S, et al. Gender, 619 HIV Testing and Stigma: The Association of HIV Testing Behaviors and Community-Level 620 and Individual-Level Stigma in Rural South Africa Differ for Men and Women. AIDS 621 Behav. 2017 Sep;21(9):2579-88. 622 54. Hlongwa M, Mashamba-Thompson T, Makhunga S, Hlongwana K. Barriers to HIV testing 623 uptake among men in sub-Saharan Africa: a scoping review. Afr J AIDS Res. 2020 Jan 624 2;19(1):13-23. 625 55. Gumbi K. South Africa's unemployment rate rises amid power crisis [Internet]. Reuters. 626 2023 [cited 2023 Jun 26]. Available from: https://www.reuters.com/world/africa/south-627 africas-unemployment-rate-rises-329-q1-2023-05-16/ 628 56. Peer N, Lombard C, Stevn K, Levitt N. A high burden of adverse life events and poor 629 coping mechanisms experienced by urban-dwelling black South Africans. Annunziato RA, 630 editor. PLOS ONE. 2020 Sep 10;15(9):e0238320. 631 57. Hatcher AM, Lemus Hufstedler E, Doria K, Dworkin SL, Weke E, Conroy A, et al. 632 Mechanisms and perceived mental health changes after a livelihood intervention for HIV-633 positive Kenyans: Longitudinal, qualitative findings. Transcult Psychiatry. 2020 634 Feb;57(1):124–39. 635 58. MenStar Coalition. MenStar Strategy [Internet]. [cited 2023 Jun 26]. Available from: 636 https://www.state.gov/wp-content/uploads/2020/06/MenStar-Strategy.pdf

637 638 639	59.	Martinez P, Lien L, Landheim A, Kowal P, Clausen T. Quality of life and social engagement of alcohol abstainers and users among older adults in South Africa. BMC Public Health. 2014 Dec;14(1):316.
640 641 642	60.	Hatcher AM, Gibbs A, McBride RS, Rebombo D, Khumalo M, Christofides NJ. Gendered syndemic of intimate partner violence, alcohol misuse, and HIV risk among peri-urban, heterosexual men in South Africa. Soc Sci Med. 2022 Feb;295:112637.
643 644 645 646	61.	Rabie S, Tomlinson M, Almirol E, Stewart J, Skiti Z, Weiss RE, et al. Utilizing Soccer for Delivery of HIV and Substance Use Prevention for Young South African Men: 6-Month Outcomes of a Cluster Randomized Controlled Trial. AIDS Behav. 2023 Mar;27(3):842–54.
647 648 649	62.	Huis In 'T Veld D, Ensoy-Musoro C, Pengpid S, Peltzer K, Colebunders R. The efficacy of a brief intervention to reduce alcohol use in persons with HIV in South Africa, a randomized clinical trial. De Socio GV, editor. PLOS ONE. 2019 Aug 20;14(8):e0220799.
650 651	63.	Mullainathan S, Shafir E. Freeing Up Intelligence. Scientific American Mind. 2014 Feb;57–63.
652 653 654	64.	Crielaard L, Nicolaou M, Sawyer A, Quax R, Stronks K. Understanding the impact of exposure to adverse socioeconomic conditions on chronic stress from a complexity science perspective. BMC Med. 2021 Dec;19(1):242.
655 656	65.	Vélez-Coto M, Rute-Pérez S, Pérez-García M, Caracuel A. Unemployment and general cognitive ability: A review and meta-analysis. J Econ Psychol. 2021 Dec;87:102430.
657 658 659	66.	Tieosapjaroen W, Zhang Y, Fairley CK, Zhang L, Chow EPF, Phillips TR, et al. Improving access to oral pre-exposure prophylaxis for HIV among international migrant populations. Lancet Public Health. 2023 Aug;8(8):e651–8.
660 661 662	67.	Shommu NS, Ahmed S, Rumana N, Barron GRS, McBrien KA, Turin TC. What is the scope of improving immigrant and ethnic minority healthcare using community navigators: A systematic scoping review. Int J Equity Health. 2016 Dec;15(1):6.
663 664 665 666	68.	Cultural Navigators to Liaise Between Communities and Public Health [Internet]. University of Minnesota National Resource Center for Refugees, Immigrants, and Migrants. Available from: https://nrcrim.org/cultural-navigators-liaise-between- communities-and-public-health
667 668 669	69.	A human step to equal health care in South Africa's rural hospitals [Internet]. The Conversation. 2015 [cited 2023 Jun 27]. Available from: https://theconversation.com/a-human-step-to-equal-health-care-in-south-africas-rural-hospitals-41648
670 671 672 673	70.	Treves-Kagan S, Steward WT, Ntswane L, Haller R, Gilvydis JM, Gulati H, et al. Why increasing availability of ART is not enough: a rapid, community-based study on how HIV-related stigma impacts engagement to care in rural South Africa. BMC Public Health. 2015 Dec;16(1):87.

- 674 71. Hennink MM, Cooper P, Diamond I. Seasonal Work and Sexual Behavior. J Sex Res. 2000
  675 May;37(2):175–83.
- Muhumuza R, Ssemata AS, Kakande A, Ahmed N, Atujuna M, Nomvuyo M, et al.
  Exploring Perceived Barriers and Facilitators of PrEP Uptake among Young People in
  Uganda, Zimbabwe, and South Africa. Arch Sex Behav. 2021 May;50(4):1729–42.
- 73. Nabunya R, Karis VMS, Nakanwagi LJ, Mukisa P, Muwanguzi PA. Barriers and
  facilitators to oral PrEP uptake among high-risk men after HIV testing at workplaces in
  Uganda: a qualitative study. BMC Public Health. 2023 Feb 20;23(1):365.
- 682 74. Beesham I, Milford C, Smit J, Joseph Davey DL, Baeten JM, Heffron R, et al. Post-trial
  683 access to and use of pre-exposure prophylaxis in Durban, South Africa. BMC Public
  684 Health. 2023 Jun 22;23(1):1210.