

AORTA Gene: Polygenic prediction improves detection of thoracic aortic aneurysm

James P. Pirruccello MD^{1,2,3}, Shaan Khurshid MD, MPH^{4,5,6,7}, Honghuang Lin PhD^{8,9}, Lu-Chen Weng PhD^{5,6}, Siavash Zamirpour MS¹⁰, Shinwan Kany MD^{6,11}, Avanthi Raghavan MD^{4,6}, Satoshi Koyama MD, PhD^{5,6,12}, Ramachandran S. Vasani MD^{8,13,14}, Emelia J. Benjamin MD, ScM^{8,13,14}, Mark E. Lindsay MD, PhD^{4,5,6,7,15}, Patrick T. Ellinor MD, PhD^{4,5,6,7}

1 Division of Cardiology, University of California San Francisco, San Francisco, California, USA, 2 Institute for Human Genetics, University of California San Francisco, San Francisco, California, USA, 3 Bakar Computational Health Sciences Institute, University of California San Francisco, San Francisco, California, USA, 4 Cardiology Division, Massachusetts General Hospital, Boston, Massachusetts, USA, 5 Cardiovascular Research Center, Massachusetts General Hospital, Boston, Massachusetts, USA, 6 Cardiovascular Disease Initiative, Broad Institute of MIT and Harvard, Cambridge, Massachusetts, USA, 7 Harvard Medical School, Boston, Massachusetts, USA, 8 Framingham Heart Study, Boston University and National Heart, Lung, and Blood Institute, Framingham, Massachusetts, USA, 9 Department of Medicine, University of Massachusetts Chan Medical School, Worcester, Massachusetts, USA, 10 School of Medicine, University of California San Francisco, San Francisco, California, USA, 11 Department of Cardiology, University Heart and Vascular Center Hamburg-Eppendorf, Hamburg, Germany, 12 Laboratory for Cardiovascular Genomics and Informatics, RIKEN Center for Integrative Medical Sciences, Kanagawa, Japan, 13 Department of Medicine, Cardiology and Preventive Medicine Sections, Boston Medical Center, Boston University Chobanian and Avedisian School of Medicine, Boston, Massachusetts, USA, 14 Epidemiology Department, Boston University School of Public Health, Boston, Massachusetts, USA, 15 Thoracic Aortic Center, Massachusetts General Hospital, Boston, Massachusetts, USA

Running Title: AORTA Gene

Word Count: 4,053

Keywords: Ascending aorta, polygenic score, UK Biobank, *All of Us*, Framingham Heart Study, Mass General Brigham

Correspondence:

James P. Pirruccello, MD
University of California San Francisco
Division of Cardiology
555 Mission Bay Blvd South #3118
San Francisco, CA 94158
james.pirruccello@ucsf.edu

Patrick T. Ellinor, MD, PhD
Massachusetts General Hospital
Cardiology Division
55 Fruit Street
Boston, MA 02114
ellinor@mgh.harvard.edu

42 Abstract

43 **Background:** Thoracic aortic disease is an important cause of morbidity and mortality in
44 the US, and aortic diameter is a heritable contributor to risk. Could a polygenic
45 prediction of ascending aortic diameter improve detection of aortic aneurysm?

46
47 **Methods:** Deep learning was used to measure ascending thoracic aortic diameter in
48 49,939 UK Biobank participants. A genome-wide association study (GWAS) was
49 conducted in 39,524 participants and leveraged to build a 1.1 million-variant polygenic
50 score with *PRScs-auto*. Aortic diameter prediction models were built with the polygenic
51 score (“AORTA Gene”) and without it. The models were tested in a held-out set of 4,962
52 UK Biobank participants and externally validated in 5,469 participants from Mass
53 General Brigham Biobank (MGB), 1,298 from the Framingham Heart Study (FHS), and
54 610 participants from *All of Us*.

55
56 **Results:** In each test set, the AORTA Gene model explained more of the variance in
57 thoracic aortic diameter compared to clinical factors alone: 39.9% (95% CI 37.8-42.0%)
58 vs 29.2% (95% CI 27.1-31.4%) in UK Biobank, 36.5% (95% CI 34.4-38.5%) vs 32.5%
59 (95% CI 30.4-34.5%) in MGB, 41.8% (95% CI 37.7-45.9%) vs 33.0% (95% CI 28.9-
60 37.2%) in FHS, and 34.9% (95% CI 28.8-41.0%) vs 28.9% (95% CI 22.9-35.0%) in *All*
61 *of Us*. AORTA Gene had a greater AUROC for identifying diameter ≥ 4 cm in each test
62 set: 0.834 vs 0.765 ($P=7.3E-10$) in UK Biobank, 0.808 vs 0.767 in MGB ($P=4.5E-12$),
63 0.856 vs 0.818 in FHS ($P=8.5E-05$), and 0.827 vs 0.791 ($P=7.8E-03$) in *All of Us*.

64

65 **Conclusions:** Genetic information improved estimation of thoracic aortic diameter when
66 added to clinical risk factors. Larger and more diverse cohorts will be needed to develop
67 more powerful and equitable scores.

68

69

70 Introduction

71 Thoracic aortic disease is an important cause of morbidity and mortality^{1,2}. Ascending
72 thoracic aortic enlargement is a well-established risk factor for ascending aortic
73 dissection^{3,4}. Indeed, the majority of dissections occur in individuals with aortic diameter
74 ≥ 4 cm, including more than 90% in the International Registry of Aortic Dissection and
75 100% in Kaiser^{4,5}. Contemporary guidelines do not describe a role for population
76 screening of thoracic aortic aneurysm⁶, and universal imaging would likely be
77 impractical.

78
79 These limitations have spurred the development of clinical risk scores to identify
80 individuals who are likely to have aortic enlargement on confirmatory imaging⁷⁻⁹. These
81 scores have been shown to explain approximately 30% of the variance in ascending
82 aortic diameter, with an area under the receiver operator characteristic curve (AUROC)
83 near 0.78 for detecting individuals with diameter ≥ 4 cm. Population genetics studies
84 suggest that ascending aortic diameter is highly heritable, with a proportion of variance
85 attributable to common genetic factors of over 60%¹⁰. We therefore sought to assess
86 whether incorporating polygenic risk might improve estimation of ascending aortic
87 diameter over clinical factors alone.

88

89 Methods

90 Study design

91 Model development and internal validation were conducted in UK Biobank, with external
92 validation within the *All of Us* cohort, the Mass General Brigham Biobank (MGB), and
93 the Framingham Heart Study (FHS). This study was reported in accordance with the
94 transparent reporting of a multivariable prediction model for individual prognosis or
95 diagnosis (TRIPOD) statement¹¹.

96
97 Study protocols complied with the tenets of the Declaration of Helsinki. All UK Biobank
98 participants provided written informed consent¹², and only those participants who had
99 not withdrawn consent as of March 7, 2023 were analyzed. The UK Biobank analyses
100 were considered exempt by the UCSF Institutional Review Board (IRB), #22-37715. UK
101 Biobank analyses were conducted under application #41664. Each *All of Us* biobank
102 participant provided written informed consent¹³. The *All of Us* analyses were considered
103 exempt by the UCSF IRB, #22-37715. The MGB study protocols were approved with a
104 waiver of informed consent by the Mass General Brigham IRB. All FHS participants
105 provided written informed consent, and the FHS imaging analyses were previously
106 approved by the IRB of the Boston University Medical Center¹⁴. The FHS analyses were
107 also approved by the MGB IRB.

108 Study populations

109 In UK Biobank participants, ascending aortic diameter was measured from magnetic
110 resonance imaging (MRI) using a previously described deep learning model¹⁵. Model

111 development and internal validation were conducted in UK Biobank. External validation
112 of the models was pursued in MGB participants with ascending aortic diameter
113 measured from TTE for clinical indications¹⁶; in FHS participants with ascending aortic
114 diameter measured for research purposes from non-contrast computed tomography
115 (CT)¹⁴; and in *All of Us* biobank participants who had clinical aortic diameter
116 measurements from transthoracic echocardiography (TTE)¹³. In MGB, race or ethnicity
117 was extracted from the electronic health record¹⁶ and the validation analyses were
118 repeated in individuals identified as Black.

119 **Modality-specific differences**

120 The clinical approach to measuring ascending aortic diameter differs by modality. UK
121 Biobank MRI-based aortic diameter measurements were derived from the blood pool
122 diameter only, incorporating neither aortic wall. For TTE, used in the MGB and *All of Us*
123 cohorts, standard measurements use the “leading edge to leading edge” approach,
124 incorporating one wall. For non-contrast CT measurements, used in FHS, both walls are
125 incorporated into the diameter.

126 **Splitting the UK Biobank for training and validation**

127 49,939 UK Biobank participants with MRI measurements of aortic diameter were
128 randomly assigned a number from 0-999. Those from 0-799 contributed to the genome-
129 wide association study (GWAS) used to produce the polygenic score (N=39,524). Those
130 from 800-999 related within 3 degrees of kinship to the 0-799 group were excluded to
131 avoid polygenic score overfitting (N=557). Those remaining from 0-899 contributed to
132 the development of non-genetic models (N=44,420). Those from 800-899 were included

133 in the derivation of weights for the polygenic score component of the AORTA Gene
134 combined clinical and genetic model (N=4,896). Those ≥ 900 were analyzed for
135 validation (N=4,962; **Figure 1**).

136 **Thoracic aortic aneurysm in *All of Us***

137 Within *All of Us*, thoracic aortic aneurysm was defined as the first occurrence of
138 SNOMED codes 433068007, 74883004, or 426948001; ICD10CM codes of I71.01,
139 I71.1, I71.2, or I77.810; ICD9CM codes of 441.01, 441.1, 441.2, or 447.71; ICD10PCS
140 codes of 02RX0JZ or 02RW0JZ; the ICD9Proc code 39.73; or CPT4 codes 33863,
141 75957, 33880, 75956, or 33881. Prevalent analyses were conducted based on the
142 presence of disease labels prior to the time of enrollment. Time-to-event analyses were
143 conducted beginning at the time of enrollment and censored on July 1, 2022 using the R
144 *survival* package, excluding participants whose diagnoses occurred prior to enrollment.
145 Any tabular data with an *All of Us* sample count < 20 was modified in accordance with *All*
146 *of Us* policies.

147 **Genome-wide association study and polygenic scores**

148 Common genetic contributions to ascending aortic diameter in the 39,524 GWAS
149 participants were discovered using REGENIE v3.2.7¹⁷ with imputed variants provided by
150 UK Biobank^{18,19}. GWAS covariates included age, age², sex, the MRI serial number, the
151 genotyping array, and the first ten principal components of ancestry. The summary
152 statistics were trimmed to the ~ 1.1 million UK Biobank-specific variants within HapMap3
153 identified by the *PRSCs* authors, and underwent Bayesian weighting by *PRSCs-auto* on
154 its default settings to produce polygenic score weights²⁰.

155

156 The polygenic score was applied to all UK Biobank participants with imputed genetic
157 data¹⁹. Variants with imputation quality <0.3 were removed.

158

159 The polygenic score was applied to all MGB participants with imputed genetic data^{16,21}
160 with imputation into the TOPMed panel²².

161

162 The polygenic score was applied to FHS participants with CT and genetic data
163 available. For FHS, genotyping was done on the Affymetrix GeneChip 500K Array Set &
164 50K Human Gene Focused Panel (Affymetrix, Santa Clara, CA, USA). Variants with call
165 rate <97% or Hardy–Weinberg equilibrium $P < 1E-06$ were excluded. The remaining
166 variants were then imputed to the 1000 Genomes Project phase I release 3 panel by
167 MACH version 1.0^{23,24}. Variants with imputation quality <0.3 were removed.

168

169 The polygenic score was applied to *All of Us* participants with whole genome
170 sequencing data available in release “R7”²⁵.

171

172 Within each set of participants, the polygenic score was residualized for the first 20
173 principal components of ancestry (except for FHS, where 10 were used due to
174 availability). The polygenic score in each group was zero-centered and scaled by the
175 standard deviation from the UK Biobank training set: for studies that applied the
176 average polygenic score (UK Biobank, *All of Us*) the standard deviation was 3.4E-07,

177 and for those that applied the summed polygenic score (MGB, FHS) the standard
178 deviation was 0.38.

179 **Ascending aortic diameter model training**

180 Hierarchical group least absolute shrinkage and selection operator models were built to
181 estimate ascending aortic diameter using the R package *glinternet*²⁶. Three models
182 were constructed with *glinternet* v1.0.11 using the following independent variables: (a)
183 age, age², and sex; (b) age, age², sex, and polygenic score; (c) age, age², body mass
184 index (BMI), heart rate, systolic and diastolic blood pressure, height, weight, sex, and a
185 history of diabetes, hypertension, or hyperlipidemia. A fourth model (d) was constructed
186 by taking a linear combination of the model (c) and the polygenic score. The weighting
187 for model (c) was estimated in all participants randomly assigned to 0-899, while the
188 weighting for model (b) and the linear combination for model (d) were estimated in those
189 assigned to 800-899 to avoid overlap with the GWAS cohort. For ease of reference, we
190 label model (c) “AORTA Score” (AORTA: optimized regression for thoracic aneurysm)
191 and model (d) “AORTA Gene”.

192 **Statistical analysis**

193 All statistical significance tests were two-sided, with significance defined as $P < 0.05$,
194 and performed in R 4.2.3 unless otherwise stated. For visualization, scores were plotted
195 against measured thoracic aortic diameter.

196 Outcomes

197 The primary outcome was correlation between the scores and ascending aortic
198 diameter using linear models, expressed as R^2 (variance explained). Secondary
199 outcomes included tests of calibration, and the performance of the scores for identifying
200 ascending aortic diameter ≥ 4.0 cm; the clinical diagnosis of aneurysm was also tested in
201 *All of Us*.

202
203 To assess calibration, each score was used as an independent variable in a linear
204 model predicting aortic diameter that also had an intercept, permitting significance
205 testing of both the slope and the intercept. Mean absolute error was also computed.

206
207 To assess the performance for identifying diameter ≥ 4 cm, the AUROC was computed.
208 Confusion matrices and their derived statistical measures were produced based on the
209 presence or absence of aortic diameter ≥ 4 cm and the presence or absence of predicted
210 diameters above a fixed threshold determined in the training set for each score. The
211 thresholds for the confusion matrix analysis were defined in the 4,896 participants in the
212 training set.

213
214 AUROCs were compared using the DeLong test²⁷. The continuous net reclassification
215 index (NRI) was calculated following the formula of Pencina, *et al*²⁸.

216 **Data and code availability**

217 Computed scores are returned to UK Biobank, where data are made available by UK
218 Biobank to researchers from research institutions with genuine research inquiries,

219 following IRB and UK Biobank approval. FHS data are made available to researchers
220 with approved research applications. The dbGAP study accession number used for FHS
221 validation was #phv00076329.v1.p5 for ascending aortic diameter. MGB data are
222 available to MGB investigators; external collaboration requests can be initiated through
223 <https://biobank.massgeneralbrigham.org/for-researchers>. The complete set of AORTA
224 Score covariates and their weights are available as R programs at
225 github.com/carbocation/genomisc and the polygenic score weights will be available on
226 the Polygenic Score Catalog (pgscatalog.org).

227 Results

228 Clinical characteristics

229 Among the 4,896 UK Biobank participants whose data contributed to training the
230 genetic models, 2,489 were women and 2,407 were men. The ascending aortic
231 diameter was 3.05 ± 0.31 cm for women and 3.34 ± 0.34 cm for men. The 4,962 internal
232 validation set participants had similar measurements (**Table 1; Figure 1**). The MGB
233 validation set consisted of 5,469 participants with genetic data and aortic diameter
234 measured from TTE. The FHS validation set consisted of 1,298 participants with genetic
235 data and aortic diameter measured from non-contrast CT. The *All of Us* validation set
236 consisted of 610 participants 40 years and older with genetic data and aortic diameter
237 measured from TTE; 60% identified as female and 88% as having non-Hispanic white
238 ethnicity.

239 Model validation for aortic diameter

240 Testing calibration, the slope of the AORTA Gene model was statistically consistent with
241 one in UK Biobank ($P=0.2$), MGB ($P=1.0$), FHS ($P=0.6$), and *All of Us* ($P=0.4$) (**Figure**
242 **2**). The intercept was consistent with zero in UK Biobank ($P=0.2$), MGB ($P=0.4$), and *All*
243 *of Us* ($P=0.8$), but not FHS (intercept estimate 0.26cm, $P=0.01$). The same pattern was
244 observed for the clinical AORTA Score. In contrast, the model incorporating only age,
245 sex, and the polygenic score had a calibration slope inconsistent with one in most
246 cohorts outside of UK Biobank (**eTable AA**).

247

248 The AORTA Gene model's mean absolute error (MAE) for ascending aortic diameter
249 was 0.212cm (95% CI 0.207-0.217cm) in UK Biobank, 0.282cm (95% CI 0.275-
250 0.289cm) in MGB, 0.344cm (95% CI 0.331-0.358cm) in FHS, and 0.293cm (95% CI
251 0.273-0.313cm) in *All of Us*. The clinical AORTA Score had an inferior MAE in all
252 cohorts: 0.232cm (95% CI 0.227-0.237cm) in UK Biobank, 0.291cm (95% CI 0.284-
253 0.298cm) in MGB, 0.351cm (95% CI 0.337-0.365cm) in FHS, and 0.306cm (95% CI
254 0.285-0.327cm) in *All of Us*. The age/sex/genetics model had an MAE of 0.226cm (95%
255 CI 0.221-0.231cm) in UK Biobank, 0.293cm (95% CI 0.286-0.300cm) in MGB, 0.346cm
256 (95% CI 0.332-0.360cm) in FHS, and 0.318cm (95% CI 0.296-0.340) in *All of Us*.

257
258 The AORTA Gene model explained 39.9% of the variance in ascending aortic diameter
259 (99% CI 37.8-42.0%; $P=2.6E-551$) in the UK Biobank validation set, 36.5% (95% CI
260 34.4-38.5%, $P=2.4E-541$) in MGB, 41.8% (95% CI 37.7-45.9%; $P=3.5E-154$) in FHS,
261 and 34.9% (95% CI 28.8-41.0%; $P=1E-58$) in *All of Us* (**Figure 3**). In comparison, the
262 clinical AORTA Score explained 29.3% of the variance in UK Biobank (99% CI 27.1-
263 31.4%), 32.5% in MGB (95% CI 30.4-34.5%), 33.0% (95% CI 28.9-37.2%) in FHS, and
264 28.9% (95% CI 22.9-35.0%) in *All of Us*. The age/sex/genetics model explained 31.8%
265 (95% CI 29.6-33.9%) of variance in UK Biobank, 33.2% (95% CI 31.2-35.3%) in MGB,
266 33.3% (95% CI 29.1-37.5%) in FHS, and 31.0% (95% CI 24.9-37.1%) in *All of Us*.

267

268 **Model validation for a 4cm aortic diameter threshold**

269 113 of the 4,962 internal validation set participants in UK Biobank had aortic diameter
270 ≥ 4 cm (2.3%), as did 335 of 5,469 MGB participants (6.1%), 116 of 1,298 FHS
271 participants (8.9%), and 65 of 610 *All of Us* participants (10.7%).

272
273 The AUROC for detecting diameter ≥ 4 cm for the AORTA Gene model was 0.834 (95%
274 CI 0.798-0.869) in UK Biobank, 0.808 (95% CI 0.783-0.832) in MGB, 0.856 (95% CI
275 0.821-0.891) in FHS, and 0.827 (95% CI 0.776-0.878) in *All of Us* (**Figure 4; eTable**
276 **BB**).

277
278 For the clinical AORTA Score, the AUROC was 0.765 (95% CI 0.723-0.807) in UK
279 Biobank, 0.767 (95% CI 0.741-0.792) in MGB, 0.818 (95% CI 0.776-0.859) in FHS, and
280 0.791 (95% CI 0.735-0.846) for *All of Us*. The AUROC of AORTA Gene was
281 significantly greater in all cohorts: $P=7.3E-10$ against a null hypothesis of no difference
282 in UK Biobank, $P=4.5E-12$ in MGB, $P=8.5E-05$ in FHS, and $P=7.8E-03$ in *All of Us*
283 (**eTable CC**).

284
285 The age/sex/genetics model had an AUROC of 0.805 (95% CI 0.767-0.843) in UK
286 Biobank, 0.805 (95% CI 0.781-0.829) in MGB, 0.819 (95% CI 0.781-0.856) in FHS, and
287 0.813 (95% CI 0.757-0.870) in *All of Us*. The AORTA Gene AUROC was significantly
288 greater in UK Biobank ($P=0.03$) and FHS ($P=3.9E-03$), but not in MGB ($P=0.7$) or *All of*
289 *Us* ($P=0.5$) (**eTable CC**).

290

291 The continuous NRI was positive for AORTA Gene over the clinical AORTA Score and
292 the age/sex/genetics model in all cohorts (**eTable CC**). Decomposed into its NRI(event)
293 and NRI(nonevent) components, AORTA Gene had a greater NRI(event) in all cohorts
294 compared to the AORTA Score and the age/sex/genetics model. However, its
295 NRI(nonevent) value was lower than that of the AORTA Score in MGB and UK Biobank,
296 and lower than that of the age/sex/genetics model in MGB and *All of Us*.

297 **Score threshold demonstrations**

298 To evaluate the consequences of various score thresholds whereby a portion of the
299 population could be brought forward for confirmatory thoracic imaging, two thresholds
300 were defined within the UK Biobank training set and applied in the validation sets: one
301 targeted at the top 10% of the population, and the other designed to have a sensitivity of
302 90% for aortic diameter $\geq 4\text{cm}$ (**eTable BB**).

303
304 For the target top 10% AORTA Gene cutoff, 485 (9.8%) of 4,962 participants were
305 above this threshold in UK Biobank, 640 (11.7%) of 5,469 in MGB, 99 (7.6%) of 1,295 in
306 FHS, and 97 (15.9%) of 610 in *All of Us*. 59 of 113 UK Biobank participants with
307 diameter $\geq 4\text{cm}$ were correctly classified as enlarged (sensitivity 52.2%), as were 162 of
308 335 in MGB (48.4%), 47 of 116 in FHS (40.5%), and 35 of 65 in *All of Us* (53.8%). The
309 precision was 0.122 in UK Biobank, 0.253 in MGB, 0.475 in FHS, and 0.361 in *All of Us*,
310 yielding respective F1 scores of 0.197, 0.332, 0.437, and 0.432 (**eTable BB**). A
311 comparable threshold for the clinical AORTA Score flagged 484 (9.8%) of 4,962 UK
312 Biobank participants, 722 (13.2%) of 5,469 in MGB, 92 (7.1%) of 1,295 in FHS, and 110
313 (18.0%) of 610 in *All of Us*. 41 of 113 UK Biobank participants with diameter $\geq 4\text{cm}$ were

314 correctly classified as enlarged (sensitivity 36.3%) as were 143 of 335 in MGB (42.7%),
315 38 of 116 in FHS (32.8%), and 33 of 65 in *All of Us* (50.8%). The precision was 0.85 in
316 UK Biobank, 0.198 in MGB, 0.413 in FHS, and 0.300 in *All of Us*, yielding respective F1
317 scores of 0.137, 0.271, 0.365, and 0.377 (**eTable BB**).

318
319 For the target 90% sensitive AORTA Gene cutoff, 1,737 (35.0%) of 4,962 UK Biobank
320 participants were above this threshold, 1,943 (35.5%) of 5,469 in MGB, 336 (25.9%) of
321 1,295 in FHS, and 247 (40.5%) of 610 in *All of Us*. 92 of 113 UK Biobank participants
322 with diameter ≥ 4 cm were correctly classified as enlarged (sensitivity 81.4%), as were
323 258 of 335 in MGB (77.0%), 88 of 116 in FHS (75.9%), and 56 of 65 in *All of Us*
324 (86.2%). The precision was 0.053 in UK Biobank, 0.133 in MGB, 0.262 in FHS, and
325 0.227 in *All of Us*, yielding respective F1 scores of 0.099, 0.227, 0.389, and 0.359
326 (**eTable DD**). A comparable threshold for the clinical AORTA Score flagged 2,006
327 (40.4%) of 4,962 UK Biobank participants, 2,194 (40.1%) of 5,469 in MGB, 405 (31.3%)
328 of 1,295 in FHS, and 266 (43.6%) of 610 in *All of Us*. 87 of 113 UK Biobank participants
329 with diameter ≥ 4 cm were correctly classified as enlarged (sensitivity 77.0%), as were
330 264 of 335 in MGB (78.8%), 86 of 116 in FHS (74.1%), and 52 of 65 in *All of Us*
331 (80.0%). The precision was 0.043 in UK Biobank, 0.120 in MGB, 0.212 in FHS, and
332 0.195 in *All of Us*, yielding respective F1 scores of 0.082, 0.209, 0.330, and 0.314
333 (**eTable DD**).

334 **Thoracic aortic aneurysm diagnosis in *All of Us***

335 In the 164,789 *All of Us* participants aged 40 or older with genetic data, 1,904 had an
336 electronic health record (EHR)-based diagnosis of thoracic aortic aneurysm prior to

337 enrollment. AORTA Gene's AUROC was 0.760 (95% CI 0.750-0.771) compared to
338 0.739 (95% CI 0.728-0.750) for the AORTA Score (**eTable EE**), $P=2.4E-16$ against a
339 null hypothesis of no difference (**eTable FF**). For incident disease (1,632 cases
340 diagnosed after enrollment), AORTA Gene's AUROC was 0.748 (95% CI 0.737-0.759)
341 compared to 0.729 (95% CI 0.718-0.741) for the AORTA Score, $P=9.5E-10$ against a
342 null hypothesis of no difference (**eTables EE-FF**).

343 **Analysis in Black individuals in MGB**

344 In MGB, 340 individuals were identified as Black through the electronic health record¹⁶.
345 In these individuals, AORTA Gene had an intercept consistent with zero ($P=0.5$) and a
346 slope consistent with one ($P=0.5$). The MAE was 0.297cm (95% CI 0.272-0.321cm),
347 and the model explained 34.8% of variance in ascending aortic diameter (95% CI 26.7-
348 43.0%). In comparison, for the clinical AORTA Score these values were 0.294cm (95%
349 CI 0.269-0.319cm) and 34.3% (95% CI 26.2-42.5%) (**eTable AA**). The AUROC for
350 detecting diameter ≥ 4 cm for the AORTA Gene model was 0.858 (**eTable BB**), nominally
351 greater than that of the clinical AORTA Score (AUROC 0.820, $P=0.04$ for the difference,
352 **eTable CC**), with a positive continuous NRI (0.473).

353

354 Discussion

355 Ascending thoracic aortic diameter has a SNP heritability estimated to be greater than
356 60%¹⁰, which makes the incorporation of common-variant genetic data for the
357 presymptomatic detection of individuals with aortic enlargement or aneurysm
358 conceptually appealing. But whether aortic polygenic scores encode information not
359 already latently captured through correlation with clinical covariates was unknown.
360 Here, we observed that AORTA Gene, which incorporated both an aortic polygenic
361 score and clinical covariates, improved estimation of ascending aortic diameter and
362 identification of individuals with diameter ≥ 4 cm in four cohorts—UK Biobank, MGB,
363 FHS, and *All of Us*—when compared to a previously validated model for aortic diameter
364 built from the same clinical covariates.

365
366 For the estimation of aortic diameter as a quantitative measure, a clinical model built
367 from the same covariates as the AORTA Score⁹ explained 29.3% of the variance in
368 ascending aortic diameter in UK Biobank, 28.9% in *All of Us*, 32.5% in MGB, and 33.0%
369 in FHS. The AORTA Gene model that additionally incorporated the aortic diameter
370 polygenic score increased those values to 39.9% in UK Biobank, 34.9% in *All of Us*,
371 36.5% in MGB, and 41.8% in FHS. These represent a 10.6% absolute improvement in
372 explained variance (36% relative improvement) in UK Biobank, 4% (12.3% relative
373 improvement) in MGB, 8.8% (27% relative improvement) in FHS, and 6% (21% relative
374 improvement) in *All of Us*. These substantive improvements from the polygenic score
375 over a clinical model are comparable in magnitude to the addition of key clinical
376 predictors over age and sex alone⁹.

377
378 Incorporation of the polygenic score also improved identification of individuals with
379 clinically relevant aortic enlargement (i.e., diameter ≥ 4 cm). The clinical model had an
380 AUROC of 0.765 in UK Biobank, 0.791 in *All of Us*, 0.767 in MGB, and 0.818 in FHS;
381 these respectively improved to 0.834, 0.827, 0.808, and 0.856 for AORTA Gene. The
382 AUROC is independent of the choice of score threshold, so we also demonstrated two
383 example thresholds: one based on the goal of being 90% sensitive for diameter ≥ 4 cm,
384 and another based on a scenario where there are resources to image up to 10% of the
385 population. In both scenarios, for all four cohorts, sensitivity and specificity were
386 modestly improved with the incorporation of the polygenic score. Interestingly, we also
387 observed that, compared to the comprehensive clinical score, a simpler model
388 consisting only of age, sex, and the polygenic score performed similarly. Because such
389 a model can be estimated for any future time from birth, this may have implications for
390 attempts at early risk stratification before the onset of clinical risk factors. We also noted
391 that, for the task of identifying *All of Us* participants with an EHR-based diagnosis of
392 aneurysm, all models had attenuated performance—likely in part due to limited
393 ascertainment of thoracic aortic disease in current clinical practice. Nevertheless, even
394 for this task, the AORTA Gene model remained superior to the clinical score ($P=9.5E-$
395 10).

396
397 We also note two limitations of the current model that point to future directions: first,
398 despite a heritability of nearly 60%, the current polygenic score for thoracic aortic
399 diameter derived from a GWAS of 39,524 participants added approximately 10% to the

400 variance already explained by a clinical model, suggesting that it accounted for
401 approximately 17% of the heritability of aortic diameter. In contrast, polygenic scores for
402 height—built from GWAS of millions of participants—are nearly saturated²⁹. Larger
403 GWAS sample sizes are required to increase the variance explained by polygenic
404 estimates of ascending aortic diameter. Second, polygenic scores derived in largely
405 European-ancestry populations do not capture the full range of human genetic diversity,
406 limiting their utility in individuals with more diverse genetic identities³⁰. To examine this,
407 we evaluated 340 individuals in MGB identified as Black in the electronic health
408 record¹⁶. In these individuals, the addition of the polygenic score did not improve the
409 continuous estimation of aortic diameter compared to the clinical score (R^2 34.8% for
410 AORTA Gene vs 34.3% for the clinical AORTA Score), although it did nominally
411 improve identification of individuals with diameter ≥ 4 cm (AUROC 0.858 vs 0.820,
412 $P=0.04$). We expect a key problem from incorporating genetic data to be one of inequity
413 due to differential accrual of benefits to individuals with European genetic identities.
414 Larger and more diverse GWAS are required to improve genetic predictions for people
415 across diverse genetic ancestries.

416

417 Anticipation of the relevant clinical scenario for the application of polygenic scores is
418 important for placing the current findings into context. The cost of acquiring a targeted
419 set of images of the ascending thoracic aorta may be comparable that of obtaining
420 genetic information. Therefore, we expect that the value of polygenic scores for primary
421 prevention will be realized within the context of a healthcare system that incorporates
422 genetic information as part of the prior probability across many diseases and risk

423 factors, where the cost of one-time genotyping would be amortized over all downstream
424 uses. In such a system, might there be a role, in principle, for the incorporation of a
425 genetic estimator of thoracic aortic diameter into a comprehensive model to identify
426 people who are likely to have ascending aortic enlargement? The results from the
427 present analysis suggest so.

428 **Limitations**

429 While all participants were incorporated into analyses without exclusion for race,
430 ethnicity, or ancestry, the participants in UK Biobank, MGB, FHS, and *All of Us* with
431 thoracic imaging predominantly had genetic identities similar to that of Europeans. None
432 of the models described in this manuscript are anticipated to be effective for identifying
433 individuals with aortic enlargement driven by rare pathogenic variants, which are not
434 captured in polygenic scores constructed from common variants. The models were
435 derived and tested in individuals over the age of 40 years. The attenuation in variance
436 explained in the external cohorts compared to UK Biobank may be, in part, a
437 consequence of cryptic relatedness causing overfitting to the UK Biobank cohort;
438 reducing this discrepancy is an area of interest for future efforts. Future efforts are also
439 needed to understand whether aortic prediction models can improve outcomes.

440 **Conclusion**

441 Integrating genetic information into a validated clinical model improved estimation of
442 ascending aortic diameter and the identification of individuals with thoracic aortic
443 aneurysm.

444 Acknowledgments

445 JPP had full access to all the data in the study and takes responsibility for the integrity
446 of the data and the accuracy of the data analysis. JPP and PTE conceived of the study.
447 JPP conducted the genetic analyses, developed the models, validated the models, and
448 conducted bioinformatic analyses in the UK Biobank, MGB, and *All of Us*. HL validated
449 the models in FHS. LCW applied the polygenic score in MGB. S Koyama performed
450 PCA in MGB. JPP and PTE wrote the paper. All other authors contributed to the
451 analysis plan or provided critical revisions.

452 Sources of funding

453 Organizations that provided financial support had no role in the design and conduct of
454 the study; collection, management, analysis, and interpretation of the data; preparation,
455 review, or approval of the manuscript; or decision to submit the manuscript for
456 publication.

457
458 Dr. Pirruccello is supported by National Institutes of Health (NIH) K08HL159346. Dr. Lin
459 is supported by the NIH grant U01AG068221. Dr. Kany is supported by the Walter-
460 Benjamin Fellowship from the Deutsche Forschungsgemeinschaft (521832260). Dr.
461 Raghavan is supported by the John S. LaDue Memorial Fellowship in Cardiovascular
462 Medicine from Harvard Medical School. Dr. Benjamin is supported by NIH grants
463 R01HL128914, R01HL092577, R01HL141434, and U54HL120163; and American Heart
464 Association (AHA) grant 18SFRN34110082. Dr. Lindsay is supported by the Fredman
465 Fellowship for Aortic Disease and the Toomey Fund for Aortic Dissection Research and

466 has received salary support from Bayer AG. Dr. Ellinor is supported by grants from the
467 NIH (R01HL092577, 1R01HL157635), from the AHA Strategically Focused Research
468 Networks (18SFRN34110082), and from the European Union (MAESTRIA 965286).

469
470 The *All of Us* Research Program is supported by the National Institutes of Health, Office
471 of the Director: Regional Medical Centers: 1 OT2 OD026549; 1 OT2 OD026554; 1 OT2
472 OD026557; 1 OT2 OD026556; 1 OT2 OD026550; 1 OT2 OD 026552; 1 OT2
473 OD026553; 1 OT2 OD026548; 1 OT2 OD026551; 1 OT2 OD026555; IAA #: AOD
474 16037; Federally Qualified Health Centers: HHSN 263201600085U; Data and Research
475 Center: 5 U2C OD023196; Biobank: 1 U24 OD023121; The Participant Center: U24
476 OD023176; Participant Technology Systems Center: 1 U24 OD023163;
477 Communications and Engagement: 3 OT2 OD023205; 3 OT2 OD023206; and
478 Community Partners: 1 OT2 OD025277; 3 OT2 OD025315; 1 OT2 OD025337; 1 OT2
479 OD025276. In addition, the *All of Us* Research Program would not be possible without
480 the partnership of its participants.

481
482 From the Framingham Heart Study of the National Heart Lung and Blood Institute of the
483 National Institutes of Health and Boston University School of Medicine, this project has
484 been funded in whole or in part with Federal funds from the National Heart Lung and
485 Blood Institute, Department of Health and Human Services, under Contract No.
486 75N92019D00031.

487 Disclosures

488 Dr. Ellinor has received sponsored research support from Bayer AG, IBM Health, Bristol
489 Myers Squibb and Pfizer. Dr. Ellinor has also served on advisory boards or consulted
490 for Bayer AG, MyoKardia and Novartis. The Broad Institute has filed for a patent on an
491 invention from Drs. Ellinor, Lindsay, and Pirruccello related to a previous genetic risk
492 predictor for aortic disease. Remaining authors report no disclosures.

493 References

- 494 1. Howard DPJ, Banerjee A, Fairhead JF, Perkins J, Silver LE, Rothwell PM, Oxford
495 Vascular Study. Population-based study of incidence and outcome of acute aortic
496 dissection and premorbid risk factor control: 10-year results from the Oxford
497 Vascular Study. *Circulation*. 2013;127:2031–2037.
- 498 2. Yamaguchi T, Nakai M, Yano T, Matsuyama M, Yoshino H, Miyamoto Y, Sumita Y,
499 Matsuda H, Inoue Y, Okita Y, Minatoya K, Ueda Y, Ogino H. Population-based
500 incidence and outcomes of acute aortic dissection in Japan. *European Heart*
501 *Journal Acute Cardiovascular Care*. 2021;10:701–709.
- 502 3. Kim JB, Spotnitz M, Lindsay ME, MacGillivray TE, Isselbacher EM, Sundt TM. Risk
503 of Aortic Dissection in the Moderately Dilated Ascending Aorta. *J Am Coll Cardiol*.
504 2016;68:1209–1219.
- 505 4. Solomon MD, Leong T, Sung SH, Lee C, Allen JG, Huh J, LaPunzina P, Lee H,
506 Mason D, Melikian V, Pellegrini D, Scoville D, Sheikh AY, Mendoza D, Naderi S,
507 Sheridan A, Hu X, Cirimele W, Gisslow A, Leung S, Padilla K, Bloom M, Chung J,
508 Topic A, Vafaei P, Chang R, Miller DC, Liang DH, Go AS, Kaiser Permanente
509 Northern California Center for Thoracic Aortic Disease. Association of Thoracic
510 Aortic Aneurysm Size With Long-term Patient Outcomes: The KP-TAA Study.
511 *JAMA Cardiology* [Internet]. 2022 [cited 2022 Oct 11]; Available from:
512 <https://doi.org/10.1001/jamacardio.2022.3305>
- 513 5. Pape LA, Tsai TT, Isselbacher EM, Oh JK, O'gara PT, Evangelista A, Fattori R,
514 Meinhardt G, Trimarchi S, Bossone E, Suzuki T, Cooper JV, Froehlich JB,
515 Nienaber CA, Eagle KA, International Registry of Acute Aortic Dissection (IRAD)
516 Investigators. Aortic diameter \geq 5.5 cm is not a good predictor of type A aortic
517 dissection: observations from the International Registry of Acute Aortic Dissection
518 (IRAD). *Circulation*. 2007;116:1120–1127.
- 519 6. Isselbacher EM, Preventza O, Hamilton Black J, Augoustides JG, Beck AW, Bolen
520 MA, Braverman AC, Bray BE, Brown-Zimmerman MM, Chen EP, Collins TJ,
521 DeAnda A, Fanola CL, Girardi LN, Hicks CW, Hui DS, Schuyler Jones W, Kalahasti
522 V, Kim KM, Milewicz DM, Oderich GS, Ogbechie L, Promes SB, Gyang Ross E,
523 Schermerhorn ML, Singleton Times S, Tseng EE, Wang GJ, Woo YJ. 2022
524 ACC/AHA Guideline for the Diagnosis and Management of Aortic Disease: A
525 Report of the American Heart Association/American College of Cardiology Joint
526 Committee on Clinical Practice Guidelines. *Circulation*. 2022;146:e334–e482.
- 527 7. Mori M, Gan G, Deng Y, Yousef S, Weininger G, Daggula KR, Agarwal R, Shang
528 M, Assi R, Geirsson A, Vallabhajosyula P. Development and Validation of a
529 Predictive Model to Identify Patients With an Ascending Thoracic Aortic Aneurysm.
530 *Journal of the American Heart Association*. 2021;10:e022102.
- 531 8. Obel LM, Diederichsen AC, Steffensen FH, Frost L, Lambrechtsen J, Busk M,
532 Urbonaviciene G, Egstrup K, Karon M, Rasmussen LM, Gerke O, Bovling AS,
533 Lindholt JS. Population-Based Risk Factors for Ascending, Arch, Descending, and
534 Abdominal Aortic Dilations for 60–74-Year-Old Individuals. *Journal of the American*
535 *College of Cardiology*. 2021;78:201–211.
- 536 9. Pirruccello JP, Lin H, Khurshid S, Nekoui M, Weng L-C, Ramachandran VS,
537 Isselbacher EM, Benjamin EJ, Lubitz SA, Lindsay ME, Ellinor PT. Development of
538 a Prediction Model for Ascending Aortic Diameter Among Asymptomatic

- 539 Individuals. *JAMA*. 2022;328:1935–1944.
- 540 10. Pirruccello JP, Chaffin MD, Chou EL, Fleming SJ, Lin H, Nekoui M, Khurshid S,
541 Friedman SF, Bick AG, Arduini A, Weng L-C, Choi SH, Akkad A-D, Batra P, Tucker
542 NR, Hall AW, Roselli C, Benjamin EJ, Vellarikkal SK, Gupta RM, Stegmann CM,
543 Juric D, Stone JR, Vasani RS, Ho JE, Hoffmann U, Lubitz SA, Philippakis AA,
544 Lindsay ME, Ellinor PT. Deep learning enables genetic analysis of the human
545 thoracic aorta. *Nat Genet*. 2021;
- 546 11. Collins GS, Reitsma JB, Altman DG, Moons KGM. Transparent Reporting of a
547 multivariable prediction model for Individual Prognosis Or Diagnosis (TRIPOD):
548 The TRIPOD Statement. *Annals of Internal Medicine*. 2015;162:55.
- 549 12. Collins R. UK Biobank Protocol [Internet]. 2007; Available from:
550 <https://www.ukbiobank.ac.uk/media/gnkeyh2q/study-rationale.pdf>
- 551 13. Denny JC, Rutter J, Goldstein DB, Philippakis A, Smoller J, Jenkins G, Dishman E.
552 The “All of Us” Research Program. *New England Journal of Medicine*.
553 2019;381:668–676.
- 554 14. Rogers IS, Massaro JM, Truong QA, Mahabadi AA, Kriegel MF, Fox CS,
555 Thanassoulis G, Isselbacher EM, Hoffmann U, O’Donnell CJ. Distribution,
556 determinants, and normal reference values of thoracic and abdominal aortic
557 diameters by computed tomography (from the Framingham Heart Study). *Am J*
558 *Cardiol*. 2013;111:1510–1516.
- 559 15. Pirruccello JP, Rämö JT, Choi SH, Chaffin MD, Kany S, Nekoui M, Chou EL,
560 Jurgens SJ, Friedman SF, Juric D, Stone JR, Batra P, Ng K, Philippakis AA,
561 Lindsay ME, Ellinor PT. The Genetic Determinants of Aortic Distention. *Journal of*
562 *the American College of Cardiology*. 2023;81:1320–1335.
- 563 16. Khurshid S, Reeder C, Harrington LX, Singh P, Sarma G, Friedman SF, Di Achille
564 P, Diamant N, Cunningham JW, Turner AC, Lau ES, Haimovich JS, Al-Alusi MA,
565 Wang X, Klarqvist MDR, Ashburner JM, Diedrich C, Ghadessi M, Mielke J, Eilken
566 HM, McElhinney A, Derix A, Atlas SJ, Ellinor PT, Philippakis AA, Anderson CD, Ho
567 JE, Batra P, Lubitz SA. Cohort design and natural language processing to reduce
568 bias in electronic health records research. *NPJ Digit Med*. 2022;5:47.
- 569 17. Mbatchou J, Barnard L, Backman J, Marcketta A, Kosmicki JA, Ziyatdinov A,
570 Benner C, O’Dushlaine C, Barber M, Boutkov B, Habegger L, Ferreira M, Baras A,
571 Reid J, Abecasis G, Maxwell E, Marchini J. Computationally efficient whole-
572 genome regression for quantitative and binary traits. *Nat Genet*. 2021;53:1097–
573 1103.
- 574 18. Sudlow C, Gallacher J, Allen N, Beral V, Burton P, Danesh J, Downey P, Elliott P,
575 Green J, Landray M, Liu B, Matthews P, Ong G, Pell J, Silman A, Young A,
576 Sprosen T, Peakman T, Collins R. UK biobank: an open access resource for
577 identifying the causes of a wide range of complex diseases of middle and old age.
578 *PLoS medicine*. 2015;12:e1001779.
- 579 19. Bycroft C, Freeman C, Petkova D, Band G, Elliott LT, Sharp K, Motyer A, Vukcevic
580 D, Delaneau O, O’Connell J, Cortes A, Welsh S, Young A, Effingham M, McVean
581 G, Leslie S, Allen N, Donnelly P, Marchini J. The UK Biobank resource with deep
582 phenotyping and genomic data. *Nature*. 2018;562:203.
- 583 20. Ge T, Chen C-Y, Ni Y, Feng Y-CA, Smoller JW. Polygenic prediction via Bayesian
584 regression and continuous shrinkage priors. *Nature Communications*.

- 585 2019;10:1776.
- 586 21. Karlson EW, Boutin NT, Hoffnagle AG, Allen NL. Building the Partners HealthCare
587 Biobank at Partners Personalized Medicine: Informed Consent, Return of Research
588 Results, Recruitment Lessons and Operational Considerations. *J Pers Med*
589 [Internet]. 2016 [cited 2020 Jul 14];6. Available from:
590 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4810381/>
- 591 22. Kowalski MH, Qian H, Hou Z, Rosen JD, Tapia AL, Shan Y, Jain D, Argos M,
592 Arnett DK, Avery C, Barnes KC, Becker LC, Bien SA, Bis JC, Blangero J,
593 Boerwinkle E, Bowden DW, Buyske S, Cai J, Cho MH, Choi SH, Choquet H,
594 Cupples LA, Cushman M, Daya M, Vries PS de, Ellinor PT, Faraday N, Fornage M,
595 Gabriel S, Ganesh S, Graff M, Gupta N, He J, Heckbert SR, Hidalgo B, Hodonsky
596 C, Irvin MR, Johnson AD, Jorgenson E, Kaplan R, Kardia SL, Kelly TN,
597 Kooperberg C, Lasky-Su JA, Loos RJJ, Lubitz SA, Mathias RA, McHugh CP,
598 Montgomery C, Moon J-Y, Morrison AC, Palmer ND, Pankratz N, Papanicolaou
599 GJ, Peralta JM, Peyser PA, Rich SS, Rotter JI, Silverman EK, Smith JA, Smith NL,
600 Taylor KD, Thornton TA, Tiwari HK, Tracy RP, Wang T, Weiss ST, Weng LC,
601 Wiggins KL, Wilson JG, Yanek LR, Zöllner S, North KN, Auer PL, Consortium NT-
602 O for PM (TOPMed), Hematology & Topm, Group HW, Raffield LM, Reiner AP,
603 Li Y. Use of >100,000 NHLBI Trans-Omics for Precision Medicine (TOPMed)
604 Consortium whole genome sequences improves imputation quality and detection of
605 rare variant associations in admixed African and Hispanic/Latino populations.
606 *bioRxiv*. 2019;683201.
- 607 23. Altshuler DM, Gibbs RA, Peltonen L, Altshuler DM, Gibbs RA, Peltonen L,
608 Dermitzakis E, Schaffner SF, Yu F, Peltonen L, Dermitzakis E, Bonnen PE,
609 Altshuler DM, Gibbs RA, de Bakker PIW, Deloukas P, Gabriel SB, Gwilliam R, Hunt
610 S, Inouye M, Jia X, Palotie A, Parkin M, Whittaker P, Yu F, Chang K, Hawes A,
611 Lewis LR, Ren Y, Wheeler D, Gibbs RA, Marie Muzny D, Barnes C, Darvishi K,
612 Hurles M, Korn JM, Kristiansson K, Lee C, McCarroll SA, Nemesh J, Dermitzakis
613 E, Keinan A, Montgomery SB, Pollack S, Price AL, Soranzo N, Bonnen PE, Gibbs
614 RA, Gonzaga-Jauregui C, Keinan A, Price AL, Yu F, Anttila V, Brodeur W, Daly MJ,
615 Leslie S, McVean G, Moutsianas L, Nguyen H, Schaffner SF, Zhang Q, Ghorri MJR,
616 McGinnis R, McLaren W, Pollack S, Price AL, Schaffner SF, Takeuchi F,
617 Grossman SR, Shlyakhter I, Hostetter EB, Sabeti PC, Adebamowo CA, Foster
618 MW, Gordon DR, Licinio J, Cristina Manca M, Marshall PA, Matsuda I, Ngare D,
619 Ota Wang V, Reddy D, Rotimi CN, Royal CD, Sharp RR, Zeng C, Brooks LD,
620 McEwen JE, The International HapMap 3 Consortium, Principal investigators,
621 Project coordination leaders, Manuscript writing group, Genotyping and QC,
622 ENCODE 3 sequencing and SNP discovery, Copy number variation typing and
623 analysis, Population analysis, Low frequency variation analysis, Linkage
624 disequilibrium and haplotype sharing analysis, et al. Integrating common and rare
625 genetic variation in diverse human populations. *Nature*. 2010;467:52–58.
- 626 24. Li Y, Willer CJ, Ding J, Scheet P, Abecasis GR. MaCH: Using Sequence and
627 Genotype Data to Estimate Haplotypes and Unobserved Genotypes. *Genet*
628 *Epidemiol*. 2010;34:816–834.
- 629 25. Venner E, Muzny D, Smith JD, Walker K, Neben CL, Lockwood CM, Empey PE,
630 Metcalf GA, Kachulis C, All of Us Research Program Regulatory Working Group,

- 631 Mian S, Musick A, Rehm HL, Harrison S, Gabriel S, Gibbs RA, Nickerson D, Zhou
632 AY, Doheny K, Ozenberger B, Topper SE, Lennon NJ. Whole-genome sequencing
633 as an investigational device for return of hereditary disease risk and
634 pharmacogenomic results as part of the All of Us Research Program. *Genome*
635 *Med.* 2022;14:34.
- 636 26. Lim M, Hastie T. Learning Interactions via Hierarchical Group-Lasso
637 Regularization. *Journal of Computational and Graphical Statistics.* 2015;24:627–
638 654.
- 639 27. DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or
640 more correlated receiver operating characteristic curves: a nonparametric
641 approach. *Biometrics.* 1988;44:837–845.
- 642 28. Pencina MJ, Steyerberg EW, D’Agostino RB. Extensions of net reclassification
643 improvement calculations to measure usefulness of new biomarkers. *Stat Med.*
644 2011;30:11–21.
- 645 29. Yengo L, Vedantam S, Marouli E, Sidorenko J, Bartell E, Sakaue S, Graff M,
646 Eliassen AU, Jiang Y, Raghavan S, Miao J, Arias JD, Mukamel RE, Spracklen CN,
647 Yin X, Chen S-H, Ferreira T, Ji Y, Karedera T, Lull K, Lin K, Malden DE, Medina-
648 Gomez C, Machado M, Moore A, Rueger S, Group GC-HW, Team 23andMe
649 Research, Program VMV, Initiative) D (DiscovEHR and MCH, Network) eMERGE
650 (Electronic MR and G, Study LC, Center RG, Consortium TP, Group USS,
651 Chasman DI, Cho YS, Heid IM, McCarthy MI, Ng MCY, O’Donnell CJ, Rivadeneira
652 F, Thorsteinsdottir U, Sun YV, Thai ES, Boehnke M, Deloukas P, Justice AE,
653 Lindgren CM, Loos RJF, Mohlke KL, North KE, Stefansson K, Walters RG, Winkler
654 TW, Young KL, Loh P-R, Yang J, Esko T, Assimes TL, Auton A, Abecasis GR,
655 Willer CJ, Locke AE, Berndt SI, Lettre G, Frayling TM, Okada Y, Wood AR,
656 Visscher PM, Hirschhorn JN. A Saturated Map of Common Genetic Variants
657 Associated with Human Height from 5.4 Million Individuals of Diverse Ancestries
658 [Internet]. 2022 [cited 2022 Jan 11]. Available from:
659 <https://www.biorxiv.org/content/10.1101/2022.01.07.475305v1>
- 660 30. Gurdasani D, Carstensen T, Fatumo S, Chen G, Franklin CS, Prado-Martinez J,
661 Bouman H, Abascal F, Haber M, Tachmazidou I, Mathieson I, Ekoru K, DeGorter
662 MK, Nsubuga RN, Finan C, Wheeler E, Chen L, Cooper DN, Schiffels S, Chen Y,
663 Ritchie GRS, Pollard MO, Fortune MD, Mentzer AJ, Garrison E, Bergström A,
664 Hatzikotoulas K, Adeyemo A, Doumatey A, Elding H, Wain LV, Ehret G, Auer PL,
665 Kooperberg CL, Reiner AP, Franceschini N, Maher D, Montgomery SB, Kadie C,
666 Widmer C, Xue Y, Seeley J, Asiki G, Kamali A, Young EH, Pomilla C, Soranzo N,
667 Zeggini E, Pirie F, Morris AP, Heckerman D, Tyler-Smith C, Motala AA, Rotimi C,
668 Kaleebu P, Barroso I, Sandhu MS. Uganda Genome Resource Enables Insights
669 into Population History and Genomic Discovery in Africa. *Cell.* 2019;179:984-
670 1002.e36.
- 671

672

673 Figure Legends

674 **Figure 1: Study cohorts**

675 The sample diagram depicts participant subsets. UK Biobank participants with MRI data
676 were split by random ID into a GWAS group (blue, N=39,524) and a residual group.
677 Individuals in the residual group related within 3 degrees of kinship were removed
678 (N=557), and the residual group was then split into a training group (purple, N=4,896)
679 and an internal validation group (yellow, N=4,962). As depicted in the legend, the
680 GWAS and PRS development was conducted in the GWAS group (blue, N=39,524).
681 Clinical model development was conducted in the GWAS and training groups (blue and
682 purple, N=44,420). The PRS was incorporated into the clinical score using a linear
683 model to produce the AORTA Gene model in the training group (purple, N=4,896). All
684 models were validated in the internal validation set (yellow, N=4,962) and the external
685 validation sets (orange): MGB, FHS, and *All of Us*. MGB: Mass General Brigham
686 Biobank. FHS: Framingham Heart Study. MRI: magnetic resonance imaging. GWAS:
687 genome-wide association study.

688 **Figure 2: Score calibration curves in the UK Biobank validation set**

689 Counterclockwise from top left: AORTA Gene, followed by the clinical AORTA Score,
690 the age and sex model, and the age/sex/genetics model. The x axis represents the
691 predicted ascending aortic diameter; the y axis represents the measured ascending
692 aortic diameter; both are truncated at the same points so that the x and y axes have the
693 same span. Each point represents one of the 4,962 UK Biobank internal validation set
694 participants; orange points represent women while navy points represent men. The blue

695 line shows the smoothed average value, and only extends along the x axis to the limits
696 of the observed data. The black line shows the line of ideal calibration, where a 1 cm
697 greater score would be met by a 1 cm greater aortic diameter. For all plotted scores, the
698 slopes were statistically indistinguishable from one and the intercepts were statistically
699 indistinguishable from zero as detailed in the Results.

700 **Figure 3: Variance in aortic diameter explained by the models**

701 For each cohort, the variance in ascending aortic diameter explained by the model
702 derived from age and sex (navy blue), the clinical model (AORTA Score, slate gray),
703 and the clinical model incorporating the polygenic score (AORTA Gene, orange) is
704 depicted for the internal validation set and the three external validation sets. MGB: Mass
705 General Brigham Biobank. FHS: Framingham Heart Study.

706 **Figure 4: Receiver operator characteristic curves**

707 Receiver operator characteristic (ROC) curves for the AORTA Gene model (red) and the clinical
708 AORTA Score model (blue) in the validation cohorts. The dashed diagonal line represents the
709 no-information baseline. Counterclockwise from top left: UK Biobank internal validation, MGB,
710 FHS, and *All of Us*. The AUROC for detecting aortic diameter ≥ 4 cm for the AORTA Gene model
711 in these cohorts was, respectively, 0.834, 0.808, 0.856, and 0.827. The respective AUROC for
712 the clinical AORTA Score was 0.765, 0.767, 0.818, and 0.791. MGB: Mass General Brigham
713 Biobank. FHS: Framingham Heart Study.

714 Tables

715 **Table 1: Participant characteristics**

	UK Biobank MRI Training	UK Biobank MRI Internal Validation	MGB TTE External Validation	FHS CT External Validation	All of Us TTE External Validation	All of Us Aneurysm External Validation
N	44,420	4,962	5,469	1,298	610	164,789
Men	21,053 (47.4%)	2,344 (47.2%)	2,580 (47.2%)	617 (47.5%)	246 (40.3%)	70,861 (43.0%)
Women	23,367 (52.6%)	2,618 (52.8%)	2,889 (52.8%)	681 (52.5%)	364 (59.7%)	93,928 (57.0%)
Age (years)	65.2 (7.7)	65.4 (7.7)	61.1 (14.4)	59.8 (9.0)	66.1 (11.0)	60.3 (10.9)
Medical history						
Hypertension	13,299 (29.9%)	1,501 (30.2%)	4,535 (82.9%)	532 (41.0%)	426 (69.8%)	57,845 (35.1%)
Hyperlipidemia	9,644 (21.7%)	1,103 (22.2%)	942 (17.2%)	257 (19.8%)	155 (25.4%)	6,379 (3.9%)
Diabetes	1,277 (2.9%)	160 (3.2%)	1,750 (32.0%)	115 (8.9%)	210 (34.4%)	25,568 (15.5%)
Heart rate (bpm)	68 (11)	68 (11)	76 (15)	65 (11)	69 (11)	72 (12)
Systolic blood pressure (mmHg)	140 (18)	141 (18)	128 (17)	126 (18)	129 (18)	129 (18)
Diastolic blood pressure (mmHg)	79 (10)	79 (10)	75 (10)	75 (9)	77 (11)	78 (11)
Height (cm)	169 (9)	169 (9)	169 (10)	168 (9)	166 (10)	168 (10)
Weight (kg)	75.2 (14.7)	75.4 (14.9)	84.4 (21.2)	80 (17.3)	88.9 (24.9)	84.5 (21.8)
BMI (kg/m²)	26.3 (4.2)	26.3 (4.3)	29.5 (6.6)	28.3 (5.2)	32.0 (8.4)	30.0 (7.3)
Ascending aortic diameter (cm)	3.18 (0.35)	3.19 (0.35)	3.20 (0.48)	3.45 (0.36)	3.34 (0.45)	NA
Aortic diameter ≥4cm	1,040 (2.3%)	113 (2.3%)	335 (6.1%)	116 (8.9%)	65 (10.7%)	NA
Prevalent aneurysm	NA	NA	NA	NA	NA	1,904 (1.2%)
Incident aneurysm	NA	NA	NA	NA	NA	1,632 (1.0%)

716 All values are mean (standard deviation) or count (percent of total). BPM: beats per
 717 minute. MGB: Mass General Brigham. BMI: body mass index. MRI: magnetic resonance
 718 imaging. CT: computed tomography. TTE: transthoracic echocardiography.

Figures

Figure 1: Study cohorts




Figure 2: Score calibration curves in the UK Biobank validation set




Figure 3: Variance in aortic diameter explained by the models




Figure 4: Receiver operator characteristic curves

