

Population study of post-stroke dysphagia

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

Health factors associated with development and severity of post-stroke dysphagia: an
epidemiological investigation.

Brittany N Krekeler^{1,2}, Heidi J P Schieve³, Jane Khoury⁴, Lili Ding,⁴ Mary Haverbusch², Kathleen
Alwell², Opeolu Adeoye⁵, Simona Feriolo², Jason Mackey⁶, Daniel Woo², Matthew Flaherty²,
Felipe De Los Rios La Rosa^{2,7}, Stacie Demel², Michael Star⁸, Elisheva Coleman⁹, Kyle Walsh²,
Sabreena Slavin¹⁰, Adam Jasne¹¹, Eva Mistry², Dawn Kleindorfer^{2,12} Brett Kissela²

¹University of Cincinnati College of Medicine, Department of Otolaryngology-Head and Neck Surgery
²University of Cincinnati College of Medicine, Department of Neurology and Rehabilitation Medicine
³Wake Forest School of Medicine
⁴Division of Biostatistics and Epidemiology, Cincinnati Children's Hospital Medical Center, Department of
Pediatrics, University of Cincinnati College of Medicine
⁵Washington University School of Medicine, Department of Emergency Medicine
⁶Indiana University School of Medicine, Department of Neurology
⁷Baptist Health South Florida, Miami Neuroscience Institute
⁸Sorkoa Medical Center, Israel
⁹University of Chicago Medicine, Department of Neurology
¹⁰University of Kansas Medical Center, Department of Neurology
¹¹Yale School of Medicine, Department of Neurology
¹²University of Michigan, Department of Neurology

Corresponding Author:
Brittany Krekeler, PhD, CCC-SLP
231 Albert Sabin Way, ML 0528
Room 6307C
Cincinnati, OH 45267

Population study of post-stroke dysphagia

33

34 **Abstract**

35 **Background and Purpose:** Dysphagia is a common post-stroke occurrence and has been
36 shown to impact patients' morbidity and mortality. The purpose of this study was to use a large
37 population-based dataset to determine specific epidemiological and patient health risk factors
38 that impact development and severity of dysphagia after acute stroke.

39 **Methods:** Using data from the Greater Cincinnati Northern Kentucky Stroke Study, GCNKSS,
40 involving a representative sample of approximately 1.3 million people from Southwest Ohio and
41 Northern Kentucky of adults (age ≥ 18), ischemic and hemorrhagic stroke cases from 2010 and
42 2015 were identified via chart review. Dysphagia status was determined based on bedside and
43 clinical assessments, and severity by necessity for alternative access to nutrition via nasogastric
44 (NG) or percutaneous endoscopic gastrostomy (PEG) tube placement. Comparisons between
45 patients with and without dysphagia were made to determine differences in baseline
46 characteristics and pre-morbid conditions. Multivariable logistic regression was used to
47 determine factors associated with increased risk of developing dysphagia.

48 **Results:** Dysphagia status was ascertained from 4139 cases (1709 with dysphagia). Logistic
49 regression showed: increased age, Black race, higher NIHSS score at admission, having a
50 hemorrhagic stroke (vs infarct), and right hemispheric stroke increased risk of developing
51 dysphagia after stroke. Factors associated with reduced risk included history of high cholesterol,
52 lower pre-stroke mRS score, and white matter disease.

53 **Conclusions:** This study replicated many previous findings of variables associated with
54 dysphagia (older age, worse stroke, right sided hemorrhagic lesions), while other variables
55 identified were without clear biological rationale (e.g. Black race, history of high cholesterol and
56 presence of white matter disease). These factors should be investigated in future, prospective
57 studies to determine biological relevance and potential influence in stroke recovery.

58

Population study of post-stroke dysphagia

59 **Introduction**

60 The development of swallowing impairments (dysphagia) has been shown to have an
61 impact on hospitalization timeline and comorbidities associated with acute stroke recovery.¹
62 Specifically, patients with dysphagia have poor nutrition and hydration,¹ both of which are
63 critical for neural recovery after stroke.^{1,2} Dysphagia is also associated with higher risk of
64 pneumonia, a serious and deadly complication of swallowing impairment.³⁻¹⁰

65 Prior investigations of dysphagia after ischemic or hemorrhagic stroke have focused
66 primarily on characterizing specific impairments in swallow physiology based on lesion location
67 and size,¹¹⁻¹³ or reporting on specific outcomes of these impairments in this population.^{3,14}
68 These investigations are highly relevant for assessing and treating the disorder and
69 understanding disposition of these patients. However, patient-specific factors related to
70 increased risk of developing dysphagia after stroke, and patient characteristics linked to severity
71 of dysphagia, are not well understood. These patient-specific factors include both pre-morbid
72 health conditions (e.g. heart disease) and patient demographics (e.g. sex, age, race), which in-
73 part derive social determinants of health (e.g. economic stability and health literacy/education)¹⁵
74 and are known to impact health outcomes at multiple levels.¹⁶ Gaining an understanding of
75 underlying conditions, biological, and socioeconomic risk factors that may predispose patients to
76 developing swallowing disorders after stroke would provide opportunities for preventative care.
77 To date, there have been no study populations large enough to address these gaps.

78 The goal of this work is to investigate incidence of dysphagia after stroke in a large
79 retrospective, population-based dataset to determine what patient-specific factors may influence
80 dysphagia status, or severity of impairment. These data will guide future investigations to better
81 understand how and why certain individuals develop swallowing impairments after stroke and
82 what can be done in acute clinical management of stroke to help prevent development of
83 swallowing disorders.

84 **Methods**

Population study of post-stroke dysphagia

85 The Greater Cincinnati Northern Kentucky Stroke Study (GCNKSS) has been previously
86 described in detail; ¹⁷⁻²¹ in brief, this is a population-based study using retrospective
87 ascertainment of physician adjudicated stroke cases in a sample of approximately 1.3 million
88 people across five counties in Southwestern Ohio and Northern Kentucky. All hospitalized adult
89 (age ≥ 18) ischemic and hemorrhagic stroke cases from 2010 and 2015 were identified using
90 International Classification of Diseases (ICD) stroke codes (ICD-9 430-436; ICD-10 I60-I69) at
91 15 area hospitals.

92 *Case Definition – Dysphagia vs No Dysphagia*

93 Dysphagia status was identified through Electronic Medical Record (EMR) review by
94 study team Registered Nurses. Individuals with documentation in the EMR of swallowing
95 impairment were included in the “Dysphagia” group. Given the known variability in swallowing
96 assessment practices both within and across institutions (e.g. timing of initial screening, formal
97 evaluation by a speech-language pathologist vs bedside screen by nursing staff, etc), ^{22,23}
98 documentation of dysphagia varied depending on the case, the treating institution, and clinical
99 practices at time of stroke. Assessment of dysphagia status was also documented in the EMR.
100 This could include a formal swallow assessment (bedside and/or instrumental by a speech-
101 language pathologist) or a nursing bedside screen. Only individuals who were determined not to
102 have dysphagia after one or more of these assessments were included in the “No Dysphagia”
103 group. Within the identified “Dysphagia” group, individuals were compared based on their need
104 for access to alternative nutrition during acute hospital stay. Individuals were categorized as one
105 of the following (Dysphagia PO Subgroups): 1) Dysphagia not requiring PEG/NG (i.e. dysphagia
106 management only required a modified texture and/or liquid diet), 2) Dysphagia requiring NG, 3)
107 Dysphagia requiring PEG. PO status was determined and confirmed through same chart review
108 procedures previously described. For cases where dysphagia status changed (i.e. progressed)
109 throughout inpatient stay, the most invasive intervention required was documented (i.e. PEG
110 was placed after NG).

Population study of post-stroke dysphagia

111 *Patient Characteristics - Demographics, Clinical Presentation, Pre-Morbid Conditions*

112 Patients included in the analyses were further characterized and compared (Dysphagia
113 vs No Dysphagia, and Dysphagia PO Subgroups). The following patient characteristics were
114 available for analysis, abstracted during the RN chart review process: 1) Patient demographic
115 and social determinants of health data: age, sex, race, ethnicity, and estimation of a
116 neighborhood Deprivation Index score;²⁴ 2) Clinical presentation at time of acute injury: National
117 Institutes of Health Stroke Scale [NIHSS], pre-stroke Modified Rankin Scale [mRS], Glasgow
118 Coma Score; 3) Stroke/lesion characteristics: type of case [ischemic/hemorrhagic], lesion
119 location [hemisphere], ischemic stroke subtype [small vessel, cardioembolic, large vessel]; 4)
120 Comorbidities (pre-morbid conditions): history of hypertension, diabetes, elevated cholesterol,
121 coronary artery disease, myocardial infarction [MI], congestive heart failure [CHF], prior stroke,
122 prior transient ischemic attack [TIA], dementia, brain injury or tumor, history of smoking, current
123 smoker (within the past six months), alcohol use, heavy alcohol use, and any white matter
124 disease [WMD, new and/or old]. All data regarding race and ethnicity categorization were
125 abstracted directly from the electronic medical record (EMR); as such we are unable to
126 definitively determine the source of the EMR reporting of race and ethnicity, whether it was self-
127 identification or assigned by the admitting medical professional.

128 *Statistical Analyses*

129 Study variables were summarized using descriptive statistics (mean and standard deviation or
130 median and interquartile range [IQR] and/or range for continuous variables and frequencies and
131 percentages for categorical variables) by study groups (Dysphagia status: Yes vs. Tested and
132 No; Dysphagia subgroups: Dysphagia Not Requiring PEG/NG vs. Dysphagia Requiring NG vs.
133 Dysphagia Requiring PEG; assessment: assessed vs. not assessed). Distribution was examined
134 for all continuous variables. Associations between categorical variables and study groups were
135 tested using either Chi-square or Fisher's exact tests. Associations between continuous variables
136 and study groups were tested using either two-sample t-tests, analysis of variance (ANOVA),

Population study of post-stroke dysphagia

137 Wilcoxon rank sum tests, or Kruskal-Wallis tests. Comparisons between the three Dysphagia
138 subgroups were adjusted for multiple comparisons using Bonferroni correction for categorical
139 variables, NIHSS and Glasgow Coma Score and by applying Dunnett's test, following ANOVA,
140 for age and deprivation index. Multivariable logistic regression analysis was carried out for
141 Dysphagia status (Yes vs. Assessed and No) and dichotomized Dysphagia subgroups
142 (Dysphagia requiring PEG or NG versus Dysphagia not requiring PEG or NG). P-values less or
143 equal to 0.05 were declared as statistically significant. All analysis was carried out in SAS® 9.4
144 (SAS Institute, Cary NC).

145 **Results**

146 A total of 5592 (2010 N=2644; 2015 N=2948) physician adjudicated and confirmed
147 hospitalized strokes from 15 area hospitals in the Greater Cincinnati and Northern Kentucky
148 region were reviewed.

149 *Dysphagia Status and Dysphagia Sub-Group*

150 Dysphagia status could only be ascertained from 4139 cases, which made up the final sample
151 for the analyses (**Figure 1**). Out of 4139, 1709 cases were determined to have dysphagia. Of
152 these cases, 944 did not require alternative access to nutrition (i.e. PEG/NG), 437 required NG,
153 and 328 required PEG placement (**Table 1**). In comparing the subgroups by year, there were no
154 differences between 2015 and 2010 in terms of proportion of individuals per Dysphagia
155 Subgroup.

156 *Patient Characteristics – Demographics and Clinical Presentation*

157 Patients with dysphagia were significantly older (mean 72 years vs 67.2 years
158 $p<0.0001$), a larger proportion were females (57% vs 52.2%, $p=0.002$), had higher median
159 NIHSS scores (8 vs 2; $p<0.0001$), lower median Glasgow Coma scores (14 vs 15, $p<0.0001$),
160 higher median pre-stroke mRS scores (2 vs 1, $p<0.0001$), and scored slightly higher on the
161 mean deprivation index ($p=0.01$) (**Table 2**).

Population study of post-stroke dysphagia

162 Stroke lesion characteristics between Dysphagia groups differed (**Table 3**): the
163 Dysphagia group had a larger proportion of hemorrhagic strokes as compared to those with No
164 Dysphagia (16.5% vs 11.8%, $p<0.0001$); the Dysphagia group had a greater proportion of “any
165 right hemisphere” stroke when compared to the No Dysphagia group (51.3% vs 45.8%,
166 $p=0.002$); amongst ischemic stroke cases, the Dysphagia group had a smaller proportion of
167 small vessel strokes (9.3% vs 20.7%, $p<0.0001$), a greater proportion of cardioembolic strokes
168 (38.5% vs 21.9, $p<0.0001$), and a smaller proportion of strokes where subtype could not be
169 determined (32% vs 38.2%, $p=0.0008$).

170 *Patient Characteristics - Pre-Morbid Conditions*

171 A larger proportion of individuals in the Dysphagia group as compared to the No
172 Dysphagia group had a history of hypertension (84.5% vs 81%, $p=0.004$), a history of coronary
173 artery disease (36.2% vs 32%, $p=0.0005$), history of congestive heart failure (23.5% vs 15.6%,
174 $p<0.0001$), history of prior stroke (28.4% vs 25.6%, $p=0.04$), and a history of dementia (16.7%
175 vs 8.4%, $p<0.0001$) (**Table 4**). There were a larger proportion of individuals in the No Dysphagia
176 group who had a history of TIA (16.6% vs 13%, $p=0.002$), who were current smokers (28.2% vs
177 23.8%, $p=0.002$) who reported alcohol use (at least 1-2 drinks per day; 41.7% vs 34.7%,
178 $p<0.0001$), and those who had white matter disease (71.6% vs 68.5%, $p=0.03$).

179 *Logistic Regression Model for Factors related to Dysphagia Status*

180 The variables that were considered for the logistic regression can be found in **Figure 2**.
181 Variables that were included in the final model included age (per year increase), race (Black vs
182 non-Black), baseline mRS score (0-1, i.e. functional vs 2-5), NIHSS (per point increase),
183 Hemorrhage vs Infarct, lesion on the right side, high cholesterol, coronary artery disease,
184 presence of white matter disease, deprivation index, and year of stroke (2015 vs 2010). The
185 logistic regression model (ROC= 0.78; **Figure 2**) for dysphagia status revealed an increased
186 risk of developing dysphagia after stroke for older individuals (Age/1 year OR=1.022, $p<0.0001$),
187 people of Black race (OR=1.225, $p=0.04$), cases with higher NIHSS score (per point increase

Population study of post-stroke dysphagia

188 OR=1.133, $p<0.001$), cases with a hemorrhagic stroke vs an ischemic infarct (OR=1.472,
189 $p=0.0004$), and having the stroke occur in the right hemisphere (OR=1.269, $p=0.0011$). Factors
190 showing decreased risk of dysphagia included having lower baseline mRS scores of 0-1
191 (OR=0.657, $p<0.0001$), having high cholesterol (OR=0.84, $p=0.02$), having white matter disease
192 (new or old, OR=0.801, $p=0.01$), and having been a case in 2015 (OR=0.82, $p=0.007$).

193 Discussion

194 In this large, population-based study we sought to characterize differences in patient
195 factors underlying dysphagia status for individuals after stroke and to determine which of these
196 factors were associated with an increased risk in developing dysphagia post stroke.

197 Less than half of our final sample with positive dysphagia status required alternative
198 access to nutrition via PEG or NG tube (**Table 1**). We found a greater proportion of individuals
199 with dysphagia compared to those with no dysphagia were older, female, and had higher
200 impairment scores upon stroke admission (i.e. NIHSS, GCS) and pre-stroke mRS (**Table 2**). It
201 has been well established that increased age results in age-related changes to swallow
202 movements and timing²⁵⁻²⁹ that may predispose older individuals to developing swallowing
203 impairments after stroke. Regarding our finding associating female sex with swallowing
204 impairments after stroke, a recent study showed that women had more severe outcomes after
205 ischemic stroke as compared to men, which could result in worse swallowing outcomes after
206 stroke.³⁰ However, another study of 578 patients in Seoul, South Korea found no relationship
207 between swallowing outcomes after stroke and patient sex.³¹ These alternative findings could
208 be due to regional and cultural differences. Published evidence about influence of racial and
209 ethnic background on stroke severity and outcome measures is mixed.³² However, some
210 literature suggests an ethnicity-age interaction, which could not be examined in these data, due
211 to a small number of Hispanic individuals in our study population, could be influential. Other
212 evidence points to some impact of socio-economic disparities amongst specific ethnic groups
213 could be a contributing factor,³³ which is also supported by our data. Our data also showed that

Population study of post-stroke dysphagia

214 Deprivation Index scores were slightly higher in the Dysphagia group as compared to the No
215 Dysphagia group, however, these were not found to be influential in the logistic regression
216 (**Figure 2**). These discrepancies in the literature regarding the influence of sex and ethnicity in
217 stroke-related outcomes, in general and specific to swallowing, highlight that more work is
218 needed to investigate biological drivers for these differences.

219 Lesion type differed between the Dysphagia and No Dysphagia Groups, with a greater
220 proportion of hemorrhagic, and right hemisphere strokes in the Dysphagia Group, which is
221 supported by the literature.^{13,34,35} Similarly, our findings of more dysphagia occurring with
222 increasing stroke severity (NIHSS, Glasgow Coma score,), initial pre-stroke status as measured
223 by mRS, hemorrhagic stroke, and right hemisphere lesions are also consistent with prior
224 literature.^{13,35-40} There have been some other studies, with smaller samples,⁴¹⁻⁴³ questioning
225 the influence of laterality in development of dysphagia, but our findings provide more evidence
226 towards hemispheric location of lesion being a potential risk factor for developing swallowing
227 impairments. Our findings for stroke subtype also demonstrated that there were a greater
228 proportion of cardioembolic strokes in the Dysphagia Group with ischemic stroke. This could be
229 related to the increased likelihood of cardioembolic infarcts resulting in hemorrhagic
230 transformation,⁴⁴ and as noted, hemorrhagic strokes may be more likely to present with
231 dysphagia.

232 Regarding pre-morbid factors and development of dysphagia, we found that a greater
233 proportion of individuals with a history of coronary artery disease, congestive heart failure, prior
234 stroke, dementia, and hypertension in the Dysphagia Group as compared to the No Dysphagia
235 Group.^{45,46} Alcohol use, current cigarette smoking status, high cholesterol and prior TIA were
236 also significantly different between the groups, with a higher proportion of individuals in the No
237 Dysphagia group. While prior studies don't directly study effects of alcohol consumption on
238 development of dysphagia, alcohol use is known as risk factor for stroke.^{47,48} However, studies
239 have also shown that moderate alcohol use may reduce risk of ischemic stroke,⁴⁹ and that

Population study of post-stroke dysphagia

240 alcohol use does not affect stroke severity or outcomes after stroke.⁵⁰ These mixed results
241 suggest that there may be a more complex relationship between alcohol consumption and
242 stroke-related recovery, including swallowing, that has yet to be fully discovered.

243 Prior literature clearly demonstrate that individuals with cardiac conditions and
244 dysphagia have a more difficult recovery and often poorer outcomes.^{51,52} However, it is difficult
245 to determine whether these cardiac conditions made patients more susceptible to developing
246 dysphagia after stroke simply because these individuals are more medically compromised, or if
247 there is a causal relationship between these conditions and swallowing impairments. In fact, our
248 logistic regression results did not indicate increased risk of dysphagia development for patients
249 with a history of coronary artery disease after accounting for other potential risk factors (**Table**
250 **4**), which may indicate the former hypothesis being more explanatory of these findings.

251 Other factors that were predictive of developing dysphagia after stroke included
252 increased age, Black race, higher NIHSS score at presentation, having a hemorrhagic stroke
253 (vs infarct), and right hemispheric stroke. Factors associated with decreased risk included
254 patients having a history of high cholesterol, lower pre-stroke mRS, and any white matter
255 disease detected on clinical imaging. Several findings were either surprising, or are still unclear
256 in their meaning. As discussed, literature on ethnic and racial disparities in stroke severity and
257 post-stroke care shows very mixed evidence and appears to be influenced by regional location
258 and/or size of study. Prior published data have indicated that Black individuals have more
259 severe strokes acutely than White individuals.⁵³ Current evidence provided in our study
260 provides further emphasis of need for better understanding contributions of biological vs
261 socioeconomic disparity to stroke-related outcomes in Black patients vs other racial categories.

262 ⁵⁴

263 The findings regarding hyperlipidemia and any evidence of white matter disease on
264 clinical stroke imaging being protective for risk of developing dysphagia after stroke were also
265 unexpected. One hypothesis regarding the relationship between hyperlipidemia and lesser risk

Population study of post-stroke dysphagia

266 of dysphagia could be due to use of statin medications in this population. Literature supports the
267 possibility that use of statins pre-stroke may reduce severity of stroke.⁵⁵ One study specifically
268 looking at effects of statin therapy and development of post-stroke pneumonia suggested that
269 patients who were using statins prior to admission for stroke and were treated with thrombolysis
270 during admission were less likely to develop a pneumonia.⁵⁶ This finding should be explored
271 more specifically in future studies of post-stroke dysphagia development, as this could be a
272 potential protective treatment for individuals at risk of both stroke and dysphagia. In terms of any
273 evidence of white matter disease serving as protective, this could be an incidental finding driven
274 by interdependency among the many factors explored in this study. Given that we were unable
275 to measure new white matter disease in these patients at the time of clinical imaging (i.e.
276 differentiate new WMD from existing WMD before acute admission), it is difficult to tease out the
277 meaning of this particular finding. Design of future studies of white matter disease and
278 dysphagia could potentially include a cohort of repeat stroke patients with available prior clinical
279 imaging in an attempt to better understand the relationship between existing and new WMD in
280 post-stroke dysphagia.

281 **Limitations**

282 The primary limitations of this study are due to the nature of retrospective analysis of data
283 collected as part of a large data set; therefore, several broad and more nuanced factors were
284 not available for use in our data analysis for both epidemiological factors and patient medical
285 history. Most notably, this included the type of medical professional completing dysphagia
286 evaluation and/or screening (e.g. speech-language pathologist or nursing professional). Further,
287 the source of patient demographic factors (i.e. self-report vs determination by medical
288 professional) related to race and ethnicity are not able to be determined in this type of
289 retrospective dataset. These factors may have impacted classification of patients within certain
290 groupings in our analysis. Further, the inability to differentiate new vs old WMD limits our ability
291 to fully interpret the role of WMD in development of dysphagia after stroke.

Population study of post-stroke dysphagia

292 **Conclusions**

293 This large, population-based study confirms several factors that have been frequently
294 demonstrated in the literature to be associated with developing dysphagia after stroke, including
295 older age, dementia status, more severe stroke, and lesion type and location. However, other
296 factors emerged that are not clear in their biological meaning or association, particularly findings
297 related to certain racial/ethnic groups being significant factors for dysphagia development after
298 stroke. These and other factors indicating a reduced risk, hyperlipidemia and any presence of
299 white matter disease, should be considered in future prospective work to better understand
300 these relationships and how these conditions may affect acute stroke recovery in the setting of
301 dysphagia.

302

Population study of post-stroke dysphagia

- 303 **Sources of Funding:** National Institute of Neurologic Disorders and Stroke,
304 R01NS030678.
- 305 **Disclosures:**
- 306 BN Krekeler: Reports research support from the National Institutes of Health
307 NIH/NCMRR/NICHHD
- 308 HJP Schieve: No relevant disclosures
- 309 J Khoury: Reports research support from the National Institutes of Health NIH/NINDS
- 310 L Ding: No relevant disclosures
- 311 M Haverbusch: Reports research support from the National Institutes of Health NIH/NINDS
- 312 K Alwell: No relevant disclosures
- 313 O Adeoye: Reports research support from the National Institutes of Health NIH/NINDS and
314 Equity Holder and Co-Founder – Sense Diagnostics, Inc.
- 315 S Ferioli: No relevant disclosures
- 316 J Mackey: Reports research support from the National Institutes of Health NIH/NINDS
- 317 D Woo: Reports research support from the National Institutes of Health NIH/NINDS
- 318 M Flaherty: Speaker’s bureau- CSL Behring and Alexion, Co-founder, employee, and equity
319 holder: Sense Diagnostics, Inc., Clinical trial endpoint adjudication committee member:
320 Boehringer Ingelheim and LG Chem
- 321 F De Los Rios La Rosa:
- 322 S Demel: No relevant disclosures
- 323 M Star: No relevant disclosures
- 324 E Coleman: No relevant disclosures
- 325 K Walsh: AstraZeneca: Speaker’s Bureau American Heart Association, NIH, Sense Diagnostics
326 LLC, Jan Medical Inc: Grant Funding/Research Support
- 327 S Slavin: No relevant disclosures
- 328 A Jasne: No relevant disclosures

Population study of post-stroke dysphagia

329 E Mistry: Reports research support from the National Institutes of Health NIH/NINDS

330 D Kleindorfer: No relevant disclosures

331 B Kissela: Reports research support from the National Institutes of Health NIH/NINDS

332

333

334

335

Population study of post-stroke dysphagia

336 **Table 1.** Dysphagia Subgroup Status by Year

337

Dysphagia Subgroup n = 1709	Dysphagia Not Requiring PEG/NG n =944	Dysphagia Requiring NG n = 437	Dysphagia Requiring PEG n = 328
Year			
2010 (n=782)	406 (51.9%)	223 (28.5%)	153 (19.6%)
2015 (n=927)	538 (58.0%)	214 (23.1%)	175 (18.9%)
p-value (2010 vs 2015)	0.03	0.03	1.0

*All p-values are Bonferroni adjusted

338

339

Population study of post-stroke dysphagia

340 **Table 2.** Demographics and Clinical Presentation by Dysphagia Status

	Dysphagia	No Dysphagia	p-value
N	1709	2430	
Age, year, mean (std)	72.0 (14.7)	67.2 (14.8)	<.0001
Race			0.06
White + other	1278 (74.8%)	1877 (77.3%)	
Black	431 (25.2%)	551 (22.7%)	
Unknown*	1 (0.06%)	1 (0.04%)	
Ethnicity			0.02
Hispanic	6 (0.4%)	14 (0.6%)	
Non-Hispanic	1211 (70.9%)	1804 (74.3%)	
Unknown	492 (28.8%)	611 (25.2%)	
Sex (M vs F)			0.002
Males	735 (43.0%)	1161 (47.8%)	
Females	974 (57.0%)	1268 (52.2%)	
Baseline NIHSS			<.0001
Median (25th, 75th %ile)	8 (3, 16)	2 (1, 5)	
[min, max]	[0, 40]	[0, 39]	
Glasgow Coma Score			<.0001
Median (25th, 75th %ile)	14 (11, 15)	15 (15,15)	
[min, max]	[3, 15]	[3, 15]	
Baseline mRS			<.0001
Median (25th, 75th %ile)	2 (0, 3)	1 (0, 3)	
[min, max]	[0, 5]	[0, 5]	
Baseline mRS 0-1	610 (35.8%)	1280 (52.7%)	<.0001
Deprivation Index, mean (std)	0.38 (0.13)	0.37 (0.13)	0.01

*Unknown not included in statistical comparison

341

342

343

Population study of post-stroke dysphagia

344 **Table 3.** Lesion Characteristics by Dysphagia Status

	Dysphagia	No Dysphagia	p-value ³⁴⁵
N	1709	2430	
Type of Case			<.0001*
Infarct	1426 (83.4%)	2142 (88.1%)	
Hemorrhagic (ICH/SAH)	281 (16.5%)	287 (11.8%)	
Unknown†	2 (0.1%)	1 (0.03%)	
Lesion Location(s)			<.0001*
Left Hemisphere only	681 (39.8%)	1017 (41.8%)	
Any Left Hemisphere	946 (55.4%)	1292 (53.2%)	0.64**
Right Hemisphere only	610 (35.7%)	854 (35.1%)	
Any Right Hemisphere	877 (51.3%)	1113 (45.8%)	0.002**
Brainstem only	4 (0.2%)	7 (0.3%)	
Any Brainstem	153 (9.0%)	187 (7.7%)	0.57**
Mixed Hemispheric (L+R only)	171 (10.0%)	165 (6.8%)	
Mixed (L and/or R + brainstem)	149 (8.7%)	180 (7.4%)	
Unknown	94 (5.5%)	207 (8.5%)	
Stroke subtype Infarct only			<.0001*
Small vessel (1)	132 (9.3%)	444 (20.7%)	<.0001**
Cardioembolic (2)	549 (38.5%)	470 (21.9%)	<.0001**
Large vessel (3)	189 (13.3%)	294 (13.7%)	0.97**
Other identified (5)	99 (7.0%)	116 (5.4%)	0.24**
Undetermined (8)	456 (32.0%)	818 (38.2%)	0.0008**

*overall significance; **after correcting for multiple testing; †unknown not included in the analysis

Population study of post-stroke dysphagia

346 **Table 4.** Pre-Morbid Conditions by Dysphagia Status

	Dysphagia	No Dysphagia	p-value
N	1709	2430	
History of Hypertension	1444 (84.5%)	1969 (81.0%)	0.004
History of Diabetes	611 (35.8%)	870 (35.8%)	0.97
History of Elevated Cholesterol	944 (55.2%)	1405 (57.8%)	0.10
Coronary Heart Disease	794 (46.5%)	938 (38.6%)	<.0001
History of Coronary Artery Disease	618 (36.2%)	777 (32.0%)	0.005
History of Myocardial Infarction	255 (14.9%)	341 (14.0%)	0.42
History of Congestive Heart Failure	401 (23.5%)	379 (15.6%)	<.0001
Prior Stroke	486 (28.4%)	621 (25.6%)	0.04
Prior TIA	223 (13.0%)	404 (16.6%)	0.002
History of Dementia	286 (16.7%)	204 (8.4%)	<.0001
History of Brain Injury/Tumor	12 (0.7%)	15 (0.6%)	0.84
History of Smoking (Cigarettes)	949 (55.5%)	1400 (57.6%)	0.18
Current Smoker (Cigarettes)	407 (23.8%)	686 (28.2%)	0.002
Alcohol Use	593 (34.7%)	1013 (41.7%)	<.0001
Heavy Alcohol use	159 (9.3%)	212 (8.7%)	0.52
White Matter Disease (any)	1171 (68.5%)	1739 (71.6%)	0.03

347


348

349

Population study of post-stroke dysphagia


350 **Figure 1.** Number of Strokes included from Greater Cincinnati Northern Kentucky Stroke Study
351 in 2010 and 2015

352



Population study of post-stroke dysphagia

353 **Figure 2.** Logistic Regression by Dysphagia Status



354

Variable	Adjusted Odds Ratio (aOR)	Lower 95% CI	Upper 95% CI	p-value
Age / 1 year	1.022	1.016	1.028	<.0001
Black Race	1.225	1.013	1.481	0.04
Baseline mRS 0-1	0.657	0.562	0.769	<.0001
NIHSS / 1 point increase	1.133	1.120	1.147	<.0001
Hemorrhage vs Infarct	1.472	1.189	1.822	0.0004
Right hemispheric stroke	1.269	1.100	1.463	0.0011
High Cholesterol	0.840	0.721	0.978	0.02
Coronary Artery Disease	1.153	0.990	1.343	0.07
White Matter Disease (any)	0.801	0.676	0.949	0.01
Deprivation Index	1.673	0.904	3.096	0.10
Year 2015	0.820	0.710	0.947	0.007

355

Population study of post-stroke dysphagia

356 References

- 357 1. Crary MA, Humphrey JL, Carnaby-Mann G, Sambandam R, Miller L, Silliman S.
358 Dysphagia, nutrition, and hydration in ischemic stroke patients at admission and
359 discharge from acute care. *Dysphagia*. 2013;28:69-76.
- 360 2. Ojo O, Brooke J. The use of enteral nutrition in the management of stroke. *Nutrients*.
361 2016;8:827.
- 362 3. Martino R, Foley N, Bhogal S, Diamant N, Speechley M, Teasell R. Dysphagia after
363 stroke: incidence, diagnosis, and pulmonary complications. *stroke*. 2005;36:2756-2763.
- 364 4. Perry L, Love CP. Screening for dysphagia and aspiration in acute stroke: a systematic
365 review. *Dysphagia*. 2001;16:7-18.
- 366 5. Eslick GD, Talley N. Dysphagia: epidemiology, risk factors and impact on quality of life—a
367 population-based study. *Alimentary pharmacology & therapeutics*. 2008;27:971-979.
- 368 6. Humbert IA, Robbins J. Dysphagia in the elderly. *Physical medicine and rehabilitation
369 clinics of North America*. 2008;19:853-866.
- 370 7. Altman KW, Yu G-P, Schaefer SD. Consequence of dysphagia in the hospitalized
371 patient: impact on prognosis and hospital resources. *Archives of Otolaryngology–Head &
372 Neck Surgery*. 2010;136:784-789.
- 373 8. Cabre M, Serra-Prat M, Palomera E, Almirall J, Pallares R, Clavé P. Prevalence and
374 prognostic implications of dysphagia in elderly patients with pneumonia. *Age and ageing*.
375 2010;39:39-45.
- 376 9. Cichero JA, Altman KW. Definition, prevalence and burden of oropharyngeal dysphagia:
377 a serious problem among older adults worldwide and the impact on prognosis and
378 hospital resources. In: *Stepping stones to living well with dysphagia*. Karger Publishers;
379 2012:1-11.
- 380 10. Cohen DL, Roffe C, Beavan J, Blackett B, Fairfield CA, Hamdy S, Havard D, McFarlane
381 M, McLaughlin C, Randall M. Post-stroke dysphagia: A review and design considerations
382 for future trials. *International Journal of Stroke*. 2016;11:399-411.
- 383 11. Wilmskoetter J, Bonilha L, Martin-Harris B, Elm JJ, Horn J, Bonilha HS. Mapping acute
384 lesion locations to physiological swallow impairments after stroke. *NeuroImage: Clinical*.
385 2019;22:101685.
- 386 12. Wilmskoetter J, Martin-Harris B, Pearson Jr WG, Bonilha L, Elm JJ, Horn J, Bonilha HS.
387 Differences in swallow physiology in patients with left and right hemispheric strokes.
388 *Physiology & behavior*. 2018;194:144-152.
- 389 13. Daniels SK, Pathak S, Mukhi SV, Stach CB, Morgan RO, Anderson JA. The relationship
390 between lesion localization and dysphagia in acute stroke. *Dysphagia*. 2017;32:777-784
391 @ 1432-0460.
- 392 14. Mann G, Hankey GJ, Cameron D. Swallowing function after stroke: prognosis and
393 prognostic factors at 6 months. *Stroke*. 1999;30:744-748.
- 394 15. Kunitz SJ. Sex, race and social role—history and the social determinants of health.
395 *International Journal of Epidemiology*. 2006;36:3-10. doi: 10.1093/ije/dyl296
- 396 16. U.S. Department of Health and Human Services. Social Determinants of Health; Healthy
397 People 2030. Retrieved from [https://health.gov/healthypeople/objectives-and-](https://health.gov/healthypeople/objectives-and-data/social-determinants-health)
398 [data/social-determinants-health](https://health.gov/healthypeople/objectives-and-data/social-determinants-health)
- 399 17. Broderick J, Brott T, Kothari R, Miller R, Khoury J, Pancioli A, Gebel J, Mills D, Minneci
400 L, Shukla R. The Greater Cincinnati/Northern Kentucky Stroke Study: preliminary first-
401 ever and total incidence rates of stroke among blacks. *Stroke*. 1998;29:415-421.
- 402 18. Zakaria T, Lindsell CJ, Kleindorfer D, Alwell K, Moomaw CJ, Woo D, Szaflarski JP,
403 Khoury J, Miller R, Broderick JP. Age accounts for racial differences in ischemic stroke

Population study of post-stroke dysphagia

- 404 volume in a population-based study. *Cerebrovascular Diseases*. 2008;26:376-380 %@
405 1015-9770.
- 406 19. Kissela B, Broderick J, Woo D, Kothari R, Miller R, Khoury J, Brott T, Pancioli A, Jauch
407 E, Gebel J. Greater Cincinnati/Northern Kentucky Stroke Study: volume of first-ever
408 ischemic stroke among blacks in a population-based study. *Stroke*. 2001;32:1285-1290.
- 409 20. Madsen TE, Khoury JC, Leppert M, Alwell K, Moomaw CJ, Sucharew H, Woo D, Ferioli
410 S, Martini S, Adeoye O. Temporal trends in stroke incidence over time by sex and age in
411 the GCNKSS. *Stroke*. 2020;51:1070-1076 %@ 0039-2499.
- 412 21. Madsen TE, Cummings OW, De Los Rios La Rosa F, Khoury JC, Alwell K, Woo D,
413 Ferioli S, Martini S, Adeoye O, Khatri P. Substance Use and Performance of Toxicology
414 Screens in the Greater Cincinnati Northern Kentucky Stroke Study. *Stroke*.
415 2022;53:3082-3090.
- 416 22. Chen BJ, Suolang D, Frost N, Faigle R. Practice Patterns and Attitudes Among Speech-
417 Language Pathologists Treating Stroke Patients with Dysphagia: A Nationwide Survey.
418 *Dysphagia*. 2022;37:1715-1722.
- 419 23. Fairfield CA, G. Smithard D. Assessment and management of dysphagia in acute stroke:
420 An initial service review of international practice. *Geriatrics*. 2020;5:4.
- 421 24. Brokamp C, Beck AF, Goyal NK, Ryan P, Greenberg JM, Hall ES. Material community
422 deprivation and hospital utilization during the first year of life: an urban population-based
423 cohort study. *Annals of epidemiology*. 2019;30:37-43.
- 424 25. Robbins J, Humpal NS, Banaszynski K, Hind J, Rogus-Pulia N. Age-related differences
425 in pressures generated during isometric presses and swallows by healthy adults.
426 *Dysphagia*. 2016;31:90-96.
- 427 26. Azzolino D, Damanti S, Bertagnoli L, Lucchi T, Cesari M. Sarcopenia and swallowing
428 disorders in older people. *Aging Clinical & Experimental Research*. 2019;31:799-805.
429 doi: 10.1007/s40520-019-01128-3
- 430 27. Wakabayashi H. Presbyphagia and Sarcopenic Dysphagia: Association between Aging,
431 Sarcopenia, and Deglutition Disorders. *J Frailty Aging*. 2014;3:97-103. doi:
432 10.14283/jfa.2014.8
- 433 28. Robbins J, Levine R, Wood J, Roecker EB, Luschei E. Age effects on lingual pressure
434 generation as a risk factor for dysphagia. *J Gerontol A Biol Sci Med Sci*. 1995;50:M257-
435 262.
- 436 29. Robbins JA. Normal swallowing and aging. Paper/Poster presented at: Seminars in
437 neurology; 1996;
- 438 30. Phan HT, Reeves MJ, Blizzard CL, Thrift AG, Cadilhac DA, Sturm J, Otahal P, Rothwell
439 P, Bejot Y, Cabral NL. Sex differences in severity of stroke in the INSTRUCT study: a
440 meta-analysis of individual participant data. *Journal of the American Heart Association*.
441 2019;8:e010235.
- 442 31. Kim J, Oh B-M, Lee GJ, Lee SA, Chun SW, Han TR. Clinical factors associated with
443 severity of post-stroke dysphagia. *Brain & Neurorehabilitation*. 2011;4:116-120.
- 444 32. Ellis C, Hyacinth HI, Beckett J, Feng W, Chimowitz M, Ovbiagele B, Lackland D, Adams
445 R. Racial/Ethnic Differences in Poststroke Rehabilitation Outcomes. *Stroke Research
446 and Treatment*. 2014;2014:950746. doi: 10.1155/2014/950746
- 447 33. Stansbury JP, Jia H, Williams LS, Vogel WB, Duncan PW. Ethnic disparities in stroke:
448 epidemiology, acute care, and postacute outcomes. *Stroke*. 2005;36:374-386.
- 449 34. Paciaroni M, Mazzotta G, Corea F, Caso V, Venti M, Milia P, Silvestrelli G, Palmerini F,
450 Parnetti L, Gallai V. Dysphagia following stroke. *European neurology*. 2004;51:162-167.
- 451 35. Suntrup S, Kemmling A, Warnecke T, Hamacher C, Oelenberg S, Niederstadt T, Heindel
452 W, Wiendl H, Dziewas R. The impact of lesion location on dysphagia incidence, pattern
453 and complications in acute stroke. Part 1: dysphagia incidence, severity and aspiration.
454 *European journal of neurology*. 2015;22:832-838.

Population study of post-stroke dysphagia

- 455 36. Henke C, Foerch C, Lapa S. Early screening parameters for dysphagia in acute
456 ischemic stroke. *Cerebrovascular Diseases*. 2017;44:285-290.
- 457 37. Jeyaseelan RD, Vargo MM, Chae J. National Institutes of Health Stroke Scale (NIHSS)
458 as an early predictor of poststroke dysphagia. *PM&R*. 2015;7:593-598.
- 459 38. Crary MA, Carnaby-Mann GD, Miller L, Antonios N, Silliman S. Dysphagia and nutritional
460 status at the time of hospital admission for ischemic stroke. *Journal of Stroke and*
461 *cerebrovascular diseases*. 2006;15:164-171.
- 462 39. Khedr EM, Abbass MA, Soliman RK, Zaki AF, Gamea A. Post-stroke dysphagia:
463 frequency, risk factors, and topographic representation: hospital-based study. *The*
464 *Egyptian Journal of Neurology, Psychiatry and Neurosurgery*. 2021;57:1-8.
- 465 40. Robbins J, Levine RL, Maser A, Rosenbek JC, Kempster GB. Swallowing after unilateral
466 stroke of the cerebral cortex. *Archives of physical medicine and rehabilitation*.
467 1993;74:1295-1300.
- 468 41. Fernández-Pombo A, Seijo-Raposo IM, López-Osorio N, Cantón-Blanco A, González-
469 Rodríguez M, Arias-Rivas S, Rodríguez-Yáñez M, Santamaría-Nieto A, Díaz-Ortega C,
470 Gómez-Vázquez E. Lesion location and other predictive factors of dysphagia and its
471 complications in acute stroke. *Clinical nutrition ESPEN*. 2019;33:178-182.
- 472 42. Suntrup-Krueger S, Kemmling A, Warnecke T, Hamacher C, Oelenberg S, Niederstadt
473 T, Heindel W, Wiendl H, Dziewas R. The impact of lesion location on dysphagia
474 incidence, pattern and complications in acute stroke. Part 2: Oropharyngeal residue,
475 swallow and cough response, and pneumonia. *European journal of neurology*.
476 2017;24:867-874.
- 477 43. Sharma J, Fletcher S, Vassallo M, Ross I. Prognostic value of CT scan features in acute
478 ischaemic stroke and relationship with clinical stroke syndromes. *International journal of*
479 *clinical practice*. 2000;54:514-518.
- 480 44. Hornig CR, Bauer T, Simon C, Trittmacher S, Dorndorf W. Hemorrhagic transformation
481 in cardioembolic cerebral infarction. *Stroke*. 1993;24:465-468.
- 482 45. Easterling CS, Robbins E. Dementia and dysphagia. *Geriatric Nursing*. 2008;29:275-
483 285.
- 484 46. Smithard DG, O'Neill PA, England RE, Park CL, Wyatt R, Martin DF, Morris J. The
485 natural history of dysphagia following a stroke. *Dysphagia*. 1997;12:188-193.
- 486 47. Shiotsuki H, Saijo Y, Ogushi Y, Kobayashi S. Relationship between Alcohol Intake and
487 Stroke Severity in Japanese Patients: a Sex-and Subtype-Stratified Analysis. *Journal of*
488 *Stroke and Cerebrovascular Diseases*. 2022;31:106513.
- 489 48. O'donnell MJ, Xavier D, Liu L, Zhang H, Chin SL, Rao-Melacini P, Rangarajan S, Islam
490 S, Pais P, McQueen MJ. Risk factors for ischaemic and intracerebral haemorrhagic
491 stroke in 22 countries (the INTERSTROKE study): a case-control study. *The Lancet*.
492 2010;376:112-123.
- 493 49. Elkind MS, Sciacca R, Boden-Albala B, Rundek T, Paik MC, Sacco RL. Moderate
494 alcohol consumption reduces risk of ischemic stroke: the Northern Manhattan Study.
495 *Stroke*. 2006;37:13-19.
- 496 50. Fekete K, Szatmári S, Szócs I, Szekeres C, Szász J, Mihálka L, Smolanka V, Kardos L,
497 Csiba L, Bereczki D. Prestroke alcohol consumption and smoking are not associated
498 with stroke severity, disability at discharge, and case fatality. *Journal of Stroke and*
499 *Cerebrovascular Diseases*. 2014;23:e31-e37.
- 500 51. Matsuo H, Yoshimura Y, Fujita S, Maeno Y. Dysphagia is associated with poor physical
501 function in patients with acute heart failure: a prospective cohort study. *Aging Clinical*
502 *and Experimental Research*. 2020;32:1093-1099.
- 503 52. Matsuo H, Yoshimura Y, Fujita S, Maeno Y. Incidence of Dysphagia and Its Association
504 With Functional Recovery and 1-Year Mortality in Hospitalized Older Patients With Heart

Population study of post-stroke dysphagia

- 505 Failure: A Prospective Cohort Study. *Journal of Parenteral and Enteral Nutrition*.
506 2021;45:372-380.
- 507 53. Jones MR, Horner RD, Edwards LJ, Hoff J, Armstrong SB, Smith-Hammond CA,
508 Matchar DB, Oddone EZ. Racial variation in initial stroke severity. *Stroke*. 2000;31:563-
509 567.
- 510 54. Howard VJ. Reasons underlying racial differences in stroke incidence and mortality.
511 *Stroke*. 2013;44:S126-S128.
- 512 55. Hong K-S, Lee JS. Statins in acute ischemic stroke: a systematic review. *Journal of*
513 *stroke*. 2015;17:282.
- 514 56. Scheitz JF, Endres M, Heuschmann PU, Audebert HJ, Nolte CH. Reduced risk of
515 poststroke pneumonia in thrombolized stroke patients with continued statin treatment.
516 *International Journal of Stroke*. 2015;10:61-66.
- 517