- 1 Title Page
- 2
- 3 **Title:** Characterizing HIV status documentation among cancer patients at regional cancer centers in Malawi,
- 4 Zimbabwe, and South Africa
- 5
- 6 **Authors:** Michalina A Montaño^{1,2,3}, Takudzwa Mtisi⁴, Ntokozo Ndlovu⁴, Margaret Borok⁴, Agatha Bula⁵,
- 7 Maureen Joffe⁶, Rachel Bender Ignacio^{2,3*}, Maganizo B Chagomerana^{5,7*}
- 8
- 9 ¹University of Washington, Department of Global Health
- ²University of Washington, Department of Medicine, Division of Allergy and Infectious Diseases
- ³Fred Hutchinson Cancer Center, Vaccine and Infectious Diseases Division
- ⁴University of Zimbabwe, Faculty of Medicine and Health Sciences
- 13 ⁵UNC Project Malawi
- ⁶University of the Witwatersrand, Faculty of Health Sciences, Strengthening Oncology Services Research Unit
- 15 ⁷University of North Carolina at Chapel Hill, Department of Medicine
- 16
- 17 *Contributed equally
- 18
- 19 **Corresponding author:**
- 20 Michalina A Montaño
- 21 micham@uw.edu
- 22

23 Abstract

24 Introduction

25 In East and Southern Africa, people with HIV (PWH) experience worse cancer-related outcomes and are at

- 26 higher risk of developing certain cancers. Siloed care delivery pathways pose a substantial barrier to co-
- 27 management of HIV and cancer care delivery.

28 Methods

29 We conducted cross-sectional studies of adult cancer patients at public radiotherapy and oncology units in

30 Malawi (Kamuzu Central Hospital), Zimbabwe (Parirenyatwa Group of Hospitals), and South Africa (Charlotte

- 31 Maxeke Hospital) between 2018-2019. We abstracted cancer- and HIV-related data from new cancer patient
- 32 records and used Poisson regression with robust variance to identify patient characteristics associated with
- 33 HIV documentation.

34 Results

35 We included 1,648 records from Malawi (median age 46 years), 1,044 records from South Africa (median age

36 55 years), and 1,135 records from Zimbabwe (median age 52 years). Records from all three sites were

37 predominately from female patients; the most common cancers were cervical (Malawi [29%] and Zimbabwe

38 [43%]) and breast (South Africa [87%]). HIV status was documented in 22% of cancer records from Malawi,

39 92% from South Africa, and 86% from Zimbabwe. Patients with infection-related cancers were more likely to

40 have HIV status documented in Malawi (adjusted prevalence ratio [aPR]: 1.92, 95% confidence interval [CI]:

41 1.56-2.38) and Zimbabwe (aPR: 1.16, 95%CI: 1.10-1.22). Patients aged ≥60 years were less likely to have HIV

42 status documented (Malawi: aPR: 0.66, 95% CI: 0.50-0.87; Zimbabwe: aPR: 0.76, 95%CI: 0.72-0.81) than

patients under age 40 years. Patient age and cancer type were not associated with HIV status documentation
in South Africa.

45 Conclusion

Different cancer centers have different gaps in HIV status documentation and will require tailored strategies to
improve processes for ascertaining and recording HIV-related information in cancer records. Further research
by our consortium to identify opportunities for integrating HIV and cancer care delivery is underway.

49

50

51 Keywords:

52 HIV, oncology, epidemiology, Malawi, South Africa, Zimbabwe

54 INTRODUCTION

55 In East and Southern Africa, cancer disproportionately impacts people with HIV (PWH). PWH are at higher risk of developing certain cancers (1, 2), and experience worse cancer-related outcomes, including 56 57 higher risk of treatment complications, more rapid progression, and higher cancer-specific mortality (2-7). Since 2016, the World Health Organization has recommended that all PWH begin lifelong antiretroviral therapy 58 (ART) as soon as possible after diagnosis. Immediate ART reduces overall morbidity and mortality from HIV-59 60 associated malignancies and other diseases (8-10). Cancer treatment guidelines for PWH recommend co-61 management of HIV and cancer care to ensure uninterrupted ART during cancer therapy (11). However, the guidelines lack specific recommendations beyond ensuring standard of care HIV therapy, and there is a lack of 62 63 evidence-based guidance for co-management of HIV and cancer care delivery in resource-constrained 64 settings. Siloed care delivery pathways and limited facilities for cancer treatment make co-management of HIV and cancer care delivery difficult (12), with evidence that engagement and barriers to HIV care impact receipt 65 of timely cancer care in this setting (13). 66

67 Ascertaining the HIV status of patients being treated for cancer is a critical step in successful cancer 68 treatment. Stable and continuous access to ART throughout cancer treatment is associated with better 69 tolerance of cancer therapy, higher response rates, and improved survival (14, 15). Conversely, PWH with 70 moderate to severe immunosuppression have an increased risk of morbidity and mortality from radiation and 71 cytotoxic chemotherapy, likely due to a compounding effect of these therapies on existing HIV-related 72 immunosuppression (16, 17). ART decreases immune exhaustion and restores cell-mediated immunity among PWH (18, 19), which may improve cancer treatment outcomes by preventing further immunosuppression. 73 74 Ascertaining HIV and antiretroviral treatment status for all patients being treated for cancer is necessary to 75 avoid poor cancer-related outcomes, as well as to decrease risk of infections and all-cause mortality among 76 people who may be unaware of their HIV status or not receiving ART (10).

Integration of HIV and cancer care delivery could improve cancer outcomes for people with comorbid cancer and HIV. Critical first steps to integrating HIV and cancer care delivery in East and Southern Africa include understanding both the burden of HIV at cancer treatment centers, and the scope of HIV-related documentation in cancer center records. Identifying gaps in HIV status ascertainment is the most basic component of cancer care teams being able to optimize clinical decision-making for PWH who are being

treated for cancer. HIV testing and treatment cascades continue to improve in the region, and integrating HIV and cancer-related service delivery will require that these improvements be reflected in cancer-related records. We formed a new consortium across cancer referral centers in East and Southern Africa to study and implement improved HIV care for cancer patients. In this first evaluation, we measured HIV status documentation in medical records at national referral cancer treatment centers in Malawi, South Africa, and Zimbabwe, and assessed demographic and clinical characteristics associated with HIV status documentation in cancer records.

- 89
- 90 METHODS
- 91 Study design, setting, and population

We conducted cross-sectional studies of adult cancer patients at three cancer treatment centers 92 participating in our consortium from Malawi (Kamuzu Central Hospital (KCH)), South Africa (Charlotte Maxeke 93 Johannesburg Academic Hospital (CMJAH)), and Zimbabwe (Parirenyatwa Group of Hospitals (PGH)). KCH is 94 95 one of two national teaching hospitals in Malawi and provides oncology services to half of the country's population. Radiotherapy services were not available in Malawi, and patients requiring radiation to treat their 96 cancer during the study period were referred to institutions outside the country (20). CMJAH is a guaternary-97 care teaching hospital located in Johannesburg, affiliated with the University of the Witwatersrand, and 98 99 provides complete diagnostic, surgical, radiation and medical oncology services for Johannesburg and national referrals. PGH is a tertiary-care teaching hospital in Harare affiliated with the University of Zimbabwe and is 100 one of two public cancer treatment facilities in the country; PGH provides radiation, medical and surgical 101 oncology services to two thirds of Zimbabwe's population (21). We limited analysis to all new patients 102 registered during 2018 and 2019, and excluded patients who were under age 18 years, those with a known 103 recurrent cancer diagnosis, and those with benign diagnoses. 104

105

106 Data sources and measures

At KCH, we obtained data from the KCH Cancer Registry, a hospital-based registry which collects data on demographic and clinical characteristics for all persons with cancer presenting to KCH during 2018-2019, including cancer diagnosis and HIV status (22). KCH Cancer Registry data are abstracted directly from hospital

110 records. At CMJAH, we abstracted data from patient medical records from the Department of Medical 111 Oncology and research records from a large breast cancer cohort. At PGH, we abstracted data from patient 112 medical records from the Radiotherapy and Oncology Centre (ROC). PGH has a Kaposi Sarcoma Clinic 113 separate from the ROC, and KS patients were therefore not included among the reviewed records unless they 114 were receiving treatment at the ROC.

We collected data on age, sex, district or province of residence, cancer diagnosis, and HIV status 115 (positive/negative/not documented). Based on residence data, we categorized participants as either residing 116 outside or in the same province/district as the cancer center at which they were receiving cancer treatment. 117 Due to the strong association between oncogenic viral infections and HIV, we categorized cancer diagnoses as 118 either infection-related or not infection-related. Cancers that are most likely or universally resultant from 119 oncogenic viral infection (Epstein-Barr virus, Hepatitis B and C viruses, human papillomavirus (HPV), Kaposi 120 sarcoma-associated herpesvirus) were categorized as infection-related cancers: Hodgkin and non-Hodgkin 121 lymphomas, liver cancers, anal, cervical, penile, and vulvovaginal cancers, and Kaposi sarcoma (23). We did 122 not consider cancers of mixed pathogenesis (e.g., head and neck or gastric, which may or may not be 123 attributable to HPV and H. pylori, respectively), as infection-related, due to inability to confirm histology and 124 infectious etiology in all cases. 125

126

127 Statistical analysis

Descriptive statistics for demographic and clinical characteristics of all participants included in the 128 analysis were summarized separately for each study site. We calculated the proportion of participants with HIV 129 status documented in their medical records, and the HIV prevalence among those with documented HIV 130 131 status. We used generalized linear models with Poisson distribution and robust variance to identify participant characteristics associated with documented HIV status. We selected covariates a priori based on 132 characteristics previously documented to be associated with either HIV testing or barriers to HIV care 133 134 engagement in these regions (24-27). The covariates of interest included age, gender, residence proximity to 135 cancer treatment center, and infection-related cancer diagnosis. We developed separate models for each study site to understand the context of HIV status documentation separately for each cancer center and assist 136

- with developing site-specific solutions to improve status documentation. Two-sided statistical tests were
 performed at an alpha level of 0.05. We used Stata version 17.0 (College Station, Texas) for all analyses.
- 139

This study was approved by the University of Washington Institutional Review Board, University of North Carolina Institutional Review Board, Malawi National Health Sciences Research Committee, University of the Witwatersrand Human Research Ethics Committee (Medical), and the Joint Research Ethics Committee for the University of Zimbabwe Faculty of Medicine and Health Sciences and Parirenyatwa Group of Hospitals. Patients were not involved in the design or conduct of this research.

- 145
- 146 **RESULTS**
- 147 Malawi

We included 1,648 records from KCH, Malawi. Participants had a median age of 46 years and were 148 predominantly female (Table 1). Most participants (72.3%) reported residing in Lilongwe district, where KCH is 149 located. Infection-related cancers were common and accounted for over half of all cancer diagnoses. HIV 150 status was documented in 22% of available cancer records. Among patients with documented HIV status, 47% 151 were HIV-positive. Data on ART status was not available in cancer records from Malawi. The most common 152 cancers were cervical, breast, Kaposi sarcoma, and non-Hodgkin lymphoma, among which the proportion of 153 records with no HIV status documented was 80.3%, 84.9%, 75.3%, and 46.9%, respectively (Table 2; Figure 154 1). We observed differences in HIV status documentation by age, gender, residence proximity to KCH, and 155 infection-related cancer status (Table 3). Patients aged 60 years or older were less likely to have their HIV 156 status documented (adjusted prevalence ratio [aPR]: 0.66, 95% confidence interval [CI]: 0.50-0.87). Male 157 patients were more likely to have HIV status documented compared to female patients (aPR: 1.45, 95% CI: 158 1.21-1.74). Similarly, patients residing outside Lilongwe district were more likely to have HIV status 159 documented compared to patients residing within Lilongwe district (aPR: 1.68, 95% CI: 1.41-2.01). Patients 160 with infection-related cancers were also more likely to have HIV status documented than patients with non-161 162 infection-related cancers (aPR: 1.92, 95%CI: 1.56-2.38).

163

164 South Africa

We included 1,044 records from CMJAH, South Africa. Over 90% of participants from this site were 165 166 female, and the median age was 55 years. Almost all participants (97.0%) reported residing in Gauteng Province, where CMJAH is located. Infection-related cancers were not common among available data from 167 CMJAH; data were obtained primarily from patients with breast cancer (86.9% of available records at the 168 sampled clinic). HIV status was documented among 91.7% of available cancer records, and HIV prevalence 169 was 24.2% among those with documented HIV status. Among patients with documented HIV, 96.9% were on 170 ART at the time they registered as patients at CMJAH. Among patients with breast cancer, 7.3% had no HIV 171 status documented in their cancer record. Patient age, gender, residence, and cancer type were not 172 associated with HIV status documentation in South Africa. 173

- 174
- 175 Zimbabwe

We included 1,135 records from PGH, Zimbabwe. Participants had a median age of 52 years, and 176 three guarters were female. Most participants (65.4%) resided outside Harare Province, where PGH is located. 177 Infection-related cancers were common, accounting for 49.8% of cancer diagnoses. HIV status was 178 documented in 86% of available cancer records. Among patients with documented HIV status, HIV prevalence 179 was 37%. Among patients with documented HIV, 95.6% were on ART at the time they registered at PGH. The 180 most common cancers were cervical, head and neck, and breast, among which 3.5%, 14.7%, and 16.8%, 181 182 respectively, had no HIV status documented in cancer records (Table 2; Figure 1). We observed differences in HIV status documentation by age and infection-related cancer status. Patients aged 60 years or older were 183 less likely to have their HIV status documented compared to patients under age 40 years (aPR 0.76, 95%CI: 184 0.72-0.81). Patients with infection-related cancers were more likely than those with non-infection-related 185 cancers to have their HIV status documented (aPR: 1.16, 95%CI:1.10-1.22) (Table 4). Gender and residence 186 proximity to cancer center were not significant predictors of HIV status documentation in multivariable models. 187 188

189 DISCUSSION

The objectives of this project were to document the burden of HIV among patients with cancer, measure HIV status documentation in cancer records, and identify characteristics associated with undocumented HIV status. The goal of these activities was to provide context-specific recommendations for

improvement of HIV status documentation, and guide future strategies to integrate components of HIV care 193 194 delivery into cancer care for PWH. Amongst the three sites studied in this new consortium focused on implementing coordinated HIV and cancer care, we found variable needs in achieving universal HIV 195 documentation at each center. While Zimbabwe and South Africa had high rates of HIV documentation with 196 near complete documentation of ART receipt for those with documented HIV, a minority of cancer patients in 197 Malawi had HIV status available in their records. Among cancer patients with documented status, the 198 prevalence of HIV was much higher than in the general population, even considering the older age distribution 199 of cancer patients compared to the general population in each country. 200

201

202 Malawi

In Malawi, HIV status was documented in 22% of cancer records. In this study, HIV prevalence among 203 patients with documented HIV status was almost five times the general population HIV prevalence, and over 204 three times the prevalence for adults over age 50 years (28). Over three guarters of cancer records did not 205 206 have HIV status documented. Malawi has made considerable progress toward meeting the first and second Joint United Nations Programme on HIV/AIDS (UNAIDS) 95-95-95 targets; 98% of women and 89% of men 207 over age 15 years and living with HIV in Malawi are aware of their HIV status, and 97% of PWH aware of their 208 status are on ART (29). The high proportion of patients with undocumented HIV status that we observed may 209 be due to failure to document HIV-related information in cancer records rather than lack of HIV testing in this 210 population. While HIV testing rates are higher among women in Malawi (26, 28), men were more likely than 211 women to have their HIV status documented in cancer records, which may further support that those tested in 212 outside settings were less likely to have those results reported than those tested for indications associated with 213 214 their cancer diagnosis.

The high HIV prevalence we observed among patients with their HIV status documented may be due to indication bias. Patients with infection-related cancers (signs and symptoms possibly indicative of HIV) may be more likely to be referred for HIV testing during cancer care delivery and would therefore be more likely to have their HIV status documented than patients with other cancers. Patients with infection-related cancers were more likely to test positive for HIV than patients with non-infection-related cancers. The HIV prevalence among

patients with their HIV status documented may therefore not accurately represent the HIV prevalence among
 the overall population of patients with cancer.

Knowing patients' ART status and regimen likely also improves cancer care decision-making, but clinical care teams are at a disadvantage if this information is not recorded and available to them. Malawi's progress meeting the 95-95-95 targets suggests that changes to the way HIV-related information is ascertained by cancer care teams and documented in cancer records may result in substantial improvements in HIV care coordination among cancer patients, even while most cancer patients are likely already aware of their status. Strategies to improve HIV testing can then be targeted to the small proportion of patients with cancer whose HIV status is truly unknown rather than those whose HIV status is known but not documented.

229

230 South Africa

In South Africa, HIV status was documented in 92% of cancer records. We did not find characteristics 231 associated with HIV status documentation among this patient population. A high proportion of PWH had their 232 233 ART status recorded, and were on ART. However, the majority of available data were from patients with breast cancer due to the population at the specific clinic sampled, and the results may not generalize to other 234 cancers. Specifically, there has been concerted effort at this institution to create algorithms for breast cancer 235 providers to ensure HIV testing and care are secured prior to cancer treatment (30, 31). This may have 236 237 impacted our overall ability to measure associations between patient characteristics and HIV status documentation, and these associations should be explored among patients with other types of cancer in this 238 and similar institutions. Recent data from UNAIDS shows 94% of PWH in South Africa know their status, which 239 is reflected in the high proportion of cancer records with HIV status documented at CMJAH (29). Further along 240 241 the HIV care cascade, however, only 78% of South African women and 68% of men with HIV over age 15 years are on ART. Among records included in our analysis, we found that 97% of PWH being treated for 242 cancer at CMJAH were on ART, a higher proportion than the general population of PWH. This may be due in 243 part to specific protocols for cancer treatment at CMJAH, especially for people with breast cancer, and possible 244 245 higher engagement in clinical care among people with cancer.

- 246
- 247 Zimbabwe

In Zimbabwe, HIV status was documented in 86% of cancer records. People over age 60 years and 248 249 those with non-infection related cancers were less likely to have their HIV status documented, and increasing HIV status ascertainment for these groups will be necessary to meet the goal of universal HIV status 250 ascertainment for cancer patients. Among patients with documented HIV status, the HIV prevalence was 37%, 251 which is triple the general population prevalence, and double the HIV prevalence for adults over age 50 years 252 (32). Among patients with a documented positive HIV status, the vast majority had their ART status 253 254 documented, and were on ART at the time they entered care at PGH. This suggests that almost all patients with cancer at PGH who know their HIV status are already established in HIV care when they enter cancer 255 care. Demographic data show that two thirds of PGH cancer patients reside in a different province from where 256 257 the cancer center is located. Since most PWH access HIV care near where they live, a substantial proportion of PWH accessing cancer therapy at PGH may experience disruptions to HIV care access due to traveling to 258 receive cancer therapy, as previously demonstrated in Uganda by members of our consortium (33). Any 259 integration of HIV and cancer service delivery at PGH should include strategies to ensure these patients can 260 261 reliably access their ART during cancer treatment. Finally, like Malawi, Zimbabwe has met the UNAIDS 90-90-90 goals and is progressing toward the new 95-95-95 targets (29). Among PWH, 98% of women and 95% of 262 men aged 15 years or older know their status, and 96% of those who know their status are on ART. If HIV 263 testing and ART use among Zimbabweans with cancer are similar to the general population, this suggests that 264 265 improvements to HIV status documentation, supplemented by targeted HIV testing for people with unknown status, is likely to bridge the gap to achieving universal HIV status ascertainment for people with cancer being 266 treated at PGH, especially for those with malignancies not typically considered HIV-associated. 267

268

Data on the burden of HIV and HIV status documentation among patients with cancer from other countries in East and Southern Africa are sparse. A cross-sectional study conducted in Uganda in 2015 reported that 29% of patients with cancer at the Uganda Cancer Institute had no recorded HIV status (34). Among patients with HIV status documented, 32% had HIV, more than four-fold higher than the general population prevalence (7%) at the time. Another study, conducted between 2014 and 2016 in Malawi, was able to ascertain the HIV status of 34% of registered cancer patients at KCH, of whom 58% had HIV (22). This proportion is substantially higher than the general population HIV prevalence, but like our findings from KCH

reported above, may not accurately reflect the prevalence of HIV among all cancer patients due to the high 276 277 proportion of missing data. Importantly, the timelines of these two studies coincided with early scale up of strategies to meet the newly established 90-90-90 goals but largely preceded removal of CD4 thresholds for 278 ART initiation by the WHO (35). Since then, the proportion of people who know their HIV status has improved 279 across East and Southern Africa (29). It seems unlikely that people with cancer are much less likely to receive 280 testing and know their status than the general population, and gaps in HIV status ascertainment for 281 patients with cancer may be partly due to siloing of information and care rather than lack of testing; stigma 282 283 around both HIV and cancer and use of hand-carried HIV records in many locations may contribute to lack of HIV documentation(18). 284

Our findings suggest that different cancer centers will require tailored strategies to address gaps in HIV 285 286 and HIV-treatment status documentation for patients with cancer as a first step in providing coordinated HIV and cancer care. The proportions of cancer patients without documented HIV status reported above include 287 both people with cancer who are unaware of their HIV status, and those who know their status, but whose 288 289 status may not be available to their cancer care team. The high proportion of PWH in Malawi, South Africa, and Zimbabwe who know their status suggests that under-ascertainment of HIV status for patients with cancer may 290 be primarily a result of siloed care delivery systems in these locations. Improving cancer care for PWH will 291 require that cancer providers and clinical care teams also know the HIV and ART status of their patients. 292 293 Improving ascertainment will likely require a combination of approaches that are site-specific. Changes to the 294 process for ascertaining and recording HIV status and other HIV-related information in cancer-related medical records may result in marked improvement in HIV status documentation overall, with HIV testing to supplement 295 these medical record system changes focused on patients with unknown HIV status. Further research to 296 297 identify and apply context-specific opportunities for integrating HIV and cancer care delivery may be necessary 298 to improve outcomes for patients with comorbid HIV and cancer in East and Southern African settings. Work by our consortium is ongoing at these three sites plus the Uganda Cancer Institute, to identify opportunities to 299 300 integrate HIV care into cancer care, starting with developing site-specific process maps and institutional 301 barriers and facilitators to providing coordinated HIV treatment initiation and continuity for those receiving 302 cancer care.

303

- 304 Abbreviations
- 305 aPR: adjusted prevalence ratio
- 306 ART: antiretroviral therapy
- 307 CI: confidence interval
- 308 CMJAH: Charlotte Maxeke Johannesburg Academic Hospital
- 309 HIV: Human Immunodeficiency Virus
- 310 HPV: Human papillomavirus
- 311 KCH: Kamuzu Central Hospital
- 312 KS: Kaposi Sarcoma
- 313 PGH: Parirenyatwa Group of Hospitals
- 314 PWH: People with HIV
- 315 ROC: Radiation and Oncology Centre

317 **DECLARATIONS**

318

319 Ethics Approval

320	All procedures performed in this study involving human participants were in accordance with the ethical
321	standards of the institutional and/or national research committees, and with the 1964 Helsinki declaration and
322	its later amendments or comparable ethical standards. The study received ethical approval from the University
323	of Washington Institutional Review Board, University of North Carolina Institutional Review Board, Malawi
324	National Health Sciences Research Committee, University of the Witwatersrand Human Research Ethics
325	Committee (Medical), and the Joint Research Ethics Committee for the University of Zimbabwe Faculty of
326	Medicine and Health Sciences and Parirenyatwa Group of Hospitals.
327	
328	Consent for Publication
329	Not applicable
330	
331	Availability of Data and Materials
332	The data that support the findings of this study are not publicly available due to privacy or ethical restrictions.
333	Data are available from the authors upon reasonable request and permission from each participating hospital
334	and institutional review board.
335	
336	Competing Interests
337	All authors declare they have no competing interests.
338	
339	Funding
340	This study was funded by the National Institutes of Health: National Institute of Allergy and Infectious Diseases
341	(R21 AI155055) and National Cancer Institute (T32 CA080416).
342	
343	Author Contributions

- Conception and study design: MB, MBC, MJ, MAM, NN, RBI; data collection: all authors; data analysis: MBC,
- MAM, TM; review and interpretation of results: all authors; manuscript preparation: MAM; manuscript revision
- and final approval of the submitted manuscript: all authors.
- 347

348 Acknowledgements

- The authors wish to thank the staff at PGH, University of Zimbabwe, CMJAH, Wits Health Consortium, KCH,
- and UNC Project Malawi who assisted with data collection for this study.
- 351
- 352

REFERENCES

1. Yarchoan R, Uldrick TS. HIV-Associated Cancers and Related Diseases. The New England journal of medicine. 2018;378(11):1029-41.

2. Chinula L, Moses A, Gopal S. HIV-associated malignancies in sub-Saharan Africa: progress, challenges, and opportunities. Curr Opin HIV AIDS. 2017;12(1):89-95.

3. Reddy P, Ebrahim S, Singh B, Ramklass S, Buccimazza I. Breast Cancer and HIV: a South African perspective and a critical review of the literature. South African journal of surgery Suid-Afrikaanse tydskrif vir chirurgie. 2017;55(1):10-5.

4. Simonds HM, Botha MH, Neugut AI, Van Der Merwe FH, Jacobson JS. Five-year overall survival following chemoradiation among HIV-positive and HIV-negative patients with locally advanced cervical carcinoma in a South African cohort. Gynecologic oncology. 2018;151(2):215-20.

5. Dryden-Peterson S, Bvochora-Nsingo M, Suneja G, Efstathiou JA, Grover S, Chiyapo S, et al. HIV Infection and Survival Among Women With Cervical Cancer. Journal of clinical oncology : official journal of the American Society of Clinical Oncology. 2016;34(31):3749-57.

6. Ngidi S, Magula N, Sartorius B, Govender P, Madiba TE. Incidence of chemotherapy-induced neutropenia in HIV-infected and uninfected patients with breast cancer receiving neoadjuvant chemotherapy. S Afr Med J. 2017;107(7):595-601.

7. Camandaroba MPG, de Araujo RLC, Silva VSE, de Mello CAL, Riechelmann RP. Treatment outcomes of patients with localized anal squamous cell carcinoma according to HIV infection: systematic review and meta-analysis. J Gastrointest Oncol. 2019;10(1):48-60.

8. Consolidated guidelines on the use of antiretroviral drugs for treating and preventing HIV infection: what's new.: World Health Organization; 2015.

9. Group TAS, Danel C, Moh R, Gabillard D, Badje A, Le Carrou J, et al. A Trial of Early Antiretrovirals and Isoniazid Preventive Therapy in Africa. The New England journal of medicine. 2015;373(9):808-22.

 Lundgren JD, Babiker AG, Gordin F, Emery S, Grund B, Sharma S, et al. Initiation of Antiretroviral Therapy in Early Asymptomatic HIV Infection. The New England journal of medicine. 2015;373(9):795-807.
 Reid E, Suneja G, Ambinder RF, Ard K, Baiocchi R, Barta SK, et al. Cancer in People Living With HIV,

Version 1.2018, NCCN Clinical Practice Guidelines in Oncology. Journal of the National Comprehensive Cancer Network : JNCCN. 2018;16(8):986-1017.

12. Koh WJ, Anderson BO, Carlson RW. NCCN resource-stratified and harmonized guidelines: A paradigm for optimizing global cancer care. Cancer. 2020;126 Suppl 10:2416-23.

13. Low DH, Phipps W, Orem J, Casper C, Bender Ignacio RA. Engagement in HIV Care and Access to Cancer Treatment Among Patients With HIV-Associated Malignancies in Uganda. J Glob Oncol. 2019;5:1-8.

14. Hessol NA, Pipkin S, Schwarcz S, Cress RD, Bacchetti P, Scheer S. The impact of highly active antiretroviral therapy on non-AIDS-defining cancers among adults with AIDS. Am J Epidemiol. 2007;165(10):1143-53.

15. Montaño MA, Chagomerana MB, Borok M, Painschab M, Uldrick TS, Bender Ignacio RA. Impact of Antiretroviral Therapy on Cancer Treatment Outcomes among People Living with HIV in Low- and Middle-Income Countries: a Systematic Review. Curr HIV/AIDS Rep. 2021;18(2):105-16.

16. Chiao EY, Dezube BJ, Krown SE, Wachsman W, Brock MV, Giordano TP, et al. Time for oncologists to opt in for routine opt-out HIV testing? Jama. 2010;304(3):334-9.

17. Deeken JF, Pantanowitz L, Dezube BJ. Targeted therapies to treat non-AIDS-defining cancers in patients with HIV on HAART therapy: treatment considerations and research outlook. Curr Opin Oncol. 2009;21(5):445-54.

18. Hunt PW, Deeks SG, Rodriguez B, Valdez H, Shade SB, Abrams DI, et al. Continued CD4 cell count increases in HIV-infected adults experiencing 4 years of viral suppression on antiretroviral therapy. AIDS (London, England). 2003;17(13):1907-15.

19. Landay A, da Silva BA, King MS, Albrecht M, Benson C, Eron J, et al. Evidence of ongoing immune reconstitution in subjects with sustained viral suppression following 6 years of lopinavir-ritonavir treatment. Clinical infectious diseases : an official publication of the Infectious Diseases Society of America. 2007;44(5):749-54.

20. Kamuzu Central Hospital - National Cancer Center [Available from: <u>https://www.esmo.org/for-patients/esmo-designated-centres-of-integrated-oncology-palliative-care/esmo-accredited-designated-centres/kamuzu-central-hospital-national-cancer-center.</u>

21. The National Cancer Prevention And Control Strategy For Zimbabwe 2014 - 2018. Zimbabwe Ministry of Health and Child Care; 2014.

22. Horner MJ, Salima A, Chilima C, Mukatipa M, Kumwenda W, Kampani C, et al. Frequent HIV and Young Age Among Individuals With Diverse Cancers at a National Teaching Hospital in Malawi. J Glob Oncol. 2018;4:1-11.

23. Krump NA, You J. Molecular mechanisms of viral oncogenesis in humans. Nat Rev Microbiol. 2018;16(11):684-98.

24. Croome N, Ahluwalia M, Hughes LD, Abas M. Patient-reported barriers and facilitators to antiretroviral adherence in sub-Saharan Africa. Aids. 2017;31(7):995-1007.

25. Hakim AJ, Tippett Barr BA, Kinchen S, Musuka G, Manjengwa J, Munyati S, et al. Progress Toward the 90-90-90 HIV Targets in Zimbabwe and Identifying Those Left Behind. J Acquir Immune Defic Syndr. 2021;88(3):272-81.

26. Maman D, Ben-Farhat J, Chilima B, Masiku C, Salumu L, Ford N, et al. Factors associated with HIV status awareness and Linkage to Care following home based testing in rural Malawi. Trop Med Int Health. 2016;21(11):1442-51.

27. West CA, Chang GC, D WC, Bray R, Kinchen S, Behel S, et al. Unawareness of HIV Infection Among Men Aged 15-59 Years in 13 Sub-Saharan African Countries: Findings From the Population-Based HIV Impact Assessments, 2015-2019. J Acquir Immune Defic Syndr. 2021;87(Suppl 1):S97-s106.

28. Malawi Population-based HIV Impact Assessment 2020-2021 (MPHIA 2020-2021): Final Report. Lilongwe: MOH Malawi November 2022.

29. UNAIDS Data 2022. Geneva: Joint United Nations Programme on HIV/AIDS; 2022.

30. Chasimpha S, McCormack V, Cubasch H, Joffe M, Zietsman A, Galukande M, et al. Disparities in breast cancer survival between women with and without HIV across sub-Saharan Africa (ABC-DO): a prospective, cohort study. The Lancet HIV. 2022;9(3):e160-e71.

31. Cubasch H, Joffe M, Hanisch R, Schuz J, Neugut AI, Karstaedt A, et al. Breast cancer characteristics and HIV among 1,092 women in Soweto, South Africa. Breast cancer research and treatment. 2013;140:177-86.

32. Zimbabwe Population-based HIV Impact Assessment 2020 (ZIMPHIA 2020): Final Report. Harare December 2021.

33. Low DH, Phipps W, Orem J, Casper C, Bender Ignacio RA. Engagement in HIV Care and Access to Cancer Treatment Among Patients With HIV-Associated Malignancies in Uganda. Journal of Global Oncology. 2019(5):1-8.

34. Bender Ignacio R, Ghadrshenas M, Low D, Orem J, Casper C, Phipps W. HIV Status and Associated Clinical Characteristics Among Adult Patients With Cancer at the Uganda Cancer Institute. J Glob Oncol. 2018;4:1-10.

35. 90-90-90: An ambitious treatment target to help end the AIDS epidemic. Joint United Nations Programme on HIV/AIDS (UNAIDS); 2014.

Table 1. Demographic and Clinical Characteristics of Adult Cancer Patients at Regional											
Cancer Treatment Centres in Malawi, South Africa, and Zimbabwe											
	Ma	lawi	South	Africa	Zimbabwe						
	N=1	648	N=1	044	N=1135						
Characteristic	n	%	n	%	n	%					
Age at Diagnosis		~~ /	470	40.0							
<40	534	32.4	173	16.6	175	15.4					
40-49	429	26.0	255	24.4	288	25.4					
50-59	314	19.2	293	28.1	215	18.9					
60 or older	368	22.3	323	30.9	457	40.3					
Gender											
Female	1040	63.1	964	92.3	829	73.0					
Male	605	36.7	80	7.7	306	27.0					
Residence Proximity to Hospital											
Resides in same province as hospital	1191	72.3	1013	97.0	393	34.6					
Resides outside hospital province	446	27.1	11	1.1	742	65.4					
Infection-Related Cancer Status											
Not infection-related	715	43.4	994	95.2	570	50.2					
Infection-related	933	56.6	50	4.8	565	49.8					
HIV Status Documentation											
Not Documented	1286	78.0	87	8.3	155	13.7					
Documented	362	22.0	957	91.7	980	86.3					
	001			•							
HIV Status ^a											
HIV-Negative	191	52.8	725	75.8	614	62 7					
HIV-Positive	171	47.2	232	24.2	366	37.3					
	17.1	-1.2	202	27.2	000	07.0					
ART Status ^b											
			210	96 9	350	95.6					
Not on ART			213	27	7	1 0					
ART status not documented			0	0.1	7 Q	2.5					
			1	0.4	5	2.0					

Due to missing values, percentages may not sum to 100%. 3 participants from Malawi missing age and gender; 11 participants from Malawi and 20 from South Africa missing residence.

^a Of records with documented HIV status. Denominators are 362 (Malawi), 957 (South Africa), and 980 (Zimbabwe). ^b Among participants with HIV. Denominators are 232 (South Africa), and 366 (Zimbabwe). ART status was not

available in records from Malawi.

Table 2. Cancer Diagnoses and Documented HIV Status of Adult Cancer Patients at Regional Cancer Treatment Centers in Malawi, South Africa, and Zimbabwe

	Malawi (N=1648)						South Africa					Zimbabwe (N=1135)						
Diagnosis	HIV-		(N=1646) HIV+		Unk	Unknown		HIV-		HIV+ Un		nown	iown HIV-		HIV+		Unknown	
	n	%	n	%	n	%	п	%	n	%	п	%	n	%	n	%	n	%
Anal	0	0.0	1	20.0	4	80.0	0	0.0	1	100.0	0	0.0	2	15.4	10	76.9	1	7.7
Bladder	1	14.3	0	0.0	6	85.7	4	66.7	1	16.7	1	16.7	9	56.3	0	0.0	7	43.8
Bone or soft tissue	1	14.3	1	14.3	5	71.4	3	75.0	1	25.0	0	0.0	7	53.9	4	30.8	2	15.4
Breast	22	8.7	16	6.4	214	84.9	644	70.9	198	21.8	66	7.3	73	68.2	16	15.0	18	16.8
Cervical	58	12.3	35	7.4	379	80.3	1	50.0	0	0.0	1	50.0	246	50.0	229	46.5	17	3.5
CNS (primary, incl. brain)	0	0.0	2	14.3	12	85.7	1	100.0	0	0.0	0	0.0	1	50.0	0	0.0	1	50.0
Colorectal	1	3.5	3	10.3	25	86.2	24	82.8	1	3.5	4	13.8	37	72.6	8	15.7	6	11.8
Conjunctival SCC	0	0.0	0	0.0	1	100.0												
Esophageal	4	5.6	1	1.4	67	93.1	1	50.0	1	50.0	0	0.0	17	56.7	9	30.0	4	13.3
Gastric	1	10.0	0	0.0	9	90.0	6	66.7	1	11.1	2	22.2	24	72.7	3	9.1	6	18.2
Head and Neck	3	6.4	2	4.3	42	89.4	2	100.0	0	0.0	0	0.0	53	48.6	40	36.7	16	14.7
Hodgkin Lymphoma	20	35.7	15	26.8	21	37.5	3	27.3	7	63.6	1	9.1	2	66.7	0	0.0	1	33.3
Kaposi Sarcoma	13	5.6	44	19.1	174	75.3	3	27.3	8	72.7	0	0.0	0	0.0	1	100.0	0	0.0
Leukemia (Acute)							0	0.0	0	0.0	2	100.0	2	100.0	0	0.0	0	0.0
Leukemia (Chronic)	2	20.0	0	0.0	8	80.0	1	50.0	1	50.0	0	0.0						
Lung	0	0.0	0	0.0	14	100.0	1	100.0	0	0.0	0	0.0	13	52.0	7	28.0	5	20.0
Liver	2	15.4	0	0.0	11	84.6	6	60.0	2	20.0	2	20.0	20	57.1	6	17.1	9	25.7
Melanoma	1	20.0	1	20.0	3	60.0	1	50.0	0	0.0	1	50.0	6	60.0	1	10.0	3	30.0
Non-Hodgkin Lymphoma	38	34.2	21	18.9	52	46.9	7	50.0	7	50.0	0	0.0	0	0.0	5	100.0	0	0.0
Ovarian	0	0.0	0	0.0	10	100.0	1	50.0	1	50.0	0	0.0	9	69.2	0	0.0	4	30.8
Pancreatic or Biliary	0	0.0	1	20.0	4	80.0	2	66.7	0	0.0	1	33.3	2	33.3	1	16.7	3	50.0
Penile	0	0.0	5	22.7	17	77.3	0	0.0	1	100.0	0	0.0	1	16.7	5	83.3	0	0.0
Prostate	4	18.2	0	0.0	18	81.8	9	100.0	0	0.0	0	0.0	38	50.0	6	7.9	32	42.1
Renal	1	25.0	0	0.0	3	75.0	1	50.0	0	0.0	1	50.0	3	75.0	0	0.0	1	25.0
Vulvovaginal	1	4.4	8	34.8	14	60.9							2	20.0	8	80.0	0	0.0
Unknown	15	10.7	6	4.3	119	85.0							7	41.2	3	17.6	7	41.2
Other	3	4.6	9	13.6	54	81.8	6	60.0	1	10.0	3	30.0	40	71.4	4	7.1	12	21.4
CNS: Central nervous system; SCC: Squamous cell carcinoma																		

Table 3. Demographic and Clinical Characteristics by HIV Status Documentation among Adult Cancer Patients at Regional Cancer Treatment Centres in Malawi South Africa, and Zimbabwe											
	Undocument	ed HIV Statu	s Documente	d HIV Stat	us						
MALAWI	n	%	n	%	р						
Age at Diagnosis											
<40	404	75.7	130	24.3	o o o -						
40-49	323	75.3	106	24.7	0.007						
50-59	247	18.1	6/ 59	21.3							
	310	04.2	00	15.0							
Gender											
Female	844	81.2	196	18.8	<0.001						
Male	440	12.1	165	27.3							
Residence Proximity to Hospital											
Resides in same province as hospital	978	82.1	213	17.9	<0.001						
Resides outside hospital province	301	67.5	145	32.5							
Infection-related Cancer ^a											
Not infection-related	614	85.9	101	14.1	<0.001						
Infection-related	672	72.0	261	28.0							
		0 (A (
SOUTH AFRICA	n	%	n	%	р						
Age at Diagnosis	10	6.0	161	02.1							
40	12	0.9	101	93.1	0.75						
50-59	20	8.2	269	92.2	0.75						
60 or older	31	9.6	200	90.4							
	01	0.0	202	00.1							
Gender	70	0 0	995	01.0	0.57						
Male	19	0.2	000 72	91.0 QA A	0.57						
	0	10.0	12	30.0							
Residence Proximity to Hospital	05	0.4		04.0	0.00						
Resides in same province as hospital	85	8.4	928	91.6	0.93						
Resides outside hospital province	I	9.1	10	90.9							
Infection-related Cancer											
Not infection-related	83	8.4	911	91.6	0.44						
Infection-related	4	8.0	46	92.0							
	n	0/	n	0/	n						
Age at Diagnosis		/0		/0	þ						
<40	5	2.9	170	97.1							
40-49	14	4.9	274	95.1	<0.001						
50-59	11	5.1	204	94.9							
60 or older	125	27.4	332	72.6							
Gender											
Female	84	10.1	745	89.9	< 0.001						
Male	71	23.2	235	76.8							
Residence Provimity to Heapitel											
Resides in same province as bospital	57	14 5	226	85 5	0 55						
Resides outside hospital province	98	13.2	644	86.8	0.00						
	50	10.2	0-1-1	50.0							
Intection-related Cancer	407	00 0	110	77 7	~0.001						
Infection-related	۱ <i>۲۱</i> ۵۵	22.3 5.0	443 527	11.1 95.0	∼ 0.001						
	20	0.0		33.0							
^a We considered the following cancers to be inf Sarcoma, liver, penile, and vulvovaginal	ection-related: a	nal, cervical, H	lodgkin and non-H	odgkin lymp	ohomas, Kaposi						

Table 4. Predictors of Documented HIV Status among Adult Cancer Patients at Regional Cancer Treatment Centres in Malawi, South Africa, and Zimbabwe												
MALAWI	PR	95% CI	a	aPR	95% CI	p						
Age at Diagnosis <40 40-49 50-59 60 or older	Ref. 1.01 0.88 0.65	(0.81 – 1.27) (0.68 – 1.14) (0.49 – 0.86)	 0.9 0.3 0.002	Ref. 0.98 0.88 0.66	(0.79 – 1.22) (0.69 – 1.14) (0.50 – 0.87)	 0.9 0.3 0.003						
Gender Female Male	Ref. 1.45	 (1.21 – 1.73)	 <0.001	Ref. 1.45	 (1.21 – 1.74)	 <0.001						
Residence Proximity to Cancer Centre Resides in same province as hospital Resides outside hospital province	Ref. 1.82	 (1.52 – 2.18)	 <0.001	Ref. 1.68	 (1.41 – 2.01)	 <0.001						
Infection-related Cancer Not infection-related Infection-related	Ref. 1.98	 (1.61 – 2.44)	 <0.001	Ref. 1.92	 (1.56 – 2.38)	 <0.001						
SOUTH AFRICA	PR	95% CI	р	aPR	95% CI	р						
Age at Diagnosis <40 40-49 50-59 60 or older	Ref. 0.99 0.99 0.97	 (0.94 – 1.05) (0.94 – 1.04) (0.92 – 1.03)	 0.7 0.6 0.3	Ref. 0.99 0.99 0.97	 (0.94 – 1.05) (0.94 – 1.05) (0.92 – 1.03)	 0.7 0.8 0.4						
Gender Female Male	Ref. 0.98	 (0.91 – 1.06)	 0.6	Ref. 0.98	 (0.90 – 1.07)	 0.7						
Residence Proximity to Hospital Resides in same province as hospital Resides outside hospital province	Ref. 0.99	 (0.82 – 1.20)	 0.9	Ref. 0.99	 (0.82 – 1.19)	 0.9						
Infection-related Cancer Not infection-related Infection-related	Ref. 1.00	 (0.92 – 1.09)	 0.9	Ref. 1.01	 (0.91 – 1.12)	 0.8						
ZIMBABWE	PR	95% CI	р	aPR	95% CI	р						
Age at Diagnosis <40 40-49 50-59 60 or older	Ref. 0.98 0.98 0.75	 (0.94 – 1.02) (0.94 – 1.02) (0.70 – 0.80)	 0.3 0.3 <0.001	Ref. 0.96 0.96 0.76	 (0.92 – 1.00) (0.92 – 1.00) (0.72 – 0.81)	 0.04 0.06 <0.001						
Gender Female Male	Ref. 0.85	 (0.80 – 0.91)	 <0.001	Ref. 0.97	 (0.90 – 1.04)	 0.4						
Residence Proximity to Hospital Resides in same province as hospital Resides outside hospital province	Ref. 1.02	 (0.97 – 1.07)	 0.6	Ref. 0.99	 (0.95 – 1.04)	 0.8						
Infection-related Cancer Not infection-related Infection-related	Ref. 1.22	 (1.17 – 1.28)	 <0.001	Ref. 1.16	 (1.10 – 1.22)	 <0.001						

aPR: adjusted prevalence ratio; CI: confidence interval; Ref: reference category; PR: prevalence ratio;

Multivariable models include age, gender, residence proximity to hospital, and infection-related cancer diagnosis ^a We considered the following cancers to be infection-related: anal, cervical, Hodgkin and non-Hodgkin lymphomas, Kaposi

Sarcoma, liver, penile, and vulvovaginal

Figure 1. Cancer diagnoses and documented HIV status of adult cancer patients at regional cancer treatment centers in Malawi, South Africa, and Zimbabwe. Boxes indicate infection-related cancers.

