

1 Full length Title: Predictors of voluntary uptake of modern
2 contraceptive methods in rural Sindh, Pakistan

3 **Short Title: Predictors of modern contraceptive methods**
4

5 Authors:

6 Zahid Memon^{1¶*}, Wardah Ahmed^{1¶}, Abeer Mian^{1¶}, Muhammad Jawwad^{1¶}, Shah Muhammad^{1¶},
7 Abdul Qayyum Noorani^{3¶}, Zulfiqar Bhutta^{1¶} and Hora Soltani^{2£}

8 Affiliations:

9 1 The Aga Khan University

10 2 Sheffield Hallam University

11 3 Aga Khan Foundation

12 Country of Residence:

13 ¶ Pakistan

14 £ United Kingdom

15 *Corresponding author email address: zahid.memon@aku.edu
16
17
18
19
20
21
22
23
24
25
26
27
28
29

30

31 **Abstract**

32 Introduction: The use of modern contraceptive methods (MCMs) in Pakistan has been stagnant for
33 the last decade. In Sindh, current contraceptive use is at 28.9%, of which 25% is MCMs use. Such
34 a low uptake translates into high unmet need 17% amongst married women. To bridge the gap
35 between the health system and beneficiaries, there is a need to assess predictors that influence
36 voluntary uptake of MCMs among women, at the health services and individual levels.

37 Methods: A cross-sectional household survey was conducted in two districts of Sindh, Pakistan
38 namely Matiari and Badin .In total, 1684 Married Women of Reproductive Age (MWRA) 15-49
39 years were interviewed. For the selection of eligible respondents, a two- stage stratified cluster
40 sampling strategy was used. Univariate and multivariable logistic regression was used to determine
41 the predictive factors for the increase in the use of MCM.

42 Results: Mean age was $32.3 \pm SD 7.1$ years. Average number of children per woman was 4.0 ± 2.0 .
43 Use of modern methods of contraceptive was 26.1% [n=441].Statistically significant socio
44 demographic predictors of MCM included: Number of children 4 or more (AOR 5.234 95%CI
45 2.78-9.84), Mother having primary education (AOR 1.730 95% CI 1.26-2.36), and Husband
46 having middle education [AOR 1.69 95% CI 1.03 – 2.76].Maternal health services indicators
47 included postnatal checkup of mother (AOR 1.46 95% CI 1.09 – 2.05); women who were visited
48 by Lady Health Workers in their postnatal period and were counseled on family planning
49 (AOR 1.83 95% CI 1.386 - 2.424).

50 Conclusion: Voluntary uptake of modern contraceptive methods is higher in women having 2 or
51 more children, having primary education and husband having middle education. Significantly,
52 receiving post-natal checkup at facility, and Lady Health Worker visit after delivery have more

53 likelihood to opt for contraception. Additionally, young couple counseling on family planning is
54 imperative to bridge the gap between knowledge and its translation into practice. There is also a
55 need to focus on the provision of integrated family planning and maternal, newborn, and child
56 health services through facility-based and community engagement platforms.

57 Keywords: Modern contraceptive methods, Predictors, Family planning, maternal health,
58 Reproductive health, Lady health worker

59

60

61

62

63

64

65

66

67

68

69

70

71 Introduction

72 According to the 7th population and housing census 2023, the population of Pakistan stood at 241.5
73 million, with an annual growth rate of 2.5% making it 5th populous country in the world [1]. This
74 growth rate is double compared to South Asia's average rate (1.2%). It is forecasted to reach 403
75 million inhabitants in 2050 [2]. This rapid and for the most part, uncontrolled population growth
76 has contributed to the economic and social crisis. Pakistan faces and affects the health of women
77 and children disproportionately. Overall, 63% of the population lives in rural areas with low
78 accessibility to health care services.

79 Global evidence indicates that such rampant population growth rates are also associated with poor
80 coverage of family planning programs and unmet needs for contraception. Investments in social
81 determinants and programs for supporting and scaling up modern contraception rates play a key
82 role in reducing maternal and infant mortality and improving human, social, and economic
83 development on an individual and national level [3, 4, 5].

84 Despite the benefits of family planning, the use of modern contraceptive methods (MCMs) in
85 Pakistan has been stagnant for the last decade. The country-wide contraceptive prevalence rate
86 (CPR) stands at 34%, with MCMs comprising 26% of this rate [6]. Within Pakistan, the second
87 largest province of Sindh reports contraceptive use at 28.9%, of which 25% is attributed to MCM
88 use. Such low uptake rates translate into high unmet need 17% amongst married women who are
89 fecund and desire birth spacing or limit the number of children they have. Despite these needs,
90 they are currently not accessing or utilizing any methods of contraception. [7].

91 The reasons for this low MCM access and utilization are varied and multifaceted. A major
92 contributing factor is the prevalence of early marriages, which constitutes 16.3% of all marriages
93 and primarily involves those between the ages 15 and 19 years [7].

94 In keeping with the need to improve these rates, previous studies have attempted to explore various
95 predictors of modern contraceptive uptake. Predictors of uptake covered in the local literature
96 include a woman's age, education, wealth status, and positive attitude towards contraception at an
97 individual level [8]. Further studies have reported that interventions encompassing elements of
98 social franchising, mid-level health care providers (community midwives) engagement, and mass
99 media participation substantially increase the prevalence of contraceptive use [9, 10]. Thus, these
100 determinants provide an important evidence base to effectively design, implement and further
101 evaluate interventions aimed at increasing contraceptive uptake.

102 Addressing these determinants requires a distinct strategy, intervention, or program. However,
103 introducing numerous new programs external to the existing healthcare infrastructure is unfeasible
104 in light of the country's economic instability combined with a lack of political will and a heavily
105 underfunded healthcare system [11]. Therefore, the main aim of this paper is to assess the
106 predictors associated with the uptake of MCMs in the domains of sociodemographic, reproductive,
107 and maternal health with corresponding knowledge, attitude, and use amongst married women of
108 reproductive age (MWRA) between 15-49 years in two districts of rural Sindh. Protocol paper has
109 been published for more details on methodology [12].

110 **Methods and Materials**

111 This was a cross-sectional study with face-to-face interviews of married women of reproductive
112 age at household level, using a structured questionnaire, as the pregnancies in Pakistan primarily
113 occur following marriage [13]. This survey was conducted in two districts of the province of Sindh:
114 Matiari and Badin (Figure 1).

115 **Figure 1. Study Sites – Badin and Matiari Districts of Province Sindh, Pakistan**

116 The selection of districts was based on propensity matching score further details could be obtained
117 from protocol paper [12]. Data was collected on all relevant program indicators, including the
118 study population's modern contraceptive prevalence rate (mCPR), antenatal and postnatal care
119 services, skilled birth attendance, and socio-economic and demographic status.

120 Sample size estimation

121 As per the Multiple Indicator Cluster Survey for Sindh 2018-2019 [7], the CPR reported for Sindh
122 province was 28.9%. Based on the CPR, with 95% CI and 80% power, an assumed design effect
123 of 1.5 and a 7% non-response rate, a sample size of 880 MWRA (15-49 years) per district was
124 estimated.

125 The targeted population for this study consisted of MWRA between 15-49 years of age residing in
126 the study districts within the catchment population of public sector health facilities. The exclusion
127 criteria for the sample included: (i) families who had moved into the project area within three
128 months with no intent to stay for a prolonged period in the project area and (ii) women of
129 reproductive age with mental disorders.

130 A two-stage stratified cluster sampling strategy was used to select eligible respondents. Clusters
131 were created with at least 100-150 households and were established as primary sampling units
132 (PSUs). In the first sampling stage, clusters were randomly selected from each district's list of
133 clusters (sampling frame). Subsequently, a line listing operation was conducted to determine
134 eligible households assigned as secondary sampling units (SSUs). In the second stage, 20 eligible
135 households with at least one MWRA were selected using systematic sampling to reach the
136 calculated sample size (covering 44 clusters in each district).

137 Survey questionnaire

138 The household questionnaire was adopted from the Pakistan Demographic and Health Survey
139 (PDHS) questionnaire for women of reproductive age [6]. The questionnaire was then pre-tested
140 in 5% of the total sample population, then translated into the local Sindhi language. A customized
141 digital application and software were developed for digital data collection.

142 Survey training

143 Before data collection, a five-day training workshop on the data collection process was conducted
144 for field teams (directly involved in field-based data collection). All sections of the questionnaires
145 were discussed in detail, followed by a mock field-based data collection activity for data collectors
146 supervised by the senior research staff. In each district, there are three teams consisting of four
147 senior research assistants and one research associate who have received 16 years of education in
148 social or medical allied sciences. The teams were carefully formed based on educational
149 qualifications and job descriptions to effectively conduct the survey. The last training session was
150 dedicated to feedback and troubleshooting based on issues reported from the mock exercise. The
151 sessions involved extensive training on informed consent, manual and digital data collection
152 processes. Training manuals were developed and distributed to the data collectors and
153 questionnaires for pilot testing in the field.

154 Data collection and data confidentiality

155 The study was conducted in each district by the assigned data collection team consisting of one
156 team leader and four data collectors. Before the data collection process commenced, line listing
157 was undertaken in selected clusters. Verbal informed consent was taken from married respondents
158 and details of the risks and benefits of their participation were explained in the presence of an

159 impartial witness and guardian if respondent was under 18 years of age. They were given an
160 opportunity to ask any questions they had regarding the study, after which their consent was
161 obtained (except for those refusing to participate).

162 Data was collected between 1st October 2020 and 31st December 2020, on personal digital
163 assistants (Model Samsung SM T289) through an Android application designed for field-based
164 data collection. Data collection duration was 45- 60 minutes. All data was uploaded electronically,
165 analyzed, and displayed on the virtual project dashboard. To maintain confidentiality, coded data
166 was stored in a password-protected and encrypted system, ensuring that only authorized
167 individuals had access to it.

168 Data analysis

169 The collected data was then analyzed in STATA version 17 (Stata Corp, Texas). Descriptive
170 statistics, bivariate statistical tests, chi-square tests, and independent sample t-tests were used to
171 summarize the data. Categorical variables were presented in frequency and percentages, while
172 mean and standard deviation were used for continuous variables. Furthermore, univariate and
173 multivariable logistic regression was used to determine the predictive factors associated with
174 increase in the use of modern methods of contraceptives amongst married women of reproductive
175 age. Variables significant at a p-value of less than 0.20 in the univariate analysis were included
176 in multivariable logistic regression to measure association at 95% confidence interval [14]. The
177 final model was constructed using backward elimination; variables were retained if the p-value
178 was less than 0.05 [15].

179 Ethical approval

180 Ethical approval was received from the Ethical Review Committee (ERC) of the Aga Khan
181 University on July 16, 2020 (2020-3606-18261). The study protocol was approved by the National
182 Bioethics Committee, Pakistan (4-87/NBC-514/22/857).

183 Results

184 From the total sample size of 1,760, 96% (n=1,684) of MWRA were interviewed. There was a
185 high response rate of 96%, as projected in the interview-based studies at the household level.

186 Sociodemographic characteristics of MWRA

187 In terms of sociodemographic characteristics in table 1, the mean age of the study participants
188 was 32.3years, with a standard deviation (SD) of 7.1 years. Among respondents, 27.0% (n=453)
189 having formal education. The average number of children per woman was 4.0 ± 2.0 , and the
190 average number of members living in the household was 7.2 ± 2.7 .

191 **Table 1: Socio demographic characteristics of MWRA(n=1,684)**

Variable	Overall	
	mean	SD
Household Size	7.2	± 2.7
Respondent's age	32.3	± 7.1
Husband's Age	36.4	± 8.5
Number of Children (Parity)	4.0	± 2.0
Number of Pregnancies	4.2	± 2.3
	n	%
Mother's Education		
No education	1231	73.0
Primary	224	13.3
Middle	58	3.4
Secondary	68	4.0
Intermediate or above	103	6.1
Wealth Quantile		
Poorest	335	19.8
Poor	336	19.9
Middle	339	20.1
Rich	338	20.0
Richest	336	19.9

192

193 Current use of contraceptives

194 Table 2 showed that from a total of 1,684 married women, 76.2% (n=1255) reported that they
195 heard at least one method of family planning to delay pregnancy. The difference in knowledge of

196 contraceptives was found to be significant in current users compared to non-users. Almost half of
197 the women who reported having knowledge of contraceptives had used family planning methods
198 in the past. Ever use of any contraceptive method to delay or avoid pregnancy was observed to be
199 significant amongst respondents.

200 The contraceptive prevalence rate was 27.4 % (n=462) in the sampled population. Furthermore,
201 the overall use of modern contraceptive methods was observed to be slightly lower at 26.1%
202 (n=441). Among different methods, the most common methods reported by the respondents were
203 condoms (n=87; 5.1%) followed by pills (n=83; 5.0%) and injectables (n=63; 3.7%). Out of all the
204 long-acting reversible methods, less than two percent of the sample population reported using
205 implants (n=41; 2.4%) and IUDs (n=12; 0.7%). The use of implants was seen as preferential to the
206 use of IUDs across both districts.

207 Moreover, the lactation amenorrhea method (LAM) use was reported by 6 women. However, one
208 woman reported the use of standard days methods (SDM). Moreover, among permanent methods
209 of contraception, 6.5% (n=109) of the respondents reported female sterilization and only one
210 respondent reported male sterilization method use.

211 The prevalence of traditional methods was less than two percent amongst all surveyed women
212 (n=22; 1.3%). Among such practices, 1.1% (n=19) of observations were recorded for the
213 withdrawal method, which was found to be the most common, followed by the rhythm method
214 (n=3; 0.1%). Additionally, distribution of modern contraceptive methods among socio
215 demographic characteristics attached as supporting information (S1_File).

216 **Table 2: Current use of contraceptives reported by MWRA (n=1,684)**

Characteristics	Overall	
	n	%

217	Ever heard of any method to delay pregnancy	1255	76.2
218	Ever used any method to delay/avoid pregnancy	625	38.5
219	Current use of contraceptive	463	27.4
220	Any modern method	441	26.1
220	Pills	83	5.0
221	Injectables	63	3.7
222	Condoms	87	5.1
222	Intra Uterine Devices(IUD)	12	0.7
223	Implant	41	2.4
224	Standard Days Method (SDM)	1	0.1
224	Lactational Amenorrhea Method (LAM)	6	0.3
225	Female sterilization	109	6.5
226	Male sterilization	1	0.1
226	Traditional method	22	1.3
227	Rhythm method	3	0.1
228	Withdrawal	19	1.1

229
230

231 Predictors for use of modern contraceptive methods (MCM)

232 Table 3 shows the univariate and multivariate analysis of the predictors associated with the use of
233 modern contraceptives.

234 Univariate Analysis

235 The odds of using contraceptives among women with more than four children were four times
236 higher than those with only one child (OR=3.739; 95% CI=2.274 – 6.149). The likelihood of use
237 of contraceptives was two times higher for women aged 35-39 years compared to women aged
238 below 19 years (OR 2.556 95% CI 1.345-4.857). Women who had completed middle-level
239 education were at least twice as likely to use modern contraceptives to delay pregnancy than those
240 without education (OR 1.629 95% CI 0.970 – 2.737). Similarly, husbands who had completed their
241 middle-level education were at least two times more likely to use contraceptives compared to those

242 who were uneducated (OR 1.879 95% CI 1.201 - 2.939). The odds of modern contraceptive use
243 among women who belonged to the rich wealth index were two times higher than women in the
244 poorest wealth index category (OR = 1.722. 95% CI 1.051 - 2.).

245 Women who were using health facilities for maternal health services showed a significant increase
246 in the odds of using contraceptives in their postpartum period compared to women who were not
247 using health facilities. Mothers who sought antenatal care services had higher odds of adopting
248 modern contraceptive methods (OR = 1.259 95% CI 0.956-1.778) as compared to those who didn't
249 have ANC visits. Similarly, mothers who had visited health facilities four or more times for
250 antenatal care checkups were 1.259 times more likely to use contraceptives in their postpartum
251 period than those who had visited less than four times during their pregnancy.

252 Moreover, mothers who delivered at a health facility showed higher odds of using contraceptives
253 than those who delivered at home (OR = 1.335 95% CI 0.996 - 1.790). The odds of use of
254 contraceptives amongst women with cesarean birth were 1.4 times the odds of use of
255 contraceptives amongst women who had either normal or assisted delivery (95% CI 1.041 - 1.888).
256 Women who had a postnatal checkup also showed higher odds of using postpartum contraceptives
257 than those who didn't (OR = 1.330 95% CI 1.008 - 1.754). Lady Health Worker visits after delivery
258 and for PNC were associated with a significant increase in the odds of the use of contraception
259 (OR = 2.028 95% CI 1.569- 2.622). Similarly, family planning counseling during antenatal care
260 visits also showed a significant increase in the odds of the use of postpartum contraceptives
261 compared to those who had not received any counseling (OR = 1.604 95% CI 0.746-3.446).

262 Multivariate Analysis

263 The multivariate logistic model shows that women with four or more children are four times more
264 likely to use contraceptives than those with only one child (AOR = 5.234 95% CI 2.784 - 9.842).
265 Compared to the bivariate model, after controlling for other covariates, the odds of use of
266 contraceptives were significantly higher among women who had completed primary education
267 compared to uneducated women (AOR = 1.730 95%CI 1.264 – 2.367).The odds ratio remained
268 significant for husbands who had attended middle level education in the multivariate model
269 (AOR = 1.695 95%CI 1.039- 2.762).The odds of a postnatal checkup of the mother were 1.446
270 times higher amongst mothers who had used contraceptives (AOR = 1.466 95% CI 1.093-2.052)
271 as well as women who were visited by the Lady Health Worker in their postnatal period and were
272 counseled on family planning (AOR = 1.833 95%CI 1.386- – 2.424).

Table 3: Predictors for use of modern contraceptive method (MCM) among Married Women of Reproductive age (MWRA- 15-49years)

Variable	Modern Contraceptive Use		Univariate Analysis		Multivariate Analysis	
	Yes	No	Crude Odd Ratio	95% CI	Adjusted Odds Ratio	95% CI
Number of children						
1	30 (6.8%)	230 (18.7%)	1	1	1	1
2	62 (14.3%)	210 (16.4%)	2.395*	1.425 – 4.024	2.641*	1.445 - 4.825
3	60 (12.4%)	203 (15.9%)	2.140*	1.284 – 3.567	2.644*	1.401 - 4.990
4 or more children	289 (66.6%)	600 (49.0%)	3.739*	2.274 - 6.149	5.234*	2.784 - 9.842
Mother's age [in years]						
15-19	7 (2.2%)	44 (3.9%)	1	1		
20-24	35 (8.1%)	143 (12.0%)	1.186	0.562 - 2.505	-	-
25-29	58 (12.7%)	257 (19.1%)	1.171	0.551 - 2.487	-	-
30-34	117 (24.9%)	310 (25.0%)	1.747	0.806 - 3.787	-	-
35-39	119 (28.5%)	245 (19.6%)	2.556*	1.345 - 4.857	-	-
40 or above	105 (23.7%)	244 (20.5%)	2.025*	0.983 - 4.170	-	-
Mother's education						
No education	289 (67.3%)	942 (76.0%)	1	1	1	1
Primary	79 (17.4%)	145 (12.3%)	1.599*	1.211 - 2.112	1.730*	1.264 - 2.367
Middle	20 (4.4%)	38 (3.0%)	1.629*	0.970 – 2.737	1.671	0.950 - 2.940
Secondary	20 (4.2%)	48 (3.8%)	1.240	0.716 - 2.150	1.534	0.820 - 2.867
Intermediate or above	33 (6.7%)	70 (4.8%)	1.579	0.931 - 2.677	1.819	1.017 - 3.254
Husband's education						
No education	158 (38.0%)	501 (39.9%)	1	1	1	1
Primary	72 (15.6%)	253 (21.1%)	0.777	0.527 - 1.146	0.713	0.481 - 1.055
Middle	42 (8.9%)	67 (5.0%)	1.879*	1.201 – 2.939	1.695*	1.039 – 2.762
Secondary	61 (12.8%)	152 (12.2%)	1.097	0.739 – 1.629	0.953	0.631-1.439
Intermediate or above	107 (24.6%)	250 (20.0%)	1.277	0.919 – 1.774	0.964	0.657- 1.416
Wealth Index						
Poorest	69 (14.4%)	266 (19.5%)	1	1	1	1
Poor	70 (16.8%)	266 (21.4%)	1.067	0.675 - 1.689	-	-
Middle	93 (21.8%)	246 (21.1%)	1.400	0.895 - 2.188	-	-

Rich	104 (24.7%)	234 (19.4%)	1.722*	1.051 - 2.820	-	-
Richest	105 (22.3%)	231 (18.7%)	1.620	0.976 – 2.689	-	-
Sought ANC	408 (92.6%)	1077 (87.4%)	1.790*	1.073 - 2.985	1.736*	1.078-2.796
4 or more ANC Visits	213 (48.8%)	521 (43.1%)	1.259*	0.956 - 1.778	-	-
FP counseling during ANC Check-up	27 (5.9%)	50 (3.8%)	1.604*	0.746 – 3.446	-	-
LHW Visited during pregnancy	304 (69.0%)	865 (69.2%)	0.988	0.742 - 1.37	-	-
Facility births	361 (81.3%)	949 (76.5%)	1.335*	0.996 - 1.790	-	-
Skilled Birth Attendant	360 (81.1%)	952 (76.7%)	1.308*	0.973- 1.758	-	-
Type of Delivery						
Normal/Assisted	347 (80.0%)	1046 (84.9%)	1	1		
C-Section	94 (20.0%)	197 (15.1%)	1.402*	1.041 - 1.888	-	-
PNC checkup of mother	287 (63.0%)	707 (56.2%)	1.330*	1.008 - 1.754	1.466*	1.093 – 2.052
LHW Visited after delivery	300 (65.7%)	741 (57.9%)	1.397*	1.076 - 1.813	-	-
LHW counseling for family planning as a part of PNC visit	150 (33.6%)	253 (20.0%)	2.028*	1.569 - 2.622	1.833*	1.386- 2.424

273 Discussion

274 This study assessed the predictors for opting for voluntary modern contraceptive methods to
 275 inform future strategies and decisions. This was achieved through univariate and multivariate
 276 analysis for specific variables, including knowledge, practices, access, and use of reproductive and
 277 maternal health services. The results showed the number of children, mother’s education,
 278 husband’s education, postnatal checkup of mother, and LHW counseling for family planning as
 279 part of the PNC visit; were the most identified predictors of modern contraceptive methods. The
 280 results provide an insight into the design of family planning and maternal health programs and the
 281 required modifications as per the local context.

282 It was noted that the average number of live births was at least four children per MWRA; women
 283 with more than four children were more likely to use modern contraceptives. This showed a
 284 significant inclination of the sample population towards limiting their pregnancy rather than

285 spacing between desired pregnancies, particularly in women aged more than 35 years. The
286 majority of them opted for sterilization, a surgical procedure, while younger respondents tend to
287 prefer short-acting methods like condoms, pills, and injections. The utilization of long-acting
288 reversible contraceptives (LARC), such as intrauterine devices (IUDs) and implants, was not
289 widely observed. These patterns among participants align with the results of similar studies
290 conducted in similar contexts. [16, 17, 18].

291 Conversely, contraception needs related to desire for spacing are predicted to be greater among
292 younger women; however, the results show the reverse whereby contraceptive use amongst
293 younger woman was lower. This dilemma of high need but low use discerned in our results is in
294 line with the findings of previous studies in similar settings [19, 8].

295 Additionally, the prevalence of contraceptive use among the respondents was almost similar to
296 that reported in the national and regional statistical records [6, 7]. However, the current study has
297 highlighted gaps between the awareness and practice of modern contraceptive methods. Though
298 majority of the participants were aware of contraceptive methods, this did not translate into
299 practice as consistent with other studies [20]. It also highlights the intention to use contraceptives
300 and its translation into practice as based on the willingness motivated by the presence and
301 fulfilment of favorable conditions [21]. This could be indicated as an unmet need for family
302 planning in married couples [22]. Another significant element contributing to narrowing the gap
303 between knowledge and application is the accessibility of contraceptive methods along with their
304 consistent availability. The uninterrupted provision of these methods plays a pivotal role in
305 addressing the issue of unmet needs and fostering the adoption of MCM [23]. Pakistan, as a
306 signatory to the Sustainable Development Goals (SDGs) and a participant in the FP 2020
307 commitments, has taken a firm stance on ensuring comprehensive access to family planning

308 resources, services, and commodities. Diverse initiatives have been initiated within the country in
309 collaboration with international and domestic non-governmental organizations, with sustained
310 support from National, Provincial, and District governments. A recent development in this context
311 is the introduction of the Costed Implementation Plan (CIP) by the Sindh government under the
312 FP2020-30 initiative [24] which holds the potential to expedite advancements towards achieving
313 the desired mCPR targets while focusing on the management of supply continuity.

314 Moreover, other negative attitudes and misconceptions, including fear of side effects and a
315 preference for a male baby, continue to contribute to low uptake of contraceptives [22, 25, 8, 18,
316 26]. Among subgroups in the sample population, the contraceptive prevalence rate was higher in
317 women from the rich quintile, similar to other studies in the local context [8, 27]. Interestingly,
318 women educated at the primary level were more likely to opt for modern contraceptive methods
319 than women who had attained comparatively higher education as opposed to previous studies [27,
320 28]. Furthermore, in univariate analysis, contraceptive use showed an increase with increasing
321 levels of males' education. However, this growth was more significant for husbands who attained
322 education until the middle level. These findings are compounded by the literature that describes a
323 comprehensive range of studies on family planning and its association with male engagement [29,
324 10]. Improving contraceptive uptake and support for opting for methods of choice relies upon the
325 provision of couple counseling and increased male engagement [30].

326 This study's most important finding was identifying postnatal checkup (PNC) as the most
327 significant predictor for opting for family planning. Respondents who sought PNC were more
328 likely to use modern contraceptive methods. Evidence in the literature has endorsed that postnatal
329 counseling before discharge significantly improves motivation for and practice of contraceptive
330 uptake [26, 31, 32, 33]. This study also observed that Lady Health Worker's counselling delivered

331 during a PNC visit increased the likelihood of family planning utilization. The probable cause of
332 this finding can be linked to the preference of the local community for Lady Health Workers to
333 discuss family planning matters on a one-to-one basis at the household level while maintaining
334 confidentiality [18]. Moreover, Lady Health Workers also provide family planning commodities
335 like condoms and pills during their routine visits. Consistent with our results, referral to a health
336 care facility through LHW also increases postpartum utilization of family planning. Therefore, the
337 Lady Health Worker Program strongly influences women's fertility choices and their use of sexual
338 and reproductive health care services [9].

339 Our study population reported a lower prevalence of family planning counseling received during
340 ANC visit. At the same time, these visits still influenced positive behaviors towards modern
341 contraceptive methods. In addition to this, family planning exposure was seen to be associated
342 with subsequent antenatal care visits, a higher number of deliveries performed by skilled
343 professionals, and a greater chance of delivery at a health facility [9]. Earlier studies have shown
344 a positive relationship between the delivery of high-quality antenatal family planning counseling
345 services and increased contraceptive use [34, 35].

346 Another study conducted in a similar setting as our study sites, concluded that in communities
347 where a higher proportion of women received quality antenatal care with an explanation of the
348 importance of postpartum family planning and where discussion of birth spacing was more
349 common, there was significantly higher contraceptive use particularly LARC. [36, 37]. These
350 studies were short-term funded programs highlighting the need for sustainable programs and
351 scaling up existing resources. Thus, multiple determinants then affect contraceptive uptake in the
352 context of Sindh, and strategically addressing these are imperative in improving MCM prevalence
353 rates.

354 However, what is achievable and sustainable is the expansion of interventions that meet specific
355 needs and target these factors by utilizing existing healthcare delivery systems.

356 Limitations and Strengths

357 A key strength of our cross-sectional study was that we had the opportunity to explore MCM
358 uptake in the context of multiple, complex factors, unique to the context of rural Sindh. This is
359 important, given that rural Sindh is home to highly marginalized and vulnerable communities
360 situated within disaster-prone areas that adversely affect reproductive health outcomes for males
361 and females.

362 Therefore, this study provides an important evidence-bases that can inform future research,
363 particularly on the qualitative side, to understand these predictors better and further, strategically
364 plan to address them at the community and facility level.

365 The cross-sectional study design of our study has several limitations on account of self-reported
366 data for multiple factors collected at a single point in time. However, this was controlled by the
367 backward selection of significant indicators for developing a final model of predictors.

368 Furthermore, we kept sociodemographic factors as predictors for family planning practices, which
369 followed the same pattern as the national trends and were covered by other local studies. The
370 analysis was further expanded for the studied sociodemographic variables to focus mainly on the
371 underlying factors influencing contraception uptake in our study districts. Although the data
372 regarding negative attitudes, concerns about potential side effects, and a preference for male babies
373 was not specifically addressed in the interviews, these factors have been extensively documented
374 as contributing to the low adoption of modern contraceptives.

375 Another limitation of our study is that our sampled population resided within the catchment area
376 of health facilities and already had some form of access to healthcare services in this regard.

377 Despite this limitation, our study population's CPR and MCMs prevalence rate was similar to other
378 regional and national level surveys. This validated the findings that the location of residence, on
379 its own, does not affect contraceptive uptake rates [36], making our sample comparable to other
380 studies.

381 A further limitation of the cross-sectional nature of this study is that findings cannot be used to
382 analyze all behaviors with regards to contraceptive uptake, given that it has been conducted at one
383 point in time. Understanding predictors requires a more detailed inquiry into motivations,
384 specifically, at the individual level, to adopt MCMs and the risks and challenges faced, in day to
385 day life owing to this. The study tried to mitigate this issue by the use of very detailed
386 questionnaires that comprehensively covered these aspects to a certain extent.

387 Conclusion and Recommendations

388 This study showed that the voluntary uptake of modern contraceptive methods is higher in women
389 receiving skilled-based delivery, postnatal checkups at the facility, and Lady Health Workers visits
390 after delivery. Furthermore, the results showed that women used MCMs for limiting pregnancies
391 rather than spacing and that utilization increases with age. This is an important finding that can be
392 used to inform behavioral change interventions to improve outcomes.

393 In addition to this, bridging the gap between knowledge and uptake to increase uptake is necessary.
394 Hence, various recommendations have been proposed in the light of this study for providing family
395 planning and maternal, newborn, and child health integrated services through facility-based
396 platforms and community engagement. This is associated with an increased likelihood of modern
397 contraceptive method acceptance. Integrated services must include antenatal and postnatal care
398 visits that incorporate family planning services [38, 31, 39]. Providing these services, calls for the

399 capacity strengthening of Lady Health Workers on family planning counseling regarding method-
400 specific side effects.

401 A further recommendation of this study is to focus on the introduction of feasible and effective
402 interventions which include in-service training, provision of job aids, and educational materials to
403 health care providers. This will allow them to counsel patients for family planning effectively.

404 At the community level, initiating and consistently monitoring educational sessions or programs
405 highlighting the importance of MCM use to community members is important. It should focus on
406 engaging adolescents through these community-level platforms to educate them about MCM at an
407 appropriate age to gradually reduce social resistance to uptake. A further focus should be on the
408 education of young couples on sexual and reproductive health and MCM. Such education,
409 information, and communication must be developed and delivered within the bounds of privacy
410 and respect for socio-cultural norms. Eventually, the positive impact emanating from these
411 initiatives should be used as an evidence base for advocating for policy changes and reform in
412 reproductive health service delivery and access oriented towards encouraging MCM use and
413 empowering young girls and women to reach their full potential.

414 Acknowledgment

415 We extend our appreciation to the Department of Health and the Population Welfare Department
416 of the Sindh Government for their essential partnership in project implementation. Our sincere
417 gratitude is directed towards the study participants, whose involvement and input are highly
418 valued. Additionally, we acknowledge the dedicated district-based personnel for their consistent
419 assistance throughout project activities, playing a crucial role in the effective achievement of this
420 initiative.

421 **Funding Details**

422 “This research was funded by Global Affairs Canada through United Nations Population Fund

423 (UNFPA), grant number SMK532400”

424

425 **Declaration of Interest statement**

426 The authors report there are no competing interests to declare

427 References

428

- [1] PBS, "https://www.pbs.gov.pk/content/announcement-results-7th-population-and-housing-census-2023-digital-census," Pakistan Bureau of Statistics, August 2023. [Online]. Available: <https://www.pbs.gov.pk/content/population-census>. [Accessed 7 August 2023].
- [2] UN, "United Nations, Department of Economic and Social Affairs, Population Division. World Population Prospects: The 2019 Revision, DVD Edition.," 15 1 2022. [Online]. Available: <http://data.un.org/Data.aspx?q=pakistan+population&d=PopDiv&f=variableID%3a12%3bcrID%3a586>. [Accessed 18 2 2022].
- [3] Wazir MA, Alazar YM, Kadirov B. Family planning: smartest investment for achieving the sustainable developments goals for Pakistan. *J Pak Med Assoc.* 2021 Nov 1;71(11). <https://www.researchgate.net/publication/355789536>
- [4] Chola L, McGee S, Tugendhaft A, Buchmann E, Hofman K. Scaling up family planning to reduce maternal and child mortality: the potential costs and benefits of modern contraceptive use in South Africa. *PloS one.* 2015 Jun 15;10(6):e0130077.
- [5] T Ali SA, Tikmani SS, Harrison MS, Saleem S, Goldenberg RL, McClure EM, Pasha O. Pregnancy Outcomes among Women with an Unintended Pregnancy: Findings from a Prospective Registry in Rural Pakistan. *EC Gynaecology.* 2018;7(2):57. https://ecommons.aku.edu/pakistan_fhs_mc_chs_chs/357
- [6] PDHS, "National Institute of Population Studies (NIPS) [Pakistan] and ICF.Pakistan Demographic and Health Survey slamabad`, Pakistan`, and Rockville`, Maryland`, USA: NIPS and ICF.," 2017-18.
- [7] MICS, "Multiple Indicator Cluster Survey 2018-19, Bureau of Statistics, Planning & Development Board, Government of Sindh, 2020. Sindh; Survey Findings Report. Karachi, Pakistan: Bureau of Statistics Sindh.," 2020.
- [8] Asif, M.F., Pervaiz, Z. Socio-demographic determinants of unmet need for family planning among married women in Pakistan. *BMC Public Health* 19, 1226 (2019). <https://doi.org/10.1186/s12889-019-7487-5>
- [9] Khan Q, Wang S. The effect of family planning exposure on fertility choices and reproductive health care in rural Pakistan. *Journal of Asian Economics.* 2021 Apr 1;73:101283.<https://doi.org/10.1016/j.asieco.2021.101283>
- [10] Azmat, S.K., Hameed, W., Hamza, H.B. *et al.* Engaging with community-based public and private mid-level providers for promoting the use of modern contraceptive methods in rural Pakistan: results from two innovative birth spacing interventions. *Reprod Health* 13, 25 (2016). <https://doi.org/10.1186/s12978-016-0145-9>

- [11] Hatcher, P., Shaikh, S., Fazli, H. *et al.* Provider cost analysis supports results-based contracting out of maternal and newborn health services: an evidence-based policy perspective. *BMC Health Serv Res* **14**, 459 (2014). <https://doi.org/10.1186/1472-6963-14-459>
- [12] Memon ZA, Reale S, Ahmed W, Spencer R, Lashari TH, Bhutta Z, Soltani H. Effects of integrating family planning with maternal, newborn, and child health services on uptake of voluntary modern contraceptive methods in rural Pakistan: protocol for a quasi-experimental study. *JMIR research protocols*. 2022 Mar 8;11(3):e35291.
- [13] A. a. S. M. A. Aziz, "'Proximate Determinants of Fertility in Pakistan [with Comments].'", *The Pakistan Development Review*, vol. 33, no. 4, pp. 727-42, 1994.
- [14] Teshale AB. Factors associated with unmet need for family planning in sub-Saharan Africa: A multilevel multinomial logistic regression analysis. *PloS one*. 2022 Feb 10;17(2):e0263885.
- [15] Beson, P., Appiah, R. & Adomah-Afari, A. Modern contraceptive use among reproductive-aged women in Ghana: prevalence, predictors, and policy implications. *BMC Women's Health* **18**, 157 (2018). <https://doi.org/10.1186/s12905-018-0649-2>.
- [16] Hackett K, Nausheen S, Yameen S, Hussain I, Khaneez Z, Shah I, Soofi SB. Exploring reasons for low uptake of widely available modern contraceptive methods in Karachi, Pakistan: a mixed-methods study. *J Glob Health Sci*. 2021 Dec;3(2):e12. <https://doi.org/10.35500/jghs.2021.3.e12>
- [17] Gonie, A., Wudneh, A., Nigatu, D. *et al.* Determinants of family planning use among married women in bale eco-region, Southeast Ethiopia: a community based study. *BMC Women's Health* **18**, 50 (2018). <https://doi.org/10.1186/s12905-018-0539-7>
- [18] Mustafa G, Azmat SK, Hameed W, Ali S, Ishaque M, Hussain W, Ahmed A, Munroe E. Family planning knowledge, attitudes, and practices among married men and women in rural areas of Pakistan: Findings from a qualitative need assessment study. *International journal of reproductive medicine*. 2015 Sep 1;2015. <https://doi.org/10.1155/2015/190520>
- [19] Anand BK, Singh J, Mohi MK. Study of unmet need for family planning in immunisation clinic of a teaching hospital at Patiala, India. *The Internet Journal of Health*. 2010;11(1):23-4.
- [20] Mubarik M, Jameel N, Khalil R, Gul S, Women MR. SOURCE OF KNOWLEDGE AND CHOICE OF CONTRACEPTIVE METHOD AMONG MARRIED RURAL WOMEN OF PAKISTAN." *International Journal of Recent Advances in Multidisciplinary Research*, vol. 3, no. 6, pp. 576-1580, 2016. ISSN: 2350-0743
- [21] Agha, S. Intentions to use contraceptives in Pakistan: implications for behavior change campaigns. *BMC Public Health* **10**, 450 (2010). <https://doi.org/10.1186/1471-2458-10-450>

- [22] Avinash KR, Suganthi S, Venmathi E. Prevalence of Contraceptive Methods among Eligible Couples in an Urban Area of Chennai-A Cross Sectional Study. *Journal of Pharmaceutical Research International*. 2021 Dec 13;33(55A):262-7.
- [23] Bakali Mukasa, Moazzam Ali, Madeline Farron & Renee Van de Weerd (2017) Contraception supply chain challenges: a review of evidence from low- and middle-income countries, *The European Journal of Contraception & Reproductive Health Care*, 22:5, 384-390, DOI: [10.1080/13625187.2017.1394453](https://doi.org/10.1080/13625187.2017.1394453)
- [24] CIP, "Costed Implementation Plan on Family Planning for Sindh (2015–2020). Pakistan: Population Welfare Department," *Family Planning 2030*, Karachi, 2016.
- [25] Channon, M. D. (2017). Son Preference and Family Limitation in Pakistan: A Parity- and Contraceptive Method-Specific Analysis. *International Perspectives on Sexual and Reproductive Health*, 43(3), 99–110. <https://doi.org/10.1363/43e4317>
- [26] Memon Z, Zaidi S, Riaz A. Residual Barriers for Utilization of Maternal and Child Health Services: Community Perceptions From Rural Pakistan. *Glob J Health Sci*. 2015 Nov 3;8(7):47-57. doi: [10.5539/gjhs.v8n7p47](https://doi.org/10.5539/gjhs.v8n7p47). PMID: 26925902; PMCID: PMC4965661
- [27] Azmat, S.K., Ali, M., Ishaque, M. *et al.* Assessing predictors of contraceptive use and demand for family planning services in underserved areas of Punjab province in Pakistan: results of a cross-sectional baseline survey. *Reprod Health* 12, 25 (2015). <https://doi.org/10.1186/s12978-015-0016-9>
- [28] Aslam SK, Zaheer S, Qureshi MS, Aslam SN, Shafique K. Socio-economic disparities in use of family planning methods among Pakistani women: findings from Pakistan demographic and health surveys. *PloS one*. 2016 Apr 7;11(4):e0153313.
- [29] Agha S, Fitzgerald L, Fareed A, Rajbhandari P, Rahim S, Shahid F, Williams E, Javed W, Currie S. Quality of labor and birth care in Sindh Province, Pakistan: Findings from direct observations at health facilities. *PloS one*. 2019 Oct 17;14(10):e0223701.
- [30] Ashfaq, Seemin and Maqsood Sadiq. 2015. "Engaging the missing link: Evidence from FALAH for involving men in family planning in Pakistan—Case study." Washington, DC: Population Council, The Evidence Project. doi: [10.31899/rh5.1001](https://doi.org/10.31899/rh5.1001)
- [31] Baqui AH, Ahmed S, Begum N, Khanam R, Mohan D, Harrison M, Al Kabir A, McKaig C, Brandes N, Norton M, Ahmed S; Projahnmo Study Group in Bangladesh. Impact of integrating a postpartum family planning program into a community-based maternal and newborn health program on birth spacing and preterm birth in rural Bangladesh. *J Glob Health*. 2018 Dec;8(2):020406. doi: [10.7189/jogh.08.020406](https://doi.org/10.7189/jogh.08.020406). PMID: 30023053; PMCID: PMC6036944.
- [32] Udgiri R, Sorganvi V. Knowledge attitude and practices of family planning methods among postnatal mothers-a hospital based study. *Age*. 2016;18(20):21-3. ISSN 0974-1143

- [33] Abbasi Y, Shaikh SR, Memon KN. Barriers and missed opportunities towards immediate and early post-partum family planning methods in Pakistan. *The Professional Medical Journal*. 2020 Jul 10;27(07):1448-53. <https://doi.org/10.29309/TPMJ/2020.27.07.4273>
- [34] León FR, Monge R, Zumarán A, García I, Ríos A. Length of counseling sessions and the amount of relevant information exchanged: a study in Peruvian clinics. *International Family Planning Perspectives*. 2001 Mar 1:28-46. <https://doi.org/10.2307/2673802>
- [35] Sebastian MP, Khan ME, Kumari K, Idnani R. Increasing postpartum contraception in rural India: evaluation of a community-based behavior change communication intervention. *International perspectives on sexual and reproductive health*. 2012 Jun 1:68-77. <https://www.jstor.org/stable/41585361>
- [36] Tappis H, Kazi A, Hameed W, Dahar Z, Ali A, Agha S. The role of quality health services and discussion about birth spacing in postpartum contraceptive use in Sindh, Pakistan: a multilevel analysis. *PloS one*. 2015 Oct 20;10(10):e0139628.
- [37] Ismail A, Ashraf M, Idris IB, Thaver I, Ahmad AM, Siddique SJ. IDENTIFYING FACTORS ASSOCIATED WITH THE USE OF A SCALED-UP POSTPARTUM FAMILY PLANNING PROJECT IMPLEMENTED IN PUNJAB, PAKISTAN. *Malaysian Journal of Public Health Medicine*. 2021 Aug 28;21(2):466-74. <https://doi.org/10.37268/mjphm/vol.21/no.2/art.1091>
- [38] Brickley DB, Chibber K, Azman H, Spaulding A, Lindegren ML, Kennedy C. Systematic Review of Integration between Maternal, Neonatal and Child Health and Nutrition and Family Planning, Final Report. USAID, Cochrane HIV/AIDS Group and University of California, Washington, DC: USAID, Cochrane HIV/AIDS Group and University of California. 2011.
- [39] Cooper CM, Wille J, Shire S, Makoko S, Tsega A, Schuster A, Hausi H, Gibson H, Tappis H. Integrated family planning and immunization service delivery at health facility and community sites in Dowa and Ntchisi districts of Malawi: a mixed methods process evaluation. *International Journal of Environmental Research and Public Health*. 2020 Jun;17(12):4530. <https://doi.org/10.3390/ijerph17124530>
- [40] Caliendo M, Kopeinig S. Some practical guidance for the implementation of propensity score matching. *Journal of economic surveys*. 2008 Feb;22(1):31-72. <https://doi.org/10.1111/j.1467-6419.2007.00527.x>
- [41] Jabeen, S., Rathor, A., Riaz, M. *et al.* Demand- and supply-side factors associated with the use of contraceptive methods in Pakistan: a comparative study of demographic and health surveys, 1990–2018. *BMC Women's Health* **20**, 265 (2020). <https://doi.org/10.1186/s12905-020-01112-4>
- [42] Nadeem, M., Malik, M.I., Anwar, M. *et al.* Women Decision Making Autonomy as a Facilitating Factor for Contraceptive Use for Family Planning in Pakistan. *Soc Indic Res* **156**, 71–89 (2021). <https://doi.org/10.1007/s11205-021-02633-7>

[43] Khowaja F, Pervaiz M, Khatoon N. Male Involvement in Family Planning among Squatter Settlement Residents of Karachi. *Journal of Asian Midwives (JAM)*. 2019;6(1):26-42.

[44] Saleem, S., Rizvi, N., Shahil Feroz, A. *et al.* Perceptions and experiences of men and women towards acceptability and use of contraceptives in underserved areas of Karachi, Pakistan: a midline qualitative assessment of Sukh initiative, Karachi Pakistan. *Reprod Health* **17**, 95 (2020). <https://doi.org/10.1186/s12978-020-00946-3>

429

430 [Supporting Information](#)

431 S1_File PDF

Pakistan

medRxiv preprint doi: <https://doi.org/10.1101/2023.09.04.23295034>; this version posted September 5, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).



Figure