perpetuity. It is made available under a CC-BY 4.0 International license .

Cognition, physical function, and life purpose in the rural elderly population: a systematic review protocol

- 4 Hércules Lázaro Morais Campos1; Elisa Brosina De Leon2; Ingred Merllin Batista
- 5 de Souza3; Anna Quialheiro4; Elizabete Regina Araújo de Oliveira5
- 1 Professor at the Physiotherapy Department of the Federal University of
- 7 Amazonas/ Institute of Health and Biotechnology (UFAM/ISB). 2 PhD student in
- 8 Collective Health at the Department of Collective Health at the Federal University
- 9 of Espírito Santo (UFES).

1

2

3

- 2 Professor at the Universidade Federal do Amazonas, Stricto Sensu Post-
- 11 graduation Program in Human Movement Sciences, Faculdade de Educação
- 12 Física e Fisioterapia, Manaus, Amazonas, Brazil.
- 13 3 Department of Physical Therapy, Speech-Language Pathology, and Audiology
- and Occupational Therapy, Faculty of Medicine at University of São Paulo,
- 15 Cipotânea Street, Vila Universitária, 51, São Paulo 05360-000, SP, Brazil; e-
- 16 mail:ingredmerllin@usp.br
- 4 Health and Human Movement Unit, Polytechnic Institute of Health of the North
- 18 CESPU, Portugal.
- 5 Professor of the Post-graduate Program in Collective Health of the Federal
- 20 University of Espírito Santo.

Abstract

21

22

23

- 24 **Introduction:** Aging in the rural setting world from the perspective of cognition,
- 25 physical function, and life purpose, essential constructs for a prosperous old age,
- still needs to be better discussed. Thus, this systematic review protocol highlights
- the prevalence of cognitive decline, physical functioning, and life purpose in older
- adults aging in rural community settings. Methods and analysis: We will include
- 29 NOTTOS Sectiones studies obtains the dentities of the contraction o

perpetuity.

It is made available under a CC-BY 4.0 International license .

MEDLINE, LILACS, PsycINFO, Scopus, SciELO, and Web of Science. For the 30 31 first selection of studies, the Ryyan software will be used, and to check the methodological quality and the risk of bias, we will use COSMIN. For the primary 32 analysis, the titles and abstracts available in the search engine will be analyzed 33 by the following MeSH descriptors "Physical functioning,"; "Cognition,"; 34 "Cognitive function,": "Life purpose,": Personal satisfaction: Subjective well-35 being; "Elderly"; "Older"; "Rural aging"; "Rural population"; "Communities, rural"; 36 "Distribution, rural spatial"; "Medium communities"; "Rural settlement"; "Small 37 community". If necessary, the secondary analysis will adopt a complete reading 38 of the selected articles by two blinded reviewers and confirmed by a third person. 39 Publication bias will be assessed using cross-sectional analytical study quality. 40 Sensitivity analyses will be performed by retrieving one article at a time and 41 42 analyzing their endpoints with the proposal to identify the manuscripts that significantly influenced the combined prevalence of the endpoints. 43

Strengths and limitations of this study:

- 46 It will present worldwide epidemiological data on aging in rural settings from the
- 47 perspective of cognition, physical functioning, and life purpose.
- 48 It is the first systematic review that involves the theme of life purpose worldwide
- 49 in rural elderly.

44

45

53

54

55

- It is a cross-sectional study that although it cannot describe clinical outcomes,
- 51 presents sociodemographic, cognition, physical functioning, and life purpose data
- that can help in public health decision-making for this population.

INTRODUCTION

- Aging within the rural context takes place in a variety of ways around the world,
- and increasingly the literature has pointed out that people aging in this unique
- setting have different epidemiological and health indicators than seniors aging in
- the urban context (1)
- There are controversies about the elderly aging in the rural context, sometimes
- the challenges are pointed as independence and active participation in
- community life, as well as attention to safety, choice, and location of housing,
- 63 loneliness, social isolation, difficulty accessing services, leisure, food,

perpetuity.

It is made available under a CC-BY 4.0 International license.

transportation, agriculture itself until retirement. On the other hand, seniors who age in rural settings may have good health, and quality of life especially in the cognitive(1) aspect, and have better access to health, services, and healthier living and eating habits although they are at constant risk of frailty (2) Well-conducted cross-sectional and observational studies have relevance within aging and measure the prevalence of health outcomes as well as help understand the social determinants of health and above all describe the characteristics of a given population (3)

There is still no synthesis of what aging in the rural world context looks like from

There is still no synthesis of what aging in the rural world context looks like from the perspective of cognition, physical function, and life purpose. Thus, this systematic review protocol was designed with the following question: (1) what is the prevalence of cognition, physical functioning, and life purpose in older adults aging in rural community settings?

METHODS AND ANALYSIS

This systematic review protocol followed the PICO strategy and was registered with the International prospective register of systematic reviews (PROSPERO) under number 2022 CRD42022311053.

Eligibility Criteria

Databases and Search Strategy

The search will include cross-sectional studies until April/2022 in journals indexed in the following health databases: Embase, Medical Literature Analysis and Retrieval System Online (MEDLINE), Latin American and Caribbean Literature in Health Sciences (LILACS), PsycINFO, SciVerse Scopus (Scopus), SciELO (Scientific Electronic Library Online) and Web of Science. Published until December 2021, studies that epidemiologically evaluated the outcomes of cognition, functionality, and life purpose within rural aging and that were published in Portuguese, English or Spanish will be accepted. The protocol will follow the Preferred Reporting Items for Systematic Reviews (PRISMA) quidelines (4)

The search methods will use 3-step Boolean operators with the search strategies following the uniqueness of each database. The search strategy will use: "rural aging", "elderly", "old", "elderly", "physical functioning", "cognition", and "life purpose" according to Boolean operators. Two independent HLMC and EBDL reviewers will search the studies, and if necessary, a third reviewer will be included to decide on tie-breaking criteria.

Study selection and data extraction

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

After evaluation of the titles and abstracts retrieved from the searches, potential full texts will be assessed for eligibility by two independent reviewers. Authors of potential full texts will be contacted for questions about the eligibility criteria and studies will be excluded when there is no response. Studies meeting the eligibility criteria will be included. For studies with the same sample, only those with the most representative sample of that population will be considered. Data relevant to the theme in question will be extracted from the selected articles.

The selection of articles will contemplate the following eligibility criteria: (a) inclusion criteria: studies that presented a prevalence of functioning conditions, cognitive decline, and life purpose in rural elderly; and (b) exclusion criteria: systematic review studies; methodological studies; instrument validation studies; and qualitative studies. The following information will be recorded: year of publication of the study, authors, characteristics of the institutions in which the study was conducted, sampling strategy, type of sample, prevalence of outcomes, and their associated factors. The information will be organized and presented in tables.

Study Evaluation

To assess the methodological quality of each study, i.e., to assess the risk of bias in a study's results, the corresponding COSMIN Risk of Bias box must be filled in. To determine the overall quality of a study, the lowest rating of any standard in the box is taken (i.e. the "worst score counts" principle). For example, if for a reliability study, one item in a box is rated as 'inadequate', the

overall methodological quality of that reliability study is rated as 'inadequate' which will generate a quality score at the end of the systematic review (5).

Strategy for Data Overview

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

The synthesis of outcome study data will be organized and presented according to the recommendations of the Meta-analysis of Observational Studies in Epidemiology (MOOSE) statement (6). When two or more articles are reported as results from the outcome database, only the most comprehensive one (with more details on sampling strategy and response rate) will be included in the metaanalysis. Results will be presented in Forest Plots, showing 95% confidence intervals and p-values. Heterogeneity will be assessed using the I² statistic, which will be considered high when I2 is equal to or greater than 75%. Statistical procedures will be performed in STATA 14.0, with the significance level set at 5% for two-tailed tests (6).

Patient and Public Involvement

There is no direct patient or public involvement in this study.

ETHICS AND DISSEMINATION

Ethical approval is not required for this study since no personal or private

information of the individuals will be involved. It is intended to submit this study

to a peer-reviewed academic journal.

Cross-sectional studies are more economically feasible but easy to conduct and

can establish preliminary evidence so that further advanced studies can be

conducted. Prevalence and incidence are established and can generate

important hypotheses, they can be analytical or descriptive, and depending on

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

revision of the manuscript.

whether the outcome variable is assessed possible associations with exposures or risk factors can be established (3). It seems that the functional physical capacity of elderly people aging in rural settings needs to be better understood, elderly women with good income are more likely to be sedentary as well as elderly men (7). Studies have demonstrated the factors associated with frailty in rural elderly such as age, gender, health status variables and that included self-perception of health and number of chronic conditions, functional covariates variables such as disability in basic activities of daily living (ABVD), disability in instrumental ADLs, length of stay in the chair and psychosocial problems, it seems that depressive symptoms and cognitive impairment, greater comorbidity and disability was found in this population (8). As for cognition, there is divergence in the literature related to the place of birth and where the rural elderly live. Sometimes the rural environment is pointed out as positive for aging (2) and at other times not (9) and (10). The Purpose of Life in rural elderly is still little investigated and very little is known about the impact of this condition on rural aging until this moment the literature points out that the elderly with a purpose of life are more resilient, have less risk for developing dementia, feel less pain and are happier and more functional (11). There are many questions to be answered about the way of aging in the rural context and some gaps still need to be filled and understood to understand rural elder aging. After all, is rural aging positive or negative on the cognition, physical function, and life purpose of the elderly population? **Contributions:** HLMC- Writing the review protocol, reviewing the search strategies, recording the review, writing the manuscript, and final revision of the manuscript. EBDL- Writing of the review protocol, review of the search strategies, and final revision of the manuscript. IMBS- Writing of the review protocol, review of the search strategies and final

- AQ- Writing of the review protocol, review of the search strategies and final review 196
- 197 of the manuscript.
- ERAO- Writing of the review protocol, review of the search strategies and final 198
- review of the manuscript. 199

201 **Funding Statement**

200

204

205

206

209

210

211

- Research scholarship from the Amazonas State Research Support Foundation 202
- 203 - Brazil (FAPEAM).

Competing Interest Statement 207

208 No competing interest.

REFERENCES

- 1. Winterton R BM. A decade of rural ageing research in the Australasian Journal 212
- 213 on Ageing. 2020 [cited 2023 Jan 14]; Available from:
- https://onlinelibrary.wiley.com/doi/10.1111/ajag.12879 214
- 2. Garbaccio JL, Tonaco LAB, Estêvão WG, Barcelos BJ. Aging and quality of 215
- life of elderly people in rural areas. Rev Bras Enferm [Internet]. 2018 [cited 2023] 216
- 14];71:724–32. Available from: 217 Jan
- http://www.scielo.br/j/reben/a/pC3sjdGyJnPbyC9PXygQRrF/abstract/?lang=en 218
- 3. Wang X, Cheng Z. Cross-Sectional Studies: Strengths, Weaknesses, and 219
- Recommendations. Chest [Internet]. 2020 Jul [cited 220
- 14];158(1S):S65-71. Available from: https://pubmed.ncbi.nlm.nih.gov/32658654/ 221
- 4. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et 222
- al. The PRISMA 2020 statement: an updated guideline for reporting systematic 223
- reviews. BMJ [Internet]. 2021 Mar 29 [cited 2023 Jan 14];372. Available from: 224
- https://pubmed.ncbi.nlm.nih.gov/33782057/ 225
- 5. Cosmin. Systematic reviews of measurement properties. 2017. 6. DF Stroup 1 226
- JBSMEOGWDRDMBJBTSST. Metanálise de estudos observacionais em 227

- epidemiologia: uma proposta de relato. Grupo de meta-análise de estudos 228
- observacionais em epidemiologia (MOOSE). JAMA. 2000;283(15):2008–12. 229
- 230 7. Ethisan P, Somrongthong R, Ahmed J, Kumar R, Chapman RS. Factors
- 231 Related to Physical Activity Among the Elderly Population in Rural Thailand. J
- Prim Care Community Health [Internet]. 2017 Apr 1 [cited 2023 Jan 14];8(2):71-232
- 6. Available from: https://pubmed.ncbi.nlm.nih.gov/27799413/ 233
- 234 8. Curcio CL, Henao GM, Gomez F. Frailty among rural elderly adults. BMC
- Geriatr [Internet]. 2014 Jan 10 [cited 2023 Jan 14];14(1). Available from: 235
- https://pubmed.ncbi.nlm.nih.gov/24405584/ 9. Silva JNMA da, Leite MT, 236
- Gaviraghi LC, Kirsten VR, Kinalski S da S, Hildebrandt LM, et al. Predicting 237
- dimensions of clinical-functional conditions and cognition in the elderly. Rev Bras 238
- Enferm [Internet]. 2020 Jul 13 [cited 2023 Jan 15];73:e20190162. Available from: 239
- http://www.scielo.br/j/reben/a/YdR9HPnBdgDXPR6wgKCZY5m/?lang=en 240
- 10. Jing Z, Li J, Wang Y, Ding L, Tang X, Feng Y, et al. The mediating effect of 241
- psychological distress on cognitive function and physical frailty among the 242
- elderly: Evidence from rural Shandong, China. J Affect Disord [Internet]. 2020 243
- 244 May 1 [cited 2023 Jan 14];268:88–94. Available from:
- https://pubmed.ncbi.nlm.nih.gov/32158011/ 245

248

249

- 11. Cristina Cristóvão Ribeiro MSYALNeri. Propósito de vida na vida adulta e na 246
- terceira idade: revisão integrativa. Cien Saude Colet. 2020;25(6):2127-42. 247