

1 **Prevalence and predictors of self-medication for COVID-19 among slum dwellers in Jinja City,**
2 **Uganda**

3

4 **Short title: Self medication for covid-19**

5

6 **Authors**

7 Prossy Nakito^{1*}, Angela N. Kisaky^{2,3}, Abel Wilson Walekhwa^{4,5}, Gloria Tumukunde³ Charity Mutesi³,

8 Nicholas Muhumuza³, Carolyne Nyamor³, David Musoke⁵, Geoffrey Musinguzi⁵, Dathan M.

9 Byonanebye^{1,6}

10 Affiliations:

11 ¹Department of Community Health and Behavioural Sciences, School of Public Health, Makerere
12 University, Kampala, Uganda

13 ²African Field Epidemiology Network Lugogo House, Plot 42 Lugogo By-Pass, Kampala

14 ³Makerere University School of Public Health, Department of Health Policy Planning and Management.

15 ⁴Diseases Dynamics Unit, Department of Veterinary Medicine, University of Cambridge,
16 United Kingdom

17 ⁵Department of Disease Control and Environmental Health, University School of Public Health.

18 ⁶Kirby Institute, University of New South Wales, Sydney, Australia

19

20 ***Corresponding author and address:** Prossy Nakito

21 Department of Community Health and Behavioural Sciences

22 School of Public Health, Makerere University, Kampala Uganda

23 Email: prossynakito@gmail.com

24

25

26 **Abstract**

27

28 **Introduction:** Self-medication is a serious public health concern globally and is more prevalent in
29 underserved populations, especially in resource limited settings. The lack of effective treatment for COVID-
30 19 and poor access to healthcare were drivers of self-medication. We investigated the prevalence and
31 associated factors with self-medication for COVID-19 among slum dwellers in a Ugandan slum.

32 **Methods and materials:** We conducted a cross-sectional study using randomly selected respondents from
33 slums in Jinja city, Uganda. Households were proportionately selected from the slums and one participant
34 with confirmed or self-reported COVID-19 during 2021 was recruited. Poisson regression with robust
35 standard errors was used to determine the crude (CPR) and adjusted prevalence ratios (APR) (95% CI) of
36 factors associated with self-medication. Variables were selected apriori and backward elimination approach
37 used to fit the final multivariate model in which variables with a $P \leq 0.05$ were included.

38 **Results:** Overall, 517 respondents were recruited, median age (years) was 31 (26-40), and 59% were male.
39 The prevalence of self-medication for COVID-19 was 87.23% (451/517), 95% CI: [84.00%-90.00%] and
40 56% knew that self-medication was dangerous. Age ≥ 50 years, compared to 20-29 years [APR: 1.12, 95%
41 CI: 1.05, 1.20], being female [APR: 1.07, 95% CI: 1.02, 1.13], minor [APR: 1.62, 95% CI: 1.25, 2.11], and
42 severe symptoms [APR: 1.51, 95% CI: 1.16, 1.96], access to internet [APR: 1.13, 95% CI: 1.07, 1.20].
43 Having medical insurance [APR: 0.63, 95% CI: 0.46, 0.87] and awareness about laws against self-
44 medication [APR: 0.89, 95% CI: 0.81, 0.97] were associated with a lower risk of self-medication.

45 **Conclusion:** The prevalence of self-medication in slum dwellers in Uganda was high despite high
46 awareness about its dangers. Self-medication was common in those with severe symptoms and those access
47 to internet. There is need to control infodemia and improve health insurance cover in informal settlements
48 within Uganda.

49 **Keywords:** *Self-medication, COVID-19, Slum dwellers, Confirmed COVID-19, Self-reported COVID-19*

50 **Introduction**

51
52 While Africa did not registered very high incidence of Coronavirus disease 2019 (COVID-19) compared
53 to the rest of the world, the disease had significant impact on people living in informal settlements. As of
54 February 2023, the World Health Organization (WHO) estimated that there were more than 9.4 million
55 cases of COVID-19, with approximately 173,295 deaths [1, 2]. Uganda experienced a first wave and a more
56 serious second wave between November to December 2020 and April to June 2021, respectively [3]. The
57 second wave of the pandemic was fuelled by five different variants and a high mortality rate, estimated at
58 36% [4]. Cumulatively, the country had reported 170,409 confirmed cases and 3,630 deaths by February
59 28th, 2023 [5]. Amidst this burden, effective treatments for COVID-19 were not accessible in Africa, and
60 the government treatment guidelines focused on symptom management [2].

61 COVID-19 impacted nearly every sector of development the world over. The health sector was not spared
62 and it experienced monumental challenges in trying to cope with and respond to the pandemic [6]. Several
63 countries instituted strategies such as restrictions on travel to reduce disease transmission and the burden
64 on the health [7, 8]. However, in resource limited settings, these measures exacerbated the existing
65 challenge of limited access to health services, especially in underserved communities. The infodemia
66 around COVID-19 infection, overstretched health systems and limitations in access to health services
67 exacerbated self-medication [9]. The World Health Organization (WHO) identified self-medication as a
68 significant public health issue with risks if not used properly [10]. It defines self-medication as the selection
69 and utilization of medicines to treat self-recognized symptoms or ailments without consulting a physician
70 [11]. It also includes the usage or re-usage of previously prescribed or unused drugs; direct purchasing of
71 prescription drugs without consultation; and irrational use of over-the-counter medications [12].

72 The effects of self-medication includes antimicrobial resistance, drug interactions, addiction or dependence,
73 and drug toxicity [13]. Potential risks of self-medication include incorrect self-diagnosis, adverse reactions,
74 dangerous drug interactions, incorrect administration, incorrect dosage, incorrect therapy selection,

75 masking of a severe disease, and the risk of drug dependence and abuse [14, 15]. Although the prevalence
76 and correlates of self-medication have been widely discussed in many countries [16], there is limited data
77 on the prevalence and factors associated with self-medication in underserved populations in informal
78 settlements. We estimated the prevalence, and factors associated with self-medication for COVID-19
79 amongst urban slum dwellers in Jinja City.

80 **Methods**

81

82 **Study design**

83 This was a cross-sectional study among adult urban slum dwellers aged 18 years and above who reported
84 having had confirmed or self-reported COVID-19 during the study period from January to December 2021.

85 **Study Area**

86 This study was carried out in Jinja city in Eastern Uganda, a region with an estimated population of 331,079
87 people [17]. Jinja City has eight slum but this study was conducted in five randomly selected slums. The
88 city also houses several educational institutions, commercial centres, industries, and factories, hence a
89 growing sprawl of unplanned settlements. This has given rise to several slum communities around the city,
90 which, among others, face a myriad of challenges, including poverty, poor health service delivery, and
91 illiteracy. Slum dwellers sought treatment from drug shops or stores, public hospitals [18], local pharmacy
92 [19], road-side herbalists [20] and traditional healers [20]. In Uganda, health care is provided in both private
93 and public health centres. Uganda provides primary health care through its National Minimum Health Care
94 Package at the lowest level health centres, while tertiary care is provided at the highest level health centres
95 [21]. The largest public health facility in Jinja district is Jinja Regional Referral Hospital, which provides
96 tertiary services, while several lower-level health facilities serve the majority of the population.

97

98

99

100 **Study population**

101 Participants who were non-residents, and adults not capable of giving consent for any reason (mentally ill,
102 critically sick, and those not able to respond), were excluded from the study as they did not represent the
103 true residents of the slums.

104 **Sample size determination**

105 The minimum sample size was estimated using the Kish Leslie sample size estimation formula for cross-
106 sectional studies [22]. In this study, we used a self-medication prevalence of 72% previously reported in
107 Namuwongo, a slum in Kampala [23] to estimate a sample size of 314. Through multistage sampling, the
108 estimated sample size was adjusted with a design effect of 1.5. Therefore, with a sample size of 518
109 respondents, this study had 80% power to determine the prevalence of self-medication at 95% confidence
110 interval.

111 **Sampling Procedures and participant recruitment**

112
113 A multi-stage sampling strategy was employed to select study respondents. Five slums were randomly
114 selected for this study out of eight slums in the study area, three slums were randomly selected in Southern
115 Division and two slums in Northern Division.

116 In the first stage, the researchers used a stratified sampling method in which five of Jinja's eight slums were
117 randomly selected and proportionate to their size. In the second stage, the researcher selected households
118 from the five selected slums. Later, a systematic sampling technique was further used, where every 41st
119 household (derived when we divided 21,284 total households by the 518-sample size calculated) was
120 selected to reach the desired number of respondents per slum. This was done with the guidance of the local
121 council leaders and/or Village Health Team (VHT) members. The local council or VHT members assisted
122 in guiding the researchers to locate the boundaries of each slum, provided a list of households in each slum,
123 and introduced the researchers to the community and the selected households. For the third stage, simple
124 random sampling was used to select a participant in each household where more than one person met the

125 inclusion criteria. The Kish Grid was used to select the required participant to avoid selection bias [24, 25].

126 Where there was no eligible person, the next household with an eligible respondent was selected. This was

127 continued until the desired number of respondents of 518 was reached.

128 **Data collection methods and tools**

129

130 A questionnaire was used to obtain quantitative data from the study respondents. The questionnaire used

131 was adapted from previous studies [26-28] and modified to fit our study. The tool was translated into the

132 local '*Lusoga*' by language expert and pre-tested in Polota slum, which is similar to the Jinja city slums.

133 The Cronbach's Alpha coefficient reliability test was used to assess the data collection tool's validity

134 (coefficient alpha=0.83). The questionnaire was then uploaded onto KoBo Toolbox tool version 2022.1.2,

135 a mobile phone application for data collection [29]. Data were collected by research assistants who were

136 conversant with both the local language and English. Study respondents received consent from all

137 respondents prior to the interviews. Data were collected using semi-structured questions installed on the

138 KoBo Toolbox application tool version 2022.1.2.

139 Participants were asked (Yes/No) if they used any medications (medicines and herbs) for self-medication

140 during COVID illness without a prescription or consultation with a medical practitioner/physician.

141 Participants were also interviewed on several demographic and social characteristics. The characteristics

142 were categorised as either predisposing, enabling, or need factors. Predisposing factors included: age, sex,

143 ethnicity, occupation, education level, cultural beliefs, and societal influences. Enabling factors refer to

144 availability constraints and included household income, medical insurance, proximity to medical services,

145 advertisements, sources of advice, and legislation. Need factors are those which trigger the action to self-

146 medicate, such as type of illness, duration of illness, the severity of illness, history of illness, and health

147 status.

148 **Data management and analysis**

149
150 The data collected were transferred into Microsoft Excel software and then STATA version 14.0 for further
151 statistical analysis. Data cleaning was done by checking for variables with missing data and duplicates as
152 well as transforming variables from string to numerical for analysis purposes to come up with a complete
153 dataset. Outliers were identified using geographical methods and a Q–Q plot (quantile-quantile plot). The
154 prevalence of self-medication for COVID-19 and symptoms suffered by slum dwellers was estimated.

155 **Statistical analysis**

156
157 The factors associated with self-medication for COVID-19 were determined using Poisson regression with
158 robust standard errors. We determined the crude prevalence ratios (cPR) and adjusted prevalence ratios
159 (APR) and 95% confidence intervals (CIs) for several covariates associated with self-medication for
160 COVID-19. The backward elimination method was used and variables with a P-value ≤ 0.20 were
161 considered at multivariable stage modelling stage. Only independent variables with a p-value ≤ 0.05 were
162 considered significant. All estimates were reported with a 95% confidence interval. Multi-collinearity was
163 checked using scatter plots and no variables had correlation coefficient greater than 0.4. The model with
164 the least Akaike information criterion (AIC) and Bayesian information criterion (BIC) values was selected
165 as the final model. The goodness of fit test was used to test the fitness of the final model.

166 **Ethical Considerations**

167
168 We obtained ethical approval to conduct this study from Makerere University, School of Public Health
169 Higher Degrees Research and Ethics Committee (HDREC), approval number FWA00011353.
170 Administrative clearance was also obtained from the Jinja city authorities. Written informed consent was
171 obtained from all respondents before the interview. Confidentiality was maintained throughout the study.

172 **Results**

173 **Characteristics of study participants.**

174 A total of 517 respondents were interviewed representing 99.80% response rate. The median age of
 175 respondents (IQR) was 31 (26-40) years and 40.42% (209/517) were within the age group of 30-39. In
 176 addition, 58.99% (305/517) were males. Among the respondents, 52.80% (273/517) had attained
 177 secondary-level education, 71.76% (371/517) were married while 50.68% (262/517) had a family monthly
 178 income of less than 100,000 Uganda shillings (approximately \leq 26.32 U.S. dollars), median income (IQR)
 179 being 100,000 (50,000-300,000) **Table 1.**

180 **Table 1: Socio-demographic information of the respondents**

Variable	Frequency (n=517)	Percentage (%)
Age (years)		
20-29	125	24.18
30-39	209	40.43
40-49	105	20.31
\geq 50	78	15.09
Median age (IQR) years	31 (26-40)	
Sex		
Male	305	59.99
Female	212	41.01
Religion		
Anglican	112	21.66
Catholic	135	26.11
Muslim	159	30.75
Pentecostal	90	17.41
Other	21	4.06
Education		
No formal	47	9.09
Primary	137	26.50
Secondary	273	52.80
Tertiary	60	11.61
Marital Status		
Never Married	97	18.76
Married	371	71.76
Separated	26	5.03
Widower/widow	23	4.45
Monthly Income (UGX)		
<100, 000	262	50.68
100,001-200,000	89	17.21
200,001-300,000	68	13.15
300,001-400,000	34	6.58
>400,000	64	12.38
Median income (IQR)	100,000 (50,000-300,000)	

181 **IQR:** Inter-quartile range

182 **Prevalence of self-medication for COVID-19**

183 Overall, the prevalence of self-medication for COVID-19 was 87.23% (451/517), 95% CI: [84-90] among
184 slum dwellers in Jinja. Of the 451 respondents who self-medicated for COVID-19, 56.76% (256/451)
185 agreed that they were aware of the risks associated with self-medication, while 15.74% (71/451) disagreed.
186 The most common symptoms suffered were difficulty in breathing, sore throat, loss of appetite, chest pain,
187 fever and other (dizziness, vomiting, bloating of the stomach and hearing loss) symptoms. **Fig 1 Symptoms**
188 **of COVID-19 among respondents who self-medicated**

189 **Factors associated with self-medication for COVID-19**

190 After controlling for potential confounders, being old (>50 years) [APR: 1.12, 95% CI:1.05, 1.20],
191 female sex [APR: 1.07, 95% CI: 1.02, 1.13], symptoms severity: minor symptoms [APR: 1.62, 95% CI:
192 1.25, 2.11]; severe symptoms [APR: 1.51, 95% CI: 1.16, 1.96], having medical insurance [APR: 0.63, 95%
193 CI: 0.46, 0.87], access to internet [APR: 1.13, 95% CI: 1.07, 1.20], family/friends influence [APR: 1.27,
194 95% CI: 1.12, 1.42], and anxiety/fear of being quarantined [APR: 1.35, 95% CI: 1.14, 1.60] were
195 significantly associated with higher rates of self-medication for COVID-19. Being aware of laws against
196 self-medication was also found to be associated with a lower risk of self-medication [APR: 0.89, 95% CI:
197 0.81, 0.97]. On the other hand, having had a prior COVID-19 test, previous experience with the disease,
198 health facility being far, long waiting time at the health facility, short duration of the disease or its
199 symptoms, poor present health status, highest education levels, marital status and household monthly
200 income were all found not statistically significant. **Table 2**

201

202

203

204

205 **Table 2: Bivariate and multivariable factors associated with self-medication among slum dwellers**
 206 **in Jinja City**

Variables	Have you ever self-medicated (n=517)		CPR (95% CI)*	APR (95%CI)**	p-value
	No n (%) n=66	Yes n (%) n=451			
Sex					
Male	37 (17.45)	175 (82.55)	Ref	Ref	
Female	29 (9.51)	276 (90.49)	1.07 (1.02, 1.13)	1.1 (1.0, 1.1)	0.006
Age (years)					
20-29	19(15.20)	106(84.80)	Ref	Ref	
30-39	29(13.88)	180(86.12)	1.02(0.93, 1.11)	1.03(0.97, 1.09)	0.384
40-49	13(12.38)	92(87.62)	1.03(0.93, 1.15)	1.05(0.98, 1.12)	0.133
50+	5(6.41)	73(93.59)	1.10(1.00, 1.21)	1.12(1.05, 1.20)	0.001
Anxiety and or fear of quarantine					
No	21(42.86)	28(57.14)	Ref	Ref	
Yes	9(2.82)	310(97.18)	1.70(1.33, 2.17)	1.35(1.14, 1.60)	<0.001
Have access to medical insurance					
No	46(9.52)	437(90.48)	Ref	Ref	
Yes	20(58.82)	14(41.18)	0.46(0.30, 0.68)	0.63(0.46, 0.87)	0.005
Internet/ advertisement					
No	47(21.17)	175(78.83)	Ref	Ref	
Yes	18(6.12)	276(93.88)	1.19(1.11, 1.28)	1.13(1.07, 1.20)	0.001
Influenced by my family or friends					
No	48(46.15)	56(53.85)	Ref	Ref	
Yes	18(4.36)	395(95.64)	1.78(1.48, 2.13)	1.27(1.12, 1.42)	<0.001
Aware of any law on medicine use					
No	39(9.11)	389(90.89)	Ref	Ref	
Yes	27(30.34)	62(69.66)	0.77(0.67, 0.88)	0.89(0.81, 0.97)	0.008
Symptom severity					
No symptoms	39 (67.24)	19 (32.76)	Ref	Ref	
Minor symptoms	11(5.14)	203(94.86)	2.89(2.00, 4.19)	1.62(1.25, 2.11)	<0.001
Severe symptoms	16(6.53)	229(93.47)	2.85(1.97, 4.13)	1.51(1.16, 1.96)	0.002
Prior COVID-19 test					
No	22(7.12)	287(92.88)	Ref	Ref	
Yes	18(11.69)	136(88.31)	0.95(0.89,1.01)	1.01(0.96,1.06)	0.736
Previous experience with the disease					
No	49(11.53)	376(88.47)	Ref	Ref	
Yes	17(18.48)	75(81.52)	0.92(0.83, 1.02)	0.96(0.89,1.03)	0.236
The health facility is far					
No	49(11.64)	372(88.36)	Ref	Ref	
Yes	17(17.71)	79(82.29)	0.93(0.84, 1.03)	0.93(0.85,1.01)	0.091
Long waiting time at the health facility					
No	45(14.61)	263(85.39)	Ref	Ref	
Yes	21(10.05)	188(89.95)	1.05(0.99, 1.12)	0.01(0.96,1.06)	0.679
The short duration of the disease or its symptoms/signs					
No	51(14.37)	304(85.63)	Ref	Ref	
Yes	15(9.26)	147(90.74)	1.06(0.99, 1.13)	1.02(0.97, 1.07)	0.375

Poor present health status					
No	41(17.23)	197(82.77)	Ref	Ref	
Yes	25(8.96)	254(91.04)	1.10(1.03, 1.18)	0.97(0.92, 1.02)	0.214
Education					
No education	5(10.64)	42(89.36)	Ref	Ref	
Primary	15(10.95)	122(89.05)	1.00(0.89, 1.12)	0.98(0.89, 1.08)	0.733
Secondary	30(10.99)	243(89.01)	1.00(0.89, 1.12)	1.03(0.93, 1.14)	0.548
Tertiary	16(26.67)	44(73.33)	0.82(0.68, 0.98)	1.00(0.87, 1.14)	0.992
Marital Status					
Never Married	20(20.62)	77(79.38)	Ref	Ref	
Married	41(11.08)	330(88.92)	0.89(0.81, 0.99)	1.00(0.93, 1.08)	0.987
Separated	2(7.69)	24(92.31)	1.04(0.92, 1.17)	0.97(0.85, 1.11)	0.626
Widower/widow	3(13.04)	20(87.98)	0.98(0.83, 1.15)	1.03(0.92, 1.15)	0.605
Monthly income (UGX)[¶]					
0-100,000	42(16.03)	220(83.97)	Ref	Ref	
100001-200,000	8(8.99)	81(91.01)	1.08(0.99, 1.18)	1.04(0.97, 1.11)	0.245
200,001-300,000	8(11.76)	60(88.24)	1.05(0.95, 1.16)	0.99(0.93, 1.05)	0.700
300,001-400,000	3(8.82)	31(91.17)	1.09(0.97, 1.22)	0.97(0.88, 1.06)	0.504
400001-max	5(7.81)	59(92.19)	1.10(1.00, 1.20)	0.99(0.92, 1.06)	0.781

207 *Note: CPR= *Crude Prevalence Ratio, APR= **Adjusted Prevalence Ratio. [¶]1USD=3800 Uganda shillings*
 208 *(UGX)*

209 Discussion

210 Our findings show that there is a high prevalence of COVID-19 self-medication among urban slum
 211 dwellers. This was demonstrated by the fact that 87.2% of urban slum dwellers in Jinja city reported using
 212 self-medication. The high prevalence of self-medication may be a result of the challenges related to
 213 healthcare access brought about by the pandemic (high health costs, travel restrictions, and a shortage of
 214 health staff). Studies conducted in Bangladesh and Uganda found almost a similar prevalence to this current
 215 study of 88.3% [30] and 88% [31] respectively. Further, a study conducted among the general public found
 216 a decrease in self-medication practices during the COVID-19 pandemic lockdown compared to the practice
 217 before the lockdown [31]. This disparity could be explained by the fact that during the COVID-19
 218 lockdown, the public was not moving freely, making it difficult for them to obtain drugs. The prevalence
 219 of self-medication in our study, however, was way higher than that reported among residents of urban slums
 220 [32]. This discrepancy can be explained by the fact that both of these studies were conducted way before
 221 the COVID-19 pandemic which started almost three years later.

222 Factors found to be associated with self-medication for COVID-19 included age, sex, anxiety and or fear
223 of quarantine, access to medical insurance, internet/advertisement, awareness of the laws regarding
224 medicine use, family/ friend influence, and symptom severity. Our study revealed that respondents who
225 were 50 years and older showed a 12% increase in the likelihood of self-medication compared to those in
226 the 20-29 age bracket. This could be explained by the higher COVID-19 mortality among the elderly, hence
227 creating a higher perceived risk among this population. In addition, older people are prone to chronic
228 infections, thus being susceptible to infections like COVID-19 [33]. In addition, studies conducted in
229 Uganda, Eritrea, Ethiopia, Nigeria, Belgrade and Cameroon have reported that as one gets older, they
230 acquire accumulated knowledge due to disease experience [33-40]. Contrary studies have been done in
231 Jabalpur [32] and Syrian [41], where individuals aged 18 to 40 years were more likely to self-medicate.
232 This age difference could be due to the population dynamics of the respondents.

233 Our study showed that females were two times more likely to self-medicate than males. The reason for the
234 association between being a female and self-medication is not known, but in the context of the COVID-19
235 outbreak, greater anxiety among women, as described in Iran and Italy, cannot be excluded [42, 43]. In
236 addition, this might have been due to the exacerbated stress among women associated with the COVID-19
237 pandemic and efforts to mitigate its spread [44]. This is consistent with results in studies conducted in Spain
238 and elsewhere that found that more females self-medicated compared to their counterparts [45, 46].
239 However, there are controversial data on the relationship between sex and self-medication as studies
240 conducted in Iran and the United Arab Emirates [47, 48] respectively did not report any association. This
241 difference could be due to these studies having been conducted among different study populations.

242 In the current study, respondents who reported anxiety and fear of being quarantined had a 35% increase in
243 the likelihood of self-medication compared to their counterparts. It is important to note that during the
244 period under study, COVID-19-related morbidity and mortality were very high in Uganda and could have
245 increased anxiety levels and fear of infection. Anxiety could have exacerbated self-medication as people
246 tried their level best to prevent the infection, treat any present signs and symptoms, as well as boost

247 immunity to fight off disease. In addition, the mass fear of COVID-19, termed “corona phobia” [49-51]
248 coupled with stringent infection prevention and control measures, was reported to have caused a lot of
249 depression, confusion, fear, anger, and despair among people. This consequently resulted in substance use,
250 which is self-medication [49]. Similar findings have been reported by other scholars [52-54].

251 This study found that having access to medical insurance was more protective compared to those who did
252 not have medical insurance. At the peak of the second wave of COVID-19 in Uganda, the public healthcare
253 system could not ably serve all confirmed and suspected COVID-19 cases effectively, since treatment was
254 expensive in private facilities, people who had insurance accessed treatment easily in these private facilities,
255 and those offered better services [55, 56]. The majority of those who accessed private facilities were likely
256 to have medical insurance, as supported by findings of a study conducted in Iran [57].

257 The findings of this study show that individuals who reported access to the internet and advertisements
258 showed a 13% increase in the likelihood of self-medication compared with those who did not. Comparable
259 studies have been conducted in Iran and India [58, 59]. There was infodemia about the COVID-19
260 pandemic due to much information in both digital and physical environments. This resulted in confusion
261 and risk-taking behaviours that could harm health, foster distrust in health authorities or systems, and
262 undermine the public health response, leading to self-medication [60, 61]. This information was used by
263 the public as a replacement for medical consultations [59, 62]. In this response, the WHO designed a
264 taxonomy to address infodemia, which was piloted in individual countries and WHO regions to generate
265 localized insights and actions [60]. Furthermore, these findings indicated that the likelihood of self-
266 medication for COVID-19 increased by 27% due to influence from family and friends. This finding reflects
267 the familial clustering of the disease by the respondents since some reported self-medication for the
268 prevention of COVID-19 infection. A possible reason is that family and friends tend to live together and
269 often communicate with each other. Therefore, individuals who tested positive for COVID-19 might have
270 taken the initiative to inform their recent acquaintances, who, under the scare of acquiring infection,
271 resorted to self-medication in the hope of prevention [62-66].

272 Respondents who experienced minor and severe symptoms had a higher likelihood of 62% and 51%,
273 respectively, of self-medicating compared to those who had no symptoms. These results mean that there
274 was a high likelihood of self-medication among slum dwellers who either experienced minor or severe
275 symptoms. Individuals with minor symptoms ignored the disease and thought that since it was not severe,
276 they would easily recover in a short time. This is supported by studies conducted in Nigeria, and Ethiopia
277 and a systematic review [36, 38, 67] respectively. On the other hand, those with severe symptoms viewed
278 themselves as having the severe form of the disease that requires immediate attention and cannot be delayed
279 due to issues of health facility access, affordability, drug stock-outs, and long waiting times. This is
280 comparable to studies conducted in Brazil [68] and the USA [69]. Respondents who knew laws regarding
281 medicine use were more likely to self-medicate, which is in line with a study conducted in Colombia [70].
282 During the COVID-19 era, the greater access of people to the media, the internet, and their ability to
283 understand information about the medicine and its related laws on use found on social networks may explain
284 this finding. Since many patients get knowledge about drugs from previous prescriptions [71], physicians
285 should limit superfluous prescriptions of antibiotics and implement guideline-based practices.

286 **Limitations of the study**

287 The study had potential limitation. First, this study based on self-reporting, hence more likely to lead to
288 social desirability bias because of stigma, anxiety, and fear. However, this was minimized by assuring the
289 respondents before the study about information confidentiality, reviewing respondents' medical records
290 while conducting data collection, and checking the Jinja City health database on COVID-19 test results.
291 This study was conducted during the COVID-19 pandemic when people had fear and stigma about freely
292 interacting with anyone for fear of contracting the disease. However, this was addressed by ensuring all the
293 research assistants observed COVID-19 SOPs while in the field. These findings can be used to plan and
294 manage future epidemics or any emergencies of public health concern and also guide the engagement of
295 slum dwellers towards a reduction in the prevalence of self-medication. Therefore, these results could be
296 generalizable to the general population of urban slum dwellers.

297 **Conclusion**

298 This study reports a very high prevalence of self-medication for COVID-19 among urban slum dwellers in
299 Jinja City. This is a signal for potential future antimicrobial resistance outbreaks, especially in vulnerable
300 groups like urban slum dwellers. Self-medication has been known to result in unintended healthcare
301 outcomes such as antimicrobial resistance, adverse effects, drug interactions, and drug dependence, among
302 others. According to WHO, the COVID-19 pandemic situation is likely to be prolonged for years and will
303 have a huge effect on the social-economic and psychosocial impact on people's lifestyles and behaviour.
304 This poses a more serious threat that might push people to self-medicate. Therefore, MoH and NDA should
305 design stringent measures for the prescription, sale, and use of medicine, not forgetting to regulate
306 advertisements in addition to sensitization of the slum dwellers.

307 **Recommendations**

308 Based on the findings, the researchers suggest the following recommendations to the Ugandan government
309 and other stakeholders. The Ministry of Health needs to adopt WHO-approved guidance on the detection
310 of info-demic signals and implement medical insurance policy to benefit the most vulnerable populations
311 like slum dwellers. The National Drugs Authority (NDA) should put in place stringent mechanisms to
312 regulate the prescription, sale, hawking, and use of conventional medicines, especially antibiotics. Also,
313 NDA should liaise with herbal medicine practitioners to establish active elements in socially acceptable
314 and effective herbal medicines. The City Health Team (CHT) should control and manage the sale of any
315 medicine by conducting routine inspections, routine support supervision of all sellers of medicines and
316 mass sensitisation of all stakeholders. During health crises, such as the COVID-19 pandemic, slum dwellers
317 and the general public ought to practice good health-seeking behaviours.

318

319

References

1. WHO, *World Health Organization situation report (Corona virus disease)*. 2023.
2. WHO. *WHO recommends two new drugs to treat COVID-19, 14 January 2022 Statement, Geneva, Accessed 5th March 2022*. 2022.
3. Elayeete, S., et al., *Comparative epidemiologic analysis of COVID-19 patients during the first and second waves of COVID-19 in Uganda*. IJID Regions, 2022. **3**: p. 160-167.
4. Bongomin, F., et al. *High mortality during the second wave of the coronavirus disease 2019 (COVID-19) pandemic in Uganda: experience from a national referral COVID-19 treatment unit*. in *Open Forum Infectious Diseases*. 2021. Oxford University Press US.
5. MOH, *Corona virus pandemic, Covid-19 [Online]*. Available: <https://www.health.go.ug/covid/> [Accessed 07/03 2023]. 2023.
6. Kaye, A.D., et al., *Economic impact of COVID-19 pandemic on healthcare facilities and systems: International perspectives*. Best Practice & Research. Clinical Anaesthesiology, 2020.
7. Liu, F., M. Wang, and M. Zheng, *Effects of COVID-19 lockdown on global air quality and health*. Science of the Total Environment, 2021. **755**: p. 142533.
8. Chu, I.Y.-H., et al., *Social consequences of mass quarantine during epidemics: a systematic review with implications for the COVID-19 response*. Journal of travel medicine, 2020. **27**(7): p. taaa192.
9. Sindhuja, P., et al., *COVID-19 AND HEALTH CARE SERVICES IN RURAL INDIA: CHALLENGES AND CONSEQUENCES*. Turkish Online Journal of Qualitative Inquiry, 2021. **12**(4).
10. WHO, *World Health Organization. (2000). Guidelines for the regulatory assessment of medicinal products for use in self-medication*. World Health Organization. 2000.
11. WHO, *Guidelines for the regulatory assessment of medicinal products for use in self-medication*. 2000, World Health Organization.
12. Eticha, T. and K. Mesfin, *Self-medication practices in Mekelle, Ethiopia*. PloS one, 2014. **9**(5): p. e97464.
13. Bashige CV, M.M., Bakari AS, Sangwa KG, Kahumba BJ, Pierre D, Lumbu SJB, *Prévalence et caractéristiques de l'automédication chez les étudiants de 18 à 35 ans résidant au campus de la Kasapa de l'Université de Lubumbashi*. Pan Afr Med, 21, 77. . 2015. **12**(2): p. 007-020.
14. Chouhan, K. and S.B. Prasad, *Self-medication and their consequences: a challenge to health professional*. Asian J Pharm Clin Res, 2016. **9**(2): p. 314-317.
15. De Sanctis, V., et al., *Prevalence, attitude and practice of self-medication among adolescents and the paradigm of dysmenorrhea self-care management in different countries*. Acta Bio Medica: Atenei Parmensis, 2020. **91**(1): p. 182.
16. Selvaraj, K.K., S.G.; Ramalingam, A. , *Prevalence of self-medication practices and its associated factors in Urban Puducherry, India*. Perspect. Clin. Res. 2014, 5, 32–36. 2014.
17. UBOS, *Statistical Abstract: Uganda Bureau of statistics 2020*.
18. Adams, A.M., et al., *Healthcare seeking for chronic illness among adult slum dwellers in Bangladesh: A descriptive cross-sectional study in two urban settings*. PloS one, 2020. **15**(6): p. e0233635.
19. van der Heijden, J., et al., *'Working to stay healthy', health-seeking behaviour in Bangladesh's urban slums: a qualitative study*. BMC Public Health, 2019. **19**(1): p. 1-13.
20. Bawah, A.T., et al., *Prevalence of diabetes among homeless and slum dwellers in Accra, Ghana: a survey study*. BMC research notes, 2019. **12**(1): p. 1-5.
21. Mijumbi-Deve R, K.I., Nankya E, Sewankambo N. , *Primary health care systems (PRIMASYS): case study from Uganda*. World Health Organization; 2017 [cited 2023 March 12]. Available from: <https://www.who.int/alliance-hpsr/projects/primasys/en/>. 2021.
22. Kish, L., *Survey sampling*. 1965.

23. Oniba, O.R., *Factors Influencing Self-medication Practices Among Adults in Urban Slum Households*. 2012, International Health Sciences University.
24. Dreier, M., et al., *Development and evaluation of e-mental health interventions to reduce stigmatization of suicidality—a study protocol*. BMC psychiatry, 2019. **19**(1): p. 1-12.
25. Ahmed, S.H., et al., *The prevalence of selected risk factors for non-communicable diseases in Hargeisa, Somaliland: a cross-sectional study*. BMC Public Health, 2019. **19**: p. 1-10.
26. Quispe-Cañari, J.F., et al., *Self-medication practices during the COVID-19 pandemic among the adult population in Peru: A cross-sectional survey*. Saudi pharmaceutical journal, 2021. **29**(1): p. 1-11.
27. Fernando, A., et al., *A descriptive study of self-medication practices among Sri Lankan national level athletes*. BMC research notes, 2017. **10**: p. 1-12.
28. Ngigi, C.K., P. Mwangi, and J. Wala, *Self-medication with antibiotics prior to seeking treatment among adult patients attending outpatient department at gatundu sub-county hospital, Kiambu County, Kenya*. Imp J Interdiscip Res, 2016. **2**.
29. Lakshminarasimhappa, M., *Web-based and smart mobile app for data collection: Kobo Toolbox/Kobo collect*. Journal of Indian Library Association, 2022. **57**(2): p. 72-79.
30. Nasir, M., A. Chowdhury, and T. Zahan, *Self-medication during COVID-19 outbreak: a cross sectional online survey in Dhaka city*. Int J Basic Clin Pharmacol, 2020. **9**(9): p. 1325-30.
31. Dare, S.S., et al., *COVID-19 pandemic and behavioural response to self-medication practice in Western Uganda*. medRxiv, 2021.
32. Jain, S., et al., *Prevalence of self-medication practices among the residents of urban slums located near govt. medical college, Jabalpur*. Int J Community Med Public Health, 2018. **5**: p. 811-7.
33. Lukovic JA, M.V., Pekmezovic T, Trajkovic G, Ratkovic N, Aleksic D, Grgurevic A. , *Self-medication practices and risk factors for self-medication among medical students in Belgrade, Serbia*. PLoS One. 2014 Dec 11;9(12):e114644. doi: 10.1371/journal.pone.0114644. PMID: 25503967; PMCID: PMC4263675. 2014.
34. Niwandinda, F., et al., *Patterns and practices of self-medication among students enrolled at Mbarara University of Science and Technology in Uganda*. Integrated pharmacy research & practice, 2020. **9**: p. 41.
35. Araia, Z.Z., N.K. Gebregziabher, and A.B. Mesfun, *Self medication practice and associated factors among students of Asmara College of Health Sciences, Eritrea: a cross sectional study*. Journal of Pharmaceutical Policy and Practice, 2019. **12**(1): p. 3.
36. Kifle, Z.D., et al., *Self-medication practice and associated factors among private health sciences students in Gondar Town, North West Ethiopia. A Cross-Sectional Study*. INQUIRY: The Journal of Health Care Organization, Provision, and Financing, 2021. **58**: p. 00469580211005188.
37. Ocan, M., et al., *Patterns and predictors of self-medication in northern Uganda*. PloS one, 2014. **9**(3): p. e92323.
38. Oche, O.M., et al., *Prevalence and factors associated with self medication among people living in urban slums of Sokoto Metropolis, Sokoto State, Nigeria*. Cent African J Public Heal, 2019. **5**(6): p. 302-309.
39. Jerez-Roig, J., et al., *Prevalence of self-medication and associated factors in an elderly population: a systematic review*. Drugs & aging, 2014. **31**(12): p. 883-896.
40. Kong, J.D., et al., *SARS-CoV-2 and self-medication in Cameroon: a mathematical model*. Journal of biological dynamics, 2021. **15**(1): p. 137-150.
41. Abdelwahed, R.N., M. Jassem, and A. Alyousbashi, *Self-medication practices, prevalence and associated factors among Syrian adult patients: a cross-sectional study*. 2022.
42. Moghanibashi-Mansourieh, A., *Assessing the anxiety level of Iranian general population during COVID-19 outbreak*. Asian journal of psychiatry, 2020. **51**: p. 102076.
43. Casagrande M, F.F., Tambelli R, Forte G. , *The enemy who sealed the world: Effects quarantine due to the COVID-19 on sleep quality, anxiety, and psychological distress in the Italian population*. Sleep Med. 2020. 2020.

44. Bochicchio, L.A., et al., *Understanding alcohol and marijuana use among sexual minority women during the COVID-19 pandemic: A descriptive phenomenological study*. Journal of Homosexuality, 2021. **68**(4): p. 631-646.
45. Behzadifar, M., et al., *Prevalence of self-medication in university students: systematic review and meta-analysis*. East Mediterr Health J, 2020. **26**(7): p. 846-57.
46. Sadio, A.J., et al., *Assessment of self-medication practices in the context of the COVID-19 outbreak in Togo*. BMC public health, 2021. **21**(1): p. 1-9.
47. Abdi, A., et al., *Prevalence of self-medication practice among health sciences students in Kermanshah, Iran*. BMC pharmacology and Toxicology, 2018. **19**(1): p. 1-7.
48. Sridhar, S.B., et al., *Assessment of nature, reasons, and consequences of self-medication practice among general population of Ras Al-Khaimah, UAE*. International Journal of Applied and Basic Medical Research, 2018. **8**(1): p. 3.
49. Sarkar, A. and J.K. Rajamani, *Assessment of Psychology, Behaviour and Self-Medication Potential Among Indian People During COVID-19 Pandemic*. Biomedical and Pharmacology Journal, 2021. **14**(4): p. 2285-2295.
50. Amin, S., *The psychology of coronavirus fear: Are healthcare professionals suffering from coronavirus phobia?* International Journal of Healthcare Management, 2020. **13**(3): p. 249-256.
51. Heiat, M., et al., *Phobia and Fear of COVID-19: origins, complications and management, a narrative review*. Annali Di Igiene Medicina Preventiva E Di Comunita, 2021. **33**(4): p. 360-370.
52. Wegbom, A.I., et al., *Self-Medication Practices and Associated Factors in the Prevention and/or Treatment of COVID-19 Virus: A Population-Based Survey in Nigeria*. Frontiers in Public Health, 2021. **9**.
53. Chopra, D., et al., *Prevalence of self-reported anxiety and self-medication among upper and middle socioeconomic strata amidst COVID-19 pandemic*. Journal of education and health promotion, 2021. **10**: p. 73-73.
54. Mbazzi, F.B., et al., *The impact of COVID-19 measures on children with disabilities and their families in Uganda*. Disability & Society, 2020: p. 1-24.
55. Davalbhakta, S., et al., *Private health sector in India-ready and willing, yet underutilized in the covid-19 pandemic: a cross-sectional study*. Frontiers in Public Health, 2020. **8**: p. 571419.
56. Ilesanmi, O.S. and A.A. Afolabi, *Positioning the private health facilities towards continuous service provision during the COVID-19 pandemic*. Hospital Practices and Research, 2020. **5**(4): p. 122-125.
57. Karimy, M., et al., *Risk factors associated with self-medication among women in Iran*. BMC Public Health, 2019. **19**(1): p. 1033.
58. Soroush, A., et al., *Exploring the perceived factors that affect self-medication among nursing students: a qualitative study*. BMC nursing, 2018. **17**(1): p. 1-7.
59. Agarwal, T., et al., *Use of internet for practice of self-medication: We are heading toward an era of internet pharmacy*. Medical Journal of Dr. DY Patil Vidyapeeth, 2021. **14**(1): p. 36.
60. Purnat, T.D., et al., *Infodemic signal detection during the COVID-19 pandemic: development of a methodology for identifying potential information voids in online conversations*. JMIR infodemiology, 2021. **1**(1): p. e30971.
61. Calleja, N., et al., *A public health research agenda for managing infodemics: methods and results of the first WHO infodemiology conference*. JMIR infodemiology, 2021. **1**(1): p. e30979.
62. Malik, M., et al., *Self-medication during Covid-19 pandemic: challenges and opportunities*. Drugs & Therapy Perspectives, 2020. **36**(12): p. 565-567.
63. Lei, X., et al., *Self-medication practice and associated factors among residents in Wuhan, China*. International journal of environmental research and public health, 2018. **15**(1): p. 68.
64. Kassie, A.D., B.B. Bifftu, and H.S. Mekonnen, *Self-medication practice and associated factors among adult household members in Meket district, Northeast Ethiopia, 2017*. BMC Pharmacology and Toxicology, 2018. **19**(1): p. 1-8.

65. Zhang, A., et al., *Self-medication with antibiotics for protection against COVID-19: the role of psychological distress, knowledge of, and experiences with antibiotics*. *Antibiotics*, 2021. **10**(3): p. 232.
66. Goodwin, R., et al., *Anxiety, perceived control and pandemic behaviour in Thailand during COVID-19: Results from a national survey*. *Journal of Psychiatric Research*, 2021. **135**: p. 212-217.
67. Quincho-Lopez, A., et al., *Self-medication practices to prevent or manage COVID-19: A systematic review*. *PloS one*, 2021. **16**(11): p. e0259317.
68. Pastore, G.P., et al., *Self-medication among myofascial pain patients: a preliminary study*. *The Open Dentistry Journal*, 2018. **12**: p. 347.
69. Nishith, P., et al., *A test of self-medication hypothesis for drug use in homeless persons: the role of severe mental illness*. *Journal of Social Distress and Homelessness*, 2022: p. 1-8.
70. Machado-Alba, J.E., et al., *Social, cultural and economic factors associated with self-medication*. *Biomédica*, 2014. **34**(4): p. 580-588.
71. Nepal, G. and S. Bhatta, *Self-medication with antibiotics in WHO Southeast Asian Region: a systematic review*. *Cureus*, 2018. **10**(4).