

Investigating potential myopia risk factors, including chronotype, in Estonian adolescents

Teele Palumaa^{1,2,3}, Delis Linntam¹, Reili Rebane¹, Marika Tammaru⁴, Kadi Palumaa¹

¹ Eye Clinic, East Tallinn Central Hospital, Tallinn, Estonia

² Institute of Genomics, University of Tartu, Tartu, Estonia

³ Department of Ophthalmology, Emory University, Atlanta, USA

⁴ Research Department, East Tallinn Central Hospital, Tallinn, Estonia

For correspondence:

Teele Palumaa: teele.palumaa@ut.ee; postal address: Ravi 18, 10138 Tallinn, Estonia

Abstract

Purpose: To evaluate chronotype, lifestyle factors, and parental myopia in relation to myopia, and characterise the effect of cycloplegia on spherical equivalent refraction (SER) in Estonian secondary school students.

Methods: One hundred twenty-three students aged between 15 and 17 years from three secondary schools in Estonia participated in the study. They underwent a comprehensive ocular examination, including cycloplegic refraction and ocular biometry. Chronotype was evaluated with the Morningness–Eveningness Questionnaire. Participants also completed a questionnaire about their daily activities, including time spent outdoors, time spent on near-work and mid-working distance activities, and parental myopia. Myopia was defined as cycloplegic SER ≤ -0.50 D. Logistic regression analysis was performed to assess the association of the studied factors with myopia.

Results: In a multivariate regression model, having two myopic parents was associated with higher odds of myopia (OR 3.78, 95% CI 1.15–12.42). We found no association between myopia and chronotype. Notably, time spent outdoors, and doing near-work and mid-working distance activities, did not affect the likelihood of having myopia. Non-cycloplegic SER was significantly more myopic than cycloplegic SER on average by 0.86 D ($p < 0.0001$, Wilcoxon matched-pairs signed rank test).

Conclusion: Consistent with previous reports, we identified parental myopia as a myopia risk factor. Chronotype was not associated with myopia in our study sample. Interestingly, there was no association between myopia and time spent outdoors or near work. Using non-

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

cycloplegic refraction would lead to a significant overestimation of myopia.

Keywords

myopia, risk factors, chronotype, circadian rhythms, cycloplegia

Introduction

The prevalence of myopia, or short-sightedness, has increased markedly over the last decades [1]. Should the current trend continue, approximately half of the global population will be myopic by 2050 [1]. Myopia risk factors have been studied extensively worldwide. It has emerged that the risk of developing myopia is reduced in children who spend more time outdoors [2–5] and increased with more near-work [6], and computer use [7].

Several recent studies have raised the possibility that myopia is associated with circadian rhythms. Genes associated with refractive errors in genome-wide studies have an increased representation of genes controlling circadian rhythms [8]. It is also known that axial length and choroidal thickness show diurnal rhythmicity in humans [9, 10] and chicks [11, 12]. Furthermore, inducing myopia in chicks with form deprivation abolishes this circadian rhythmicity in axial length [11].

There is also evidence to suggest that short-sightedness may be linked to sleep. For example, myopic children have been reported to sleep later than children with no myopia [13–16], and to have worse sleep quality [17, 13]. However, the data in the literature is inconsistent and several studies have found no associations between myopia and sleep quality [18] and bedtime [19, 20]. Interestingly, though, poor sleep quality is associated with evening chronotype, or one's late sleep-wake timing preference [21–23], raising the possibility that late chronotype may be the link between myopia and poor sleep quality. Chronotype has been studied in the context of myopia in a couple of studies, and no definitive associations have been found. In one report, the circadian phases of myopes and non-myopes did not differ [24] and in another, self-report of a "morning" or "evening" type was not associated with myopia progression [15].

Some discrepancies between the findings of the studies listed above might be related to whether myopia was self-reported [16], defined with cycloplegic [14, 19, 18, 20, 15] or non-cycloplegic [13, 24] refraction measurements. It is known that refraction measurements are more myopic without cycloplegia [25, 26], which may lead to overestimating the number of myopes in studies using non-cycloplegic refraction to determine myopia.

In this study, we assessed the effect of chronotype, parental myopia status, and lifestyle factors on myopia in Estonian adolescents. We also analysed the difference between

cycloplegic and non-cycloplegic spherical equivalent refraction (SER) measurements to estimate its influence on assessing myopia prevalence in our study population.

Methods

The study was conducted at the East Tallinn Central Hospital Eye Clinic in 2019. It adhered to the principles of the Declaration of Helsinki and was approved by the Tallinn Medical Research Ethics Committee. Form 10 students from three public schools in Tallinn were invited to participate, and a total of 123 students were recruited. Written informed consent was obtained from the participants and their parents or legal guardians. All participants were European Caucasians. The schools differed slightly in their education profile, putting more emphasis on science, arts, or sports.

Participants underwent a comprehensive ocular examination, including non-cycloplegic and cycloplegic refraction, ocular biometry and biomicroscopy. Cycloplegia was achieved by applying two rounds of 1% cyclopentolate and 0.5% tropicamide with a 5-minute interval, as per [3]. Cycloplegia was evaluated 40 minutes after the application of the first round and considered adequate when the pupil was ≥ 6 mm in diameter and not responsive to light. Ocular refraction was measured with HandyRef-K autorefractor (Nidek, Tokyo, Japan), axial length and anterior corneal curvature with IOLMaster 500 (Zeiss, Oberkochen, Germany). SER was calculated as sphere + $\frac{1}{2}$ cylinder, and participants' SER was defined as the mean SER of the left and right eyes. Myopia was defined as cycloplegic SER ≤ -0.50 dioptres (D), high myopia as SER ≤ -6.00 D, astigmatism as ≤ -1.00 dioptre cylinder and hyperopia as SER ≥ 2.00 D.

Participants filled out a questionnaire about daily activities, including time spent outdoors, doing near work and mid-working distance activities, separately for weekdays and free days. Total time spent on each type of activity was calculated per week. Near-work activities were defined as reading, writing, and using a smartphone. Mid-working distance activities were defined as watching TV, working with a computer, and playing video games. The weekly total number of hours spent outdoors, doing near work and mid-working distance activities were calculated and used in the analyses. Chronotype was evaluated with the Morningness-Eveningness Questionnaire (MEQ) [27]. MEQ was translated from English into Estonian and pilot-tested on a separate cohort of thirty 16-17-year-olds to eliminate any misunderstandings in the wording. The questionnaire also contained questions about parental myopia.

Statistical analyses were performed using Stata Software for Statistics and Data Science (version 14.2). Graphs were produced in Prism (version 9, GraphPad). Continuous

variables were expressed as mean or median when appropriate, and qualitative variables were summarised as frequency and percentage. Wilcoxon signed-rank test was used to compare the SER before and after cycloplegia and Spearman rho was calculated to characterise correlations between ocular parameters. Variables associated with myopia were assessed with multivariate logistic regression analysis, and results presented as odds ratios (OR) with their 95% confidence interval (CI). A two-tailed p value of <0.05 was considered statistically significant.

Results

A total of 123 students participated in the study (57% female), and the majority were 16 years old (71%). 18% reported having two myopic parents, and 41% had one myopic parent. Half of the participants were from the science-oriented school, 28% from the arts-oriented and 22% from the sports-oriented school (see Table 1 for a demographic characterisation). The prevalence of myopia in our study population was 30.9%, high myopia was present in 2.4%, hyperopia in 3.3% and astigmatism in 11.4%.

The average time spent outdoors per week was 16.5 hours for both non-myopes and myopes (Fig. 1A). Myopes spent slightly more time on near-work activities, 54.4 hours, compared to 52.0 hours per week in non-myopes (Fig. 1B). The average time spent on mid-working distance activities was 21.3 hours for non-myopes and 17.7 hours for myopes (Fig. 1C). In MEQ, a larger score indicates an earlier chronotype [27]. In our study sample, the MEQ score was 46.7 for non-myopes and 48.8 for myopes (Fig. 1D).

Multivariate logistic regression analyses of the factors studied (Table 2) revealed that participants with two myopic parents were 4.3 times more likely to have myopia (95% CI 1.19 – 15.74). Subjects attending the sports-oriented school had lower odds of myopia than participants attending the science-oriented school (OR 0.12, 95% CI 0.03 – 0.51). We found no significant association between chronotype and refractive status. Time spent outdoors, on near work and mid-working distance activities were not associated with myopia in our study.

The SER values displayed a left-skewed distribution in our study population with the median cycloplegic SER +0.25 D and mean SER –0.38 D (Fig. 2A). Myopia exhibited both an axial and corneal component as more myopic refraction was correlated with both longer axial length ($r = -0.76$, Fig. 2B), and smaller radius of corneal curvature, or steeper corneas ($r = 0.17$, Fig. 2C).

The SER with and without cycloplegia were significantly different, non-cycloplegic SER was, on average, 0.89 D more myopic than cycloplegic SER (mean non-cycloplegic SER – 1.27 D, mean cycloplegic SER –0.38 D; median non-cycloplegic SER –0.88 D, median

cycloplegic SER +0.25 D, $p < 0.0001$, Wilcoxon matched-pairs signed rank test, Fig. 3A). Non-cycloplegic SER was ≤ -0.50 D in 83 participants (67% of the study population) while cycloplegic SER was ≤ -0.50 D in 38 participants (31% of the study population). Therefore, had we defined myopia based on non-cycloplegic SER, we would have overestimated the prevalence of myopia by more than two times. The non-cycloplegic and cycloplegic SER were highly correlated ($r = 0.74$, Spearman correlation, Fig. 3B).

Discussion

This is the first study to investigate myopia risk factors in Estonia. Chronotype, assessed with MEQ, was not associated with myopia in our sample. While later bedtime has been associated with myopia [13–16], a link between chronotype and myopia has not been established. In their study, Flanagan et al. [24] found that myopes had higher melatonin levels at all time points studied, while the circadian parameters, i.e. dim-light melatonin onset, phase of circadian activity determined with actigraphy, and MEQ score did not differ between myopes and non-myopes. It should be mentioned that non-cycloplegic refraction was used to determine myopia [24]. Furthermore, self-reported “morning” or “evening” type was also not associated with myopia progression in a study by Lee et al. [15]. Larger studies are needed to further evaluate the potential link between human circadian rhythms and refractive errors.

In agreement with other investigations [28], we determined parental myopia as a myopia risk factor. While increased time spent outdoors has been convincingly shown to protect from myopia development in multiple studies [2–5], we did not see such a correlation in our study population. The reason behind this observation may be our limited sample size and the use of a questionnaire to measure time spent outdoors. It has been demonstrated that a daily activities diary correlates better with time spent outdoors evaluated by wearable light meters than a questionnaire [29]. However, this is not the first study to report no association between time outdoors and myopia in adolescents. Schmid et al.³⁷[30] found no differences in the time spent in sunlight between 17 to 25-year-old myopes and non-myopes. Lu et al. [31] reported no differences in self-reported time spent outdoors and myopia status in Singaporean adolescents (mean age 14.7 years). Furthermore, Hagen et al. [32] saw no differences in reported outdoor time between myopes and non-myopes in 16 to 19-year-old Norwegian adolescents. Our study participants were aged between 15 to 17 years, and it may be that outdoor time at a younger age, which is more critical for refractive development, confers a protective effect, and outdoor time during adolescence may not be reflective of the previous outdoor exposure.

We assessed the SER before and after cycloplegia and determined an average difference of -0.89 D, with the non-cycloplegic values being more myopic. This is similar to a previous study reporting a -0.63 D difference [26]. Had we used non-cycloplegic SER values, we would have overestimated the prevalence of myopia in our study population by more than two-fold. This highlights the striking difference in the measured non-cycloplegic and cycloplegic autorefractometer values and emphasises the need to use cycloplegia for determining refractive status, which may be one underlying cause of conflicting results in the literature.

In conclusion, this is the first study evaluating myopia risk factors in Estonia. In agreement with published work, we found that parental myopia was a risk factor for myopia. Decreased time spent outdoors and increased time on near-work activities were not myopia risk factors in our study sample. In addition, we investigated whether chronotype affected myopia development and found that in our study sample, chronotype, assessed with MEQ, was not associated with myopia.

List of abbreviations

CI – confidence interval

D – dioptre

MEQ – Morningness-Eveningness Questionnaire

OR – odds ratio

SD – standard deviation

SER – spherical equivalent refraction

Declarations

Ethics approval and consent to participate

Tallinn Medical Research Ethics Committee gave ethical approval for this work. Written informed consent was obtained from the participants and their parents or legal guardians.

Consent for publication

Not applicable

Competing interests

The authors declare that they have no competing interests.

Funding

This research project received no external funding.

References

1. Holden BA, Fricke TR, Wilson DA, Jong M, Naidoo KS, Sankaridurg P, et al. Global Prevalence of Myopia and High Myopia and Temporal Trends from 2000 through 2050. *Ophthalmology*. 2016;123:1036–42.
2. Pärssinen O. The wearing of spectacles and occurrence of myopia. Tampere: Acta Universitatis Tamperensis, ser A vol 208, University of Tampere. 1986;208.
3. Rose KA, Morgan IG, Ip J, Kifley A, Huynh S, Smith W, et al. Outdoor Activity Reduces the Prevalence of Myopia in Children. *Ophthalmology*. 2008;115:1279–85.
4. Xiong S, Sankaridurg P, Naduvilath T, Zang J, Zou H, Zhu J, et al. Time spent in outdoor activities in relation to myopia prevention and control: a meta-analysis and systematic review. *Acta Ophthalmol (Copenh)*. 2017;95:551–66.
5. Ho C-L, Wu W-F, Liou YM. Dose–Response Relationship of Outdoor Exposure and Myopia Indicators: A Systematic Review and Meta-Analysis of Various Research Methods. *Int J Environ Res Public Health*. 2019;16:2595.
6. Huang H-M, Chang DS-T, Wu P-C. The Association between Near Work Activities and Myopia in Children—A Systematic Review and Meta-Analysis. *PLoS ONE*. 2015;10.
7. Enthoven CA, Tideman JWL, Polling JR, Yang-Huang J, Raat H, Klaver CCW. The impact of computer use on myopia development in childhood: The Generation R study. *Prev Med*. 2020;132:105988.
8. Hysi PG, Choquet H, Khawaja AP, Wojciechowski R, Tedja MS, Yin J, et al. Meta-analysis of 542,934 subjects of European ancestry identifies new genes and mechanisms predisposing to refractive error and myopia. *Nat Genet*. 2020;52:401–7.
9. Wilson LB, Quinn GE, Ying G, Francis EL, Schmid G, Lam A, et al. The Relation of Axial Length and Intraocular Pressure Fluctuations in Human Eyes. *Invest Ophthalmol Vis Sci*. 2006;47:1778–84.
10. Chakraborty R, Read S a., Collins MJ. Diurnal variations in axial length, choroidal thickness, intraocular pressure, and ocular biometrics. *Invest Ophthalmol Vis Sci*. 2011;52:5121–9.
11. Weiss S, Schaeffel F. Diurnal growth rhythms in the chicken eye: relation to myopia development and retinal dopamine levels. *J Comp Physiol A*. 1993;172:263–70.
12. Nickla DL, Wildsoet C, Wallman J. Visual Influences on Diurnal Rhythms in Ocular Length and Choroidal Thickness in Chick Eyes. *Exp Eye Res*. 1998;66:163–81.
13. Ayaki M, Torii H, Tsubota K, Negishi K. Decreased sleep quality in high myopia children. *Sci Rep*. 2016;6 May:33902–33902.
14. Liu XN, Naduvilath TJ, Wang J, Xiong S, He X, Xu X, et al. Sleeping late is a risk factor for myopia development amongst school-aged children in China. *Sci Rep*. 2020;10:17194.
15. Lee SS-Y, Mackey DA. Prevalence and Risk Factors of Myopia in Young Adults: Review of Findings From the Raine Study. *Front Public Health*. 2022;10:861044.

16. Xu S, Zong Z, Zhu Y, Zhang X, Zhang Y, Wang X, et al. Association between sleep-wake schedules and myopia among Chinese school-aged children and adolescents: a cross-sectional study. *BMC Ophthalmol*. 2023;23:135.
17. Zhou Z, Morgan IG, Chen Q, Jin L, He M, Congdon N. Disordered Sleep and Myopia Risk among Chinese Children. *PloS One*. 2015;10:e0121796.
18. Stafford-Bell N, McVeigh J, Lingham G, Straker L, Eastwood PR, Yazar S, et al. Associations of 12-year sleep behaviour trajectories from childhood to adolescence with myopia and ocular biometry during young adulthood. *Ophthalmic Physiol Opt J Br Coll Ophthalmic Opt Optom*. 2022;42:19–27.
19. Wei S-F, Li S-M, Liu L, Li H, Kang M-T, Sun Y-Y, et al. Sleep Duration, Bedtime, and Myopia Progression in a 4-Year Follow-up of Chinese Children: The Anyang Childhood Eye Study. *Invest Ophthalmol Vis Sci*. 2020;61:37–37.
20. Li M, Tan C-S, Xu L, Foo L-L, Yap F, Sun C-H, et al. Sleep Patterns and Myopia Among School-Aged Children in Singapore. *Front Public Health*. 2022;10:828298.
21. Sun J, Chen M, Cai W, Wang Z, Wu S, Sun X, et al. Chronotype: implications for sleep quality in medical students. *Chronobiol Int*. 2019;36:1115–23.
22. Rique GLN, Fernandes Filho GMC, Ferreira ADC, de Sousa-Muñoz RL. Relationship between chronotype and quality of sleep in medical students at the Federal University of Paraiba, Brazil. *Sleep Sci*. 2014;7:96–102.
23. Lima ÁMA, Varela GCG, Silveira HA da C, Parente RDG, Araujo JF. Evening chronotypes experience poor sleep quality when taking classes with early starting times. *Sleep Sci*. 2010;3:45–8.
24. Flanagan SC, Cobice D, Richardson P, Sittlington JJ, Saunders KJ. Elevated Melatonin Levels Found in Young Myopic Adults Are Not Attributable to a Shift in Circadian Phase. *Invest Ophthalmol Vis Sci*. 2020;61:45–45.
25. Flitcroft DI, He M, Jonas JB, Jong M, Naidoo K, Ohno-Matsui K, et al. IMI – Defining and Classifying Myopia: A Proposed Set of Standards for Clinical and Epidemiologic Studies. *Invest Ophthalmol Vis Sci*. 2019;60:M20–30.
26. Sankaridurg P, He X, Naduvilath T, Lv M, Ho A, Smith E, et al. Comparison of noncycloplegic and cycloplegic autorefraction in categorizing refractive error data in children. *Acta Ophthalmol (Copenh)*. 2017;95:e633–40.
27. Horne JA, Ostberg O. A self-assessment questionnaire to determine morningness-eveningness in human circadian rhythms. *Int J Chronobiol*. 1976;4:97–110.
28. Zhang X, Qu X, Zhou X. Association between parental myopia and the risk of myopia in a child. *Exp Ther Med*. 2015;9:2420–8.
29. Moafa MAM. Quantifying Risk Factors for Myopia: Instrument Validation and Establishing Dosage for Light Intensity and Duration. The University of Sydney; 2019.
30. Schmid KL, Leyden K, Chiu Y, Lind S-R, Vos DJ, Kimlin M, et al. Assessment of Daily Light and Ultraviolet Exposure in Young Adults. *Optom Vis Sci*. 2013;90:148–55.

31. Lu B. Associations Between Near Work, Outdoor Activity, and Myopia Among Adolescent Students in Rural China: The Xichang Pediatric Refractive Error Study Report No. 2. Arch Ophthalmol. 2009;127:769.

32. Hagen LA, Gjelle JVB, Arnegard S, Pedersen HR, Gilson SJ, Baraas RC. Prevalence and Possible Factors of Myopia in Norwegian Adolescents. Sci Rep. 2018;8:13479.

Figure 1




Figure 1. Distribution of factors analysed in non-myopes and myopes. (A) Time spent outdoors, (B) on near work and (C) mid-working distance activities in non-myopes and myopes. (D) Distribution of chronotype, measured with MEQ, in non-myopes and myopes. Data are presented as mean \pm SD. MEQ, Morningness-Eveningness Questionnaire; SD, standard deviation.

Figure 2




Figure 2. Characterisation of SER of the study population. (A) Distribution of the participants' mean cycloplegic SER grouped into one dioptre bins. Correlations between (B) SER and axial length, and (C) SER and radius of corneal curvature in individual eyes (Spearman correlation, individual values fitted with simple linear regression to illustrate the relationship). SER, spherical equivalent refraction; D, dioptres.

Figure 3




Figure 3. Effect of cycloplegia on SER. (A) Difference between cycloplegic and non-cycloplegic SER ($p < 0.0001$, Wilcoxon matched-pairs signed rank test; boxplot presents median with minimum and maximum values, “+” denotes the mean, individual participant values presented). (B) Correlation between cycloplegic and non-cycloplegic SER (Spearman correlation, individual values fitted with simple linear regression to illustrate the relationship). SER, spherical equivalent refraction; D, dioptres.

Table 1. Demographic characteristics of the study population.

	All participants		Myopes	
	N	%	N	%
Total	123	100	38	30.9
Characteristic				
Sex				
Female	70	56.9	24	34.3
Male	53	43.1	14	26.4
Age				
15 years	5	4.1	1	20.0
16 years	87	70.7	26	29.9
17 years	31	25.2	11	35.5
Parental myopia				
None	43	35.0	8	18.6
One myopic parent	58	47.2	19	32.8
Both parents myopic	22	17.9	11	50.0
School				
Science	61	49.6	27	42.3
Arts	35	28.5	8	22.9
Sports	27	22.0	3	11.1

The demographics are characterised for the entire study population. The number and proportion of myopes in each demographic group are presented.

Table 2. Effect of demographic and lifestyle factors and chronotype on myopia status.

Characteristic	Odds ratio	p value	95% CI
Sex			
Female	1.00		
Male	1.83	0.213	0.71 – 4.76
Parental myopia			
None	1.00		
One myopic parent	1.50	0.459	0.52 – 4.33
Both parents myopic	4.32	0.026	1.19 – 15.74
School			
Science	1.00		
Arts	0.41	0.103	0.14 – 1.20
Sports	0.12	0.004	0.03 – 0.51
Chronotype (MEQ score)			
I tercile (evening types)	1.00		
II tercile	2.29	0.158	0.73 – 7.26
III tercile (morning types)	2.69	0.074	0.91 – 7.98
Time spent outdoors (hours per week)			
I tercile (4.5-11.0)	1.00		
II tercile (11.25-17.75)	1.17	0.794	0.35 – 3.90
III tercile (18.0-43.5)	1.00	0.996	0.32 – 3.14
Time spent on near work (hours per week)			
I tercile (14.0-45.0)	1.00		
II tercile (46.0-59.0)	0.58	0.363	0.18 – 1.86
III tercile (60.0-95.0)	1.00	0.996	0.32 – 3.15
Time spent on middle-distance work (hours per week)			
I tercile (3.0-15.0)	1.00		
II tercile (15.5-24.5)	1.88	0.270	0.61 – 5.76
III tercile (25.0-68.0)	0.97	0.961	0.30 – 3.19

A multivariate logistic regression model was fitted with myopia status as the dependent variable. Sex, parental myopia, school, chronotype tercile, outdoor time tercile, near work tercile and mid-working distance activity time tercile were included as independent variables. A *p* value of <0.05 was considered statistically significant and marked in bold. CI, confidence interval; MEQ, Morningness-Eveningness Questionnaire.