

1 **Capturing recent *Mycobacterium tuberculosis* infection by tuberculin skin test vs.**  
2 **interferon-gamma release assay**

3 **Jesús Gutierrez<sup>1\*</sup>, Mary Nsereko<sup>2</sup>, LaShaunda L. Malone<sup>3</sup>, Harriet Mayanja-Kizza<sup>2</sup>, Hussein**  
4 **Kisingo<sup>2</sup>, W. Henry Boom<sup>3</sup>, Charles M. Bark<sup>4¶</sup>, Catherine M. Stein<sup>1,3¶</sup>**

5  
6 <sup>1</sup>Department of Population and Quantitative Health Science, Case Western Reserve University,  
7 Cleveland, OH, United States

8 <sup>2</sup>Uganda-CWRU Research Collaboration and Department of Medicine, School of Medicine,  
9 Makerere University, Kampala, Uganda

10 <sup>3</sup>Department of Medicine, Case Western Reserve University and University Hospitals Cleveland  
11 Medical Center, Cleveland, OH, United States

12 <sup>4</sup>Division of Infectious Diseases, MetroHealth Medical Center, Cleveland, OH, United States

13  
14 \* Corresponding author

15 E-mail: [jesus.gutierrez@case.edu](mailto:jesus.gutierrez@case.edu) (JG)

16  
17 ¶ These authors contributed equally as senior authors of this work.

18

19

## 20 Abstract

21 Reductions in tuberculosis (TB) incidence and mortality require identification of individuals  
22 at high risk of developing active disease, such as those with recent *Mycobacterium tuberculosis*  
23 (*Mtb*) infection. Current tests used for *Mtb* infection cannot distinguish recent from remote  
24 infection. Using a longitudinal, prospective household contact (HHC) study in Kampala, Uganda,  
25 we diagnosed new *Mtb* infection using both the tuberculin skin test (TST) and interferon-gamma  
26 release assay (IGRA) to determine if these tests differentiated or added value as measures of  
27 recent *Mtb* infection. We identified 13 HHCs who only converted the IGRA (QFT-only converters),  
28 39 HHCs who only converted the TST (TST-only converters), and 24 HHCs who converted both  
29 tests (QFT/TST converters). Univariate analysis revealed that TST-only converters were older  
30 than the other groups. This result was confirmed by our logistic regression model where increased  
31 odds of TST-only conversion was associated with age ( $p=0.02$ ), in addition to crowdedness  
32 ( $p=0.025$ ). The odds of QFT conversion increased with a higher epidemiologic risk score  
33 ( $p=0.038$ ), which suggests it is a better measure of *Mtb* exposure and infection. QFT/TST  
34 converters had higher QFT quantitative values at conversion than QFT-only converters and a  
35 bigger change in TST quantitative values at conversion than TST-only converters. Collectively,  
36 these data indicate that TST conversion alone likely overestimates *Mtb* infection. Its correlation  
37 to older age suggests boosting of BCG responses and/or longer environmental mycobacterial  
38 exposure. This result also suggests that QFT/TST conversion may be associated with a more  
39 robust immune response, which should be considered when planning vaccine studies.

## 40 Introduction

41 Tuberculosis (TB) continues to be a major public health problem globally. In 2019,  
42 approximately 10 million people became ill with TB and over 1.4 million succumbed to the disease.  
43 The WHO's End TB Strategy aims to achieve a 95% reduction in TB mortality and an 90%  
44 reduction in the TB incidence rate by 2035 (1). Although some regions in the world are on track  
45 to achieve these milestones, the African continent has fallen behind (2). Moreover, 2020 marked  
46 the first year that TB deaths increased worldwide due to the COVID-19 pandemic (3). Faster  
47 reductions in TB incidence and deaths require improvements in multiple facets, including  
48 identification of those at risk of progressing to TB. Several studies have demonstrated that of  
49 those with latent *Mycobacterium tuberculosis* (*Mtb*) infection (LTBI) who progress to TB, most do  
50 so within the first two years post-infection (4). Furthermore, individuals with a recent  
51 *Mycobacterium tuberculosis* (*Mtb*) exposure and conversion of their tuberculin skin test (TST)  
52 and/or interferon-gamma release assay (IGRA) have a higher risk for progression to TB (5-7).  
53 Identifying and treating individuals with recent *Mtb* infection could be an important step in  
54 preventing TB.

55 Today, we rely on the TST or IGRA to diagnose *Mtb* infection. These tests have important  
56 limitations. First, the tests only infer the presence of *Mtb* based on a person's T-cell response to  
57 *Mtb* antigens and are poor predictors of progression to TB (8). Also, given that both tests use  
58 different stimuli and measure different immunologic responses, discordance rates are substantial  
59 (9). Since T-cell responses persist, a positive result cannot distinguish between recent vs. remote  
60 *Mtb* infection (10, 11), and some with a remote infection may have cleared the infection (4, 7, 10).  
61 Lately, the IGRA has become the preferred test of choice to diagnose LTBI over the TST due to  
62 improved specificity and requiring only one visit to perform the test (12-16). In a meta-analysis,  
63 the pooled sensitivity for the QFT reached 78% and the specificity among BCG-vaccinated and  
64 non-BCG-vaccinated persons was 96% and 99%, respectively (15). This preference is

65 demonstrated in the recommendations of many national TB programs in low-endemic countries  
66 (15, 17) and has expanded to clinical TB research (18, 19).

67 Despite the increased specificity for *Mtb* of IGRA, TST is still widely used in TB endemic  
68 areas for cost and simplicity reasons in the evaluation of latent *Mtb* infection (LTBI). Longitudinal  
69 studies (9), such as household contact (HHC) studies, with at least 6 months of follow-up and  
70 repeat testing can capture recent and/or new *Mtb* infection and offer a unique opportunity to  
71 evaluate both tests of sensitization to *Mtb* antigens (20). In this study, we aimed to determine if  
72 the TST adds additional information and/or value to the characterization of IGRA converters.  
73 Optimal characterization of converters could help maximize the capture of recent *Mtb* infection as  
74 well as the design of clinical studies to prevent new and treat recent *Mtb* infection (21). Given the  
75 discordance between TST and IGRA, understanding the impact of this discordance in the  
76 identification of recent or new *Mtb* infection, i.e., TST/IGRA converters is important. Finally, if  
77 identification of recent converters is practical, focused preventive drug treatment could impact the  
78 incidence of TB.

## 79 **Materials and methods**

### 80 **Study setting**

81 The study is set in the greater Kampala metropolitan area, Uganda. The prevalence of  
82 active TB in Uganda is high with recent estimates reaching 401 per 100,000 for individuals 15  
83 years of age and older (22). The prevalence in Kampala has been estimated to be even higher at  
84 764 per 100,000 (23).

### 85 **Study population**

86 Individuals suspected to have active pulmonary TB at participating health centers in  
87 Kampala were referred to the Uganda-Case Western Reserve University Research Collaboration

88 clinic starting in 2015. The referred TB suspects underwent counseling and consent for a standard  
89 TB assessment under a separately approved protocol for ongoing TB studies within the  
90 Collaboration. Diagnosis of TB for the index case was established through a standard  
91 assessment, based on chest radiograph, history, physical examination, and sputum smear for  
92 acid-fast bacilli (AFB) or MTB/RIF GeneXpert® testing. If the assessment confirmed disease,  
93 these individuals with pulmonary TB (“index cases”) were asked for permission to approach their  
94 household contacts (HHCs) who were at least 15 years old to determine their interest in study  
95 participation. Index cases were not enrolled in this study. Upon receiving documented permission  
96 from the TB index case, HHCs were contacted to undergo informed consent within 28 days (or 3  
97 months of HIV+ HHCs) of the confirmed TB diagnosis of the index case. Consenting HHCs who  
98 met eligibility criteria (see below) were enrolled if the index case of their household was sputum  
99 smear positive for AFB, and subsequently confirmed to have active TB by sputum culture or  
100 MTB/RIF GeneXpert® positivity. Recruitment for this study occurred from June 29<sup>th</sup>, 2016 until  
101 March 10<sup>th</sup>, 2020.

102 The study protocol was reviewed and approved by The Uganda National Council on  
103 Science and Technology, and the institutional review board at the University Hospitals Cleveland  
104 Medical Center. Written consent was obtained from all participants ages 18 and older. Assent  
105 was obtained prior to screening from individuals ages 15 to 17, as well as written informed consent  
106 from their parent or guardian.

## 107 **Inclusion criteria**

108 In order to be enrolled in the study as a HHC, potential participants needed to meet the  
109 following criteria: 1. Informed Consent or assent (where indicated) 2. Adults and minors aged 15  
110 years of age and older; 3. Individuals living in the same building (house, hut, or apartment), or a  
111 portion of the building, as the index case, thereby sharing air-space with the index case, for at  
112 least one week during the three-month period immediately preceding the diagnosis of TB in the

113 index case; 4. Negative urine pregnancy test in women of child-bearing potential. Breastfeeding  
114 women were permitted to enroll in this study.

## 115 **Exclusion criteria**

116 In addition, potential participants were excluded from the study due to the following criteria:

117 1. TB or other febrile illness or uncontrolled disease; 2. Peripheral blood CD4 lymphocyte count  
118 < 200/mm<sup>3</sup> for HIV+ participants; 3. Chest radiograph consistent with TB; 4. Expected to be  
119 unavailable for the 12-month follow-up period.

## 120 **Study design**

121 This is an ongoing prospective longitudinal cohort study with a household contact design.  
122 HHCs age 15 years and older who were living with the index cases and met eligibility criteria as  
123 outlined above were enrolled. At baseline, each HHC underwent TB symptom screening, HIV  
124 testing, a pregnancy test for females, an IGRA test and TST placement to be checked within 3  
125 days (S1 Fig). The IGRA test used was the QuantiFERON®-TB Gold (QFT) or the QuantiFERON®-  
126 TB Gold Plus (QFT-Plus), depending on availability during study follow-up. The QFT and QFT-  
127 Plus results were interpreted according to the manufacturer's recommendations. The TST test  
128 was performed by the Mantoux method (5 tuberculin units per 0.1 ml of purified protein derivative,  
129 Tubersol; Connaught Laboratories Limited, Willowdale, Ontario Canada). TST positivity was  
130 defined by a maximum induration of 10 mm or greater, or an induration of 5 mm or greater in HIV+  
131 individuals. In addition, each HHC underwent an individual risk assessment, a review of medical  
132 history, a complete physical examination, and a chest radiograph. Households were evaluated to  
133 determine if the house was a muzigo (the traditional muzigo in Uganda is a one-room dwelling in  
134 which all activities are done apart from cooking) (24), whether the cooking took place inside or  
135 outside the house, number of inhabitants, number of rooms, and number of windows. All HHC  
136 were followed for twelve months. Initially, follow-up visits occurred at 3, 6, and 12 months after  
137 enrollment. Subsequently, a 9-month visit was added to the protocol. During each follow -up visit,

138 HHCs underwent a health check, a TB symptom screening, and a repeat IGRA test for HHCs who  
139 had not become pregnant. A repeat TST was performed at 12 months for those HHCs who had  
140 a negative TST at enrollment. Data collection and management for this paper was performed  
141 using OpenClinica open-source software, version 3.16. Copyright OpenClinica LLC and  
142 collaborators, Waltham, MA, USA, [www.OpenClinica.com](http://www.OpenClinica.com).

## 143 **Definitions**

144 Once the HHCs completed follow-up, they were classified based on the following  
145 definitions (Fig 1):

146 **Fig 1. CONSORT diagram for this study.** Out of 383 household contacts enrolled in our study,  
147 69 were excluded. 314 household contacts were classified using both TST and QFT. Those  
148 who were defined as concordant LTBI (both TST and QFT results were consistently positive)  
149 and discordant “resisters” (either 2 negative TST or all negative QFT) were excluded from  
150 analysis (n=166). The remaining 148 household contacts were defined as follows: 39 TST-only  
151 converters (negative baseline TST with positive TST at 12 months), 13 were QFT-only  
152 converters (negative QFT at baseline followed by positive QFT during follow-up), 24 were  
153 QFT/TST converters (negative baseline TST and QFT followed by positive TST and QFT during  
154 follow-up), and 72 were concordant short-term “resisters” (all TST and QFT tests remained  
155 negative).

- 156 • A person with LTBI was defined as a HHC with all positive QFT and TST results.
- 157 • A “short-term” resister (“resister”) was defined as a HHC with all negative QFT and TST results  
158 throughout the course of the study.
- 159 • A QFT-only conversion was based on at least 3 QFT results, requiring a negative QFT at  
160 baseline followed by at least one positive QFT result during follow-up, accompanied by TST  
161 results that did not change throughout the study.

- 162 • A TST-only conversion was based on two TST results (baseline and month 12) and was  
163 defined as an original TST of <10 mm (or <5 mm for HIV+ individuals) followed by a TST  
164 reaction of ≥10 mm (or ≥5 mm for HIV+ individuals) on subsequent testing with an increase  
165 of ≥6 mm (25), accompanied by QFT results that remained positive or negative throughout  
166 the study.
- 167 • A QFT/TST conversion is defined as a conversion of both the QFT and TST as described  
168 above. There were 6 QFT converters (5 QFT/TST converters and one QFT-only converter)  
169 who had just one positive QFT result because it occurred at month 12.

170 Those HHCs with indeterminate or inconsistent QFT results were classified by consensus  
171 into one of these categories based on the preponderance of the data, when possible. Otherwise,  
172 these HHCs were excluded from the statistical analyses described below. There were also 5  
173 individuals with one positive QFT in the middle of the observation period who were excluded from  
174 the analysis. Based on a single QFT result, it is unclear if these reflect a reversion event vs. a  
175 false-positive QFT. Serial and consistent QFT results are required to reliably detect definite  
176 conversion events, i.e., recent *Mtb* infection. Variability in QF responses over time has been  
177 observed in another study (26).

## 178 **Statistical analyses**

179 First, we examined the similarity between subjects classified using only TST results or  
180 only QFT results to illustrate the impact of discordance between the two tests. This analysis  
181 included all classified HHCs.

182 Second, we aimed to illustrate how TST and QFT would define conversion differently  
183 among HHCs. This analysis only included the three types of converter groups (QFT-only  
184 converters, TST-only converters, and QFT/TST converters) and the “resister” group (HHCs who  
185 remained QFT negative and TST negative during 12-month follow-up despite a high-level  
186 exposure to an infectious TB index case). “Resisters” were included in this analysis since they



187 are known to have unique and robust immune responses (27), making them an appropriate  
188 control group. Comparisons were first made at the univariate level. Analysis focused on the  
189 following variables: epidemiological risk score (ERS), HIV status, body mass index (BMI), BCG  
190 status, quantitative IGRA values, quantitative TST values, type of housing (a marker of  
191 socioeconomic status), and clinical characteristics of the index case that have been associated  
192 with higher risk of transmission. The ERS consists of variables indicative of risk for *Mtb* exposure  
193 and infection and has been previously used in other studies examining the risk of LTBI (28, 29).  
194 Further details on the ERS can be found in the Supplement. Comparisons were done using the  
195 chi-square test, student's t-test, analysis of variance, Mann-Whitney test, and the Kruskal-Wallis  
196 test. Chi-square was used for categorical variable comparison, while other tests were used to  
197 examine continuous variables. Significance was assessed using a 0.05 alpha cut-off, using the  
198 Bonferroni correction to correct for multiple comparisons. When an omnibus comparison was  
199 significant, a pairwise comparison was done to determine which pair differences were contributing  
200 to this result. This univariate analysis was then repeated combining QFT-only converters and  
201 QFT/TST converters into a single category, QFT converters.

202 Third, a cluster analysis of all available variables using Gower distance was performed to  
203 determine if the subject clustering obtained was similar to the predetermined conversion groups.  
204 The purpose of this analysis was to examine whether clinical and epidemiological variables could  
205 better define subgroups of HHCs and if these groupings aligned with a TST and/or QFT-based  
206 definition. Additional details are found in the Supplement.

207 Fourth, we evaluated the difference between TST-only conversion and QFT-conversion  
208 using a logistic regression model, where the dependent variable was a TST-only conversion event  
209 versus a QFT conversion event. The independent variables that were initially included were  
210 chosen based on the results of the univariate and cluster analyses described above. Once these  
211 were chosen, we proceeded to formulate our logistic regression models by using two different  
212 methods. The first model contained all of the chosen independent variables. The second model

213 was obtained using an automated backwards elimination process. The two models were  
214 compared using the Akaike information criterion (AIC) method and the one with the lowest AIC  
215 was chosen.

216 Finally, since previous studies have suggested different cut-off values for positivity  
217 (especially for the IGRA), we examined the quantitative changes in both the QFT and TST tests  
218 to better quantify changes associated with each of our conversion definitions. All analyses were  
219 performed using R (30).

## 220 **Results**

### 221 **Differences in classification among all classified HHCs by type** 222 **of test**

223  
224 This first comparison was performed on the 314 individual HHCs enrolled by January 2022  
225 that could be defined as converter, LTBI or “resister” (Fig 1). We focused on how similarly subjects  
226 could be classified by TST or QFT. Table 1 shows that if classification was based on TST, there  
227 would be 63 converters. By QFT, the number of converters would be 37. If both tests are used,  
228 there would be only 24 converters. The conversion rate for TST-only converters was 27.7% (39  
229 out of 141 TST negatives at baseline) and for QFT-only converters, it was 9.7% (13 out of 134  
230 QFT negatives at baseline). The conversion rate for QFT/TST converters was 22.2% (24 out of  
231 108 TST and QFT negatives at baseline). The concordance rate QFT and TST is lowest for  
232 converters (64.9%) compared to LTBI and “resisters” (82.8% and 74.2%, resp.). These analyses  
233 were restricted to subjects with consistent QFT and TST results according to our definitions.

234

235

236

237

238

239 **Table 1. Cross-tabulation table of the discordance between QFT-only and TST-only**  
240 **classifications of converters, LTBI and “resisters”.**

		QFT-based Classification			
		N (%)	Converter	LTBI	“Resister”
TST-based Classification	Converter	24 (64.9%) <sup>a</sup>	27 (15.0%) <sup>b</sup>	12 (12.4%) <sup>b</sup>	63 (20.1%)
	LTBI	11 (29.7%) <sup>c</sup>	149 (82.8%) <sup>d</sup>	13 (13.4%) <sup>d</sup>	173 (55.1%)
	“Resister”	2 (5.4%) <sup>c</sup>	4 (2.2%) <sup>d</sup>	72 (74.2%) <sup>e</sup>	78 (24.8%)
	Total	37 (100%)	180 (100%)	97 (100%)	314 (100%)

241 Of the 314 HHCs, 166 were excluded from the analysis, 39 were QFT-only converters, 13 were  
242 TST-only converters, and 24 were QFT/TST converters. In addition, 72 “resisters” were used as  
243 a control group.

244 <sup>a</sup>These participants were classified as QFT/TST converters (n=24)

245 <sup>b</sup>These participants were classified as QFT-only converters (n=39)

246 <sup>c</sup>These participants were classified as TST-only converters (n=13)

247 <sup>d</sup>These participants were excluded from the analysis (n=166)

248 <sup>e</sup>These participants were classified as concordant “resisters” (n=72)

249 **Individual, household and TB index characteristics of TST-**  
250 **only, QFT-only and TST/QFT converters**

251  
252 The main analysis was performed on the three converter groups and “resisters” (Fig 1).  
253 Table 2 summarizes the characteristics and comparisons for each converter group with  
254 “resisters”. Converter groups did not differ in terms of sex, HIV positivity, BMI, presence of BCG  
255 scar, ERS, and smoking history. They did differ in age ( $p = 0.04$ ). TST-only converters were older  
256 than “resisters”: median age 32 [20-47] vs. 23 [19.9-36.5], respectively,  $p = 0.006$ ). TST-only  
257 converters also appeared older than the other two converter groups, however, this difference did  
258 not reach statistical significance.

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275 **Table 2. Univariate analysis comparing QFT-onl, TST-only and QFT/TST converters based on individual and household**  
 276 **characteristics.**

	<b>QFT-only converters</b>	<b>TST-only converters</b>	<b>QFT/TST converters</b>	<b>“Resisters”</b>	<b>p-value (test)</b>	<b>Relevant pairwise p-value</b>
N	13	39	24	72	NA	NA
<b>Individual Characteristics</b>						
Age	26 [20-36]	32 [20-47]	26.5 [23-39.3]	23 [19.8-36.5]	0.04* (KW test)	TST-only vs. “Resister”: 0.006*
Sex (female)	8 (61.5%)	21 (53.8%)	19 (79.2%)	43 (59.7%)	0.23 (Fisher’s)	NA
BCG scar present	12 (92.3%)	31 (81.6%)	14 (58.3%)	53 (75.7%)	0.08 (Fisher’s)	NA
HIV positive	0 (0%)	3 (7.7%)	3 (12.5%)	4 (5.6%)	0.57 (Fisher’s)	NA
BMI	22.2 [19.6-28.3]	23.6 [19.5-27.0]	22.5 [20.5-28.1]	22.8 [21.0-25.2]	0.93 (KW test)	NA
TB Risk score	7 [6-8]	6 [6-7]	7 [6-7]	6 [6-7]	0.15 (KW test)	NA
Quantitative IGRA values at conversion	2.0 [0.9–3.8]	3.4 [0.3-9.3]	4.1 [1.7-10]	NA	0.15 (KW test)	NA
No history of smoking	12 (92.3%)	37 (94.9%)	23 (95.8%)	61 (84.7%)	0.09 (Fisher’s)	NA
Spouse to	4 (30.8%)	7 (17.9%)	5 (20.8%)	8 (11.1%)	0.26 ( $\chi^2$ )	NA

Index						
<b>Household Characteristics</b>						
Living in Muzigo	9 (69.2%)	20 (51.3%)	9 (37.5%)	21 (29.2%)	0.02* (Fisher's)	QFT-only vs. "Resister": 0.01* TST-only vs. "Resister": 0.04*
Cooking inside home	1 (7.7%)	10 (25.6%)	11(45.8%)	28 (38.9%)	0.05* (Fisher's)	QFT-only vs. QFT/TST: 0.03* QFT-only vs. "Resister": 0.03*
Number of windows	1 [0-2]	1 [1-3]	2.5 [1-6]	3 [1-4]	0.007* (KW test)	QFT-only vs. QFT/TST: 0.003* QFT-only vs. "Resister": 0.003*
People per room	2.5 [2-3]	2 [1.33-3]	1.67 [1-2.25]	2 [1.3-2.67]	0.08 (KW test)	NA
Sleeping in same room	11 (84.6%)	21 (53.8%)	15 (62.5%)	33 (45.8%)	0.053 (X <sup>2</sup> )	NA
Sleeping in same bed	5 (38.5%)	6 (15.4%)	6 (25%)	12 (16.7%)	0.23 (X <sup>2</sup> )	NA

277 Counts (percentages) or median [quartiles]

278 ERS: Epidemiologic risk score

279 KW: Kruskal-Wallis

280 X<sup>2</sup>: Chi-square test

281 \* Statistically significant at  $p < 0.05$

282 In comparing household characteristics by conversion group (Table 2), a greater  
283 proportion of QFT-only converters (69.2%) live in muzigos. A higher proportion of QFT-only  
284 converters (92.3%) cook outside the home when compared to the other groups. QFT-only  
285 converters lived in homes with fewer windows (1 window [0-2]) when compared to QFT/TST  
286 converters (2.5 windows [1-6]) and “resisters” (3 windows [1-4]) ( $p = 0.003$  and  $p = 0.003$ ,  
287 respectively). There were no differences in a measure of crowdedness (people per room) and  
288 sleeping location (sleeping in the same room or the same bed).

289 Next, we examined the characteristics of the index case for the different converter groups  
290 (Table 3). Index cases of TST-only converters had more advanced lung disease (61.5%) when  
291 compared to those of QFT-only converters (23.1%) ( $p = 0.04$ ). Index cases of TST-only converters  
292 had a longer duration of cough than those of QFT/TST converters: 90 days [60-142.5] vs 52.5  
293 days [30-60] ( $p = 0.007$ ). In contrast QFT/TST converters had a higher proportion of index cases  
294 with hemoptysis (33.3%) than index cases of TST-only converters (10.3%) ( $p = 0.04$ ).

295 **Table 3. Univariate analysis comparing QFT-only, TST-only and QFT/TST converters based on index case characteristics.**

	<b>QFT-only converters</b>	<b>TST-only converters</b>	<b>QFT/TST converters</b>	<b>“Resisters”</b>	<b>p-value (test)</b>	<b>Relevant pairwise p-value</b>
Sex (female)	8 (61.5%)	20 (51.3%)	17 (70.8%)	35 (48.6%)	0.26 (Fisher’s)	NA
BCG scar present	10 (76.9%)	30 (76.9%)	17 (73.9%)	36 (56.3%)	0.10 (Fisher’s)	NA
HIV positive	1 (10.0%)	4 (12.9%)	0 (0%)	12 (18.8%)	0.14 (Fisher’s)	NA
BMI	19 [17-21]	18.5 [17-21]	18 [17-20]	19 [18-20]	0.42 (KW test)	NA
Cavitary lesions present	11 (84.6%)	32 (82.1%)	22 (91.7%)	43 (60.6%)	0.006* (Fisher’s)	TST-only vs. “Resister”: 0.03* QFT/TST vs. “Resister”: 0.005*
Advanced lung disease	3 (23.1%)	24 (61.5%)	14 (58.3%)	25 (35.2%)	0.01* (Fisher’s)	TST-only vs. QFT-only: 0.04* TST-only vs. “Resister”: 0.01*
Positive Smear	8 (88.9%)	29 (100%)	20 (100%)	55 (100%)	0.08 (Fisher’s)	NA
Positive GeneXpert	13 (100%)	39 (100%)	24 (100%)	71 (98.6%)	1.00 (Fisher’s)	NA
Coughing	13 (100%)	39 (100%)	24 (100%)	71 (100%)	1.00 (Fisher’s)	NA
Cough duration	60 [60-90]	90 [60-142.5]	52.5 [30-60]	60 [30-90]	0.03* (KW)	TST-only vs. QFT/TST: 0.007*



(days)					test)	TST-only vs. "Resister": 0.017*
Fever	9 (69.2%)	30 (76.9%)	19 (79.2%)	54 (76.1%)	0.92 ( $X^2$ )	NA
Fever duration (days)	42 [14-75]	60 [30-90]	30 [27.8-60]	30 [14-56.3]	0.10 (KW test)	NA
Productive sputum	13 (100%)	36 (92.3%)	24 (100%)	68 (95.8%)	0.55 (Fisher's)	NA
Productive sputum duration (days)	30 [30-90]	60 [30-120]	30 [27.8-60]	30 [30-60]	0.11 (KW test)	NA
Purulent sputum	6 (46.2%)	22 (56.4%)	14 (58.3%)	37 (52.9%)	0.89 ( $X^2$ )	NA
Purulent sputum duration (days)	45 [30-60]	52.5 [30-120]	30 [23.3-60]	30 [15-90]	0.25 (KW test)	NA
Hemoptysis	1 (7.7%)	4 (10.3%)	8 (33.3%)	7 (10.0%)	0.04* (Fisher's)	QFT/TST vs. "Resisters": 0.02* QFT/TST vs. TST-only: 0.04*
Hemoptysis duration (days)	14 [14-14]	5 [2.5-12.8]	2 [1.5-5]	7 [2.5-10.5]	0.58 (KW test)	NA
Dyspnea	6 (46.2%)	20 (52.6%)	10 (41.7%)	38 (53.5%)	0.77 (Fisher's)	NA
Dyspnea duration (days)	60 [37.5-60]	90 [30-160]	60 [37.5-90]	30 [30-82.5]	0.18 (KW test)	NA
Weight loss	12 (92.3%)	33 (84.6%)	18 (75%)	59 (83.1%)	0.62 (Fisher's)	NA

Weight loss duration (days)	60 [27.8-78.8]	60 [30-90]	52.5 [30-71.3]	30 [30-82.5]	0.20 (KW test)	NA
-----------------------------	----------------	------------	----------------	--------------	----------------	----

296 Counts (percentages) or median [quartiles]

297 ERS: Epidemiologic risk score

298 KW: Kruskal-Wallis

299  $\chi^2$ : Chi-square test

300 \* Statistically significant at  $p < 0.05$

301 When all QFT converters were compared to TST-only converters and “resisters”, no new  
302 significant associations were identified (Supp Table 1 and Supp Table 2).

### 303 **Identifying predictors of QFT and TST conversion**

304 Next, we conducted a logistic regression analysis of QFT conversion vs. TST-only  
305 conversion to determine how they might differ epidemiologically (Table 4). Included variables  
306 were based on the univariate analyses in tables 2 and 3. We also included variables that were  
307 most influential in the cluster analysis using Gower distance (S2 Fig). Based on backwards  
308 elimination, we arrived at a model containing six predictors including age of the household  
309 contact, presence of cavitory lesions in the chest x-ray of the index case, extent TB lung disease  
310 in index cases by chest x-ray, duration of index case cough in days, ERS, and number of people  
311 per room.

312 **Table 4. Logistic regression model of predictors of TST-only conversion vs. QFT**  
313 **conversion.**

314

COVARIATES	N	ODDS RATIO	95% CI	p-VALUE
Age (years)	NA	1.05	1.01 – 1.10	0.020*
Presence of cavitory lesions				
NO	11	(Ref.)	(Ref.)	(Ref.)
YES	65	0.29	0.055 – 1.310	0.118
Index cough duration (days)	NA	1.01	1.000 – 1.018	0.099
Epidemiologic risk score	NA	0.61	0.375 – 0.956	0.038*
People per room	NA	1.559	1.085 – 2.404	0.025*
Advanced lung disease				
NOT FAR ADVANCED	35	(Ref.)	(Ref.)	(Ref.)
FAR ADVANCED	41	2.58	0.818 – 8.866	0.115

315 \* Statistically significant at  $p < 0.05$

316 CI: Confidence interval

317 We found that the odds of undergoing TST-only conversion increased by 5% (95% CI [.01-  
318 .10]) for each additional year of age. The odds of TST-only conversion also increased by 55.9%  
319 (95% CI [.08-1.4]) for each additional person per room. On the other hand, the odds of TST

320 conversion decreased by 39% (95% CI [0.38-0.96]) for each point increase on the ERS. Based  
321 on an AUC of 0.81, this logistic regression model provided good discriminatory power (Table 4).

## 322 **Quantitative QFT and TST values for TST, QFT and TST/QFT converters**

323 Next, we compared QFT quantitative values at the time of QFT conversion (Fig 2). This  
324 showed that QFT-only converters had lower IFN-gamma secretion values at conversion than  
325 QFT/TST converters (2.03 IU [0.92-3.82] vs. 4.14 IU [1.73 vs. 10.0],  $p=0.03$ ). We also compared  
326 quantitative TST values (Fig 3). At baseline, QFT/TST converters had a median zero mm baseline  
327 TST (range 0-0 mm), while TST-only converters had a higher median 2 mm baseline TST (range  
328 0-8 mm) ( $p=0.001$ ). By month 12, these 2 groups were indistinguishable: median: 15.6 mm vs.  
329 14.6 mm ( $p=0.25$ ).

330 **Fig 2. Quantitative QFT values of QFT/TST converters and QFT-only converters before and**  
331 **after QFT-conversion.** Red dotted line indicates 0.35 IU cut-off. If QFT plus was used, the  
332 highest value was used for this analysis. QFT-only converters had a lower value at conversion  
333 when compared to QFT/TST converters (2.03 [0.92-3.82] vs. 4.14 [1.73-10.0],  $p=0.03$ ).

334 **Fig 3. Quantitative TST values of QFT/TST converters and TST-only converters at baseline**  
335 **and at month 12 of follow-up.** Red dotted line indicates 10 mm. cut-off. At baseline, TST-only  
336 converters had a higher TST value compared to QFT/TST converters (2.0 [0.0-8.0] vs. 0.0 [0.0-  
337 0.0],  $p=0.001$ ). By month 12, this difference had disappeared (14.6 [12.2-16.3] vs. 15.6 [13.9-  
338 17.6],  $p=0.25$ ).

## 339 **Discussion**

340 The goal of this study was to determine the advantages of using both the TST and IGRA  
341 tests in assessing recent *Mtb* infection. Although the results do not provide a clear differentiation  
342 between TST vs. IGRA converters, they do highlight the following important conclusions.

343           When examining TST-only converters, we observe that they are more numerous than  
344 IGRA-only converters. This result is similar to what was seen in a 2014 HHC study from Brazil  
345 (31). While this could be the result of BCG cross-reactivity, BCG cross-reactivity is minimal when  
346 vaccination occurs during infancy and the TST is performed at least 10 years after (32). Given  
347 that all our HHCs were at least 15 years of age and that the median age of our TST-only  
348 converters is 32, BCG cross-reactivity is unlikely to be the only reason behind the higher TST  
349 conversion rate. TST-only converters also appear to be older than the other converter groups.  
350 This association is present in the univariate analysis and confirmed by our logistic regression  
351 model. The positive association between TST positivity and age has been detected in previous  
352 studies across different populations and age groups (33-38). All of these studies, however, were  
353 cross-sectional in nature and could not differentiate between recent TST conversion and long-  
354 standing *Mtb* infection. Therefore, our study is the first one to identify an association between  
355 recent TST conversion and increasing age that could be possibly linked to longer environmental  
356 exposure to non-tuberculosis mycobacteria (NTM) in the setting of a BCG-primed immune  
357 response to cross-reactive mycobacterial antigens.

358           Furthermore, the use of the TST in conjunction with the QFT allowed us to identify  
359 converters who appear to have distinct immunologic reactions at the time of conversion. When  
360 we examine QFT/TST converters, we see that the QFT/TST converters have a higher change in  
361 QFT quantitative values at the time of QFT conversion than QFT-only converters. This difference  
362 of values is maintained at month 12, though this result did not attain statistical significance.  
363 Similarly, change in TST increment is larger in QFT/TST converters than in TST-only converters.  
364 Overall, these results are indicative of a more robust immunologic response in HHCs who convert  
365 both tests. This idea is supported by studies in Senegal and South Africa that found that a  
366 combination of both a positive TST and a positive IGRA appeared better at predicting TB than  
367 either response alone (39, 40). Indeed, the predictive value of TB for TST and IGRAs has been  
368 the focus of several studies and continues to be a controversial subject. Some studies have found

369 no difference between these two tests with respect to predictive ability (40-43). On the other hand,  
370 other studies have found that IGRAs were better at predicting subsequent TB (44, 45). In the  
371 end, it is unclear if the TST or the QFT on their own may offer an adequate predictive value for  
372 subsequent TB disease. However, the significance of the more robust immunologic response by  
373 combined QFT/TST converters is not clear and is deserving of further investigation or  
374 development of a better biomarker for recent and/or new Mtb infection.

375 Our analysis suffered from some limitations. First, our small sample size limited the power  
376 of the study. Second, although we obtained IGRA results at different points during the follow-up,  
377 TST was only obtained at the beginning and at the end of the follow-up.

378 In conclusion, our findings indicate that TST converters are older. If this association is  
379 due to ongoing exposure to environmental mycobacteria, as we have speculated, then vaccine  
380 design studies that only use the TST to assess endpoints will likely overestimate Mtb infection.  
381 Second, our results indicate that other epidemiological variables do not distinguish TST from  
382 IGRA converters and therefore, cannot help define “true” conversion. Third, combined QFT/TST  
383 converters appear to have a more robust immune response. Consequently, neither the TST nor  
384 IGRA alone may be sufficient to provide an adequate understanding of conversion/recent  
385 infection. Future studies using both of these tests will be needed to elucidate whether dual  
386 conversion is a useful endpoint for clinical and immunological studies.

## 387 **Acknowledgements**

388 We want to acknowledge the contributions made by senior physicians, medical officers, health  
389 visitors, laboratory and data personnel: Prof. Moses Joloba, Dr. Brenda Okware, Dr. Timothy  
390 Muwanguzi, Dorcas Lamunu, Joan Tusabe, Saidah Menya, Veronica Kizza, Druscillah Ssekandi,  
391 Joan Nassuna, Ann Ritah Namuganga, Sophie Nalukwago, Michael Odie, Henry Kawoya, Keith  
392 Chervenak, and Dr. Bonnie Thiel. The data collection and management for this paper was  
393 performed using the OpenClinica open source software, version 3.16. Copyright OpenClinica LLC  
394 and collaborators, Waltham, MA, USA, [www.OpenClinica.com](http://www.OpenClinica.com). This study would not be possible  
395 without the generous participation of the Ugandan patients and families.

## 396 **References**

- 397 1. World Health Organization. Implementing the end TB strategy: the essentials. Geneva:  
398 World Health Organization; 2016.
- 399 2. World Health Organization. Global tuberculosis report 2020. Geneva: World Health  
400 Organization; 2020.
- 401 3. Bagcchi S. WHO's Global Tuberculosis Report 2022. *Lancet Microbe*. 2023;4(1):e20.
- 402 4. Behr MA, Kaufmann E, Duffin J, Edelstein PH, Ramakrishnan L. Latent Tuberculosis:  
403 Two Centuries of Confusion. *Am J Respir Crit Care Med*. 2021;204(2):142-8.
- 404 5. Reichler MR, Khan A, Sterling TR, Zhao H, Chen B, Yuan Y, et al. Risk Factors for  
405 Tuberculosis and Effect of Preventive Therapy Among Close Contacts of Persons With  
406 Infectious Tuberculosis. *Clin Infect Dis*. 2020;70(8):1562-72.
- 407 6. Sloot R, Schim van der Loeff MF, Kouw PM, Borgdorff MW. Risk of tuberculosis after  
408 recent exposure. A 10-year follow-up study of contacts in Amsterdam. *Am J Respir Crit Care*  
409 *Med*. 2014;190(9):1044-52.
- 410 7. Behr MA, Edelstein PH, Ramakrishnan L. Revisiting the timetable of tuberculosis. *BMJ*.  
411 2018;362:k2738.
- 412 8. Zellweger JP, Sotgiu G, Corradi M, Durando P. The diagnosis of latent tuberculosis  
413 infection (LTBI): currently available tests, future developments, and perspectives to eliminate  
414 tuberculosis (TB). *Med Lav*. 2020;111(3):170-83.
- 415 9. Stein CM, Mayanja-Kizza H, Hawn TR, Boom WH. Importance of Study Design and  
416 Phenotype Definition in Ongoing Studies of Resistance to Latent *Mycobacterium tuberculosis*  
417 Infection. *J Infect Dis*. 2020;221(6):1025-6.



- 418 10. Behr MA, Edelstein PH, Ramakrishnan L. Is Mycobacterium tuberculosis infection life  
419 long? *BMJ*. 2019;367:15770.
- 420 11. Boom WH, Schaible UE, Achkar JM. The knowns and unknowns of latent  
421 Mycobacterium tuberculosis infection. *J Clin Invest*. 2021;131(3).
- 422 12. Ewer K, Deeks J, Alvarez L, Bryant G, Waller S, Andersen P, et al. Comparison of T-  
423 cell-based assay with tuberculin skin test for diagnosis of Mycobacterium tuberculosis infection  
424 in a school tuberculosis outbreak. *Lancet*. 2003;361(9364):1168-73.
- 425 13. Lalvani A, Pathan AA, McShane H, Wilkinson RJ, Latif M, Conlon CP, et al. Rapid  
426 detection of Mycobacterium tuberculosis infection by enumeration of antigen-specific T cells.  
427 *Am J Respir Crit Care Med*. 2001;163(4):824-8.
- 428 14. Brock I, Weldingh K, Lillebaek T, Follmann F, Andersen P. Comparison of tuberculin  
429 skin test and new specific blood test in tuberculosis contacts. *Am J Respir Crit Care Med*.  
430 2004;170(1):65-9.
- 431 15. Pai M, Zwerling A, Menzies D. Systematic review: T-cell-based assays for the diagnosis  
432 of latent tuberculosis infection: an update. *Ann Intern Med*. 2008;149(3):177-84.
- 433 16. Lalvani A. Diagnosing tuberculosis infection in the 21st century: new tools to tackle an  
434 old enemy. *Chest*. 2007;131(6):1898-906.
- 435 17. Drobniewski F, Cobelens F, Zellweger JP, Workshop KNE. Use of Gamma-interferon  
436 assays in low- and medium-prevalence countries in Europe: a consensus statement of a Wolfheze  
437 Workshop organised by KNCV/EuroTB, Vilnius Sept 2006. *Euro Surveill*. 2007;12(7):E070726  
438 2.

- 439 18. Chang PC, Wang PH, Chen KT. Use of the QuantiFERON-TB Gold In-Tube Test in the  
440 Diagnosis and Monitoring of Treatment Efficacy in Active Pulmonary Tuberculosis. *Int J*  
441 *Environ Res Public Health*. 2017;14(3).
- 442 19. Moses MW, Zwerling A, Cattamanchi A, Denkinger CM, Banaei N, Kik SV, et al. Serial  
443 testing for latent tuberculosis using QuantiFERON-TB Gold In-Tube: A Markov model. *Sci Rep*.  
444 2016;6:30781.
- 445 20. Stein CM, Hall NB, Malone LL, Mupere E. The household contact study design for  
446 genetic epidemiological studies of infectious diseases. *Front Genet*. 2013;4:61.
- 447 21. Garcia-Basteiro AL, White RG, Tait D, Schmidt AC, Rangaka MX, Quaife M, et al. End-  
448 point definition and trial design to advance tuberculosis vaccine development. *Eur Respir Rev*.  
449 2022;31(164).
- 450 22. Aceng JR. The Uganda National Tuberculosis Prevalence Survey, 2014 - 2015.  
451 Kampala2016.
- 452 23. Sekandi JN, List J, Luzze H, Yin XP, Dobbin K, Corso PS, et al. Yield of undetected  
453 tuberculosis and human immunodeficiency virus coinfection from active case finding in urban  
454 Uganda. *Int J Tuberc Lung Dis*. 2014;18(1):13-9.
- 455 24. Reporter RM. The Deluxe "Muzigo" - Single Bedroom House: Real Muloodi News  
456 Network; 2021 [Available from: [https://realmuloodi.co.ug/the-deluxe-muzigo-single-bedroom-](https://realmuloodi.co.ug/the-deluxe-muzigo-single-bedroom-houses/)  
457 [houses/](https://realmuloodi.co.ug/the-deluxe-muzigo-single-bedroom-houses/)].
- 458 25. Menzies D. Interpretation of repeated tuberculin tests. Boosting, conversion, and  
459 reversion. *Am J Respir Crit Care Med*. 1999;159(1):15-21.
- 460 26. Kroon EE, Kinnear CJ, Orlova M, Fischinger S, Shin S, Boolay S, et al. An observational  
461 study identifying highly tuberculosis-exposed, HIV-1-positive but persistently TB, tuberculin

- 462 and IGRA negative persons with *M. tuberculosis* specific antibodies in Cape Town, South  
463 Africa. *EBioMedicine*. 2020;61:103053.
- 464 27. Lu LL, Smith MT, Yu KKQ, Luedemann C, Suscovich TJ, Grace PS, et al. IFN-gamma-  
465 independent immune markers of *Mycobacterium tuberculosis* exposure. *Nat Med*.  
466 2019;25(6):977-87.
- 467 28. Mandalakas AM, Kirchner HL, Lombard C, Walzl G, Grewal HM, Gie RP, et al. Well-  
468 quantified tuberculosis exposure is a reliable surrogate measure of tuberculosis infection. *Int J*  
469 *Tuberc Lung Dis*. 2012;16(8):1033-9.
- 470 29. Ma N, Zalwango S, Malone LL, Nsereko M, Wampande EM, Thiel BA, et al. Clinical  
471 and epidemiological characteristics of individuals resistant to *M. tuberculosis* infection in a  
472 longitudinal TB household contact study in Kampala, Uganda. *BMC Infect Dis*. 2014;14:352.
- 473 30. R Development Core Team. R: A language and environment for statistical computing.  
474 Vienna, Austria: R Foundation for Statistical Computing; 2010.
- 475 31. Ribeiro-Rodrigues R, Kim S, Coelho da Silva FD, Uzelac A, Collins L, Palaci M, et al.  
476 Discordance of tuberculin skin test and interferon gamma release assay in recently exposed  
477 household contacts of pulmonary TB cases in Brazil. *PLoS One*. 2014;9(5):e96564.
- 478 32. Farhat M, Greenaway C, Pai M, Menzies D. False-positive tuberculin skin tests: what is  
479 the absolute effect of BCG and non-tuberculous mycobacteria? *Int J Tuberc Lung Dis*.  
480 2006;10(11):1192-204.
- 481 33. Hemmati M, Ghadiri K, Rezaei M. Tuberculin Reactivity in School Age Children; Five-  
482 year Follow-up in Iran. *Iran J Pediatr*. 2011;21(1):39-44.

- 483 34. Pong AL, Anders BJ, Moser KS, Starkey M, Gassmann A, Besser RE. Tuberculosis  
484 screening at 2 San Diego high schools with high-risk populations. *Arch Pediatr Adolesc Med*.  
485 1998;152(7):646-50.
- 486 35. Martinez L, Arman A, Haveman N, Lundgren A, Cabrera L, Evans CA, et al. Changes in  
487 tuberculin skin test positivity over 20 years in periurban shantytowns in Lima, Peru. *Am J Trop*  
488 *Med Hyg*. 2013;89(3):507-15.
- 489 36. Saito M, Bautista CT, Gilman RH, Bowering A, Levy MZ, Evans CA. The value of  
490 counting BCG scars for interpretation of tuberculin skin tests in a tuberculosis hyperendemic  
491 shantytown, Peru. *Int J Tuberc Lung Dis*. 2004;8(7):842-7.
- 492 37. Mori T, Leung CC. Tuberculosis in the global aging population. *Infect Dis Clin North*  
493 *Am*. 2010;24(3):751-68.
- 494 38. Chan-Yeung M, Dai DL, Cheung AH, Chan FH, Kam KM, Tam CM, et al. Tuberculin  
495 skin test reaction and body mass index in old age home residents in Hong Kong. *J Am Geriatr*  
496 *Soc*. 2007;55(10):1592-7.
- 497 39. Lienhardt C, Fielding K, Hane AA, Niang A, Ndao CT, Karam F, et al. Evaluation of the  
498 prognostic value of IFN-gamma release assay and tuberculin skin test in household contacts of  
499 infectious tuberculosis cases in Senegal. *PLoS One*. 2010;5(5):e10508.
- 500 40. Mahomed H, Hawkrige T, Verver S, Abrahams D, Geiter L, Hatherill M, et al. The  
501 tuberculin skin test versus QuantiFERON TB Gold(R) in predicting tuberculosis disease in an  
502 adolescent cohort study in South Africa. *PLoS One*. 2011;6(3):e17984.
- 503 41. Bakir M, Millington KA, Soysal A, Deeks JJ, Efee S, Aslan Y, et al. Prognostic value of  
504 a T-cell-based, interferon-gamma biomarker in children with tuberculosis contact. *Ann Intern*  
505 *Med*. 2008;149(11):777-87.

- 506 42. Harstad I, Winje BA, Heldal E, Oftung F, Jacobsen GW. Predictive values of  
507 QuantiFERON-TB Gold testing in screening for tuberculosis disease in asylum seekers. *Int J*  
508 *Tuberc Lung Dis.* 2010;14(9):1209-11.
- 509 43. Kik SV, Franken WP, Mensen M, Cobelens FG, Kamphorst M, Arend SM, et al.  
510 Predictive value for progression to tuberculosis by IGRA and TST in immigrant contacts. *Eur*  
511 *Respir J.* 2010;35(6):1346-53.
- 512 44. Diel R, Loddenkemper R, Niemann S, Meywald-Walter K, Nienhaus A. Negative and  
513 positive predictive value of a whole-blood interferon-gamma release assay for developing active  
514 tuberculosis: an update. *Am J Respir Crit Care Med.* 2011;183(1):88-95.
- 515 45. Leung CC, Yam WC, Yew WW, Ho PL, Tam CM, Law WS, et al. T-Spot.TB  
516 outperforms tuberculin skin test in predicting tuberculosis disease. *Am J Respir Crit Care Med.*  
517 2010;182(6):834-40.

## 518 **Supporting information**

519 **S1 Fig. Study protocol for the Kampala household contact study.**

520 **S2 Fig. Visualization of cluster model based on partitioning around medoids using two**  
521 **clusters.**

522 **S1 Table. Univariate analysis comparing QFT-only, TST-only and QFT/TST converters**  
523 **based on individual and household characteristics.** For this comparison, QFT-only  
524 converters and QFT/TST converters were combined into a single category, QFT converters.

525 **S2 Table. Univariate analysis comparing QFT-only, TST-only and QFT/TST converters**  
526 **based on index case characteristics.** For this comparison, QFT-only converters and  
527 QFT/TST converters were combined into a single category, QFT converters.

**Fig 1. CONSORT diagram for this study.** Out of 383 household contacts enrolled in our study, 69 were excluded. 314 household contacts were classified using both TST and QFT. Those who were defined as concordant LTBI (both TST and QFT results were consistently positive) and discordant "resisters" (either 2 negative TST or all negative QFT) were excluded from analysis (n=166). The remaining 148 household contacts were defined as follows: 39 TST-only converters (negative baseline TST with positive TST at 12 months), 13 were QFT-only converters (negative QFT at baseline followed by positive QFT during follow-up), 24 were QFT/TST converters (negative baseline TST and QFT followed by positive TST and QFT during follow-up), and 72 were concordant short-term "resisters" (all TST and QFT tests remained negative).




Fig 1

**Fig 2. Quantitative QFT values of QFT/TST converters and QFT-only converters before and after QFT-conversion.** Red dotted line indicates 0.35 IU cut-off. If QFT plus was used, the highest value was used for this analysis. QFT-only converters had a lower value at conversion when compared to QFT/TST converters (2.03 [0.92-3.82] vs. 4.14 [1.73-10.0],  $p=0.03$ ).




Fig 2

**Fig 3. Quantitative TST values of QFT/TST converters and TST-only converters at baseline and at month 12 of follow-up.** Red dotted line indicates 10 mm. cut-off. At baseline, TST-only converters had a higher TST value compared to QFT/TST converters (2.0 [0.0-8.0] vs. 0.0 [0.0-0.0],  $p=0.001$ ). By month 12, this difference had disappeared (14.6 [12.2-16.3] vs. 15.6 [13.9-17.6],  $p=0.25$ ).




Fig 3