1	Screen-based media device use, its associated factors and sleep
2	outcomes among undergraduate health students studying in
3	colleges of Kathmandu valley, Nepal
4	Karina Maharjan ¹ , Rajan Paudel ¹ , Kshitij Kunwar ^{1*}
5	
6	
7	
8	¹ Central Department of Public Health, Institute of Medicine, Tribhuvan University,
9	Kathmandu, Nepal
10	
11	
12	*Corresponding author:

E-mail: kshitijkunwar7@gmail.com (KK)

Abstract

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

People of all age groups are engaged with various screen-based devices. Use of screen-based media devices beyond duration considered problematic has become a problem worldwide. Yet, undergraduate students are engaged in various screen-based activities for academic works and entertainment which can influence their sleep. A descriptive cross-sectional study design was used to collect information from 384 undergraduate students using a web-based questionnaire and convenience sampling technique. The data collected was analyzed using IBM SPSS Statistics 20.0. Descriptive and regression analyses were done to explore factors that influence screen-based media use. The average duration of screen-based media use by the participants was found to be 7.12 hours per day. Factors such as sex of the participants, the use of screen-based media during major mealtimes and home-specific rules limiting screen-based media use were found to be associated with screen-based media use. Only 45.3% of participants got the recommended hours of sleep (7 or more hours). Around 9.4% of the participants had long sleep onset latency (more than 45 minutes). Participants who used screen-based media devices during night before sleep were more likely to get insufficient sleep (OR (Odds Ratio) 3.668, 95% CI (Confidence Interval) (1.784-7.539)). A significant association was found between sex of the participants, use of screen-based media devices during major meal times and home rules limiting screen-based media use with the use of screen-based media for five hours or more per day. Screen-based media use and screen-based media use during night influence the quantity of sleep. Overuse of screen-based media is degrading the quality of sleep among undergraduate health students and is an alarming problem. There is a need to promote good sleeping habits through the reduction of

- 37 this modifiable behaviour, which could be done by promoting home-based rules about
- 38 screen-based media devices use, especially at night before sleep.
- 39 **Keywords:** Screen time, screen-based media use, screen-based media use and sleep, sleep
- 40 quality, sleep quantity

Introduction

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

In the last decade there has been a sharp increase in availability and use of screen-based media devices such as television, laptop, smartphones, videogame consoles and computers and as a result people are exposed to unintentional artificial light [1]. Having access to these technologies has become an important part of everyday life. Most of the adults have at least one electronic media device in their bedroom. These electronic devices play an important part to support education of students and have also become a medium of entertainment. Nowadays people spend most of their leisure time using screen-based media devices. Not only during leisure time but they also tend to use it alongside performing any other work. College students are found engaged with various screen-based devices. They are engaged in various screen-based media related activities for the purpose of entertainment or works related to academics. Also, those devices are used to attend classes remotely, complete various assignments and assignments and reading materials are also distributed in forms that can be accessed through screen-based devices. Academic works are also increasing screen time of college students. It is seen that nine out of every ten college students use laptop or phone on a regular basis [2]. Use of screen-based media devices above recommended duration has become a problem worldwide. People of all age groups including children, adolescents, adults or elderly are engaged in various screen-based activities. While waking a teenager in the morning, there is higher odds that we will find a cell phone under their pillow. It was probably the last thing that they used before sleeping and probably the first thing that they will touch after waking up [3]. Screen-based media use is affected by various factors that may be individual factors i.e. gender, physical activity, home physical factors i.e. availability, location and accessibility of devices and social factors i.e. rules and limitation, TV co-viewing [4]. Screen time usage

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

among students is due to academic requirement and need for being connected socially through electronic devices [2]. A study on technology use among college students found that students are found spending an average of 12 hours daily in some types of media among which nine and a half hours are spent with their gadgets like mobile devices, gaming devices, computers, etc [5]. Mostly people are awake 16 to 18 hours a day, among which spending 12 hours in electronic devices is a lot. Sleep is an important part of human life. It is estimated that humans spend a third of their lifetime sleeping or attempting to do so [6]. Sufficient sleep duration is required for healthy growth and development of the mind and body especially for children and adolescents. College students should get about seven to nine hours of sleep [7]. Getting required amount of sleep helps improve grades, lowers the risk of obesity, improves memory, improves mood. Active use of screen-based media among college going students especially night time use affects sleep in a negative way, it can be a factor for short sleep duration, disturbed sleep, increased sleep onset latency and delayed bed, wake up time. It causes sleep deprivation among them. College student may become sleep deprived due to various factors such as lighting, sleep environment, medications, body pain, aches, stress, shift based work [8]. Among these factors screen time and excessive use of light emitting device are common factors [9]. Surveys have shown the association of screen-based media use with sleep [6,9– 12]. Sleep quality is found negatively correlated with hours spent on electronic devices [13]. Use of media before sleep is an ascribed factor for insufficient sleep among younger people and adult [14]. Studies have shown adverse association between screen-based media use and sleep health where delayed bedtime and shorter total sleep have been found to be most consistently related to media use [15–17].

Poor sleep can negatively affect academic performance. Inadequate sleep is also associated with cardiovascular, inflammatory and metabolic consequences. Insufficient sleep to an individual is shown to be associated with impaired cellular responses, depression, anxiety and obesity [6]. Therefore, this study aimed and has determined screen-based media device use, its associated factors and sleep outcomes among undergraduate health students from Kathmandu valley, Nepal.

Methods

Study design and setting including study site, design, sample size

and sampling

The descriptive cross-sectional design was designed to assess the screen-based media device use, its associated factors and sleep outcomes among the undergraduate students of health background studying in colleges within Kathmandu valley (Kathmandu, Bhaktapur and Lalitpur districts). The respondents included Bachelor of Medicine and Bachelor of Surgery (MBBS), Bachelor of Dental Surgery (BDS), Bachelor in Public Health (BPH), Bachelor of Science in Medical Laboratory Technology (BSc. MLT), Bachelor in Pharmacy (B. Pharm), Bachelor of Science in Nursing (Bsc. Nursing), Bachelor of Nursing Science (BNS) and Bachelor in Optometry (B. optom), BPT (Bachelor in Physiotherapy), BASLP (Bachelor of Audiology & Speech Language Pathology) students. Data was collected from the undergraduates remotely by administering web-based questionnaire considering the risk of infection from COVID-19 by mobilizing the students' network online.

The sample size was calculated using formula, n=Z^2p(1-p)/e^2 and sample size was calculated to be 347, assuming 10% non-response rate it was fixed as 382.

Data collection

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

The questionnaire was adopted from previous studies Questionnaire regarding sleep was adapted based on Holly Scott [12] and questionnaire regarding screen-based media use and factors associated were adapted from Granich et.al [4]. The questionnaire had four sections: socio-demographic information, home media environment, screen time and sleep outcomes. The main outcome variable were screen time and factors affecting screen-based media use. Screen time referred to the total hours spent using screen-based media devices. Pretesting was carried out among 40 undergraduate students. The data was collected from 15-JAN-2021 to 28-FEB-2021. Ethical approval for the study was obtained from the Institutional Review Board (IRB), Maharajguni Medical Campus, Institute of Medicine, Tribhuwan University, Kathmandu, Nepal (Approval letter reference number: 201(6-11) E2 077/078). Participants were informed about the process, objective of the study, their privacy and confidentiality of the information they provide. Participants provided informed consent in the same web-based form before they could take part in the study by answering a question related to agreeing or disagreeing to take part in the study which they could voluntarily choose as "Yes" or "No". Data processing and analysis Data collected online was downloaded and Microsoft Excel was used to clean the data and was exported to IBM SPSS Statistics 20.0 for further coding and analysis. Information of independent variables were analysed using descriptive statistics namely frequency, percentage, median. Inferential statistics using Chi-square test was done to find out the association between

screen-based media use and independent variables and logistic regression was performed to

- determine the strength of association. Confidence interval of 95% was used and p-value
- 135 < 0.05 was considered statistically significant.

Results

134

136

- Table 1 shows the demographic characteristics 384 participants of the study which included
- 41.1% male and 58.9% female participants. The median age of the participants was 22 years.

Table 1: Demographic characteristics of participants (N=384)

Characteristics	Number	Percentage
Age Group (in years)		
< 20	51	13.3
20 or above	133	86.7
Median age	22	
Interquartile Range	20-23	
Sex		
Male	158	41.1
Female	226	58.9
Faculty		
ВРН	119	31.0
B. Pharm	57	14.8
MBBS	37	9.6
BDS	48	12.5
BNS or Bsc. Nursing	72	18.8
Others (Bsc MLT, B.Optom, BPT,	51	13.3
BASLP etc.)		
Study Year		
First year (1st and 2nd semester)	100	26.0
Second year (3 rd and 4 th semester)	71	18.5
Third year (5 th and 6 th semester)	85	22.1
Fourth year and above (7th semester and	128	33.3
above)		

University		
Tribhuvan University	262	68.2
Kathmandu University	46	12.0
Pokhara University	27	7.0
Purbanchal University	37	9.6
Patan Academy of Health Sciences	12	3.1
Father's Education		
Upto secondary level	114	29.7
Higher than Secondary level	270	70.3
Mother's Education		
Illiterate (can't read and write)	40	10.4
Upto secondary level	167	43.5
Higher than Secondary level	177	46.1
Monthly Income of Family (in Nepali		
Rupees)		
Less than 25,000	111	28.9
25,000 - 50,000	141	36.7
More than 50,000	132	34.4

The household media environment plays role in screen-based media behaviour. It was found that all participants come with a household of more than three media device of some kind and 79.4% participants had their own screen-based media devices as shown in Table 2. It was found that 21.6% of the participant's household had some rules limiting the screen-based media use such as allocated screen time and in 39.6% participant's homes, screen-based media devices were being viewed during major meals. Participant's household with 3 or more variety of screen-based media device among the five types taken under this study was found to be 92.4%.

140

141

142

143

144

145

146

147

Table 2: Household media environment and screen-based media use (N=384)

Household media environment and screen-	Number	Percentage
based media use		
Have internet connection	349	90.9
Have access to TV channels	320	83.3
Have own screen-based media device	305	79.4
Presence of home Rules/Regulations about media	83	21.6
use		
Use screen-based media during major meals	152	39.6
(breakfast, lunch, dinner)		
Number of types of screen-based media		
devices present in the household		
Less than 3 types	29	7.6
Equal to or more than 3 types	355	92.4
Number of types of screen-based media device		
owned by the respondent		
None	79	20.6
One type	90	23.4
Two or more types	215	56
Average screen-based media use in 24 hours		
≥5 hours	294	76.6
<5 hours	90	23.4
Average duration of screen-based media use	7.12 hours	
per 24 hours		
Screen-based media use before bedtime		
Doesn't use screen-based media	43	11.2

341	88.8	
Duration of screen-based media use before		
178	46.3	
117	30.5	
46	12.0	
	178 117	

The major study variable of this research was screen-based media use among undergraduate students of the study area. Participants were asked total hours of use of screen-based media devices in 24 hours. The responses were then classified as <5 hours and ≥ 5 hours of use, on the basis of association seen with screen-based media use for more than or equals to five hours with computer vision syndrome.

Table 2 shows that 76.6 % participants spend more than or equals to 5 hours of their time being engaged in some kind of screen-based media activity in 24 hours. The average hours of screen-based media device use among the study participants was 7.12 hours per day (24 hours). Majority (69.5%) participants responded that they prefer to use smartphones over other screen-based medias. As for the purpose of use, majority (46.1%) of the participants reported using screen-based media devices for entertaining themselves.

Table 3 shows the sleep related outcomes of the participants with average sleep of 6.72 hours. Sleep onset latency refers to the time required for an individual to fall asleep, it was found that 9.4% participants have long SOL (Sleep Onset Latency) (more than 45 minutes).

Table 3: Sleep outcomes of participants (N=384)

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

Sleep Outcomes	Number	Percentage
Sleep time		
Before 11 PM	214	55.7

11 PM and after	170	44.3
Sleep Onset Latency (SOL)		
≤ 45 minutes	348	90.6
> 45 minutes	36	9.4
Total sleep duration		
<7 hours	210	54.7
≥ 7 hours	174	45.3
Average hours of sleep	6.72 hours	
Range of total hours of sleep	2-11 hours	

Results from the study showed that screen-based media use was associated with sex of the participants, mealtime viewing of the screen-based media and rules limiting screen-based media use as shown in Table 4.

Table 4: Association between variables and screen-based media use

167

168

169

170

171

Characteristics	Screen-based media use		p-value
	≥ 5 hours (%)	< 5 hours (%)	
Age in years			
<20	38 (74.5)	13 (25.5)	0.710
20 and above	256 (76.9)	77 (23.1)	
Sex			
Male	138 (87.3)	20 (12.7)	<0.001
Female	156 (69)	70 (31)	
Own personal device			
Yes	237 (77.7)	68 (22.3)	0.300
No	57 (72.2)	22 (27.8)	
Has internet Access			
Yes	264 (75.6)	85 (24.4)	0.187
No	30 (85.7)	5 (14.3)	
Availability of TV channels			
Available	240 (75)	80 (25)	0.110

Not available	54 (84.4)	10 (15.6)	
Uses screen-based media			
device during major meal tin	ne		
Yes	126 (82.9)	26 (17.1)	0.019
No	168 (72.4)	64 (27.6)	
Presence of Rule about using	5		
screen-based media device a	t		
household			
Yes	56 (67.5)	27 (17.1)	0.029
No	238 (79.1)	63 (20.9)	
Uses screen-based media dev	vice in bedroom		
Yes	156 (77.2)	46 (22.8)	0.746
No	138(75.8)	44 (24.2)	
Number of types of devices			
owned by the respondent			
Less than three types	25 (86.2)	4 (13.8)	0.210
Three or more types	269 (75.8)	86 (24.2)	

This study found that there was no association of personal media ownership and electronic media device in bedroom with night time screen-based media use.

173

174

175

176

177

178

179

180

181

The results of logistic regression analyses are shown in Table 5. Female participants were less likely to have more than or equal to 5 hours screen time (AOR (Adjusted Odds Ratio) 0.356, 95% CI (1.200-0.635)) than male participants. Participant who do not view screenbased media devices while having meal was less likely to have screen time of more than or equal to 5 hours (AOR 0.474, 95% CI (0.277 - 0.812)) then those who don't. Likewise, participants not having home rules limiting screen time were more likely to have screen time

182 of more than or equal to 5 hours (AOR 1.788, 95% CI (1.013-3.155)) than those who have 183 rules.

Table 5: Logistic regression for factors of screen use for more than recommended hours (≥ 5 hours)

Characteristics U	(nadjusted OR (95% CI)	Adjusted OR (95%
		CI)
Age in years		
< 20	1	1
20 and above	1.137 (0.577-2.244)	1.175 (0.578-2.389)
Sex		
Male	1	1
Female	0.323 (0.187-0.558)	0.356 (0.200-0.635)
Owns personal device		
Yes	1	1
No	0.743 (0.424 – 1.303)	0.687 (0.373 – 1.267)
Has internet access		
Yes	1	1
No	1.932 (0.727-5.136)	1.363 (0.462 – 4.016)
Availability of TV channels		
Yes	1	1
No	1.800 (0.876-3.700)	1.203 (0.548-2.639)
Use of screen-based media device du	ıring	
major meal time		
Yes	1	1
No	0.542 (0.325-0.903)	0.474 (0.277 - 0.812)
Presence of rule at home about usin	g	
screen-based media device		
Yes	1	1
No	1.821 (1.065-3.115)	1.788 (1.013-3.155)

Uses screen-based media device in bedroom

Yes	1	1
No	0.925 (0.577-1.483)	1.015 (0.611 – 1.686)

Number of types of devices owned by the

respondent

186

187

188

189

190

191

192

193

194

Less than three types 1 1 Three or more types 0.748 (0.318-1.761) 6.202 (2.267-16.964)

Regression analysis presented in Table 6 showed that participants who use screen-based media devices for ≥ 5 hours a day were more likely to get inadequate sleep (AOR 3.898, 95%) CI (2.325-6.534)) than those who use screen-based media for less than 5 hours a day. Likewise, participants who view screen-based media device before sleeping were more likely to get inadequate sleep (AOR 3.668, 95% CI (1.784-7.539)) than those who do not view screen-based media device before sleep.

Table 6: Logistic regression for factors of inadequate sleep (<7 hours)

Characteristics	Unadjusted OR (95% CI)	Adjusted OR (95% CI)
Duration of use of screen-based		
media		
\geq 5 hours	3.847(2.313 - 6.398)	3.898 (2.325-6.534)
< 5 hours	1	1
Night time screen-based media		
use		
Yes	3.567 (1.776-7.205)	3.668 (1.784-7.539)
No	1	1

Discussion

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

This study was carried out to find out the screen-based media use, factors associated with it and sleep related outcomes among health undergraduate students studying in colleges of Kathmandu valley. The study has assessed the present scenario of screen-based media use among undergraduate students and their sleep outcomes. This study found that undergraduate students spend an average of 7.2 hours in a day using some form of screen-based media device which is similar to findings from a study done by Kaiser Family Foundation that showed average use screen-based media device for 7.5 hours per day among young adolescents [3]. Similar study done by Yen-Sen Lee et al, found that the students spend an average of 8.3 hours in a day using screen-based devices [13]. This study showed that most of the participants are engaged in screen-based activity during evening time. A study by Angela Lynn Sargent on screen time and sleep condition among selected college students showed that 75% students reported always use of electronic media device one hour before bedtime [18]. Similarly, this study showed that 88.8% participants reported using screen-based media device before sleeping. A study by Granich and Kaiser Family Foundation studied the relationship of household screen-based media and screen-based media use among adolescents. Both study have concluded that there is positive association between availability and accessibility of media devices with time spent on screen-based media use [3,4]. In contrast to this study, the current study showed no association between accessibility and availability of screen-based media device with its use. Though, significant association was seen between having home rules limiting screen-based media use and meal time viewing of screen-based media devices with screen-based media use among the study participants. The percentage of individuals with an internet connection was recorded to be 36.7% in 2021 [19]. In contrast to this, the household with internet access were measured to be 90.9% in this study. This could be due to the fact

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

that the sample was based on the urban population of Kathmandu valley. The entire participant's household owned more than three screen-based media devices of any kind and 79.4% participants had their own screen-based media device, the most commonly owned device was smartphones. A study by Angela Lynn Sargent on screen time and sleep condition among selected college students showed that students on average get 6.9 hours of sleep. It also showed that 65.5% of the participants reported getting recommended 7 hours or more sleep. The study found no statistically significant difference regarding screen use before bedtime and participants hours of sleep [18]. A study carried out to see the association between portable screen-based media use and sleep outcome showed that participants who had access to media device at night were more likely to have inadequate sleep quantity, OR 1.79, 95% CI 1.39-2.31 [11]. Similarly, the current study found out that the participants got an average of 6.72 hours of sleep and also found that only 45.3% participants got recommended hours of sleep. Similar to the previous study, significant association was found between night time screen-based media use and participants sleep. A study by Nowell SB et al studied the relationship between social media use and sleep quality of undergraduate nursing students showed that sleep duration among students ranges from 4 to 10 hours, the study also showed that electronic media use correlates with poor sleep quality [20]. This study shows that the sleep duration of participants ranges from 2 to 11 hours. The data presented in the study were self-reported and partly depend on the participant recall ability thus they may subject to recall bias. This study is limited to factors affecting screenbased media use among undergraduates, sleep outcomes and it does not address any health effects due to sleep inadequacy. There are many physical, psychological, behavioural and emotional aspects of screen time as well but the variables that were studied in this study were focused only on sleep related outcomes. Also, there are many other factors affecting sleep outcomes of an individual beside screen-based media use but only screen-based media was

studied as a factor affecting sleep.

Conclusion

This study showed that the participants on average use screen-based media devices for 7.2 hours per day and only 45.3 % participants got recommended hours of sleep (7 or more hours). Around 9.4% of the participants had long sleep onset latency (more than 45 minutes). Participants who use screen-based media devices during night time before sleep were more likely to get insufficient sleep (AOR 3.668, 95% CI (1.784-7.539)).

Significant association was found between screen-based media use and variables such as sex of participant, use of screen-based media device during major meal time and home rules limiting screen-based media. Significant association was found between sleep adequacy with screen-based media use and night time screen-based media use. Thus, this research suggests the need to limit the use of screen-based media devices to promote quality of sleep. Overuse of screen-based media stands as an alarming problem among undergraduates and this can have direct influence on their physical and mental health. Limiting the use of screen-based media devices through the promotion of home-specific rules about screen-based media use especially about the use at night time before sleep can help undergraduate students get the recommended amount of sleep.

Acknowledgement

We would like to thank all respondents for their participation and academics of Institute of

medicine for their academic guidance and support for this study.

Conflict of interest

None declared. 267

266

268

Author's contribution

- 269 Conceptualization: KM,
- 270 Data curation: KM, KK
- 271 Formal analysis: KM,
- 272 Investigation: KM,
- 273 Methodology: KM, RP,
- 274 Project administration: KM,
- 275 Resources: KM,
- 276 Supervision: RP,
- 277 Validation: RP, KK,
- 278 Writing – original draft: KM,
- 279 Writing – reviewing & editing: KK, KM, RP

References 280

- 1. Green A, Cohen-Zion M, Dagan Y. Evening light exposure to computer screens 281
- 282 disrupts human sleep, biological rhythms and attention abilities. Chronobiol Int. 2017;
- DOI: 10.1080/07420528.2017.1324878 283
- 284 2. 2015-Pearson-Student-Mobile-Device-Survey-College.pdf
- 285 https://www.pearson.com/content/dam/one-dot-com/one-dot-com/ped-blogs/wp-
- content/pdfs/2015-Pearson-Student-Mobile-Device-Survey-College.pdf 286

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

3. Generation M2, Media in the lives of 8 to 18 years old, A Kaiser family foundation study, January 2010. Generation M2: Media in the Lives of 8- to 18-Year-Olds -Report (kff.org) 4. Granich J. Rosenberg M. Knuiman MW. Timperio A. Individual, social, and physical environment factors associated with electronic media use among children: sedentary behavior at home. J Phys Act Health. 2011 Jul;8(5):613–25, DOI: 10.1123/jpah.8.5.613 5. Like, Totally Wired. https://www.mediapost.com/publications/article/117012/liketotally-wired.html 6. Mireku MO, Barker MM, Mutz J, Dumontheil I, Thomas MSC, Röösli M, et al. Night-time screen-based media device use and adolescents' sleep and health-related quality of life. Environ Int. 2019;124:66–78. DOI: 10.1016/j.envint.2018.11.069 7. Writers S. Expert Sleep Advice for College Students & Teens. https://www.affordablecollegesonline.org/college-resource-center/guide-to-sleepingfor-college-students-and-teens/ 8. Factors that Interfere with Healthy Sleep - ResMed India. https://www.resmed.co.in/healthy-sleep/factors-that-interfere 9. Hysing M, Pallesen S, Stormark KM, Jakobsen R, Lundervold AJ, Sivertsen B. Sleep and use of electronic devices in adolescence: results from a large population-based study. BMJ Open. 2015 Feb 2;5(1):e006748. DOI: 10.1136/bmjopen-2014-006748 10. Arora T, Broglia E, Thomas GN, Taheri S. Associations between specific technologies and adolescent sleep quantity, sleep quality, and parasomnias. Sleep Med. 2013 Jan 1;15. DOI: 10.1016/j.sleep.2013.08.799 11. Carter B, Rees P, Hale L, Bhattacharjee D, Paradkar MS. Association Between Portable Screen-Based Media Device Access or Use and Sleep Outcomes: A

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

Systematic Review and Meta-analysis. JAMA Pediatr. 2016 01;170(12):1202-8. DOI: 10.1001/jamapediatrics.2016.2341 12. Woods HC, Scott H. Social media use in adolescence is associated with poor sleep quality, anxiety, depression and low self-esteem. DOI: 10.1016/j.adolescence.2016.05.008 13. Lee Y-S, Low B-S. Sleep pattern among electronic device users and its relationship with users' practice in Malaysia university community. Sleep Biol Rhythms. 2016 Jul 1;14(3):271-7. DOI: 10.1007/s41105-016-0052-z 14. Suganuma N et.al Using electronic media before sleep can curtail sleep time and result in self-perceived insufficient sleep. DOI:10.1111/j.1479-8425.2007.00276.x 15. Cain N, Gradisar M. Electronic media use and sleep in school-aged children and adolescents: A review. Sleep Med. 2010 Sep;11(8):735-42. DOI: 10.1016/j.sleep.2010.02.006 16. Hale L, Kirschen GW, LeBourgeois MK, Gradisar M, Garrison MM, Montgomery-Downs H, et al. Youth Screen-based media Habits and Sleep: Sleep-Friendly Screen Behavior Recommendations for Clinicians, Educators, and Parents. Child Adolesc Psychiatr Clin N Am. 2018;27(2):229–45. DOI: 10.1016/j.chc.2017.11.014 17. Asaoka S, Kamada Y et.al. Exploring the Daily Activities Associated with Delayed Bedtime of Japanese University Students. https://www.jstage.jst.go.jp/article/tjem/221/3/221 3 245/ pdf 18. Sargent Angela Lynn. Screen Time and Sleep Condition among Selected College Students. https://cornerstone.lib.mnsu.edu/cgi/viewcontent.cgi?article=1679&context=etds 19. Digital in Nepal: All the Statistics You Need in 2021. DataReportal – Global Digital Insights. https://datareportal.com/reports/digital-2021-nepal

20. Nowell SB, Thompson K. Relationship between Social Media Use and Sleep Quality 337

of Undergraduate Nursing Students at a Southeastern University.

Supporting information

- 340 S1 Dataset. Study dataset and data labels
- 341 S2 Appendix. Study questionnaire

- 342 S3 Appendix. Results of regression analysis
- 343 S4 File. Institutional Review Committee approval letter