

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

Social risk factors for pediatric asthma exacerbations: A systematic review

Shriya Vinjimoor BS^{1,3} | Caroline Vieira BA^{2,3} | Colin Rogerson MD, MPH^{1,3} | Arthur Owora MPH, PhD^{3,4} | Eneida A Mendonca MD, PhD^{1,3}

¹Indiana University School of Medicine, ²Indiana University Bloomington, ³Regenstrief Institute for Biomedical Informatics, ⁴Indiana University School of Public Health

Abstract

Objective. This systematic review aims to identify social risk factors that influence pediatric

asthma exacerbations. **Methods.** Cohort studies published between 2010 and 2020 were

systematically searched on the OVID Medline, Embase, and PsycInfo databases. Using our

established phased inclusion and exclusion criteria, studies that did not address a pediatric

population, social risk factors, and asthma exacerbations were excluded. Out of a total of 707

initially retrieved articles, 3 prospective cohort and 6 retrospective cohort studies were included.

Results. Upon analysis of our retrieved studies, two overarching domains of social determinants,

as defined by Healthy People 2030, were identified as major risk factors for pediatric asthma

exacerbations: Social/Community Context and Neighborhood/Built Environment.

Social/Community factors including African American race and inadequate caregiver

perceptions were associated with increased risk for asthma exacerbations. Patients in high-risk

neighborhoods, defined by lower levels of education, housing, and employment, had higher rates

of emergency department readmissions and extended duration of stay. Additionally, a synergistic

interaction between the two domains was found such that patients with public or no health

insurance and residence in high-risk neighborhoods were associated with excess hospital

utilization attributable to pediatric asthma exacerbations. **Conclusion.** Social risk factors play a

significant role in influencing the frequency and severity of pediatric asthma exacerbations.

Keywords: socioeconomic factor, respiratory, neighborhood characteristics, child, social

determinants of health

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

3-Question Summary Box:

1. What is the current understanding of this subject?

The individual impact of social factors such as insurance, neighborhood, and ethnicity on pediatric asthma exacerbations has previously been explored.

2. What does this report add to the literature?

This review systematically identifies the relative importance of individual sociodemographic factors and interactions between them. Race, neighborhood risk, insurance status, caregiver perceptions, and a synergistic interaction between health insurance status and neighborhood risk were found to be contributory.

3. What are the implications for public health practice?

It is important for providers to educate patients on how their surroundings impact their respiratory health and advocate for increased healthcare access for at-risk populations.

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

Introduction

Social factors have a major influence on public health outcomes among children. Asthma is the most prevalent chronic illness among children globally, with around 1 in 12 American children under the age of 18 affected¹. In 2019 alone, over 44% of pediatric asthma patients reported experiencing one or more acute asthma exacerbations². Although asthma often presents with a relapsing-remitting pattern, the frequency of asthma attacks typically decreases by mid-childhood for most pediatric asthma patients³. However, for some patients, acute asthma exacerbations are persistent throughout childhood, resulting in frequent physician visits and medication costs. Asthma patients face serious financial burdens, with an average of over \$3,266 of out-of-pocket healthcare expenditures per patient annually. In addition, these patients often face frequent school absences, mental health challenges, and a diminished quality of life⁴.

In a retrospective cohort study by Keet et al. investigating over 1 million children with asthma enrolled on Medicaid, it was found that children residing in inner-city, urban, and poorer neighborhoods, and those with Hispanic and non-Hispanic black ethnicities had a higher risk of asthma-related emergency department visits⁵. Similarly, Harrington et al found that lower parent health literacy was associated with worse pediatric asthma control⁶.

While there is an abundance of such studies evaluating the influence of individual social determinants of health on asthma, little work has been done to synthesize the relative importance of these factors in altering the risk of acute pediatric asthma exacerbations. According to Healthy People 2030, social determinants of health can be synthesized under 5 overarching categories: economic stability, education access and quality, healthcare access and quality, neighborhood and built environment, and social and community context⁷.

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

This systematic review synthesizes social factors that are associated with the risk of asthma exacerbations among pediatric patients aged 2-18 years and categorizes them based on the Healthy People 2030 classifications.

Methods

This systematic review follows the 2020 PRISMA Guidelines for Systematic Reviews.

Search strategy

Development Process: In collaboration with a medical librarian, seven known relevant seed studies were used to identify records within databases. Preliminary search terms were gathered through identifying keywords in the titles and abstracts of the seven seed studies. A pilot search strategy was developed using these keywords and additional search terms were added based on relevance to these first set of terms. The search was limited to English-language only articles. The search strategy was validated by its ability to identify the seven seed studies on the PubMed database. All seven studies were identified.

Final Search Strategy: Databases were systematically searched as of November 2, 2021. The search was limited to English-only manuscripts published between 2010 and 2020. Following a pilot PubMed search, keywords, and MeSH terms for the 3 overarching concepts of asthma, pediatrics, and social factors were searched in the following databases: OVID Medline, Embase, and PsycInfo.

The complete line-by-line search strategy with all searched keywords and MeSH terms in each database is accessible in the Appendix. The retrieved studies were uploaded onto the

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

Covidence software, a tool used for duplicate removal and manual screening of articles based on title, abstract and full text.

Eligibility Criteria

This study used phased eligibility criteria associated with each step of the screening process. During the database search, limits for English language only articles and articles published between 2010-2020 were added. No restrictions on study design or publication type were instituted at this stage.

Phase 1: Duplicates removed

Duplicate studies were automatically removed by the Covidence Software.

Phase 2: Title and abstract screening

Title and abstract screening was manually performed using the Covidence Software. Studies investigating social determinants of health, as categorized by the Healthy People 2030 definition, including race, ethnicity, income, family stability, or neighborhood safety, were included. Any studies looking solely at environmental or physiological risk factors were excluded. Studies not related to asthma, outside the participant age range of 2-18 years of age, full text unavailable/abstract-only papers, non-English manuscripts, studies solely looking at non-social variables (environmental, biological, etc.), and those not published between 2010-2020 were excluded.

Phase 3: Full text screening

Full text screening was performed manually using the Covidence Software. In addition to meeting all phase 2 criteria, studies investigating the outcome of asthma exacerbations measured

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

as inpatient or outpatient physician visits, ED visits for asthma, and patients with asthma refractory to inhalers or nebulizers were included. Associations were most commonly reported as odds ratios, risk ratios, or frequency of ED visits. Studies with designs other than randomized controlled trials, cohorts, and those investigating ambulatory care of asthma were excluded.

Selection process

Upon retrieval of studies from the implementation of the search strategy, duplicate articles were automatically excluded by the Covidence software in Phase 1. In Phase 2, two researchers independently reviewed the title and abstract of all remaining studies based on pre-established eligibility criteria. Conflicts were resolved via discussion between the two researchers till a consensus was reached. In Phase 3, full-text screening was performed by two researchers independently and conflicts were resolved in the same manner.

Data extraction

Articles meeting full text eligibility criteria in Phase 3 went through data extraction. The following factors were collected from each eligible study: Title, Author, Year of Publication, Social Determinant investigated, Outcome measure, Study Design, Statistical Analysis, Setting or study location, Sample Size, Age Group, Data and Data source, Study Aim, Key Results, and Statistical Significance of results.

Results

Study selection

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

Figure 1 shows the process we used to screen and select all included studies for this systematic review. Following the initial database search, 717 studies were retrieved for the systematic review. After 102 duplicates were removed during Phase 1, the remaining 615 studies proceeded to Phase 2. In Phase 2, 566 studies were excluded during the abstract and title screening. Upon resolving conflicts, the remaining 49 studies proceeded to full text screening in Phase 3. During full text review, 40 studies were excluded and 9 proceeded to data extraction. The data sources used by each of our 9 studies varied widely and can be identified on Table 1 [Table 1 near here]. All included studies were cohort studies, with 3 prospective and 6 retrospective designs. Additional study characteristics can be identified on Table 1 and key findings of each study can be found on Table 2 [Table 2 near here].

SDoH Domain 1: Social and community context

Six studies explored the role of social and community context in pediatric asthma exacerbations. The influence of three overarching variables were analyzed through the studies retrieved: sociodemographic factors, caregiver perceptions, and patient lifestyle.

Sociodemographic factors

The influence of sociodemographic factors on pediatric asthma exacerbations has been explored by five studies retrieved from this systematic review: Butz et al., Daniel et al., Guilbert et al., Rangachari et al., and Washington et al.

Three studies found no significant influence of sociodemographic factors on pediatric asthma ED visits. Butz et al conducted a prospective cohort study that collected information on patient demographics and asthma status via caregiver interviews at an initial ED visit and 6 months after. Age and gender were found to have no statistically significant influence on healthcare

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

utilization for very poorly controlled (VPC) asthma, as defined by the NAEPP Guidelines. Similarly, Daniel et al. conducted a retrospective cohort study with self-identified African American, Latino, and Non-Latino White pediatric patients that found no significant association between race and ethnicity and missed sleep and ED visits among pediatric asthma patients when controlling for poverty status. Finally, Rangachari et al. conducted a retrospective cohort study to explore the association between sociodemographic factors and unscheduled healthcare usage for persistent pediatric asthma patients. This study, too, concluded no statistically significant influence of age, gender, and race. Two additional studies found contrary results. Guilbert et al. conducted a prospective cohort study with 6-11 year old asthma patients that either self-identified as Black or White. This study found that Black children were more likely to have an emergency department visit for an asthma exacerbation than White children in the study period (calculated OR 4.42, $p < .0001$, $n = 348$). Further, the asthma exacerbations of Black patients were significantly more severe than those of White patients, as measured on the exacerbation hierarchy score (1.2 vs 0.6, $p < .001$). Similarly, Washington et al conducted a prospective cohort study and found that among 8-15 year-old asthma patients ($n = 544$), African American patients (RR 1.26, 95% CI: 1.14, 1.40) and Hispanic/Latino patients (RR 1.15, 95% CI: 1.01, 1.31) had a significantly higher risk of uncontrolled asthma, measured by the same NAEPP guidelines as Butz et al. This study further found that when controlled for health literacy and education, only African American patients still had a significantly higher risk than White patients for uncontrolled asthma.

Caregiver perceptions

The influence of caregiver perceptions on pediatric asthma exacerbations was the focus of two studies: Butz et al and Diep et al.

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

Diep et al conducted a retrospective cohort study using phone interviews on caregivers of 3 to 10 year-old children with persistent asthma. Caregivers' subjective social status was measured using a self-reported relative perspective from worse off or better off than several groups, including all Americans, their neighbors, and members of their race or ethnicity. The patient's asthma control was assessed by a number of ED visits for asthma in the past year, a survey of symptoms in the past two weeks, as well as the patient's score on the standardized Childhood Asthma Control Test. Results indicated no statistically significant difference in the number of pediatric asthma-related ED visits in the past year between high and low caregiver subjective social status (calculated OR 1.44, adjusted $p = 0.18$, $n = 230$).

The prospective cohort study conducted by Butz et al did find caregiver perceptions to be a significant factor. Caregiver worry about medication side effects increased the odds (OR 2.13, $p = 0.032$, $n = 222$) of a patient having very poorly controlled asthma, as diagnosed based on NAEPP guidelines. Consequently, these VPC asthma patients then had increased healthcare utilization compared to patients with well-controlled asthma (OR 1.26, $p = 0.003$, $n = 222$), including 1.26 times more frequent primary care visits with no significant difference in ICU admissions. Additionally, around 40% of caregivers of VPC asthma patients rated asthma control as higher than measured by the NAEPP guidelines, showing a disparity between caregivers' perception of health status and its reality.

Lifestyle

One retrospective cohort study, by Daniel et al., explored the impact of lifestyle practices. It evaluated the influence of missed sleep on pediatric asthma morbidity, including number of ED visits for asthma as measured on the AFSS asthma morbidity scale. It was found that more

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

frequent missed sleep in the past 12 months was significantly correlated with more frequent ED usage in the past 12 months (OR 1.67, $p < .05$, $n = 147$).

SDoH domain 2: Neighborhood and built environment

Three studies examined the role of neighborhood characteristics and built environment in relation to pediatric asthma. The two main variables pertaining to this domain were as follows: characteristics of neighborhood and effects of change in location.

Neighborhoods: High and Low Risk

Two studies – Nkoy et al. 2018 and Gjelsvik et al. 2019 – specifically focused on comparing low- and high-risk neighborhoods in relation to the number of hospital visits after an asthma diagnosis.

In Gjelsvik et al., information on patients' neighborhood risk levels was obtained through data extraction from the 2010-2014 American Community Survey and the 2010 US Census. The study found that children living in the highest poverty risk quintile were more likely to be readmitted to the hospital due to complications of asthma (OR 1.18, 95% CI 1.06,1.30, $n = 146,889$).

Nkoy et al found similar results. Through data extraction from the 2013 US census block group data and the enterprise data warehouse (EDW), patients' neighborhood risk was classified using the area of deprivation index (ADI) metric, with high ADIs relating to more deprived populations in terms of education, housing, and employment. It was found that patients residing in higher-risk neighborhoods were more likely to experience higher rates of hospital readmission (OR 1.15, 95% CI 1.06-1.24, $n = 2270$), along with a longer hospital length of stay compared to their lower-risk counterparts ($B = 1.04$, $p < 0.001$, $n=2270$).

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

Change in neighborhoods

Faison et al 2020 focused on exploring if housing instability, assessed as frequent changes in address in the 12 months prior to the study, was associated with increased rates of return ED visits. It was found that patients with at least one address change in a 12-month period were more likely to have severe asthma and more recent visitations. However, address changes were not significantly associated with an increase in return visits to the emergency department for pediatric asthma patients. Moreover, when conducting a multivariate logistical regression of the 90-day revisits, it was found that changes in address were not significantly associated with increased revisitation (OR: 1.49, 95% CI: 0.91, 2.46, $p = 0.114$, $n = 632$).

SDoH Interaction Effects: Insurance and asthma

Several studies investigated the role of patients' health insurance modality, distinguished as public or private, on pediatric asthma exacerbations. Nkoy et al. 2018, Faison et al. 2020, and Washington et al. 2018 demonstrated a significant association between public health insurance and hospital revisitation due to asthma. Nkoy et al. 2018 and Faison et al. 2020 related this increased rate of revisitation and type of insurance to place of residence, as patients who lived in high-risk areas and possessed public health insurance such as Medicaid were more likely to revisit the hospital due to asthma complications (Nkoy - OR: 1.81; 95% CI:1.46-2.24, $n = 2270$; Faison - OR: 1.94, 95% CI: 1.14-3.31, $p = 0.014$, $n = 632$) than their low-risk neighborhood counterparts. Similarly, Washington et al. 2018 found that patients with non-private health insurance were more likely to experience uncontrolled asthma (RR 1.16, 95% CI 1.08, 1.24, $p < .001$, $n = 544$) and lower quality of life (RR -0.33, 95% CI -0.43, -0.23, $p < .001$, $n = 544$) than those with private insurance.

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

Discussion

This systematic review aimed to identify social risk factors that influence pediatric asthma exacerbations. Overall, our findings suggest that race, neighborhood risk, health insurance status, and caregiver perceptions are important social factors. African American race is associated with an increased risk of asthma exacerbations compared to patients of Caucasian race. Residence in higher-risk neighborhoods was associated with higher ED readmission rates and longer hospital stays compared to lower-risk neighborhoods. Pediatric patients with public or no health insurance were also more prone to higher rates of readmission, longer hospital stays, and higher asthma severity. Finally, caregiver worry about medication side effects and inaccurate perception of patient's asthma control were associated with higher rates of very poorly controlled asthma.

Reviewed evidence suggests that social/community context and neighborhood/built environment synergistically interact to influence a child's risk of ED visits for asthma exacerbations. Specifically, race, gender, age, and ethnicity are variables that have been widely explored under social and community context. However, some studies found no significant associations while others identified that Black Race was associated with more frequent and severe asthma exacerbations that needed emergency department care. This finding was supported by previously conducted systematic reviews, including Buelo et al. 2018, in which African American children were found to have an increased risk of experiencing asthma attacks compared to non-African American populations¹⁷. Further, caregiver perceptions of the patient's asthma care were found to be significant contributors to asthma exacerbations. This result was echoed in previously conducted reviews, such as Tzeng et al. 2017, in which it was observed that

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

patients whose parents possessed low health literacy were more likely to have poor asthma control¹⁸. This finding highlights the need for improving caregiver education on asthma management and prognosis by long-term healthcare providers as it could significantly improve patient outcomes.

In exploring the influence of neighborhoods/built environment, it was found that high-risk neighborhoods were correlated with higher lengths of stay and readmission rates for pediatric asthma compared to low-risk, higher-income neighborhoods. However, the frequency of neighborhood change was not significantly associated with an increase in the rate of readmission and length of stay in pediatric asthma patients. These results are similar to those seen in Buelo et al. 2018, which determined that patients with low socioeconomic status had an increased risk for an asthma attack. These findings indicate the need for more resources, such as educational opportunities, public transportation, and access to healthcare resources, to minimize these factors from exacerbating the consequences of asthma further.

Additionally, it was found that the interaction between health insurance type and neighborhood risk had a significant influence on pediatric asthma outcomes. Overall, having public or no health insurance was associated with higher asthma severity and rates of readmission, along with longer hospital stays. These results display the necessity of implementing programs that provide financial support to patients, especially those of lower socioeconomic status.

This study does have limitations. As this topic is not well-studied, a large gap remains in the number of studies available for a review. Several articles were abstract-only publications that did not have a full text associated with them, which is a potential source of publication bias. The study screening was done by two authors. During the title and abstract screening, conflicts were

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

resolved through discussion between these two authors, which could potentially introduce bias. However, the conflicts during full-text screening were resolved by a third reviewer. Additionally, the studies included in this review were limited to the time range of 2010 to 2020, which may limit the generalization of findings to outcomes beyond those years. Despite these limitations, we worked to minimize the potential biases by following the 2020 PRISMA Guidelines for Systematic Reviews.

Conclusion

Social factors play a role in pediatric asthma. It is vital for healthcare professionals to make patients aware of how their surroundings and lifestyles affect their respiratory health, and to support social movements that enhance the social environments and resources available to at-risk patient populations. Further research using expanded data elements describing social determinants of health for pediatric asthma are needed.

Acknowledgements

We are thankful to Mirian Ramirez, from the Ruth Lilly Medical Library, for development of the search strategy. We are also grateful to our lab members, Dr. Eneida Mendonca, Dr. Colin Rogerson, Dr. Arthur Owora, and Dr. Tian He for their support with this project.

Declaration of interest

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper. This research was funded by the Indiana University Medical Student Program for Research and Scholarship and the following grants: K12HS026390, R25HL126140 (sub-contract), and R03HS029088.

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

Tables

Table 1: Information from Studies

Reference	Population age (years)	Study Design	Data Source	Social Determinant
Butz et al., 2018 ⁸	3 to 12	prospective cohort	subdata of an ongoing RCT at John Hopkins	level of caregiver worry about asthma medications
Daniel et al., 2012 ⁹	6 to 13	retrospective cohort	Rhode Island hospital, peds asthma clinic patients	missed sleep and race
Diep et al., 2019 ¹⁰	3 to 10	retrospective cohort	SB-TEAM trial in Rochester, NY.	caregivers' subjective social status
Faison et al, 2020 ¹¹	2 to 17	retrospective cohort	Vidant Medical Center	housing insecurity (measured as address changes within a 12 month period)
Gjelsvik et. al 2019 ¹²	2 to 17	retrospective cohort	Rhode Island hospitals	neighborhood risk, race, ethnicity
Guilbert et al; 2019 ¹³	6 to 11	prospective cohort	n/a	Race (black vs white)
Nkoy et al; 2018 ¹⁴	2 to 17	retrospective cohort	intermountain hospitals (from January 1 2010-June 30 2015)	neighborhood deprivation, insurance coverage
Rangachari et al, 2020 ¹⁵	8 to 17	retrospective cohort	allergy-immunology specialty outpatient peds clinic in Georgia	demographics (age, gender, race, insurance)
Washington et al, 2018 ¹⁶	8 to 15	prospective cohort	Chicago Initiative to Raise Asthma Health Equity cohort	race/ethnicity, health literacy, education, sex, age, income, employment, insurance

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

Table 2: Key Findings from Studies

Reference	Cohort Size	Measures	Key Findings
Domain 1: Social and Community Context			
Butz et al., 2018	222	OR, 95% CI	age/gender: ns; caregiver worry about medication side effects: 2.13 (1.07–4.24), p=0.032; VPC asthma patients non-urgent healthcare utilization was significantly higher than WC asthma patients(OR 1.26, SD 1.3), p = 0.003
Daniel et al., 2012	147	OR, R ²	race/ethnicity: ns; missed sleep is directly correlated with frequency of ED usage (OR 1.67, R ² 0.06, P < .05)
Diep et al., 2019	230	OR*	n.s. (OR 1.44, P = 0.18); caregiver subjective social status had no significant impact on % of children with ED visits in the past year
Guilbert et al., 2019	348	OR*	Black children had a higher prevalence and severity of ED visits for asthma exacerbations than White children (OR 4.42, p <.0001)
Rangachari et al., 2020	59	OR*	N.S. - no significant difference in unscheduled healthcare usage, such as ED visits, between African American (OR 20.93, p = 0.975) and Caucasian patients or Hispanic (OR = 11.89, p = 0.986) and Caucasian patients.
Washington et al., 2018	544	RR, 95% CI	African American patients (RR 1.26, 95% CI: 1.14, 1.40) and Hispanic/Latino patients (RR 1.15, 95% CI: 1.01,1.31) had higher risk of uncontrolled asthma, including urgent care visits than White patients
Domain 2: Neighborhood and Built Environment			
Faison et al, 2020	632	95 CI, OR	revisitation rate to ER and change in address (OR: 1.49, 95% CI: 0.91, 2.46, P = 0.114), Medicaid insurance (OR: 1.94, 95% CI: 1.14, 3.31, P = 0.014)
Gjelsvik et. al 2019	146889	95% CI, OR	annual emergency hospital visits (OR: 1.18, 95% CI: 1.06, 1.30), annual hospitalizations (OR: 0.41, 95% CI: 0.34, 0.47)


SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

Author	n	95% CI	OR, β	Outcome
Nkoy et al; 2018	2270			neighborhood deprivation: LOS (β : 1.04, $P < 0.001$), readmission (OR: 1.15, 95% CI: 1.06, 1.24), cost (β : 1.03, $P < 0.001$). Public insurance recipients: readmission (OR: 1.81, 95% CI: 1.46-2.24), cost (β : 1.147, $P < 0.001$), LOS (β : 1.171, $P < 0.001$)

*Odds ratios for the indicated articles were manually calculated using data provided in the study manuscript.

Figures

Figure 1. Process of study inclusion and exclusion



SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

References

1. Centers for Disease Control and Prevention, Vital Signs. Asthma in Children. Washington (DC): Office of the Associate Director for Communication, Digital Media Branch, Division of Public Affairs; 2018 Jun 10 [accessed 2022 Jan 4]. <https://www.cdc.gov/vitalsigns/childhood-asthma/index.html#:~:text=1%20in%2012.,50%25>.
2. Goff, S. Asthma Facts and Figures. United States: Asthma and Allergy Foundation of America; 2022 Apr [accessed 2022 Jan 5]. <https://www.aafa.org/asthma-facts/>
3. Centers for Disease Control and Prevention, National Center for Health Statistics. Asthma. Washington (DC): Office of the Associate Director for Communication, Digital Media Branch, Division of Public Affairs; 2018 Feb 1 [accessed 2022 Jan 5]. <https://www.cdc.gov/nchs/fastats/asthma.htm>
4. Johnston NW, Sears MR. Asthma exacerbations. 1: epidemiology. *Thorax*. 2006 Aug;61(8):722-8. doi: 10.1136/thx.2005.045161. Cited in: PubMed. PMID: 16877691; PMCID: PMC2104697.
5. Keet, C. A., Matsui, E. C., McCormack, M. C., & Peng, R. D. (2017). Urban residence, neighborhood poverty, race/ethnicity, and asthma morbidity among children on Medicaid. *The Journal of allergy and clinical immunology*, 140(3), 822–827. <https://doi.org/10.1016/j.jaci.2017.01.036>. Cited in: PubMed.
6. Harrington, K. F., Zhang, B., Magruder, T., Bailey, W. C., & Gerald, L. B. (2015). The Impact of Parent's Health Literacy on Pediatric Asthma Outcomes. *Pediatric allergy, immunology, and pulmonology*, 28(1), 20–26. <https://doi.org/10.1089/ped.2014.0379>. Cited in: PubMed.

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

7. Social Determinants of Health [Internet]. Social Determinants of Health - Healthy People 2030. U.S. Department of Health and Human Services; [cited 2022Oct16]. Available from: <https://health.gov/healthypeople/priority-areas/social-determinants-health>
8. Butz AM, Bellin M, Tsoukleris M, Mudd SS, Kub J, Ogborn J, Morpew T, Lewis-Land C, Bollinger ME. Very Poorly Controlled Asthma in Urban Minority Children: Lessons Learned. *J Allergy Clin Immunol Pract*. 2018 May-Jun;6(3):844-852.
<https://doi.org/10.1016/j.jaip.2017.08.007>. Epub 2017 Sep 22. PMID: 28958744; PMCID: PMC5862725.
9. Daniel LC, Boergers J, Kopel SJ, Koinis-Mitchell D. Missed sleep and asthma morbidity in urban children. *Ann Allergy Asthma Immunol*. 2012 Jul;109(1):41-6.
<https://doi.org/10.1016/j.anai.2012.05.015>. PMID: 22727156; PMCID: PMC3426919.
10. Diep J, Fagnano M, Tremblay P, Halterman JS. The relationship between caregivers' subjective social status and asthma symptoms and management for urban children. *J Asthma*. 2019 Feb;56(2):211-217. <https://doi.org/10.1080/02770903.2018.1437176>. Epub 2018 Mar 1. PMID: 29494268; PMCID: PMC6119512.
11. Faison K, Moon A, Buckman C, Cortright L, Tumin D, Campbell C, Beamon B. Change of address as a measure of housing insecurity predicting rural emergency department revisits after asthma exacerbation. *J Asthma*. 2021 Dec;58(12):1616-1622.
<https://doi.org/10.1080/02770903.2020.1818773>. Epub 2020 Sep 21. PMID: 32878515.
12. Gjelsvik A, Rogers ML, Garro A, Sullivan A, Koinis-Mitchell D, McQuaid EL, Smego R, Vivier PM. Neighborhood Risk and Hospital Use for Pediatric Asthma, Rhode Island, 2005-

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

2014. *Prev Chronic Dis*. 2019 May 30;16:E68. <https://doi.org/10.5888/pcd16.180490>. PMID: 31146802; PMCID: PMC6549429.

13. Guilbert T, Zeiger RS, Haselkorn T, Iqbal A, Alvarez C, Mink DR, Chipps BE, Szeffler SJ. Racial Disparities in Asthma-Related Health Outcomes in Children with Severe/Difficult-to-Treat Asthma. *J Allergy Clin Immunol Pract*. 2019 Feb;7(2):568-577.

<https://doi.org/10.1016/j.jaip.2018.07.050>. Epub 2018 Aug 29. Erratum in: *J Allergy Clin Immunol Pract*. 2019 Mar;7(3):1096. PMID: 30172020.

14. Nkoy FL, Stone BL, Knighton AJ, Fassel BA, Johnson JM, Maloney CG, Savitz LA. Neighborhood Deprivation and Childhood Asthma Outcomes, Accounting for Insurance Coverage. *Hosp Pediatr*. 2018 Jan 9;hpeds.2017-0032. <https://doi.org/10.1542/hpeds.2017-0032>. Epub ahead of print. PMID: 29317461.

15. Rangachari P, Griffin DD, Ghosh S, May KR. Demographic and Risk-Factor Differences between Users and Non-Users of Unscheduled Healthcare among Pediatric Outpatients with Persistent Asthma. *Int J Environ Res Public Health*. 2020 Apr 15;17(8):2704. <https://doi.org/10.3390/ijerph17082704>. PMID: 32326397; PMCID: PMC7215685.

16. Washington DM, Curtis LM, Waite K, Wolf MS, Paasche-Orlow MK. Sociodemographic Factors Mediate Race and Ethnicity-associated Childhood Asthma Health Disparities: a Longitudinal Analysis. *J Racial Ethn Health Disparities*. 2018 Oct;5(5):928-938. <https://doi.org/10.1007/s40615-017-0441-2>. Epub 2017 Nov 29. PMID: 29188553.

17. Buelo, A., McLean, S., Julious, S., Flores-Kim, J., Bush, A., Henderson, J., Paton, J. Y., Sheikh, A., Shields, M., Pinnock, H., & ARC Group (2018). At-risk children with asthma

It is made available under a [CC-BY-NC-ND 4.0 International license](#) .

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

(ARC): a systematic review. *Thorax*, 73(9), 813–824. <https://doi.org/10.1136/thoraxjnl-2017-210939>. Cited in: PubMed.

18. Tzeng, Y. F., Chiang, B. L., Chen, Y. H., & Gau, B. S. (2018). Health literacy in children with asthma: A systematic review. *Pediatrics and neonatology*, 59(5), 429–438.

<https://doi.org/10.1016/j.pedneo.2017.12.001>. Cited in: PubMed.

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

Appendix

Full Search Strategy

Database 1: Ovid MEDLINE(R)

Date run: 11/2/2021

Total results: 202

Ovid MEDLINE(R) and Epub Ahead of Print, In-Process, In-Data-Review & Other Non-Indexed Citations, Daily and Versions(R) <1946 to November 01, 2021>

No.	Query	Results
1	exp pediatrics/	61387
2	(pediatric\$ or paediatric\$ or peadiatric\$).ti,ab.	390560
3	(toddler\$ or baby\$ or baby\$ or babies).ti,ab	89315
4	(minors or minors\$).ti,ab.	4802
5	(boy or boys or boyfriend or boyhood).ti,ab.	154213
6	girl\$.ti,ab.	158970
7	(kid or kids).ti,ab,	8801
8	exp child/ or exp adolescent/	3190614
9	(child or child\$ or children\$).ti,ab,	1486711
10	(schoolchild\$ or schoolchild or "school child").ti,ab,	15244
11	(adolescenc\$ or youth\$ or teen\$).ti,ab.	376213
12	(juvenil\$ or underage).ti,ab,	88289
13	(prepubesc\$ or pubesc\$).ti,ab,	3508
14	(preschool or school or school\$).ti,ab.	327394
15	or/1-14	4034238
16	exp socioeconomic factors/	479173
17	social environment/	43848
18	social Determinants of Health/	4884
19	Social conditions/	9479
20	(social adj3 (aspect\$ or factor\$ or condition\$ or context or determinant\$ or disparit\$ or advantage\$ or disadvantage\$ or inequalit\$ or inequities or inequity or influence\$ or ecolog\$ or problem\$ or stressor\$)).ti,ab.	99465
21	((socioeconomic\$ or socio-economic\$) adj3 (aspect\$ or factor\$ or condition\$ or context or determinant\$ or disparit\$ or advantage\$ or disadvantage\$ or inequalit\$ or inequities or inequity or influence\$ or ecolog\$ or problem\$ or stressor\$)).ti,ab.	37436
22	Family Relations/	11761
23	Parent-Child Relations/	35235

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

24	((family or parent\$ or child\$) adj relat\$).ti,ab.	15347
25	exp "poverty"/ or "poverty".ti,ab.	64746
26	Social Factors/	146
27	Residence Characteristics/	36450
28	((residence or neighborhood\$) adj2 (safety or characteristic\$ or unsaf\$)).ti,ab.	2258
29	"living condition\$".ti,ab.	8528
30	or/16-29	684521
31	asthma/	131647
32	Bronchial Hyperreactivity/	7471
33	respiratory hypersensitivity/	9818
34	(asthma\$ adj3 (flare-Up or flare-ups or flaring-up or flaring-ups or flareup\$ or exaggerat\$ or acute or magnification\$ or exacerbat\$ or attack\$ or sever\$ or worse\$ or (symptom\$ adj2 increas\$)).ti,ab.	29787
35	(wheez\$ adj3 (flare-Up or flare-ups or flaring-up or flaring-ups or flareup\$ or exaggerat\$ or acute or magnification\$ or exacerbat\$ or attack\$ or sever\$ or worse\$ or (symptom\$ adj2 increas\$)).ti,ab	1240
36	((bronchial or bronchus or respiratory) adj (hyperreactivity or "hyper reactivity")).ti,ab,	2358
37	(respiratory adj (hypersensitivity or "hyper sensitivity")).ti,ab.	116
38	(airway\$ adj1 (reactive or intermittent or inflammation or hyper-responsiveness or hyperresponsiveness)).ti,ab.	21060
39	31 or 32 or 33 or 36 or 37 or 38	150168
40	Symptom Flare Up/	901
41	(flare-Up or flare-ups or flaring-up or flaring-ups or flareup\$ or exaggerat\$ or acute or magnification\$ or exacerbat\$ or attack\$ or sever\$ or worse\$ or (symptom\$ adj2 increas\$)).ti,ab.	4502460
42	40 or 41	4502762
43	34 or 35 or (39 and 42)	49833
44	15 and 30 and 43	1322
45	exp cohort studies/	2239611
46	cohort\$.tw.	703689
47	(concurrent or follow-up or incidence).tw.	1886594
48	longitudinal.tw.	277923
49	retrospective\$.tw.	864272
50	prospective\$.tw.	774379
51	or/45-50	4086049

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

52	44 and 51	505
53	limit 52 to yr=2010-2020	207
54	53 and english.la.	202

Database 2: EMBASE

Date run: 11/2/2021

Total results: 485

No	Query	Results
#38	#37 AND [embase]/lim	485
#37	#36 AND [english]/lim	512
#36	#35 AND [2010-2020]/py	518
#35	#23 AND #34	803
#34	#24 OR #25 OR #26 OR #27 OR #28 OR #29 OR #30 OR #31 OR #32 OR #33	5912712
#33	prospective*:ti,ab,kw	1198324
#32	retrospective*:ti,ab,kw	1432662
#31	longitudinal:ti,ab,kw	383417
#30	concurrent:ti,ab,kw OR 'follow up':ti,ab,kw OR incidence:ti,ab,kw	2850260
#29	cohort*:ti,ab,kw	1190885
#28	'follow up'/de	1771612
#27	'retrospective study'/de	115645

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

7		0
#2 6	'prospective study'/de	723757
#2 5	'longitudinal study'/de	163070
#2 4	'cohort analysis'/de	768639
#2 3	#3 AND #11 AND #22	2132
#2 2	#14 OR #15 OR (#18 AND #21)	108456
#2 1	#19 OR #20	618594 5
#2 0	'flare up':ti,ab OR 'flare ups':ti,ab OR 'flaring up':ti,ab OR 'flaring ups':ti,ab OR flareup*:ti,ab OR exaggerat*:ti,ab OR acute:ti,ab OR magnification*:ti,ab OR exacerbat*:ti,ab OR attack*:ti,ab OR sever*:ti,ab OR worse*:ti,ab OR 'symptoms increas*':ti,ab	612329 2
#1 9	'disease exacerbation'/exp	144064
#1 8	#12 OR #13 OR #16 OR #17	303610
#1 7	(airway* NEAR/1 (reactive OR intermittent OR inflammation OR 'hyper responsiveness' OR hyperresponsiveness)):ti,ab	31278
#1 6	((bronchial OR bronchus OR respiratory) NEXT/1 (hyperreactivity OR 'hyper reactivity')):ti,ab	3168
#1 5	(wheez* NEAR/3 ('flare up' OR 'flare ups' OR 'flaring up' OR 'flaring ups' OR flareup* OR exaggerat* OR acute OR magnification* OR exacerbat* OR attack* OR sever* OR worse* OR 'symptoms increas*')):ti,ab	2012
#1 4	(asthma* NEAR/3 ('flare up' OR 'flare ups' OR 'flaring up' OR 'flaring ups' OR flareup* OR exaggerat* OR acute OR magnification* OR exacerbat* OR attack* OR sever* OR worse* OR 'symptoms increas*')):ti,ab	48334
#1 3	'bronchus hyperreactivity'/de	12607

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

#1 2	'asthma'/exp	288992
#1 1	#4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10	851990
#1 0	'living condition*':ti,ab	10166
#9	((residence OR neighborhood*) NEAR/2 (safety OR characteristic* OR unsaf*)):ti,ab	2569
#8	'poverty'/de OR 'poverty':ti,ab	62941
#7	'family relation'/exp OR (((family OR parent* OR child*) NEXT/1 relat*):ti,ab)	127164
#6	((socioeconomic* OR 'socio economic*') NEAR/3 (aspect* OR factor* OR condition* OR context OR determinant* OR disparit* OR advantage* OR disadvantage* OR inequalit* OR inequities OR inequity OR influence* OR ecolog* OR problem* OR stressor*)):ti,ab	46027
#5	(social NEAR/3 (aspect* OR factor* OR condition* OR context OR determinant* OR disparit* OR advantage* OR disadvantage* OR inequalit* OR inequities OR inequity OR influence* OR ecolog* OR problem* OR stressor*)):ti,ab	121719
#4	'social aspects and related phenomena'/de OR 'social aspect'/de OR 'social environment'/de OR 'social status'/de OR 'social problem'/de OR 'social condition'/de OR 'social determinants of health'/de OR 'socioeconomic factors'/exp	645889
#3	#1 OR #2	490752 9
#2	pediatric*:ti,ab OR paediatric*:ti,ab OR peadiatric*:ti,ab OR toddler*:ti,ab OR baby:ti,ab OR baby*:ti,ab OR babies:ti,ab OR minors:ti,ab OR minors*:ti,ab OR boy:ti,ab OR boys:ti,ab OR boyfriend:ti,ab OR boyhood:ti,ab OR girl*:ti,ab OR kid:ti,ab OR kids:ti,ab OR child:ti,ab OR child*:ti,ab OR children*:ti,ab OR schoolchild*:ti,ab OR adolescen*:ti,ab OR youth*:ti,ab OR teen*:ti,ab OR juvenil*:ti,ab OR underage:ti,ab OR prepubesc*:ti,ab OR pubesc*:ti,ab OR preschool:ti,ab OR school:ti,ab OR school*:ti,ab	296752 7
#1	'pediatrics'/de OR 'adolescent'/de OR 'child'/exp OR 'childhood'/de OR 'juvenile'/de	404384 6

It is made available under a [CC-BY-NC-ND 4.0 International license](#) .

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

Database 3: PsycINFO 1887-current (EBSCO)

Date run: 11/2/2021

total results: 30

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

#	Query	Limiters/Expanders	Last Run Via	Results
S4 6	(S37 AND S44)	Limiters - Publication Year: 2010-2020 Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA PsycInfo	30
S4 5	(S37 AND S44)	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA PsycInfo	54
S4 4	(S38 OR S39 OR S40 OR S41 OR S42 OR S43)	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA	472,432

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

			PsycInfo	
S4 3	(TI cohort* OR AB cohort*) OR ((TI concurrent OR AB concurrent) OR (TI follow-up OR AB follow-up) OR (TI incidence OR AB incidence)) OR (TI longitudinal OR AB longitudinal) OR (TI retrospective* OR AB retrospective*) OR (TI prospective* OR AB prospective*)	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA PsycInfo	437,715
S4 2	DE "Followup Studies"	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA PsycInfo	12,307

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

S4 1	DE "Retrospective Studies"	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA PsycInfo	28,812
S4 0	DE "Prospective Studies"	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA PsycInfo	33,468
S3 9	DE "Longitudinal Studies"	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA PsycInfo	51,967

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

S3 8	DE "Cohort Analysis"	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA PsycInfo	1,557
S3 7	S7 AND S26 AND S36	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA PsycInfo	262
S3 6	S28 OR S29 OR (S32 AND S34)	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA PsycInfo	2,191

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

S3 5	S33 OR S34	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA PsycInfo	668,771
S3 4	(TI asthma* N3 (TI flare-Up OR TI flare-ups OR TI flaring-up OR TI flaring-ups OR TI flareup* OR TI exaggerat* OR TI acute OR TI magnification* OR TI exacerbat* OR TI attack* OR TI sever* OR TI worse* OR TI "symptom increas*")) OR (AB flare-Up OR AB flare-ups OR AB flaring-up OR AB flaring-ups OR AB flareup* OR AB exaggerat* OR AB acute OR AB magnification* OR AB exacerbat* OR AB attack* OR AB sever* OR AB worse* OR AB "symptom increas*"))	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA PsycInfo	665,700
S3 3	DE "Severity (Disorders)"	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA PsycInfo	19,841

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

S3 2	S27 OR S30 OR S31	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA PsycInfo	5,617
S3 1	((TI airway* OR AB airway*) N1 ((TI reactive OR AB reactive) OR (TI intermittent OR AB intermittent) OR (TI inflammation OR AB inflammation) OR (TI "hyper responsiveness" OR AB "hyper responsiveness") OR (TI hyperresponsiveness OR AB hyperresponsiveness)))	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA PsycInfo	168
S3 0	((((TI bronchial OR AB bronchial) OR (TI bronchus OR AB bronchus) OR (TI respiratory OR AB respiratory)) N1 ((TI hyperreactivity OR AB hyperreactivity) OR (TI "hyper reactivity" OR AB "hyper reactivity"))))	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA PsycInfo	13

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

S2 9	(TI wheez* N3 (TI flare-Up OR TI flare-ups OR TI flaring-up OR TI flaring-ups OR TI flareup* OR TI exaggerat* OR TI acute OR TI magnification* OR TI exacerbat* OR TI attack* OR TI sever* OR TI worse* OR TI "symptom increas*")) OR (AB wheez* N3 (AB flare-Up OR AB flare-ups OR AB flaring-up OR AB flaring-ups OR AB flareup* OR AB exaggerat* OR AB acute OR AB magnification* OR AB exacerbat* OR AB attack* OR AB sever* OR AB worse* OR AB "symptom increas*"))	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA PsycInfo	30
S2 8	(TI asthma* N3 (TI flare-Up OR TI flare-ups OR TI flaring-up OR TI flaring-ups OR TI flareup* OR TI exaggerat* OR TI acute OR TI magnification* OR TI exacerbat* OR TI attack* OR TI sever* OR TI worse* OR TI "symptom increas*")) OR (AB asthma* N3 (AB flare-Up OR AB flare-ups OR AB flaring-up OR AB flaring-ups OR AB flareup* OR AB exaggerat* OR AB acute OR AB magnification* OR AB exacerbat* OR AB attack* OR AB sever* OR AB worse* OR AB "symptom increas*"))	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA PsycInfo	1,426

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

S2 7	(DE "Asthma")	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Advanced Search Database - APA PsycInfo	5,530
S2 6	(S8 OR S9 OR S10 OR S11 OR S12 OR S13 OR S14 OR S15 OR S16 OR S17 OR S18 OR S19 OR S20 OR S22 OR S23 OR S24 OR S25)	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display
S2 5	(TI "living condition*" OR AB "living condition*")	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

S2 4	(((TI residence OR AB residence) OR (TI neighborhood* OR AB neighborhood*)) N2 ((TI safety OR AB safety) OR (TI characteristic* OR AB characteristic*) OR (TI unsaf* OR AB unsaf*)))	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display
S2 3	DE "Neighborhoods"	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display
S2 2	DE poverty OR (TI poverty OR AB poverty)	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

S2 1	((((TI family OR AB family) OR (TI parent* OR AB parent*) OR (TI child* OR AB child*)) N1 (TI relat* OR AB relat*)))	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display
S2 0	(DE "family relations") OR (DE "parent-child relations")	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

S19	(((TI socioeconomic* OR AB socioeconomic*) OR (TI "socioeconomic*" OR AB "socioeconomic*")) N3 ((TI aspect* OR AB aspect*) OR (TI factor* OR AB factor*) OR (TI condition* OR AB condition*) OR (TI context OR AB context) OR (TI determinant* OR AB determinant*) OR (TI disparit* OR AB disparit*) OR (TI advantage* OR AB advantage*) OR (TI disadvantage* OR AB disadvantage*) OR (TI inequalit* OR AB inequalit*) OR (TI inequities OR AB inequities) OR (TI inequity OR AB inequity) OR (TI influence* OR AB influence*) OR (TI ecolog* OR AB ecolog*) OR (TI problem* OR AB problem*) OR (TI stressor* OR AB stressor*)))	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display
S18	((TI social OR AB social) N3 ((TI aspect* OR AB aspect*) OR (TI factor* OR AB factor*) OR (TI condition* OR AB condition*) OR (TI context OR AB context) OR (TI determinant* OR AB determinant*) OR (TI disparit* OR AB disparit*) OR (TI advantage* OR AB advantage*) OR (TI disadvantage* OR AB disadvantage*) OR (TI inequalit* OR AB inequalit*) OR (TI inequities OR AB inequities) OR (TI inequity OR AB inequity) OR (TI influence* OR AB influence*) OR (TI ecolog* OR AB ecolog*) OR (TI problem* OR AB problem*) OR (TI stressor* OR AB stressor*)))	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

S1 7	DE "Social Ecology"	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display
S1 6	DE "social status" OR DE "Socioeconomic Status"	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display
S1 5	DE "socioeconomic factors" OR DE "Income Level" OR DE "Lower Class" OR DE "Social Class"	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

S1 4	DE "Social Issues"	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display
S1 3	DE "Family Socioeconomic Level"	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display
S1 2	DE "Social Exclusion"	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

S11	DE "Social Perception"	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display
S10	DE "Socioeconomic Status"	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display
S9	DE "Social Disadvantage" OR DE "Economic Disadvantage"	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

S8	DE "Social Environments"	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display
S7	(S1 OR S2 OR S3 OR S4 OR S5 OR S6)	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display
S6	AG "Childhood (birth-12 yrs)" OR "School Age (6-12 yrs)" OR "Adolescence (13-17 yrs)" OR "Adulthood (18 yrs & older)"	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

S5	AB (pediatric* OR paediatric* OR peadiatric* OR toddler* OR baby OR baby* OR babies OR minors OR minors* OR boy OR boys OR boyfriend OR boyhood OR girl* OR kid OR kids OR child OR child* OR children* OR schoolchild* OR adolescen* OR youth* OR teen* OR juvenil* OR underage OR prepubesc* OR pubesc* OR preschool OR school OR school*)	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display
S4	TI (pediatric* OR paediatric* OR peadiatric* OR toddler* OR baby OR baby* OR babies OR minors OR minors* OR boy OR boys OR boyfriend OR boyhood OR girl* OR kid OR kids OR child OR child* OR children* OR schoolchild* OR adolescen* OR youth* OR teen* OR juvenil* OR underage OR prepubesc* OR pubesc* OR preschool OR school OR school*)	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display
S3	SU (adolescent OR youth)	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

SOCIAL RISK FACTORS FOR PEDIATRIC ASTHMA EXACERBATIONS

S2	SU child	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display
S1	DE pediatrics	Expanders - Apply equivalent subjects Search modes - Boolean/Phrase	Interface - EBSCOhost Research Databases Search Screen - Basic Search Database - APA PsycInfo	Display