1 Main title

2 Immunization Staff Training Needs Assessment: An Overview Of Findings From Cameroon

3 Running title

4 Immunization Staff Training Needs Assessment

5 List of authors

- 6 Amani Adidja¹, Sangwe Clovis Nchinjoh^{2*}, Jessica Gu³, Fabrice Zobel Lekeumo Cheuyem¹, Nadege
- 7 Nnang Edwige², Marius Vouking⁵, Jude Nkwain Muteh⁶, Valirie Ndip Agbor⁷, Mbanga Clarence
- 8 Mvalo², Nsah Bernard², Budzi Michael Ngenge², Shalom Tchokfe Ndoula⁴, Andreas Ateke Njoh⁶,
- 9 Pietro Di Mattei⁵, Calvin Tonga⁶, Yauba Saidu^{2,8}

10 Affiliations

- 11 1. Faculty of Medicine and Biomedical Sciences, University of Yaoundé 1, Yaoundé, Cameroon
- 12 2. Clinton Health Access Initiative Inc., Yaounde P.O. Box 2664, Cameroon
- 13 3. Global Vaccine Team, Clinton Health Access Initiative Inc., Boston, MA 02127, USA;
- 14 4. Cameroon Ministry of Public Health, Yaoundé PO Box 2084, Cameroon
- 15 5. Centre pour le Développement des Bonnes Pratiques en Santé, Hôpital Central de
 Yaoundé, Henri-Dunant Avenue, Messa, Yaoundé, Cameroun.
- 17 6. Gavi, the Vaccine Alliance, Geneva, Switzerland
- Clinical Trial Service Unit and Epidemiological Studies Unit (CTSU), Nuffield Department of
 Population Health, University of Oxford, Oxford, UK
- 20 8. Institute for Global Health, University of Siena, 53100 Siena, Italy
- 21 Corresponding author*
- 22 E-mail: <u>nsclovis@clintonhealthaccess.org</u> (SCN)

23

- 24 Abstract
- 25 Background:
- 26 In recent years, Cameroon's Expanded Program on Immunization (EPI) has witnessed a sizable decline
- 27 inote: performance. Many retakeholders have a ted inversion and resources as an e cafither major

drivers for the observed trend. In a bid to better understand and redress this situation, the EPI and its partners conducted a Training Needs Assessment (TNA) among central, regional, and district staff. The assessment aimed to quantify and characterize core capability gaps and leverage the findings to design an evidence-based capacity-building plan.

Methods: A descriptive cross-sectional study was conducted using aggregated data from a survey
 carried out among EPI staff from May to September 2016 across Cameroon's health pyramid - central,
 regional, and district levels; and analysis was done using Microsoft Office Excel 2016.

35 **Results:** Over half of EPI staff had worked for less than three years in their current post, and roughly 36 three-third of them did not receive pre-service training on vaccination. Additionally, about half of them 37 had not received any form of in-service training on immunization. Supportive supervision was the most 38 frequently cited topic for training, with a surprisingly higher need at the central level (80%). Financial 39 incentive was not a primary motivating factor for learning. Approximately half of the respondents at all levels were not aware of onboarding materials. Most of the respondents identified multiple meetings, 40 high staff turnover, and competing priorities as major barriers to learning. Only about half of surveyed 41 42 staff reported participating in performance reviews, with nearly half of these reviews conducted when 43 an opportunity arose.

44 Conclusion:

45 There are still many gaps challenging the EPI goal achievement in Cameroon. Sustainably addressing 46 these issues will require a comprehensive framing of capability-building activities. High-quality EPI 47 human resources will boost the country's vaccination performance and contribute to the reduction of 48 infant mortality in the country.

49 Keywords: Training Need Assessment, Vaccination, Expanded Program on Immunization Staff

50

51

52 Background

Globally, immunization has been a remarkable tool for reducing morbidity and mortality among children since the creation of the Expanded Program on Immunization (EPI) in 1974 [1]. Many countries including Cameroon, a country in Central Africa, over the years have made significant

progress in achieving critical vaccination targets. Indeed, Cameroon reported a significant rise in vaccination coverage for the 1st and 3rd doses of the Diptheria-Tetanus-Pertussis containing vaccines (DTP) from 54% and 62% in 2000 to 88.6 % and 95% in 2013, respectively[2]. This steady increase of over 30 points has been arguably due to consistent political will and commitment towards infant health, a high social acceptance rate, financial and technical support from development partners, and the implementation of the Reach Every District (RED) strategy [3], amongst others.

Despite this promising rise in vaccination coverage, in recent years, vaccination coverage in Cameroon has stalled with occasional progressive decline in performance down to 76% and 69% for DTP-1 and DTP-3, respectively, in 2021[2]. Among other things, anecdotal evidence suggests that this may have been a direct effect of new challenges faced by the program. This includes the need to expand and sustain coverages, close equity gaps dramatically, meet the demands of a rapidly rising birth cohort and urbanization[4], introduce new vaccines [5], and aggressively pursue accelerated global and regional disease control targets[6, 7].

69 To effectively deal with the immunization challenges of the 21st century, the EPI and its partners 70 conducted several vertical vaccination system assessments and leveraged the findings to implement 71 tailored interventions to close identified gaps. Although the implementation of these actions yielded 72 positive results, the outcome is arguably limited by capacity gaps among EPI staff. Indeed, data from the 2016 national cold chain inventory showed that only 12.2% of the 9,661 surveyed staff had received 73 74 training on immunization within the previous five years, a finding which is somewhat consistent with the 16% reported in 2013 [8]. This situation, coupled with a lack of updated reference documents, high 75 staff attrition, and the absence of a robust onboarding system, have created significant capacity gaps 76 among EPI staff at all health pyramid levels. As a result, many EPI staff, including logisticians, 77 storekeepers, data managers, and vaccinators, lack the required skills to effectively do their job. 78

Although the EPI has invested significant efforts to address these challenges, five system issues have hindered progress toward building a competent workforce. First, the current training methods (notably, off-site classroom training) are mostly based on theoretical/didactical materials that are not always adapted to the Cameroonian context and that are delivered by instructors who are not sufficiently trained on the content. In addition, there are often no post-training follow-ups to ensure that training translates

84 into improved performance. Secondly, there is often poor coordination of training between partners and the government, leading to poor training quality control and sometimes duplication of efforts. Third, 85 partners often fund training activities, implying that capability-building activities can stall if partner 86 funding evaporates. Fourth, the current training system is not viewed as a continuum to the extent that 87 88 there are no clear links from pre-service to in-service to on-the-job training. Lack of this perspective often results in one-off training, which is driven by available funding for training of interest (e.g., new 89 vaccines or technology introductions). Finally, there is no monitoring and evaluation system to track 90 91 who has received/has not received training as well as knowledge utilization.

There is a need to establish a comprehensive capability program that can sustainably address current 92 and future system challenges, including the high numbers of untrained staff, high staff turnover, and 93 fast-paced and evolving technology. The first step towards establishing such a program necessitated the 94 95 conduct of a Training Needs Assessment (TNA) to quantify and characterize existing knowledge and skills gaps among EPI staff at all system tiers. In this paper, we present the findings of our study that 96 97 assessed existing capability needs among EPI staff and proposed strategies that can be leveraged to 98 address the identified gaps to inform actions or interventions from policymakers, funders, and other 99 interested immunization stakeholders.

100 Methods

101 *Study design and setting*:

A descriptive cross-sectional study was conducted using aggregated data from a survey carried out
among EPI staff from May to September 2016 across Cameroon's health pyramid - central, regional,
and district levels to establish vaccination training needs.

105 *Study population and sampling*:

Our study was based on aggregated data unveiled by the survey conducted from May to September 2016 among EPI staff from central, regional, and district levels of Cameroon's health pyramid. For the purpose of research, this data was accessed in March 7th, 2023 and authors did not have access to information that could identify individual participants. The aggregated data contained a summary outcome of interviews of staff at central and regional levels, excluding non-technical staff such as

drivers, cleaners, and security personnel. At the district level, available data was on staff considered during the survey through purposive sampling to assure proportional distribution rather than obtaining high precision. The aggregated data represents the view 28 EPI staff at central level, 52 at regional and 162 at district-level from 68 out of 200 health districts in Cameroon.

115 Inclusion and Exclusion criteria

All data from the need assessment survey conducted in 2016 representing the view of technical staff 116 working in the immunization space at the three target levels (central, regional, and districts) was 117 118 considered. At the national level, this included staff from all existing functional departments, including 119 the EPI manager and deputy, and staff from the following departments: administration and finance, 120 routine immunization and logistics, data management, monitoring and evaluation, and communication for development. At the regional level, it included the regional EPI leads, logisticians, data managers, 121 and surveillance and communication officers. At the district level, the district medical officers, the chief 122 123 of the health bureau, and EPI focal staff where included in the final analysis. At all levels, we excluded support staff such as drivers, security guards, cleaners, and those working at the secretariats without a 124 defined role as well as technical staff who did not participated in the survey. 125

126 *Data collection and procedure*

A structured excel based tool was used to extract information from aggregated data on the need assessment survey. The final tool consisted of eight sections, including basic demographic information, vaccine-specific training, core skills and competencies, the preferred format of learning, motivational factors for learning, managers' views on the current skill levels of staff under their command, human resource onboarding system at each level and skill/knowledge assessment on key immunization program concepts and practices.

133 Data processing and analysis

Once the extraction was completed, it was assess for accuracy and completeness and cleaned. Data were then exported, processed, and analyzed using Microsoft Office Excel version 2016. Results were presented using frequency and percentages for the various variables of interest.

137 Results


138 Overall, the aggregated data represents the view of 28 EPI staff at central level, 52 at regional and 162 at district-level from 68 out of 200 health districts in Cameroon. The majority of people working in EPI 139 140 had less than three years of experience in their current posts, as shown in Table 1. Moreso, our findings showed that approximately half of the respondents at all levels (central: 55%; regional: 46%, and 141 142 district: 49%) were not aware of onboarding materials for a new staff. In addition, most respondents had not received any form of pre-service training, notably 79 % at the central, 72 % at the regional, and 143 60 % at the district levels. Also, about half of the respondents at the district (50%), region (37%) and 144 central levels (58%) had not received in-service training (Table 2). 145

- 146 Table 1: Staff professional experience across the three administrative levels of the EPI in Cameroon,
- 147 2016

Professional experience*	EPI level		Tot	Total	
	n (%)				
	District	Regional	Central		
	<i>n</i> =162	<i>n</i> =52	<i>n</i> =28	<i>n</i> =242	
In all sectors					
< 3 years	10 (6)	2 (4)	2 (8)	14 (6)	
3-6 years	36 (22)	5 (10)	16 (58)	57 (24)	
>6 years	116 (72)	45 (76)	10 (34)	171 (70)	
In immunization services					
< 3 years	21 (13)	7 (14)	15 (54)	42 (18)	
3-6 years	45 (28)	16 (30)	9 (31)	70 (29)	
>6 years	96 (59)	29 (56)	4 (15)	129 (53)	

*Professional experience was defined as the total number of years an individual has worked following graduation from a training institution/university excluding internships.

148



Type of training received on	EPI level	EPI level n (%)				
immunization	n (%)					
	District	Regional	Central			
	<i>n</i> =162	<i>n</i> =52	<i>n</i> =28	<i>n</i> =242		
Pre-service						
No	97 (60)	37 (72)	22 (79)	156 (64)		
Yes	52 (32)	12 (24)	5 (17)	69 (29)		
Can't remember	13 (8)	3 (4)	1 (4)	17 (7)		
In-service						
No	81 (50)	19 (37)	15 (58)	115 (48)		
Yes	73 (45)	32 (61)	13 (42)	118 (49)		
Can't remember	8 (5)	1 (2)	0	9 (3)		

150

151 As shown in Table 3, cold chain logistics was the most covered topic during in-service training of EPI staff - 90 % of district, 83 % of regional, and 54 % of central- level EPI staff received in-service training 152 on this subject. Other important topics including but not limited to Adverse events following 153 immunization (AEFI), surveillance of vaccine-preventable diseases, conducting coverage surveys or 154 155 Supplementary Immunization Activities were less prioritized for training. On the other hand, crossdisciplinary skills necessary for day-to-day operations seem not to receive sufficient attention during 156 in-service training at all levels; for instance, less than half of district and regional staff received training 157 on planning, budgeting, and report writing. 158

Table 3: Distribution of EPI staff according to the type of topic covered during in-service training by
 159 level, 2016 160

Торіс	EPI level	Total
	n (%)	
	District Regional	Central <i>n</i> =118

<i>n</i> =73	<i>n</i> =32	<i>n</i> =13	_
66 (90)	27 (83)	7 (54)	100 (85)
57 (78)	20 (63)	0	77 (65)
55 (75)	20 (63)	0	75 (64)
38 (52)	17 (53)	4 (31)	59 (50)
44 (60)	22 (70)	0	66 (56)
32 (44)	16 (50)	0	48 (41)
38 (52)	17 (53)	4 (31)	59 (50)
54 (74)	26 (80)	1 (10)	81 (68)
51 (70)	19 (60)	0	70 (59)
31 (43)	31 (60)	0	62 (52)
23 (32)	16 (30)	0	39 (33)
17 (23)	26 (50)	0	43 (36)
38 (52)	36 (70)	0	74 (63)
21 (29)	22 (43)	0	43 (36)
	66 (90) 57 (78) 55 (75) 38 (52) 44 (60) 32 (44) 38 (52) 54 (74) 51 (70) 31 (43) 23 (32) 17 (23) 38 (52)	66 (90) 27 (83) 57 (78) 20 (63) 55 (75) 20 (63) 38 (52) 17 (53) 44 (60) 22 (70) 32 (44) 16 (50) 38 (52) 17 (53) 54 (74) 26 (80) 51 (70) 19 (60) 31 (43) 31 (60) 23 (32) 16 (30) 17 (23) 26 (50) 38 (52) 36 (70)	66 (90) $27 (83)$ $7 (54)$ $57 (78)$ $20 (63)$ 0 $55 (75)$ $20 (63)$ 0 $38 (52)$ $17 (53)$ $4 (31)$ $44 (60)$ $22 (70)$ 0 $32 (44)$ $16 (50)$ 0 $38 (52)$ $17 (53)$ $4 (31)$ $54 (74)$ $26 (80)$ $1 (10)$ $51 (70)$ $19 (60)$ 0 $31 (43)$ $31 (60)$ 0 $17 (23)$ $26 (50)$ 0 $38 (52)$ $36 (70)$ 0

161

.

.

Preferred areas of training: Core immunization topics and transferable skills 162

163 Respondents indicated the need for training in all thematic areas but with varying frequency across 164 themes and levels. Supportive supervision was the most frequently sorted topic for training, with a higher need expressed at the central level (80%). District and regional staff considered adverse events 165 following immunization (AEFI), maintenance of cold chain equipment, stock management and 166 distribution, and community engagement as critical areas where training is needed. Moreso, Over 50% 167

of staff at all levels expressed the need to be trained in preparing reports, creating work plans, delegating tasks, time management, data analysis, and preparing PowerPoint Presentations. Similarly, over half of the respondents at all levels expressed interest in gaining skills in core aspects of managing teams/working in teams. Furthermore, over 40% of respondents at all levels highlighted the need to learn how to account for project-related expenditures/expenses (Table 4).

- 173
- 174 **Table 4:** Preferred areas for training on cross-disciplinary skills among Cameroon EPI staff, 2016

Preferred transferable skill	EPI level	EPI level			
	n (%)				
	District	Regional	Central		
	<i>n</i> =152	<i>n</i> =45	<i>n</i> =30	<i>n</i> =227	
Oral communication					
Communication in public	87 (57)	30 (66)	20 (65)	137 (60)	
Negotiation and advocacy	59 (39)	11 (24)	14 (46)	84 (37)	
Mentoring	49 (32)	10 (22)	7 (23)	66 (29)	
Training/teaching skills	68 (45)	19 (43)	11 (35)	98 (43)	
Written communication					
Report writing	100 (66)	35 (78)	20 (65)	155 (68)	
Academic/research writing	35 (23)	9 (21)	13 (44)	57 (25)	
Proposal/grant writing	24 (16)	7 (15)	11 (38)	42 (19)	
Email communication	73 (48)	26 (58)	21 (69)	120 (53)	
Formal memos/Government letters	50 (33)	22 (49)	20 (65)	92 (41)	
Project management					
Creating budgets	35 (23)	17 (37)	16 (54)	68 (30)	
Monitoring budgets	38 (25)	21 (46)	14 (46)	73 (32)	
Creating workplans	90 (59)	27 (61)	19 (62)	136 (60)	

It is made available under a CC-BY 4.0 Internation	al license .
--	--------------

84 (55)	27 (60)	16 (52)	127 (56)
72 (10)			
73 (48)	13 (29)	16 (52)	102 (45)
62 (41)	20 (44)	12 (40)	94 (41)
76 (50)	22 (50)	19 (62)	117 (52)
62 (41)	16 (36)	21 (69)	99 (44)
nt 90 (59)	24 (54)	17 (58)	131 (58)
88 (58)	25 (56)	21 (69)	134 (59)
82 (54)	26 (58)	13 (42)	121 (53)
81 (53)	24 (53)	14 (46)	119 (52)
53 (35)	14 (31)	11 (35)	78 (34)
99 (65)	28 (63	16 (52)	143 (63)
71 (47)	15 (33)	16 (52)	102 (45)
74 (49)	26 (58)	15 (50)	115 (51)
27 (18)	6 (14)	11 (35)	44 (19)
49 (32)	12 (27)	1 (4)	62 (27)
-	76 (50) 62 (41) nt 90 (59) 88 (58) 82 (54) 81 (53) 53 (35) 99 (65) 71 (47) 74 (49)	76 (50) 22 (50) 62 (41) 16 (36) nt 90 (59) 24 (54) 88 (58) 25 (56) 82 (54) 26 (58) 81 (53) 24 (53) 53 (35) 14 (31) 99 (65) 28 (63) 71 (47) 15 (33) 74 (49) 26 (58)	76 (50) 22 (50) 19 (62) 62 (41) 16 (36) 21 (69) nt 90 (59) 24 (54) 17 (58) 88 (58) 25 (56) 21 (69) 82 (54) 26 (58) 13 (42) 81 (53) 24 (53) 14 (46) 53 (35) 14 (31) 11 (35) 99 (65) 28 (63 16 (52) 71 (47) 15 (33) 16 (52) 74 (49) 26 (58) 15 (50)

175

176 *Key challenges and drivers of learning*

177 Surveyed staff largely aligned on four key factors that could impact learning – multiple meetings, high

staff attrition, competing priorities, and team conflict (Table 5). With regards to motivation, financial

- incentive to learn was not among the leading motivation factors for all levels, contrary to what was previously thought. Other factors, such as obtaining a certificate of completion, being recognized by the ministry, and consideration for promotion or attending an international meeting, received better ratings than financial incentives at all levels. Classroom learning at the workplace and On-the-job mentorship and direct supervision were the top preferred learning methods in this study.
- 184

Table 5: Self-reported factors that could enhance learning among Cameroon EPI staff, 2016

187

Factor enhancing learning	EPI level			Total
	n (%)			
	District	Regional	Central	_
	<i>n</i> =162	<i>n</i> =52	<i>n</i> =28	<i>n</i> =242
Most motivational factor for learning/gaining				
new skills				
Financial incentive	68 (42)	19 (36)	8 (27)	95 (39)
Certificate of completion	91 (56)	34 (66)	8 (27)	133 (55)
Recognition from high level Ministry officials	89 (53)	29 (55)	15 (55)	133 (55)
Potential consideration for promotion	81 (50)	37 (71)	5 (18)	123 (51)
Potential for consideration for international meeting	102 (63)	37 (71)	10 (36)	149 (62)
Positive performance job review	63 (39)	21 (41)	10 (36)	94 (39)
Preferred format of learning to gain skills				
Self-paced and facilitated e-learning on a computer	13 (8)	3 (6)	0	16 (7)
Self-paced printed resource material/book	29 (18)	5 (10)	4 (14)	38 (16)
Classroom learning at the workplace	31 (19)	11 (21)	5 (17)	47 (19)
Off-site Classroom learning	26 (16)	9 (17)	4 (16)	39 (16)
On-the-job mentorship and direct supervision	24 (15)	11 (21)	6 (21)	41 (17)
On-the-job quick reference material/job aids	18 (11)	5 (10)	4 (15)	27 (11)
Key challenges staff feel could impact learning				
at the work place				
Multiple meeting	96 (59)	16 (30)	24 (87)	136 (56)
High staff attrition rate	76 (47)	19 (37)	20 (70)	115 (48)
Competing priorities	50 (31)	25 (48)	20 (70)	95 (39)

Team conflict

21 (13) 7 (13) 4 (13) 32 (13)

EPI: Expanded Program of Immunization.

188 Discussion

In this study, the majority of people working in EPI had less than three years of experience in their 189 190 current posts (Table 1), with approximately half of the respondents not aware of onboarding materials for new staff. This observation suggests high staff turnover, highlighting the need to strengthen the use 191 192 of onboarding materials which is critical to sustainability following EPI-related capacity-building efforts. Robust onboarding can dramatically improve the performance and readiness of a person who 193 194 takes on a new role in the health system. Onboarding helps build and sustain high-performing staff and maintains institutional knowledge. Therefore, onboarding materials are an essential tool of a well-195 196 functioning health system that can retain knowledge as staff changes[9]. The sub-optimal use of onboarding material despite high staff turnover arguably contributes significantly to the rising number 197 of zero-dose and under-immunized children in Cameroon. This finding is consistent with a study 198 conducted by Emily and colleagues in Tchad and Cameroon, which highlighted the role of rapid staff 199 200 turnover in implementing Gavi's health system strengthening support in both countries[10].

Moreso, the low vaccination capacity gap of EPI staff stemming from rapid staff turnover is further 201 202 compounded by sub-optimal pre-service and in-service training at all health system levels (Table 2). 203 This finding suggests that limited attention is being accorded to immunization in the curricula of nursing 204 and medical schools (pre-service training). This is consistent with a study conducted at health training 205 institutions in 12 target countries which suggested incomplete or outdated pre-service curricula, insufficient time allocation, and insufficient knowledge of the current EPI theory and practice by 206 207 lecturers[11]. Redressing this situation will, therefore, require high-level advocacy and negotiations 208 between the Ministries of Health and the ministries that govern professional education, such as the 209 ministries of higher education and vocational training [12]. However, onboarding systems within the EPI need to be strengthened to provide newcomers with the necessary skills and competencies to 210

effectively do their jobs. The finding that a sizable proportion of staff at all levels have received inservice training is reassuring in the sense that it indicates that mechanisms/systems are in place which can support in-service training. Pre-service training is fundamental in the skills acquisition of EPI staff; however, they require additional in-service training, onboarding, and supportive supervision to carry out their vaccination tasks efficiently[13].

Table 3 emphasized the knowledge gaps of EPI staff, with topics such as AEFI, surveillance, 216 Supplementary Immunization Activities, and cross-disciplinary skills such as planning, budgeting, and 217 report writing less frequently conducted. It is worth highlighting that the percentages reported for the 218 central level need to be viewed with caution for two reasons. First, some degree of functional 219 220 specialization is expected at the central level, and therefore, it will not be uncommon to see that certain 221 staff have not received in-service training on some aspects of immunization. For instance, finance staff 222 are not expected to receive in-service training on cold chain or conducting EPI coverage surveys because they may not require this knowledge for their job. Second, of the 28 central staff interviewed, 223 224 only 13 indicated that they received in-service, making it difficult to draw any meaningful conclusion 225 from the observed findings. Despite this limitation, staff at this level are expected to have a solid mastery of cross-disciplinary skills, given their role in evidence-based strategy development. And the fact that 226 227 none of the respondents from the central level reported being trained on most topics suggests that inservice training are rarely tailored to provide staff with the needed competencies to do their jobs 228 effectively. Therefore, there is a need to develop a more comprehensive and tailored capacity-building 229 plan that meets EPI staff's training needs sustainably, especially as respondents indicated the need for 230 training in all thematic areas but with varying frequency across themes and levels (Table 4). 231

232

In this study, respondents considered self-paced printed resource material/book, classroom learning at the workplace, off-site classroom learning, on-the-job mentorship and direct supervision, and on-thejob quick reference material/job aids as top learning and work aids that could be employed to improve their skills. However, based on their perspective, self-paced and facilitated e-learning on a computer was the least effective. In all, the emphasis is on repetitive practical training approaches and regular

supervision, consistent with evidence developed by Julia et al. on recommended in-service trainingtechniques based on an integrative literature review[14].

Even with a state-of-the-art platform to support in-service training, some critical factors which pose 240 serious hindrances to learning. Our study evaluated two core factors: staff motivation to learn and key 241 242 challenges they feel could impact learning. With regards to motivation, financial incentive to learn was not among the leading motivation factors for all levels. Other factors, such as obtaining a certificate of 243 completion, being recognized by the ministry, and consideration for promotion or attending an 244 245 international meeting, received better ratings than financial incentives at all levels. In addition, over 246 30% of participants at all levels indicated that positive performance reviews from a manager or 247 supervisor could motivate them to acquire new knowledge/skills. Overall, these findings suggest that 248 strategies for training should also consider non-financial incentives, such as certification, recognition by senior officials, and the institution of a system that can guarantee regular performance job reviews. 249 250 This will be important to promote (1) a culture of self-learning and (2) potentially work toward costsavings of training methods (Table 5). However, surveyed staff aligned mainly on four key factors that 251 could impact learning - multiple meetings, high staff attrition, competing priorities, and team conflict. 252 While the impact of these factors on staff professional development could not be overemphasized, it is 253 254 important to recognize that these factors could threaten the program's overall performance. Therefore, addressing them- through optimal planning and prioritization- will enhance participants' ability to learn 255 and improve the program's overall performance. 256

257

258 Conclusion

Overall, this study suggests that most EPI staff at the surveyed levels lack the requisite disciplinary and cross-disciplinary skills to effectively do their jobs. These capacity gaps, to a large extent, stem from weaknesses in pre-service and in-service training systems. According to our findings, training needs vary considerably across the different levels and within the units/ department, and only limited time is available for professional development. This variation implies that efforts need to be invested in developing a tailored capability-building program, which covers relevant immunization topics and cross-disciplinary skills for each level, and which staff sees as useful for their jobs. Also, to effectively

266	deal wi	th the diversity of staff graduates from different training institutes at all levels, there is a need to
267	institut	e a system that can guarantee targeted onboarding, mentoring, and performance review for new
268	immun	ization staff. Our study raised crucial points on preferred learning methods and showed that
269	factors	such as obtaining a certificate of course completion are more motivating than providing financial
270	incenti	ves – which are useful for effectively planning EPI staff capability building programs.
271		
272	Recom	umendation
273	Based	on our findings, we recommend immunization stakeholders:
274	1.	Establish a competency framework that clearly defines the knowledge, skills, attributes,
275		structures, roles and approaches needed for people within EPI to effectively do their job.
276	2.	Leverage the competency framework to upgrade the capability of existing staff on core
277		disciplinary and cross-disciplinary skills that are indispensable for their day-to-day operations.
278	3.	Establish a robust onboarding system for new employees to ensure that updated technical and
279		management/communication resources are available for each post/role.
280	4.	Upgrade and standardize the current performance review mechanisms at all levels, as this can
281		be a good opportunity to identify and address capability-building needs.
282	5.	Strengthen pre-service training by ensuring that core immunization aspects are sufficiently
283		integrated into the curriculum of, at least nursing and medical schools.
284	6.	Enhance in-service training, whenever it is considered, by adapting and applying adult learning
285		theories, strengthening the capacity of instructors, and appropriately selecting trainees, amongst
286		others.
287	7.	Establish a monitoring and evaluation framework to gather key information from every training
288		event and track training effectiveness. Develop a routine database to gather this key information
289		from every training event.
290	Limita	tion

This assessment had four major methodological limitations. First, data were collected from 68 health districts, representing just 36% of health districts in the country. This limitation implies that this evaluation should not be considered a comprehensive presentation of the entire scope of capability gaps

294 at the district level. Moreso, the study is based on a need assessment conducted in 2016; thus may not represent the current reality. Second, as per the objective of the TNA, this evaluation did not evaluate 295 capacity gaps at the health facility level, where capacity needs mostly revolve largely around clinical 296 practices. As such, the findings are not applicable to this level. Third, most of the findings of this 297 298 evaluation are largely based on recall and perceptions. Fourth, some staff at the central level did not respond to some questions, which implies that the results for some thematic areas may not represent the 299 variables explored. However, given the emphasis of the evaluation on pre-service and in-service 300 301 training, with a lens on disciplinary and cross-disciplinary skills, the authors are confident that these limitations have not undermined the findings – the scope of data collection allowed the assessment team 302 303 to draw out conclusions across nine thematic areas that may be applicable to central, regional and 304 districts levels.

305 Author contributions

Conception & Design: AA, FZL, MV, JNM Statistical analysis: AA, JNM, MV, FZL Initial draft
preparation: AA, SCN, JG, NNE Review and Editing: SCN, VNA, MCM, NB, BMN, STN, AAN,
PDM, CT, YS Supervision: YS All Authors read and approved submission of the final article.

309 Ethics approval and consent to participate

310 This study did not require ethical approval as it did not contain individual-level data collection.

311 Availability of data and materials

312 Data used for this research is available from the corresponding author upon reasonable request.

313 Funding

314 This research received no external funding.

315 Conflicts of Interest

316 The authors declare no conflict of interest.

317 References

perpetuity.	
It is made available under a CC-BY	4.0 International license .

318 319	1.	Chard AN, Gacic-Dobo M, Diallo MS, Sodha SV, Wallace AS (2000) Routine Vaccination Coverage — Worldwide, 2019. Morb Mortal Wkly Rep 69:1706–1710
320 321 322 323	2.	WHO Immunization Data portal. https://immunizationdata.who.int/compare.html?COMPARISON=type1WIISE/MT_AD_CO V_LONG+type2_WIISE/MT_AD_COV_LONG+option1_DTP_PLUS_coverage+option2 DTP_coverage&CODE=CMR&YEAR=. Accessed 19 Aug 2023
324 325	3.	Vandelaer J, Bilous J, Nshimirimana D (2008) Reaching Every District (RED) approach: a way to improve immunization performance. Bull World Health Organ Mar;86(3):A-B:
326 327	4.	Cameroon - urbanization 2022. In: Statista. https://www.statista.com/statistics/446865/urbanization-in-cameroon/. Accessed 19 Aug 2023
328 329	5.	Vaccine Investment Strategy 2024. https://www.gavi.org/our-alliance/strategy/vaccine- investment-strategy-2024. Accessed 19 Aug 2023
330 331	6.	Immunization Agenda 2030. https://www.who.int/teams/immunization-vaccines-and- biologicals/strategies/ia2030. Accessed 19 Aug 2023
332 333 334	7.	Immunization coverage and vaccine-preventable diseases. https://www.who.int/data/gho/data/themes/topics/sdg-target-3_b-development-assistance-and-vaccine-coverage. Accessed 19 Aug 2023
335 336	8.	Yauba S, Harmelle EE, Marius VZ, et al (2019) Availability and Status of Vaccine Cold Chain Equipment in Cameroon. J. Vaccines Vaccin.
337 338	9.	Woolforde L (2012) Onboarding Nurse Practitioners: A Healthcare System Approach to Interprofessional Education. Nurse Lead 10:32–35
339 340 341	10.	Dansereau E, Miangotar Y, Squires E, et al (2017) Challenges to implementing Gavi's health system strengthening support in Chad and Cameroon: results from a mixed-methods evaluation. Glob Health 13:83
342 343	11.	Mutabaruka E, Nshimirimana D, Goilav C, Meheus A (2005) EPI training needs assessment in 12 African countries, 2002-2004. Publ Met UAntwerpen Adres 3:14
344 345 346	12.	Juma M, Masresha B, Adekola A, Dochez C (2022) Strengthening pre-service training of healthcare workers on immunisation and effective vaccine management: the experience of Kenya Medical Training College. Pan Afr Med J 41:
347 348	13.	Nicol E, Turawa E, Bonsu G (2019) Pre- and in-service training of health care workers on immunization data management in LMICs: a scoping review. Hum Resour Health 17:92
349 350 351	14.	Bluestone J, Johnson P, Fullerton J, Carr C, Alderman J, BonTempo J (2013) Effective in- service training design and delivery: evidence from an integrative literature review. Hum Resour Health 11:51
252		