It is made available under a CC-BY-NC-ND 4.0 International license .

<u>TITLE:</u> A Retrospective Analysis of Serious Adverse Events and Deaths in US-Based Lifestyle Clinical Trials for Cognitive Health

AUTHORS: Mickeal N. Key<sup>1</sup>, Ashley R. Shaw<sup>1</sup>, Kirk I. Erickson<sup>2</sup>, Jeffrey M. Burns<sup>1</sup>, Eric D. Vidoni<sup>1</sup>

## **INSTITUTIONS:**

1) University of Kansas Alzheimer's Disease Research Center, University of Kansas Medical Center,

## Fairway, KS, USA

2) AdventHealth Research Institute, Neuroscience, Orlando, FL, USA

KEYWORDS: Death; Safety; Exercise; Diet; Research Methodology; Data Reporting

## ABSTRACT:

This retrospective analysis assessed the serious adverse events and deaths reported in lifestyle clinical trials designed to enhance cognitive health in older adults living in the United States. Data was collected from studies conducted between January 1, 2000, and July 19, 2023, using the ClinicalTrials.gov application programming interface. Our query revealed that 76% of these studies did not report trial results. The remaining studies with reported results were categorized under one of four intervention types: Cognitive/Behavioral, Exercise/Movement, Diet/Supplement, and Multi-modal. When all trial types are considered together, the results indicate that lifestyle clinical trials are safe, with no significant increase in relative risk of experiencing an SAE in an intervention group over a control group. And although the increase in relative risk of death in an intervention group over a control group was significant at 28% (X<sup>2</sup> (1, N = 36), p < 0.00688), the probability of death was not higher than the U.S. mortality rates by age. When assessing the data using intervention type, Diet/Supplement trials and Multi-modal trials both had an increase in relative risk of experiencing an SAE in the intervention over the control group, with Diet/Supplement trials at 16% (X<sup>2</sup> (1, N = 2), p < 0.0263) and Multi-modal trials at 365% (X<sup>2</sup> (1, N = 5), p < 0.00213). The Diet/Supplement trials also had an increased risk of death at 67%

It is made available under a CC-BY-NC-ND 4.0 International license .

(X<sup>2</sup> (1, N = 2), p < 0.000197). These results should be taken with careful consideration. Due to such a low reporting rate, the 36 studies included in this analysis do not accurately represent the majority of lifestyle clinical trials conducted in the U.S. This study is valuable in that it highlights the importance of reporting clinical trial results, which will improve transparency in trial results and allow for more accurate assessments of safety in the growing field of cognitive aging and lifestyle interventions for older adults.

It is made available under a CC-BY-NC-ND 4.0 International license .

### **Introduction**

Over the last two decades, many well-controlled clinical trials have improved our understanding of the potential for lifestyle modifications such as diet, exercise, meditation, and cognitive training to enhance cognitive health <sup>[1]</sup>. A recent World Health Organization analysis estimated that up to 40% of dementia cases are attributable to modifiable factors, and approximately 10% of those are potentially related to physical health behaviors <sup>[2, 3]</sup>.

Continued investigation of behavioral, non-pharmaceutical clinical trials for cognitive health warrants an assessment of their safety. Common familiarity with healthy behaviors such as walking or eating more vegetables is not equivalent to safe study conduct and should not be assumed to be safer than investigational medications. Conversely, it should not be assumed that lifestyle interventions for cognitive health, generally focused on older adults, pose an elevated risk for participants. Institutional review boards, oversight committees, and investigative teams all need reliable data to adequately assess the risk associated with these interventions.

The safety of a clinical trial is often measured in terms of the frequency, severity, and relatedness of adverse events. An adverse event (AE) is commonly defined as any physical or psychological sign, symptom, or disease experienced by a research participant during the study period. A serious adverse event (SAE) is commonly defined as any adverse event that results in death, is life-threatening, requires hospitalization, causes disability, or poses a significant hazard. A standard approach for assessing and reporting both AEs and SAEs is provided by the NIA Adverse Event and Serious Adverse Event Guidelines, Version 5.

The goal of this retrospective analysis was to evaluate the safety of reported lifestyle clinical trials for cognitive health in older adults aged 60 and older in the United States by assessing 1) serious adverse events (SAE) and 2) deaths in lifestyle clinical trials. Our hypothesis was that participants in intervention groups would experience a higher risk of SAE (including death) than participants in control It is made available under a CC-BY-NC-ND 4.0 International license .

groups. Due to the characteristics of the study population, older adults, we also hypothesized that interventions requiring exercise would experience a higher risk of SAE (including death) than any other lifestyle intervention type. Our intended outcome was to provide context for the current state of safety in lifestyle clinical trials, ideally to inform future trial design, implementation, and dissemination of trial results.

#### **Methods**

Data on the frequency of SAE and deaths in lifestyle clinical trials was extracted from ClinicalTrials.gov using their application programming interface (API). We limited our search to registered clinical trials completed between January 1, 2000 and July 19, 2023. Other search criteria included 1) minimum age of 60 years for inclusion criteria, 2) at least one explicitly identified outcome of cognitive function, 3) randomized allocation, 4) intervention with a clearly identifiable control condition, and 5) a United States-based coordinating location.

The search string supplied to the API was constructed with consideration for prior systematic reviews <sup>[1, 4-7]</sup>, and included:

Intervention/Treatment: Physical Activity OR Exercise OR Sports OR Walking OR Biking OR Bicycling OR Running OR Fitness OR Yoga OR Strength OR Resistance OR Aerobic OR Endurance OR Flexibility OR Tai Chi OR Qi Gong OR Balance OR Diet OR Dietary OR Nutrition OR Nutritional OR Food OR Supplement OR Vitamin OR Multivitamin OR Cognitive OR Cognition OR Reading OR Games OR Gaming OR Mindfulness OR Meditation OR Sleep OR Dance

Outcome measure: Cognition OR Memory OR Executive OR Thinking OR Cognitive OR Attention OR Processing

We defined lifestyle interventions as those designed to improve physical and/or cognitive health. This definition excludes interventions where a drug or device initiates the physical or cognitive change. Interventions using natural products or therapeutic compounds requiring a prescription were also

excluded. Due to the broad nature of lifestyle interventions, it was necessary to categorize similar interventions into categories. For this review, two investigators (MNK, EDV) reached consensus on assignment of each study to one of four broad lifestyle intervention categories, referred to as "Intervention Type". The "Cognitive/Behavioral" category included any intervention that involved cognitive training, brain games, sensory interventions, meditation, and assessment/screening interventions. The "Exercise/Movement" category included any intervention that involved exercise, yoga, dance, stretching/toning or similar. The "Diet/Supplement" category included any study that involved feeding, dietary changes, supplements, and bioactive natural products. Finally, the "Multimodal" category included any intervention that made use of two or more intervention categories (e.g., an intervention that had participants undergo cognitive training and exercise could be considered Multimodal).

For each included study, the number of individuals affected and at risk of experiencing an SAE, as well as the number of individuals who died during the study period, were evaluated. To provide additional context, risk and risk ratios were calculated for SAEs and deaths, both in studies overall and by intervention type. The risk of an event for intervention and control group was calculated as the number of participants affected divided by the number of participants at risk. The risk ratio was calculated as the risk of an event in the intervention divided by the risk of an event in the control group. The risk ratios are accompanied by 95% confidence intervals and a chi-square test to evaluate whether there was a statistically significant difference between intervention and control groups. The API query and statistical analyses were all performed using R Statistical Software<sup>[8]</sup>.

#### Results

Of the 6,497 lifestyle intervention studies on ClinicalTrials.gov that met the review criteria, 6461 were excluded from analysis: 4953 did not have results posted; 885 were not lifestyle interventions as described in the previous section; 589 included individuals below age 60; 19 had no cognitive outcome

It is made available under a CC-BY-NC-ND 4.0 International license .

listed; 13 were centralized outside the US; 2 did not include random allocation. Thirty-six (36) studies remained for analysis.

Of these 36 included studies, all reported SAEs and 30 studies (83%) reported deaths. The median study duration for all studies was 6 months, with the shortest intervention period lasting 2 hours (0.00274 months) and the longest intervention lasting 10 years (120 months). The percentage of participants experiencing a SAE in the included lifestyle clinical trials was 5.13%. The percentage of deaths during the study period for the included studies was 2.21%. Table 1 includes the number of studies in each of the four intervention types, the study duration, and documents the number of SAEs and deaths for each, as well as overall.

The overall risk ratio for lifestyle interventions describes a 5.8% relative decrease in risk for experiencing an SAE in the intervention group compared to the control group (0.941 times as much probability). This decrease in risk was not statistically significant, as evidenced by a chi-square test,  $X^2$  (1, N = 36), p < 0.308. When SAE risk was assessed by intervention type, Diet/Supplement and Multi-modal studies were the only ones to have statistically significant risk ratios. Both had SAE risk ratios over 1, indicating an increased relative risk of experiencing an SAE in the intervention over control, with Diet/Supplement studies at an increased relative risk of 16% (X<sup>2</sup> (1, N = 2), p < 0.0263) and Multi-modal studies at an increased relative risk of 365% (X<sup>2</sup> (1, N = 5), p < 0.000213).

When considering overall deaths during the study duration of included trials, the risk ratio describes a 28% relative increase in risk of death in the intervention group than in the control group (1.28 times as much probability). This increase in risk was statistically significant (X<sup>2</sup> (1, N = 36), p < 0.00688), and translates to a 2.98% probability of death in the intervention group and a 2.33% probability of death in the control group. This finding was likely driven by Diet/Supplement studies, which was the only intervention type to have a statistically significant risk ratio at 67% (X<sup>2</sup> (1, N = 2), p <

It is made available under a CC-BY-NC-ND 4.0 International license .

0.000197). Risk and risk ratios, along with 95% confidence intervals and chi-square p-values across all studies, and separated by intervention type, are described in Table 2.

#### Discussion

Our retrospective analysis of reported serious adverse events (SAEs) and deaths across more than 20 years of reported clinical trial results suggests a high degree of safety within lifestyle clinical trials related to cognitive health in older adults. The increased use of Consolidated Standards of Reporting Trials (CONSORT) guidelines <sup>[9]</sup> has likely improved safety reporting in the last decade, especially with the extended guidelines for safety published in 2004 <sup>[10]</sup>. Our analysis also revealed significant gaps and irregularities in reporting of SAEs (including deaths) by the field, with 76% of completed studies in our ClinicalTrials.gov query not posting results.

The overall probability of an enrollee experiencing at least one SAE was about 5%, and was consistent between intervention and control conditions. The comparable risk between SAEs occurring in the intervention group versus the control group overall is encouraging but should be interpreted with caution. We were unable to evaluate the relatedness of SAE to intervention, a common classifier in safety reporting. Rather we presumed that if lifestyle interventions conveyed significant additional risk, that risk would be identifiable in the aggregate risk ratio. A higher risk ratio of intervention was noted for both the Diet/Supplement and Multi-modal interventions, though this must be considered carefully, as both categories had 2 and 5 studies, respectively, to evaluate.

While the overall risk of death for participants in an intervention group was statistically higher than those in a control group, the probability was not higher than the U.S. mortality rate. Between 1% and 5% probability of death in a study may appear concerning, until one considers that the probability of death in a given year for a 60-year-old in the US is between 0.7% and 1.3% and steadily increases with medRxiv preprint doi: https://doi.org/10.1101/2023.09.27.23296243; this version posted September 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

age. For a 75-year-old, the probability of death in a given year is 2.77% for women and 4.06% for men,

climbing to 8.3% and 10.9% respectively by 85, a common upper age limit for trials <sup>[11]</sup>.

The data also raise the potential of reporting biases inherent in the execution of trials. For example, the Multi-modal and multi-armed Diet/Supplement studies may create more contact points for participants to report SAE to study staff. Conversely, preconceived expectations about the safety of exercise for older adults may result in closer monitoring of participants, and an artificially lower probability of SAE in Exercise interventions, due to earlier intervention modification or termination at sub-SAE threshold events.

There are limitations to this retrospective analysis. First, not all studies reported results uniformly across our analysis period. We relied on reporting in ClinicalTrials.gov, which has only been required for all drug, biologic, and device clinical trials regulated by the U.S. Federal Drug Administration since 2017<sup>[12]</sup>. This requirement also applies to all clinical trials funded by the National Institutes of Health (NIH) and the Department of Veterans Affairs (DVA). Although the database supports semistandardized reporting, we found that not all studies reported SAE and deaths. This inconsistency was recently noted in the updated CONSORT Harms 2022 guidance<sup>[13]</sup>. Broad uptake and universal implementation of these standardizing recommendations would improve our understanding of safety in our clinical research and in our interventions under investigation <sup>[14]</sup>. Another factor in the number of studies included in our analysis is the fact that most lifestyle clinical trials would not fall under FDA regulation, which ultimately means that our sample is almost exclusively NIH and DVA-funded trials. While trials with non-federal and industry funding may register in ClinicalTrials.gov, they are not obligated to comply with established policies on the submission of study results.

Regardless of the reason, insufficient reporting of clinical trial results undermines transparency in the scientific process, our commitment to the responsible conduct of research, hinders trust building with the general public, and impedes the ability to monitor trends in lifestyle clinical trials that could

It is made available under a CC-BY-NC-ND 4.0 International license .

reveal equity issues not readily apparent in other data such as intervention effectiveness or adherence. Underreporting is a source of bias for the field and a disservice to the participants who accepted risk for medical advancement, and the public who often funded the work.

Despite these limitations, our analysis has important implications for the growing field of cognitive aging and lifestyle clinical trials. Cognitive health studies have been a major focus of National Institute of Health funding in recent years. NIH requires a high degree of oversight and reporting, including data and safety monitoring committees, trial registration, and regular reporting. Additionally, cognitive health studies for people without dementia typically focus on older adults, for whom there is often greater concern regarding lifestyle modifications. Understanding the safety of these studies is and will continue to be paramount as the field grows. Though our estimates do not inform population-wide estimates of death and SAE, of which the literature is scant, the results presented are similar to the reported incidence of sudden death with physical activity in a general population of all ages <sup>[5]</sup>. As investigators, we are regularly questioned by participants, Data and Safety Monitoring Committees, Institutional Review Boards, and others, as to the safety of lifestyle interventions, without sufficient data. We strongly recommend all clinical trials fully report study results to clinicalTrials.gov and adopt the CONSORT Harms 2022 guidance for future reporting, regardless of funding source. We expect these steps will lead to greater trustworthiness and rigor in cognitive health trials going forward.

It is made available under a CC-BY-NC-ND 4.0 International license .

**Declaration of funding**: MNK was supported by fellowship funding from the National Institute on Aging of the National Institutes of Health under award number 1T32AG078114-01. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. There is no other funding to report.

**Declaration of financial/other relationships**: The authors have no financial or other relationships to declare related to this manuscript.

**Author contributions**: Conceptualization: EDV; Methodology: MNK and EDV; Writing, Review, & Editing: MNK, ARS, KIE, JMB, and EDV; Supervision & Project Administration: JMB and EDV. All authors have read and agreed to the published version of the manuscript.

Acknowledgments: No assistance in the preparation of this article is to be declared

It is made available under a CC-BY-NC-ND 4.0 International license .

Table 1. Studies Reporting SAEs and SAE Counts by Intervention Type									
	Overall	Cognitive/ Behavioral	Diet/ Supplements	Exercise/ Movement	Multi-Modal				
Number of Studies, n <sup>1</sup>	36	18	2	11	5				
Median Length of Studies in Months, n (range)	6 (0.00274-120)	6 (0.00274-120)	18.2 (2.76-33.6)	5.98 (0.23-18)	8.49 (2.76-18)				
Participants at Risk, n	21,042	8,624 10,289		1,082	1,047				
Serious Adverse Events		•			·				
Studies Reporting SAEs, n (%)	36 (100%)	18 (100%)	2 (100%)	11 (100%)	5 (100%)				
Participants Affected by SAE, n <sup>2</sup>	1,079	182	765	78	54				
Number of SAE, n <sup>3</sup>	1,111	83	942	32	54				
Deaths		•			·				
Studies Reporting Deaths, n (%)	29 (81%)	14 (78%)	2 (100%)	9 (82%)	4 (80%)				
Number of Deaths, n	466	256	201	3	6				

<sup>1</sup>Meeting inclusion criteria for analysis.

<sup>2</sup>Number of unique participants who were affected by at least one serious adverse event (SAE), not including death.

<sup>3</sup>Total Number of SAE, not including deaths. Number of SAE may be more than number of unique participants affected in cases where a participant had more than one SAE. However, we noted that two cognition intervention trials and one exercise intervention trial incorrectly underreported the total number of SAE vs the total number of participants affected resulting in the number of SAE being lower than the number of participants affected by SAE.

Table 2. Risk Ratio for Serious Adverse Events (SAEs) & Death by Intervention Type													
	SAEs						Deaths						
	Individuals	Individuals	Risk	Risk	Confidence	Chi-	Individuals	Individuals	Risk	Risk	Confidence	Chi-	
	w/ SAE	At Risk		Ratio	Interval	Square	Died	At Risk		Ratio	Interval	Square p-	
						<i>p</i> -value						value	
Cognitive/Behavioral													
Intervention	99	5143	0.0192	0.807	0.605,1.08	0.145	131	2661	0.0469	0.993	0.782,1.26	0.958	
Control	83	3481	0.0238				125	2522	0.0472				
Diet/Supplement													
Intervention	346	4262	0.0812	1.16	1.02,1.34	0.0263	109	4153	0.0255	1.67	1.27,2.21	0.000197	
Control	419	6027	0.0695				92	5935	0.0152				
Exercise/Movement													
Intervention	39	592	0.0659	0.810	0.529,1.24	0.335	1	506	0.0019	0.441	0.0402,4.85	0.492	
Control	39	480	0.0813				2	447	0.0044				
Multi-modal													
Intervention	49	710	0.0690	4.65	1.87,11.6	0.000213	5	692	0.0072	2.34	0.275,20.0	0.420	
Control	5	337	0.0148				1	325	0.0030				
Overall													
Intervention	533	10,707	0.0498	0.941	0.838,1.06	0.308	246	8006	0.0298	1.28	1.07,1.53	0.00688	
Control	546	10,325	0.0520				220	9226	0.0232				

It is made available under a CC-BY-NC-ND 4.0 International license .

# **References**

- Yu JT, Xu W, Tan CC, Andrieu S, Suckling J, Evangelou E, et al. Evidence-based prevention of Alzheimer's disease: systematic review and meta-analysis of 243 observational prospective studies and 153 randomised controlled trials. J Neurol Neurosurg Psychiatry. 2020;91(11):1201-9.
- 2. Lazar RM, Howard VJ, Kernan WN, Aparicio HJ, Levine DA, Viera AJ, et al. A Primary Care Agenda for Brain Health: A Scientific Statement From the American Heart Association. Stroke. 2021;52(6):e295-e308.
- 3. Livingston G, Huntley J, Sommerlad A, Ames D, Ballard C, Banerjee S, et al. Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. Lancet. 2020;396(10248):413-46.
- Psaltopoulou T, Sergentanis TN, Panagiotakos DB, Sergentanis IN, Kosti R, Scarmeas N.
  Mediterranean diet, stroke, cognitive impairment, and depression: A meta-analysis. Ann Neurol. 2013;74(4):580-91.
- 5. Aune D, Schlesinger S, Hamer M, Norat T, Riboli E. Physical activity and the risk of sudden cardiac death: a systematic review and meta-analysis of prospective studies. BMC Cardiovasc Disord. 2020;20(1):318.
- 6. Xu W, Wang HF, Wan Y, Tan CC, Yu JT, Tan L. Leisure time physical activity and dementia risk: a dose-response meta-analysis of prospective studies. BMJ Open. 2017;7(10):e014706.
- Shaw AR, Perales-Puchalt J, Johnson E, Espinoza-Kissell P, Acosta-Rullan M, Frederick S, et al. Representation of Racial and Ethnic Minority Populations in Dementia Prevention Trials: A Systematic Review. J Prev Alzheimers Dis. 2022;9(1):113-8.
- 8. R\_Core\_Team. R: A language and environment for statistical computing. R Core Team. Vienna, Austria: R Foundation for Statistical Computing; 2022.
- 9. Moher D, Hopewell S, Schulz KF, Montori V, Gøtzsche PC, Devereaux PJ, et al. CONSORT 2010 explanation and elaboration: updated guidelines for reporting parallel group randomised trials. Bmj. 2010;340:c869.
- 10. Ioannidis JP, Evans SJ, Gøtzsche PC, O'Neill RT, Altman DG, Schulz K, et al. Better reporting of harms in randomized trials: an extension of the CONSORT statement. Ann Intern Med. 2004;141(10):781-8.
- 11. Administration SS. Actuarial Life Table. 2020.
- 12. Zarin DA, Tse T, Williams RJ, Carr S. Trial Reporting in ClinicalTrials.gov The Final Rule. N Engl J Med. 2016;375(20):1998-2004.
- Butcher NJ, Monsour A, Mew EJ, Chan AW, Moher D, Mayo-Wilson E, et al. Guidelines for Reporting Outcomes in Trial Reports: The CONSORT-Outcomes 2022 Extension. Jama. 2022;328(22):2252-64.
- 14. Junqueira DR, Zorzela L, Golder S, Loke Y, Gagnier JJ, Julious SA, et al. CONSORT Harms 2022 statement, explanation, and elaboration: updated guideline for the reporting of harms in randomised trials. Bmj. 2023;381:e073725.