1 2	The Patient Safety Curriculum: An Interventional Study on the Effectiveness of Patient Safety Education for Jordanian Nursing Students
3	<sup>1</sup> Ahmad H. Al-Nawafleh*
4	<sup>2</sup> Sultan Musleh
5	<sup>3</sup> Nofal Nawafleh
6	
7	
8	
9	
10	
11 12	<sup>1</sup> Associate Professor of Nursing Management and Leadership, Faculty of Nursing, Mutah University, email: <u>alnawafleh@mutah.edu.jo</u>
13 14	<sup>2</sup> Associate Professor of Adult Health Nursing, Faculty of Nursing, Mutah University. Email: <u>sultanmosleh@yahoo.com</u> ; <u>s.mosleh@mutah.edu.jo</u>
15 16	<sup>3</sup> Medical Student, Faculty of Medicine, Mutah University. Email: <u>nofal.nawafleh@yahoo.com</u>
17	
18	* Corresponding author:
19	Ahmad H. Al-Nawafleh (Associate Professor of Nursing Management and Leadership,
20	Faculty of Nursing, Mutah University, Phone: +962797402080, email:
21	alnawafleh@mutah.edu.jo

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

23	The Patient Safety Curriculum: An Interventional Study on the Effectiveness of Patient
24	Safety Education for Jordanian Nursing Students
25	Abstract
26	
27	The purpose of the study was to assess the effectiveness of the World Health Organization's
28	(WHO) patient safety curriculum (PSC) in improving the patient safety education for nursing
29	students in Jordanian higher education programmes.
30	Methods: An interventional design adopting a pre-test and post-test was used. A questionnaire
31	survey was administered to 373 nursing students before and after the curriculum. Students were
32	asked to complete pre-test and post-test questionnaires on self-awareness of patient safety.
33	Results: students had moderate knowledge of factors that influence patient safety and scored as
34	the highest, (mean 3.45, SD 0.94). The greatest improvement was in the role of the health
35	organisation in error reporting (the mean difference was 0.97, P=0.001). Taking into consideration
36	the essential measurements, the results obtained after the post-test regarding the two patient safety
37	topics showed a significant increase after completing the training, demonstrating that the patient
38	safety course was effective.
39	Conclusion: the study highlights the importance of developing a curriculum in nursing schools
40	that incorporates patient safety education. The WHO PSC guide can be a great start in this domain.
41	
42	Keywords
43	
44	Nursing curriculum, nurse education; patient safety; competence; Knowledge; attitude; skills;
45	WHO
46	
47	Word Count: 3610
48	

skills;

## 50 Introduction

51 Patient safety and nursing mistakes are now a global concern for providing high quality health 52 care. In 1999, the Institute of Medicine (IOM) released a starting report on the number of deaths caused by medical errors annually in the United States (1). According to the report titled "To Err Is 53 Human: Building a Safer Health System", by the IOM, It was approximately that medical errors 54 55 caused around 98,000 fatalities on a yearly basis (1, 2, 3). The report warns policy makers, hospitals, health care staff and patients about the main problems related to patient safety. The 56 57 global focus on mitigating the negative impacts of healthcare is of utmost importance, and in the 58 last ten years, there has been significant increase in patient safety efforts due to advances in patient safety sciences (4, 5, 6, 7, 8). While, it is impossible to predict when a patient harm may occur, 59 there are measures that can be considered to prevent it from happening again. Healthcare 60 professionals and organizations should learn from previous mistakes to avoid future commitments 61 62 of them.

However, the provision of healthcare in certain situations continues to be potentially dangerous
despite earlier efforts (9, 10). Patient safety experts conclude that a major reform of nursing
education in nursing schools and resident training programmes is essential for essential
improvements in patient safety (11, 12, 13).

Despite the significance of ensuring patient safety and the necessity of raising awareness about it, only a small number of educational institutions include this subject in their syllabus (6, 11, 14). Historically, the educational program for nursing and medical students has concentrated exclusively on practical abilities; such as identifying illness, providing treatments, providing post treatment care and monitoring (9, 12, 15, 16). However, the concepts of communication skills, root cause analysis, system thinking, human factor science has been disregarded to a great extent (14, 17, 18). It is essential for patient safety that all nursing students at the undergraduate level to

possess all these essential skills and the required proficiency to reduce any harm to patients (3, 10).
It is urgent to integrate knowledge and skills about patient safety in the health care field into the nursing student curriculum.

In Jordan, nursing education has been identified as a priority and efforts have been made to improve the quality of nursing education. The PSC was introduced into undergraduate nursing programs in Jordan as a strategy to improve patient safety education for nursing students (19). This study aimed to examine the impact of PSC on nursing students' knowledge, skills, and attitudes towards patient safety. This curriculum design is based on the patient safety guide proposed by the World Health Organization (4).

### 83 Background

There is no formal educational training specifically for patient safety in Jordanian universities' nursing programs up to the time of writing this report. However, the topic of patient safety is mentioned sporadically in the nursing curriculum, often related to ensuring that the student nurses are being competent in their practice (19, 20, 21).

There is now a new drive to reform the nursing curriculum in our institution to ensure that we incorporate the most relevant teaching topics but also use the most appropriate teaching methods. Patient safety, along with other topics such as palliative care and health information systems, is on the top list of topics that we need to integrate into our curriculum. Therefore, "the Multiprofessional Patient Safety Curriculum Guide" (4) was a good opportunity to start working in this direction.

The opportunity to participate in this worldwide assessment as a "Complementary Test Site" is open to all schools and universities that specialize in nursing, dentistry, medicine, midwifery, pharmacy, and other healthcare professions (4). These institutions should be interested in incorporating topics from this guide that were adopted in similar studies (15, 17, 22, 23). These

98 universities and schools should be recognized by the national regulatory bodies of their country99 and be able to demonstrate an interest in teaching patient safety (4).

100 The university under study is now reforming its curriculum for preregistration nursing training 101 according to the University Deans Council recommendations. The new curriculum includes new 102 topics to be taught to prospective students, such as palliative care, end-of-life care and patient 103 safety. Therefore, the current guide provides an important tool for the university to utilize in 104 integrating patient safety education into our pre-registration nursing curriculum.

This study aims to evaluate the effectiveness of teaching the WHO PSC to undergraduate nursingstudents in the Jordanian context. Therefore, it has two objectives; to determine:

The value of the curriculum guide for students. The usability of the curriculum guide as an
educational tool for health education in universities and schools in Jordan.

• Impact/effect of the curriculum guide on student knowledge of patient safety in Jordan.

110

### 111 Literature Review

The WHO PSC Guide: "the multi-professional edition" provides a common platform for nurses to address their role to play in improving patient safety (4). Although nursing professionals must ensure the practice does not harm, evidence indicates that nursing education is still lagging to integrate patient safety into their curriculum (16). Therefore, this guide highlights understanding of the need to integrate the safety of patients into undergraduate nursing curriculum.

Huang and her colleagues (24) explored the factors that affect the competence of PS of nursing
students who completed a national online survey. They recruited 732 Chinese students during 2018
and 2019 across various locations (24). The main outcome factor was the rating of the patient

120 safety survey. The study found that Chinese nursing graduates showed moderate confidence in 121 their clinical safety skills, but lacked confidence in areas such as learning about socio cultural or 122 context dependent patient safety (PS) and discussing PS related topics like effective 123 communication and understanding human and environmental factors (24). Additionally, less than 124 half of the participants felt comfortable talking to individuals who engaged in unsafe practices and 125 expressing concerns about adverse reactions. Overall, these findings shed a light on the PS skills of 126 Chinese nursing graduates.

In a study conducted by Suliman (25), the level of PS literacy among nursing students in two 127 128 Jordanian universities was assessed. The study found that PS issues are not given much importance 129 in the nursing education and curricula in Jordan. The study was conducted through a cross-130 sectional survey involving 297 nursing students. The results indicated that the students had 131 moderate satisfaction with their knowledge and skills on most parameters. They were more confident in the knowledge and skills learned in the classroom compared to clinical training. 132 Additionally, senior students lacked confidence in their knowledge and skills, particularly when it 133 134 comes to writing medical error reports. The study suggests that contextual and sociocultural issues 135 related to PS should be emphasized in the nursing curriculum to enhance students' competence in 136 speaking up when they witness healthcare errors (25). To bridge the gap between theory and 137 practice, the study recommends adding PS elements to the nursing curriculum.

A study was conducted in Jordan to measure the perceptions of PS during four consecutive semesters (19). It emphasised the inclusion of PS in training programmes and nursing education. PS in the nursing curriculum will emphasise the importance of reducing medical errors, highlighting clinical excellence, and improving health outcomes. The study was a survey of 158 nursing students in their final year. It found that only 25% of the participants had used an incident report for the errors they observed. However, during their practice in clinical practice, 62% of students.

145

## 146 Justification of the Study

Patient safety is becoming a standard that is common in global and local healthcare practice. However, many nurses still do not adequately practice it. Furthermore, it is not widely taught to nursing students (16). The WHO developed a curriculum to help the nurses of the future to practice PS and thus help create a culture of safety for all patients (4).

151 It is important when adopting a training curriculum to find out whether the students can achieve 152 the competencies required after its application. Furthermore, it is important to find out how the 153 training will fit into the curriculum of the nursing program. As a result, nurses' programme 154 developers should have a list of core training objectives based on agreed core competencies and examples of training programs selected by teachers on the basis of the core training objectives and 155 target groups. When designing a training program, the developer must decide which of these skills 156 157 should be taught. Selecting a level depends on the expected level of competence and characteristics of the student (e.g. previous training in PSC, research methods, epidemiology and 158 statistics), as well as the level of training necessary to establish basic knowledge and 159 160 understanding of nursing.

161

### 162 Methods

Interventional study adopting the pre-test and post-test method. The intervention was based on the WHO PSC guide for multidisciplinary health professions (4, 13, 16, 17). We taught selected topics from this curriculum to all nursing students as part of several courses during their studies. The methodology involved administering pre-test and post-test questionnaires to students to assess their self-awareness of PS before and after the curriculum.

## 168 Study Setting and Sample

The participants who took part in the investigation were nursing students from a public university located in Jordan. Those contacted during the first semester of 2019-2020. A total of 373 students attended this PS program and their perception changes to the PS program were surveyed before and after the programme.

### **173** The Curriculum

The WHO developed training and we have chosen two topics from 11 topics because of limited funding and resources. We conducted training on topic 1: what is patient safety? and topic 9: Infection prevention and control (World Health Organization 20011)(16). The lectures were originally designed by the WHO team, but we adapted them to our Jordanian situation using contemporary health incidents as illustrative cases. These incidents were handled individually by our staff, who were able to provide detailed analysis and real management strategies.

### 180 The Study Tool:

181 The survey used in the research was adapted and modified with permission from the WHO and 182 the investigation was conducted adhering to their established protocols (4). It consisted of 23 items 183 that were organized into four groups or composites. The first composite consisted of seven items 184 and was focused on assessing the student's knowledge of PS. The second composite assessed the 185 Safety of the Healthcare System and was made up of five items. The third composite that 186 evaluated the student's ability to influence PS contained seven items. Finally, Composite 4 which was composed of four items, focused on the students' Personal Attitudes to PS. To rate the 187 students' responses, a 5-point ordinal scale was used, with "1" indicating "strongly disagree/very 188 189 poor" and "5" indicating "strongly agree/very good" (4).

190 Knowledge questions were used to determine what students have learnt about PS. Two to three 191 knowledge questions have been used for each of the topics of the curriculum guide. Students have

been asked the knowledge questions at the start of the courses on PS topics to establish baseline data on students' knowledge. Their responses to the knowledge questions at the end of the study were compared with baseline data, to quantify changes in student knowledge as a result of their learning certain PS topics.

## 196 Study Ethics and Actions

The study was approved by the university, the Scientific Research Ethics Committee, with reference number (SREC-7-2019). A voluntary self-guided survey was conducted among 373 nursing students. The survey was conducted during September 2019 and three months later during December 2019. Both were given during the entire class in the lecture room. Students have about 10-25 minutes to complete the questionnaire. The consent to take part in the study was voluntary and granted before the student responded to the questionnaire based on information given orally during the class.

204

## 205 Analysis

SPSS ver.25 was used to analyse the data. Descriptive statistics and paired t-tests were used in dataanalysis.

### 208 **Results**

The study invited 373 students to participate, and 253 (68%) questionnaires were returned for the pre-test. Two months after the last learning session or three months after the first session, 218 (58%) completed and returned the post-test questionnaire. Only forty students (16%) reported having prior exposure to PS training, while 172 (68%) acknowledged that their university had taught PS content.

### 214 Insert Figure 1 here

#### 215 Table 1: Demographic data

		Frequency	Percentage %	
Study level	2nd year	92	36.2	
	3rd year		25.6	
	4th year	93	36.6	
Has patient safety been taught in	Yes	172	67.7	
your university /school?	No	80	31.5	
Have you previously taken a patient	Yes	40	15.7	
safety course?	No	206	81.1	

216 Source: study data

#### 218 Students' attitudes before patient safety training

219 Before the training courses took place, Table 2 illustrates the knowledge, skills, and attitudes 220 (KSA) that the students had about PS. The feedback from the students indicated low level of 221 knowledge in the first composite regarding PS except for factors that contribute to human error or 222 influence PS, where they possessed moderate knowledge. The second composite is the Safety of 223 the Healthcare System evaluated by five items. Most of the students reported a neutral position 224 although some agreed and others disagree. In response to the third composite, the students agree 225 that they can influence PS. The items measure their influence, which may indicate attitude rather 226 than knowledge. This is confirmed by the student's responses to the last composite, as all students 227 indicated that they strongly agree on personal attitudes toward PS.

228

#### 229 Student attitudes after patient safety training

230 In Table 2 the results of the PS perception survey are presented. Students who had a moderate 231 understanding of the factors that affect PS achieved the highest scores (mean 3.45, SD 0.94). In 232 contrast, the lowest score was related to the role of health institutions in error reporting (mean 233 1.90, SD= 0.97). This indicates insufficient knowledge among students about the role of healthcare

<sup>217</sup> 

institutions in reporting errors. Additionally, the students received a low score in reporting error. Following the completion of the training, the students' knowledge about patient safety improved significantly, and most items receiving scores greater than three, indicating good knowledge. After the training, five out of seven items showed significant improvements, with the most improvement observed in the role of healthcare organizations in error reporting (the mean difference was 0.97, P=0.001).

Survey Item	Pre-Training		Post-Training		Correlation	<i>p</i> value
	Mean	SD	Mean	SD		
1. Different types of human error in health care	2.92	.917	3.27	.872	.034	.619
2. Factors contributing to human error	2.98	.925	3.43	.854	.175	.010
3. Factors influencing patient safety	3.45	.940	3.65	.801	.129	.057
4. Ways of speaking up about error	2.67	1.081	3.18	.877	.182	.008
5. What should happen if an error is made	2.78	1.012	3.26	1.05 3	.151	.027
6. How to report an error	2.34	1.144	2.92	1.10 5	.263	.000
7. The role of health-care organizations in error reporting	1.90	.974	2.87	1.12 9	.241	.000

# 240 Table 2: Composite 1- Knowledge of Patient Safety

241 Source: study data

247

248

Comparing the means for students' knowledge of the safety of the health care system before and after teaching the curriculum indicates an increase. For example, healthcare personnel who receive PS training scored 2.94 before and 3.13 after training with a standard deviation of (1.07 and 0.97) consequently. However, this increase is not statistically significant when we asked the students (Item 4) how often the patients received the wrong drug. (see Table 3).

Survey Item	Pre-T	raining	Post-Tra	aining	Correlation	p value
	Mean	SD	Mean	SD		
1. Most health-care workers make errors	3.33	.988	3.51	.947	.257	.000
2. In my country there is a safe system of healthcare for patients	2.94	.958	2.88	.926	.236	.000
3. Medical error is very common	3.77	.945	3.74	.906	.168	.013
4. It is very unusual for patients to be given the wrong drug	2.77	1.070	2.91	1.118	.053	.440
5. Health-care staff receive training in patient safety	2.94	1.071	3.13	.975	.211	.002

#### 249 Table 3: Composite 2- Safety of the Health care System

250 Source: study data

251 The results of students' influence on safety are not indicating a clear picture. There is a slight 252 increase in the student's ability to talk or tell others about an error they made. This result is 253 statistically significant. Students strongly believed that reporting the error will improve PS, scored 254 as highest (mean=3.93, SD1.009), students also always ensure that their PS is maintained. After 255 training, the only significant change was in the ability of the students to report their errors (items 1 256 & 7) (see Table 4).

#### 257 **Table 4: Composite 3-Personal Influence over Safety**

	Pre-Training		Post-Tr	aining	Correlatio	<i>p</i> value
Survey Item	Mea	SD	Mean	SD	n	
	n					
1. Telling others about an error I made would be easy.	3.35	1.206	3.51	1.030	.189	.005
2. It is easier to find someone to blame rather than focus on the causes of error.	2.07	1.104	2.24	1.074	.103	.131
3. I am confident about speaking to someone who is showing a lack of concern for a patient's safety.	3.48	1.153	3.42	1.005	.060	.377
4. I know how to talk to people who have made an error.	3.36	1.009	3.32	.951	.125	.067
5. I am always able to ensure that patient safety is not compromised.	3.89	1.040	3.85	.905	.037	.591
6. I believe that filling in reporting forms will help to improve patient safety.	3.93	1.009	3.90	.883	.080	.240
7. I am able to talk about my own errors.	3.60	1.026	3.76	.975	.252	.000

<sup>258</sup> Source: study data

- 259 Personal attitude towards PS did not show a significant change apart from the recognition by
- 260 students of the importance of learning how best to deal with their errors by the end of nursing
- 261 school (mean= 4.23 & SD=.8) (see Table 5). All other items were not significant, and they indicate
- 262 a decrease, too.

#### 263 Table 5: Composite 4- Personal Attitudes to Patient Safety

	Pre-Training Post-Training		Correlation	<i>p</i> value		
Survey Item	Mean	SD	Mean	SD		
1. By concentrating on the causes of incidents I can contribute to patient safety.	4.13	.775	4.04	.727	.130	.056
2. If I keep learning from my mistakes, I can prevent incidents.	4.45	.665	4.24	.836	.063	.353
<b>3.</b> Acknowledging and dealing with my errors will be an important part of my job.	4.21	.775	4.15	.826	.065	.339
4. It is important for me to learn how best to acknowledge and deal with my errors by the end of nursing school.	4.49	.687	4.23	.805	.206	.002

264 Source: study data

265

#### Discussion 266

267 The study findings demonstrated that implementing the WHO PSC had a beneficial effect on the 268 attitudes and knowledge of undergraduate nursing students regarding PS. The pre-test scores 269 revealed that the participants had a positive attitudes and moderate level of knowledge towards PS. 270 However, after the intervention, the post-test scores showed an increase in their knowledge and positive attitudes towards PS. These results are consistent with previous studies that have reported 271 272 the effectiveness of PS education in enhancing healthcare providers' knowledge and attitudes of 273 toward PS (12, 19, 26). The study results underscore the need for formal PS education and 274 highlight the importance of improving overall safety competence and health outcomes (11).

275 According to this study, students' ability to identify and talk about their mistakes improved 276 significantly after training. However, their perception towards PS showed a significant change as 277 well, as they recognized the importance of learning how to deal with their errors by the end of their 278 nursing education. This finding is consistent with Suliman (25) research which found that PS is not 279 given enough importance in Jordanian nursing education and curricula. The students had more 280 confidence in their knowledge and skills acquired in the classroom than their practical training in 281 clinical settings, particularly in writing medical error reports (24, 25). The study concludes that 282 contextual and sociocultural issues related to PS should be emphasized in nursing curriculum to 283 enable students to speak up when they observe healthcare errors. To achieve this, it is 284 recommended that PS elements be added to bridge the gap between theory and practice in the 285 nursing curriculum (25).

286

287 Study limitations:

288 Although we obtained a significant difference in the measures before/after teaching the PSC, This 289 is probably not ultimate, the curriculum probably caused the increase in KSA of PS. Other 290 variables may have also influenced this increase (16, 19). The passage of time between the pre-test and post-test without curriculum teaching could have contributed to this change. Other events may 291 292 also occur during this period that influenced the students' KSA about PS. Perhaps, the students were exposed to PS training in clinical settings or informally by the instructors. This means that 293 there is a possibility of contamination from any source that was not under the control of this study 294 (7, 27). 295

In this study, we used measures to anticipate any confounding factors during the research design. For example, we have asked whether the student has previously been exposed to PS training. However, the results of this study can be improved in future; if researchers use a control group that is not exposed to the intervention and differentiate between the intended intervention and other domains included in the curriculum. In addition, the intervention should be conducted by teachers

other than their regular course instructors to avoid using any of the materials during their normal
course. This would assist in excluding the effects of time and other factors that could affect the
study's results. This study only investigated two domains out of the 11 courses suggested by WHO
(4). Therefore, the results may not be generalised to all other topics of the curriculum suggested by
the WHO.

306

307 Implications:

308 A formal PSC adoption and its early integration into the undergraduate nursing curriculum were 309 determined to be necessary according to this study. During the study, we also anticipated that 310 teacher would need to become more involved in the creation and integration of PS in the 311 curriculum. There is a good opportunity to use the "WHO Curriculum Resource Guide" in any 312 attempt to develop an appropriate PSC. The attitude towards PS in academic culture must not only 313 include students, but there has been resistance to implementation at different levels, even from the 314 curriculum regulatory bodies. The WHO curriculum can be amended according to the culture of 315 local institutions and can be properly implemented with the support of the relevant government 316 authorities.

Future research should evaluate the usefulness of WHO's PSC in a wider and more diverse sample group to improve the universality of the results. Finally, it is recommended that more research to be done to determine how the WHO PSC has affected the patient outcomes and how it may reduce adverse reactions in clinical practice.

## 321 Conclusion

In conclusion, this study has shed important light on undergraduate nursing students' attitudes, knowledge and abilities concerning PS, both before and after training. The results indicate that the students had a low level of knowledge in some areas of PS before training, especially with regard to the role of healthcare institutions in reporting errors. However, the training using the WHO PSC

326 significantly improved the students' knowledge and attitudes in most PS areas, with the most 327 significant improvement observed in the role of healthcare organization in error reporting. The 328 curriculum was well received by the participants and they were confident in their ability to apply 329 the knowledge acquired in their future clinical practice. This indicates the need to train nursing 330 students on how to respond to healthcare errors in general, not only to provide safe bedside care to 331 the patients. Overall, the study suggests that training programs can effectively improve students' 332 KSA when included in the curriculum of nursing schools. The WHO PSC guide can be a great 333 start in this domain, which is essential to improve the quality of healthcare services.

334

## 335 Acknowledgement

336 The authors thank the University of Mutah, special thanks go to the nursing academics

337 who conducted the teaching and supported the study, and load of thanks to the study

participants who completed the questionnaires and who took the time to participate in this

339 study.

## 340 **References:**

Pierre MS, Grawe P, Bergstrom J, Neuhaus C. 20 years after To Err Is Human: A
 bibliometric analysis of 'the IOM report's' impact on research on patient safety. Safety Science.
 2022;147(20):105593.

Kohn LT, Corrigan JM, Donaldson MS. To Err Is Human : Building a Safer Health System.
 Washington, DC, USA: National Academies Press; 2000.

Sheehan P, Joy A, Fleming A, Vosper H, McCarthy S. Human factors and patient safety in
 undergraduate healthcare education: a systematic review. Human Factors in Healthcare.
 2022;2(1):100019.

World Health Organization. Patient safety curriculum guide: multi-professional edition.
 Geneva: World Health Organization; 2011. Contract No.: CC BY-NC-SA 3.0 IGO.

5. Kirwan M, Riklikiene O, Gotlib J, Fuster P, Borta M. Regulation and current status of patient safety content in pre-registration nurse education in 27 countries: Findings from the Rationing-Missed nursing care (RANCARE) COST Action project. Nurse Education in Practice. 2019;37(1):132-40.

355 6. Wu AW, Busch IM. Patient safety: a new basic science for professional education. GMS
356 journal for medical education. 2019;36(2):Doc21.

3577.Cantero-López N, González-Chordá VM, Valero-Chillerón MJ, Mena-Tudela D, Andreu-358Pejó L, Vila-Candel R, et al. Attitudes of undergraduate nursing students towards patient safety: a

perpetuity. It is made available under a CC-BY 4.0 International license .

quasi-experimental study. International journal of environmental research and public health.2021;18(4):1429.

Al-Nawafleh A, Abu-Helalah MA, Hill V, Masoud MI, Al-Mahasneh HA, Al Salti ET. Patient
 Safety Culture in Jordanian Hospitals. Health Science Journal. 2016;10(5):5.

363 9. Singh H, Mushtaq U, Marinez A, Shahid U, Huebner J, McGaffigan P, et al. Developing the
364 Safer Dx Checklist of ten safety recommendations for health care organizations to address
365 diagnostic errors. The Joint Commission Journal on Quality and Patient Safety. 2022;48(11):581366 90.

10. Evans K, Athavale N, Long S, Vincent C. Patient safety. In: Morley JE, Vellas B, Sinclair AJ,
Cesari M, Munshi M, editors. Pathy's Principles and Practice of Geriatric Medicine. 6th ed: WileyBlackwell; 2022. p. 146-64.

11. Lee SE, Morse BL, Kim NW. Patient safety educational interventions: a systematic review
with recommendations for nurse educators. Nursing open. 2022;9(4):1967-79.

Torkaman M, Sabzi A, Farokhzadian J. The effect of patient safety education on
undergraduate nursing students' patient safety competencies. Community Health Equity
Research & Policy. 2022;42(2):219-24.

37513.Ji Y, Lee H, Lee T, Choi M, Lee H, Kim S, et al. Developing an integrated curriculum for376patient safety in an undergraduate nursing program: a case study. BMC nursing. 2021;20(1):1-11.

Myung SJ, Shin J-S, Kim JH, Roh H, Kim Y, Kim J, et al. The Patient Safety Curriculum for
Undergraduate Medical Students as a First Step Toward Improving Patient Safety. Journal of
Surgical Education. 2012;69(5):659-64.

380 15. Wahlberg KJ, Pay T, Repp AB, Wahlberg EA, Kennedy AG. Effect of Patient Safety
381 Curriculum for Internal Medicine Residents on a Health Care System. ATS scholar. 2022;3(1):156382 66.

16. Mansour M, Skull A, Parker M. Evaluation of World Health Organization multiprofessional patient safety curriculum topics in nursing education: Pre-test, post-test, noneexperimental study. Journal of Professional Nursing. 2015;31(5):432-9.

17. Lawal BK, Mohammed S. Integrating patient safety education into pharmacy curriculum
 in Nigeria. Int J Pharm Pract. 2022;30(6):583-5.

Sujan MA, Furniss D, Anderson J, Braithwaite J, Hollnagel E. Resilient Health Care as the
basis for teaching patient safety – A Safety-II critique of the World Health Organisation patient
safety curriculum. Safety Science. 2019;118:15-21.

391 19. AlNawafleh AH, Al-Momani M, Alhalaiqa F. Patient Safety Education: A Perspective of
 392 Southern Jordan Nursing Students'. Mu'tah Lil-Buhuth wad-Dirasat. 2022;37(2):49-67.

393 20. Salami I, Subih M, Darwish R, Al-Jbarat M, Saleh Z, Maharmeh M, et al. Medication
394 Administration Errors: Perceptions of Jordanian Nurses. Journal of Nursing Care Quality.
395 2019;34(2):E7-E12.

Albsoul RA, Safadi RR, Alshyyab MA, FitzGerald G, Hughes JA, Ahmad M. Missed Nursing
Care in Medical and Surgical Wards in Jordan: A Cross-Sectional Study. Policy, Politics, & Nursing
Practice.0(0):15271544231155845.

Alquwez N, Cruz JP, Alshammari F, Felemban EM, Almazan JU, Tumala RB, et al. A
multi-university assessment of patient safety competence during clinical training among
baccalaureate nursing students: A cross-sectional study. Journal of clinical nursing. 2019;28(910):1771-81.

403 23. Mansour MJ, Al Shadafan SF, Abu-Sneineh FT, AlAmer MM. Integrating patient safety
404 education in the undergraduate nursing curriculum: a discussion paper. The open nursing journal.
405 2018;12:125.

406 24. Huang FF, Shen XY, Chen XL, He LP, Huang SF, Li JX. Self-reported confidence in patient
407 safety competencies among Chinese nursing students: a multi-site cross-sectional survey. BMC
408 medical education. 2020;20(1):1-10.

perpetuity. It is made available under a CC-BY 4.0 International license .

Suliman M. Measuring Patient Safety Competence Among Nursing Students in theClassroom and Clinical Settings. Nursing education perspectives. 2019;40(3):E3-E7.

411 26. Lee SE, Dahinten VS. Evaluating a patient safety course for undergraduate nursing 412 students: A quasi-experimental study. Collegian. 2023;30(1):75-83.

413 27. Kim YM, Yoon YS, Hong HC, Min A. Effects of a patient safety course using a flipped

classroom approach among undergraduate nursing students: A quasi-experimental study. Nurse

415 Education Today. 2019;79(1):180-7.

416




Figure 1: study process flow chart

