Title: EEG Based Oculographic Analysis of Epileptic Nystagmus

Aybuke Acar, MD,¹ Anthony Zampino, MD,^{1,2} Neel Fotedar, MD^{1,2}
1Epilepsy Center, Neurological Institute, University Hospitals Cle
0H, USA 44106
2Department of Neurology, Case Western Reserve University Sch
USA 44106
C

²Epilepsy Center, Neurological Institute, Offiversity Hospitals Cleveland Medical Center, Cleveland, OH, USA 44106
²Department of Neurology, Case Western Reserve University School of Medicine, Cleveland, OH,
USA 44106
 -Department of Neurology, Case Western Reserve University School of Medicine, Cleveland, OH,
USA 44106
Correspondence to: Neel Fotedar, MD
Address: 11100 Euclid Ave, Lakeside #3200, Cleveland, OH, USA 44106
Email address:

Pepartment of
USA 44106
Correspondence
Address: 11100
Email address: 1
No. of words in Correspond
Address: 11
Email addre
No. of word
No. of word
No. of figure $\begin{array}{c} 1 \\ 1 \\ 1 \\ 1 \end{array}$ Address: 11100 Euclid Ave, Lakeside #
Email address: <u>neel fotedar@uhhospit</u>
No. of words in abstract: 218
No. of words in manuscript: 1488
No. of figures: 1
No. of figures: 1
No. of tables: 2

Email address: <u>neel.fotedar@uhhospitals.org</u>
No. of words in abstract: 218
No. of words in manuscript: 1488
No. of figures: 1
No. of tables: 2
No. of videos: 2
No. of references: 11

Email and the <u>manual addression of</u> No. of words in manuscript: 1488
No. of words in manuscript: 1488
No. of figures: 1
No. of tables: 2
No. of videos: 2
No. of references: 11 No. of words in manuscript: 1
No. of figures: 1
No. of tables: 2
No. of videos: 2
No. of references: 11

No. of videos: 2
No. of reference
.

No. of figures: 1
No. of tables: 2
No. of videos: 2
No. of references: 11 No. of tables: 2
No. of videos: 2
No. of references No. of reference
:

Abstract:

|
| i c c v c
| i c in the posterior cortical regions. EN is broadly categorized into type I and type II. Type I EN consists
of contralateral repetitive saccadic eye movements alternating with post-saccadic slow drifts with
an overall contral of contralateral repetitive saccadic eye movements alternating with post-saccadic slow drifts with
an overall contralateral deviation. Type II EN is characterized by ipsilateral slow drift alternating
with contralateral co

an overall contralateral deviation. Type II EN is characterized by ipsilateral slow drift alternating
with contralateral corrective saccades. In this article, we report a method to perform oculographic
analysis of eye move with contralateral corrective saccades. In this article, we report a method to perform oculographi
analysis of eye movements using EEG only. We used this method to classify the type of EN in three
patients with parieto-occ analysis of eye movements using EEG only. We used this method to classify the type of EN in three patients with parieto-occipital seizures.
In all three patients, the ictal EEG demonstrated repetitive saccadic eye movement mationts with parieto-occipital seizures.

In all three patients, the ictal EEG demonstrated repetitive saccadic eye movements, directed

contralateral to the seizure onset zone. With prolonged time constant, we were able Facture with particle everprice examples
In all three patients, the ictal EEG demor
contralateral to the seizure onset zone. V
eye movement pattern as EN with distin
characterize the type of EN as type I and
of fast phase contralateral to the seizure onset zone. With prolonged time constant, we were able to identiffeye movement pattern as EN with distinct slow and fast phases. We were able to further characterize the type of EN as type I an

eye movement pattern as EN with distinct slow and fast phases. We were able to further
characterize the type of EN as type I and type II. In all three patients, the direction of EN (direction
of fast phase or saccades) was eharacterize the type of EN as type I and type II. In all three patients, the direction of EN (of fast phase or saccades) was contralateral to the seizure onset zone.
EN can be easily missed on video-electroencephalography of fast phase or saccades) was contralateral to the seizure onset zone.
EN can be easily missed on video-electroencephalography (vEEG) recordings because of various
reasons. Our study demonstrates a systematic method of ey Fast phase or sacritions, the contrainment is increased.

EN can be easily missed on video-electroencephalography (vEEG) recorders

reasons. Our study demonstrates a systematic method of eye moveme

be used to not only ide reasons. Our study demonstrates a systematic method of eye movement analysis on EEG, which the used to not only identify EN as seizure semiology but also classify it, without requiring additional electrodes.
Electrodes.
K reasons to not only identify EN as seizure semiology but also classify it, without requiring additional
be used to not only identify EN as seizure semiology but also classify it, without requiring additional
electrodes.
Ke be used to not only identify the semiology but also constant that η is η is an also classified electrodes.
Keywords: epileptic nystagmus; EEG; time constant; parieto-occipital

electrones.
Keywords: Keywords: epileptic nystagmus; EEG; time constant; parieto-occipital

1. Introduction:

Fractive originating in the posterior cortical regions (Kaplan and Tusa, 1993). It is characterized be
fast (saccadic) repetitive eye movements, alternating with slow movements, with the fast phase
being directed away from fast (saccadic) repetitive eye movements, alternating with slow movements, with the fast phase
being directed away from the seizure onset zone (Tusa et al., 1990).
EN is mainly classified into two types depending upon the being directed away from the seizure onset zone (Tusa et al., 1990).
EN is mainly classified into two types depending upon the underlying mechanism. Type I EN is
postulated to originate from the activation of the cortical BN is mainly classified into two types depending upon the underlyin
postulated to originate from the activation of the cortical areas respo
movements producing contraversive saccades. Type II EN is generat
areas responsibl

postulated to originate from the activation of the cortical areas responsible for saccadic eye
movements producing contraversive saccades. Type II EN is generated by the activation of cort
areas responsible for smooth purs movements producing contraversive saccades. Type II EN is generated by the activation of c
areas responsible for smooth pursuit (temporo-parieto-occipital junction), resulting in an
ipsiversive slow phase alternating with meas responsible for smooth pursuit (temporo-parieto-occipital junction), resulting in an
ipsiversive slow phase alternating with a resetting contraversive saccade (Kaplan and Tusa, 1993;
Tusa et al., 1990).
The eye acts l in since since the area are alternating with a resetting contraversive saccade (Kaplan and Tus Tusa et al., 1990).
The eye acts like an electrical dipole with the anterior being more positive and the posterior more negativ The eye acts like an electrical dipole with the anterior being more positive and the posterior being
more negative. This creates a characteristic artifact on EEG depending on the type and direction of
the eye movement (Lüd The eye acts like a
more negative. Th
the eye movemen
a large positivity a
have a characteris
baseline, which is
In this article, we
which could help
additional electro more negative. This creates a characteristic artifact on EEG depending on the type and direction of
the eye movement (Lüders and Noachtar, 2000). For example, a leftward eye movement will create
a large positivity at F7 an more the eye movement (Lüders and Noachtar, 2000). For example, a leftward eye movement will create
a large positivity at F7 and a simultaneous negativity at F8. In addition, the saccadic eye movement
have a characteristic the alarge positivity at F7 and a simultaneous negativity at F8. In addition, the saccadic eye movements
have a characteristic morphology with a rapid rise of potential, followed by a slow return to
baseline, which is dete

have a characteristic morphology with a rapid rise of potential, followed by a slow return to
baseline, which is determined by the time constant (TC) (Lüders and Noachtar, 2000).
In this article, we aim to report a systema

2. Methods:

have baseline, which is determined by the time constant (TC) (Lüders and Noachtar, 2000).

In this article, we aim to report a systematic method to analyze eye movement artifact on EE

which could help identify EN and dist baseline, which is article, we aim to report a systematic method to analyze eye movement artifact which could help identify EN and distinguish between its different types, without requided and electro-oculogram (EOG) elect In this could help identify EN and distinguish between its different types, without requiring additional electro-oculogram (EOG) electrodes.

2. Methods:

We identified three patients from our database with parietal, occip additional electro-oculogram (EOG) electrodes.

2. Methods:

We identified three patients from our database with parietal, occipital or parieto-occipital s

and reported semiology of epileptic nystagmus, who had at least 2 **2. Methods:**
2. Methods:
We identified three patients from our database vand reported semiology of epileptic nystagmus,
review, using the Neurofax EEG-1200 system by
visible in the video in any of these patients. The
ex |
| a l t
| a l

and reported semiology of epileptic nystagmus, who had at least 24 hours of vEEG data available fo
review, using the Neurofax EEG-1200 system by Nihon Kohden (Tokyo, Japan). Eyes were not
visible in the video in any of th review, using the Neurofax EEG-1200 system by Nihon Kohden (Tokyo, Japan). Eyes were not visible in the video in any of these patients. The report of EN was based on either bedside examination or from history.
First, we a visible in the video in any of these patients. The report of EN was based on either bedside
examination or from history.
First, we analyzed the EEG epoch of interest with the conventional settings of a TC of 0.1 secon
and examination or from history.

First, we analyzed the EEG epoch of interest with the conventional settings of a TC of 0.1 s

and a high-frequency filter of 70Hz in a longitudinal bipolar montage. We then analyzed 3

of pre First, we analyzed the EEG ep
and a high-frequency filter of
of pre-ictal EEG for any unust
identified epochs of unilatera
analyzed the same epochs wir
eye movements, alternating v
In addition, we also character
of prolong and a high-frequency filter of 70Hz in a longitudinal bipolar montage. We then analyzed 30 second pre-ictal EEG for any unusual eye movements, followed by the ictal EEG. In the ictal EEG, we identified epochs of unilatera of pre-ictal EEG for any unusual eye movements, followed by the ictal EEG. In the ictal EEG, we
identified epochs of unilaterally directed repetitive saccadic eye movements. Following this, we
analyzed the same epochs wit

identified epochs of unilaterally directed repetitive saccadic eye movements. Following this, we
analyzed the same epochs with a prolonged TC of ≥2 seconds to identify and characterize any s
eye movements, alternating with analyzed the same epochs with a prolonged TC of \geq 2 seconds to identify and characterize any sleve movements, alternating with the saccadic eye movements.
In addition, we also characterized a single-channel oculographi eye movements, alternating with the saccadic eye movements.
In addition, we also characterized a single-channel oculographic waveform using the same settings
of prolonged TC. By connecting F7 to an electrode relatively ind For an addition, we also characterized a single-channel oculographiof prolonged TC. By connecting F7 to an electrode relatively in
movements (e.g. Cz or Fp1), we were able to study the characte
phase.
Slow phase waveform w of prolonged TC. By connecting F7 to an electrode relatively indifferent to horizontal eye
movements (e.g. Cz or Fp1), we were able to study the characteristics of the slow phase and the fast
phase.
Slow phase waveform was

movements (e.g. Cz or Fp1), we were able to study the characteristics of the slow phase a
phase.
Slow phase waveform was characterized as linear or non-linear, depending upon the slop
waveform as it would approach the EEG phase.
Slow phase waveform was characterized as linear or non-linear, depending upon the slope of the
waveform as it would approach the EEG baseline. A negative exponential decay of the slope of the
waveform as it approach .
Slow p
wavefc
wavefc Slow phase waveform as it would approach the EEG baseline. A negative exponential decay of the slope of the waveform as it approaches the baseline is consistent with decreasing velocity as the eyes approach waveform as it waveform as it approaches the baseline is consistent with decreasing velocity as the eyes approached waveform as it approaches the baseline is consistent with decreasing velocity as the eyes approached α waveform as it approaches the baseline is consistent with decreasing velocity as the eyes approaches approaches approaches approaches approaches as the eyes approaches approaches approaches approaches approaches approache

midline or not by analyzing whether the respective waveforms would cross the EEG baseline or n
For example, if the eyes start at midline and then drift toward the left by 20 degrees, the EEG wou
show a certain positive vol For example, if the eyes start at midline and then drift toward the left by 20 degrees, the EEG would
show a certain positive voltage at F7 away from the baseline. If the eyes then come back to the right
by the same amount For example, the eyes then come back to the right show a certain positive voltage at F7 away from the baseline. If the eyes then come back to the right by the same amount, the waveform would simply return to the baseline b show the same amount, the waveform would simply return to the baseline but it would not cross it. If instead the eyes would cross the midline and actually turn to the baseline but it would not cross it. If instead the eyes by the same amound of the waveform would be twice the right side by another 20 degrees, the corresponding EEG waveform would be twice the amplitude of the initial waveform, hence crossing the EEG baseline. This analysis is

3. Results:

the corresponding EEG waveform would be twice the amplitude of the initial waveform, hence
crossing the EEG baseline. This analysis is only possible with the settings of prolonged TC.
This study was approved by the Instit crossing the EEG baseline. This analysis is only possible with the settings of prolonged TC.

This study was approved by the Institutional Review Board of University Hospitals Cleveland

Medical Center.

3. Results:

Pati This study was approved by the Institutional Review Board of University Hospitals Clevela
Medical Center.
3. **Results:**
Patient #1 and #3 were monitored for five and three days, respectively, and only one seizu
captured f The Institutional Center.

3. Results:

Patient #1 and #3 were monitored for five and three days, respectively, and only one seizure ve

captured for each of them during the recording. Patient #2 was monitored for one week 3. Results:
3. Results:
Patient #1 and #
captured for eac
initial five days,
characteristics c
Patient demogra
three patients, t
pre-ictal period.] (i ()
] t]

captured for each of them during the recording. Patient #2 was monitored for one week. During the initial five days, he had very frequent seizures at ~9-13/hour. All of these seizures had the EEG characteristics consistent initial five days, he had very frequent seizures at ~9-13/hour. All of these seizures had the EEG
characteristics consistent with nystagmus, as described below.
Patient demographics, nystagmus characteristics and EEG findi characteristics consistent with nystagmus, as described below.

Patient demographics, nystagmus characteristics and EEG findings are summarized in table 1. In

three patients, there was no specific pattern of repetitive ey Patient demographics, nystagmus characteristics and EEG findi
three patients, there was no specific pattern of repetitive eye m
pre-ictal period.
Patient #1's EEG was consistent with a right beating nystagmus
with left occ three patients, there was no specific pattern of repetitive eye movement artifact in the immediate
pre-ictal period.
Patient #1's EEG was consistent with a right beating nystagmus with leftward linear slow phases
with left pre-ictal period.

Patient #1's EEG was consistent with a right beating nystagmus with leftward linear slow phases

with left occipital seizure. With the prolonged TC, an overall initial deviation to the left was noted

wi

Patient #1's EEG
with left occipita
with rightward s
crossing the mid
with type II EN.
In patients #2 an
to the seizure on
Oculogram using
midline (Fig. 1d)
phases did show with left occipital seizure. With the prolonged TC, an overall initial deviation to the left was noted
with rightward saccadic eye movements (Fig. 1a). Oculogram using F7-Cz demonstrated the eyes
crossing the midline with with rightward saccadic eye movements (Fig. 1a). Oculogram using F7-Cz demonstrated the eyes
crossing the midline with both slow and fast phases (Fig. 1b). These characteristics are consistent
with type II EN.
In patients crossing the midline with both slow and fast phases (Fig. 1b). These characteristics are consistent with type II EN.
In patients #2 and #3, the ictal EEG demonstrated repetitive saccadic eye movements contralater.
In pati with type II EN.
In patients #2 and #3, the ictal EEG demonstrated repetitive saccadic eye movements contralaterate to the seizure onset zone with overall eye deviation in the same direction as the saccades (Fig. 1c).
Ocu In patients #2 and
to the seizure or
Oculogram using
midline (Fig. 1d)
phases did show
phase velocity (I
The frequency of EN was 2-3Hz In the seizure onset zone with overall eye deviation in the same direction as the saccades (Fig. 1c).

Oculogram using F7-Fp1 demonstrated an overall contraversive deviation without crossing the

midline (Fig. 1d). The sl Oculogram using F7-Fp1 demonstrated an overall contraversive deviation without crossing the
midline (Fig. 1d). The slow phases were very brief, becoming progressively longer. The longer slow
phases did show a waveform wit midline (Fig. 1d). The slow phases were very brief, becoming progressively longer. The longer sly
abases did show a waveform with a negative exponential slope consistent with decreasing slow
phase velocity (Fig. 1d). Thes

4. Discussion:

phases did show a waveform with a negative exponential slope consistent with decreasing slow
phase velocity (Fig. 1d). These characteristics are consistent with type I EN.
The frequency of type II EN was 0.5-2Hz with a to phase velocity (Fig. 1d). These characteristics are consistent with type I EN.
The frequency of type II EN was 0.5-2Hz with a total duration of \sim 2 minutes. The frequency of ty
I EN was 2-3Hz with a total duration of $\$ The frequency of type II EN was 0.5-2Hz with a total duration of \sim 2 minutes
I EN was 2-3Hz with a total duration of \sim 15 seconds.
4. Discussion:
Nystagmus is defined as abnormal repetitive oscillatory eye movements t The frequency of 4. Discussion:
Nystagmus is defined as abnormal repetitive oscillate
(Leigh and Zee, 2015). Epilepsy is an uncommon caus
commonly seen with seizures involving the posterior
Leigh and Zee, 2015). It is rarely associated wit |
| (c
| | (Leigh and Zee, 2015). Epilepsy is an uncommon cause of acquired nystagmus and EN is most
commonly seen with seizures involving the posterior cortical regions (Kaplan and Tusa, 1993;
Leigh and Zee, 2015). It is rarely asso (Leigh and Zee, 2015). It is rarely associated with frontal regions (Kaplan and Tusa, 1993; Leigh and Zee, 2015). It is rarely associated with frontal lobe epilepsy (Lee et al., 2014). EN car
present in isolation as seizu commonly seen with set with server with server with selection of the posterior correlation as seizure semiology and has been reported at all ages (Nicita et al., 2010; present in isolation as seizure semiology and has been present in isolation as seizure semiology and has been reported at all ages (Nicita et al.,2010;
present in isolation as seizure semiology and has been reported at all ages (Nicita et al.,2010; present in isolation as seizure semiology and has been reported at all ages (Nicita et al.,2010;

(Kaplan and Tusa, 1993; Leigh and Zee, 2015)
As demonstrated by our results, type I EN is characterized by an overall deviation of the eye:
contralateral to the seizure onset zone. A 'staircase' of contraversive saccades p As demonstrated by our results, type I EN is cha
contralateral to the seizure onset zone. A 'stairc
deviation. As mentioned above, type I EN is pos
areas responsible for saccadic eye movements s
middle frontal gyrus or the contralateral to the seizure onset zone. A 'staircase' of contraversive saccades produces this deviation. As mentioned above, type I EN is postulated to be generated by activation of the co
areas responsible for saccadic e deviation. As mentioned above, type I EN is postulated to be generated by activation of the careas responsible for saccadic eye movements such as the frontal eye field (FEF) in posterion middle frontal gyrus or the parieta areas responsible for saccadic eye movements such as the frontal eye field (FEF) in posterior middle frontal gyrus or the parietal eye field (PEF) within the banks of the intraparietal sulcus (Kaplan and Tusa, 1993; Leigh middle frontal gyrus or the parietal eye field (PEF) within the banks of the intraparietal sulcu
(Kaplan and Tusa, 1993; Leigh and Zee, 2015). The activation of these areas is known to gene
contraversive saccades with cont (Kaplan and Tusa, 1993; Leigh and Zee, 2015). The activation of these areas is known to generation traversive saccades with contraversive gaze deviation. A recent study has shown that in majority of versive seizures, the g (Contraversive saccades with contraversive gaze deviation. A recent study has shown that in
majority of versive seizures, the gaze deviation is saccadic rather than one smooth movement
(Fotedar et al., 2022) . The nystagmo majority of versive seizures, the gaze deviation is saccadic rather than one smooth movemet (Fotedar et al., 2022) . The nystagmoid appearance is due to ipsiversive slow drifts alternation the saccades, related to an addit

(Fotedar et al., 2022) . The nystagmoid appearance is due to ipsiversive slow drifts alternating
the saccades, related to an additional gaze-holding defect, which accounts for the post-saccadi
slow drift (Video 1). The vel (For exact also medial and and interpret all (MST) areas results in an injuster alternation of the post-saccedic
slow drift (Video 1). The velocity of the slow phase characteristically decreases as the eyes
approach the mi slow drift (Video 1). The velocity of the slow phase characteristically decreases as the eyes
approach the midline and the eyes never cross the midline (Kaplan and Tusa, 1993; Tusa et
al.,1990; Leigh and Zee, 2015).
Type I approach the midline and the eyes never cross the midline (Kaplan and Tusa, 1993; Tusa et
al., 1990; Leigh and Zee, 2015).
Type II EN, on the other hand, consists of an initial ipsiversive eye deviation due to a slow
movem ¹
al., 1990; Leigh and Zee, 2015).
Type II EN, on the other hand, consists of an initial ipsiversive eye deviation due to a slow
movement, which is interrupted by corrective contraversive saccades (Video 2). The initial
 Type II EN, on the other hand, c
movement, which is interrupted
ipsiversive slow movement is p
the posterior superior temporal
middle temporal (MT) and med
Experimental studies in monkey
slow eye movement (Lüders and
are m movement, which is interrupted by corrective contraversive saccades (Video 2). The initial
ipsiversive slow movement is produced by the activation of the cortical smooth pursuit re
the posterior superior temporal sulcus at ipsiversive slow movement is produced by the activation of the cortical smooth pursuit regivent per posterior superior temporal sulcus at the temporo-parieto-occipital (TPO) junction, knowledle temporal (MT) and medial sup the posterior superior temporal sulcus at the temporo-parieto-occipital (TP0) junction, known as
middle temporal (MT) and medial superior temporal (MST) areas (Leigh and Zee, 2015).
Experimental studies in monkeys have sho middle temporal (MT) and medial superior temporal (MST) areas (Leigh and Zee, 2015).
Experimental studies in monkeys have shown that activation of these areas results in an ipsiversis
slow eye movement (Lüders and Noachtar Experimental studies in monkeys have shown that activation of these areas results in an islow eye movement (Lüders and Noachtar, 2000). As demonstrated in the results, the slo
are mostly linear and both fast and slow eye m

slow eye movement (Lüders and Noachtar, 2000). As demonstrated in the results, the slow phases
are mostly linear and both fast and slow eye movements cross the midline (Kaplan and Tusa, 1993;
Tusa et al., 1990; Leigh and are mostly linear and both fast and slow eye movements cross the midline (Kaplan and Tusa, 1993).
Tusa et al., 1990; Leigh and Zee, 2015). We have summarized the characteristics of type I and II EN
in Table 2.
A third mech Tusa et al., 1990; Leigh and Zee, 2015). We have summarized the characteristics of type I and II EN in Table 2.
A third mechanism, similar to type II EN, is postulated to be related to the activation of the optokinetic re in Table 2.

A third mechanism, similar to type II EN, is postulated to be related to the activation of the

optokinetic region and can also produce a sensation of vertigo (Kaplan and Tusa, 1993; Tusa et al.,

1990; Garcí A third mee
optokinetic
1990; Garc
nystagmus
al., 2006; H
Previous st
with EN (K
5. Col optokinetic region and can also produce a sensation of vertigo (Kaplan and Tusa, 1993; Tust
1990; García-Pastor et al., 2002; Ma et al., 2015). However, in all cases, the fast phase of the
nystagmus is contralateral to th 1990; García-Pastor et al., 2002; Ma et al., 2015). However, in all cases, the fast phase of the
nystagmus is contralateral to the seizure onset zone (Lee et al., 2014; Nicita et al., 2010; Bekdik et
al., 2006; Harris et

5. Conclusions:

nystagmus is contralateral to the seizure onset zone (Lee et al., 2014; Nicita et al., 2010; Bek
al., 2006; Harris et al., 1997).
Previous studies suggest that, in awake patients, an ictal discharge frequency of ≥10Hz is nal, 2006; Harris et al., 1997).

Previous studies suggest that, in awake patients, an ictal discharge frequency of ≥ 10 Hz is associate

with EN (Kaplan and Tusa, 1993). Our study is mostly in agreement with this obse an, 2000; Harris et al., 2000;
Previous studies suggest that
with EN (Kaplan and Tusa, 19
5. **Conclusions:**
In this study, we demonstrate
could help identify and classi!
of additional EOG electrodes. Previous studies suggest that, in a matter of a matter of patients, and the BN (Kaplan and Tusa, 1993). Our study is mostly in agreement with this observation.

5. Conclusions:

In this study, we demonstrate a systematic m 5. **Conclusions:**
In this study, we demonstrate a systematic method of eye movement analysis using EEG or
could help identify and classify EN in patients with occipital and parietal seizures without
of additional EOG elect $\frac{1}{2}$ In this study, we demonstrate a systematic method of experiments with occipital and parietal seizures without the help of additional EOG electrodes.
In patients with occipital and parietal seizures without the help of addi could help in an approximation of additional EOG electrodes.
The fadditional EOG electrodes.

Tables:

Table 1: Fatient demographics, characteristics of interictal & ictal EEG, seizure semiology,
characteristics of nystagmus and relationship to seizure onset zone and neuroimaging
F: female, M: male, L: left, LPD: lateralize

F: female, M: male, L: left, LPD: lateralized periodic discharges, R: right
 $\frac{1}{2}$ F: female, M: male, L: left, LPD: lateralized periodic discharges, R: right

 Table 2: Comparison of major characteristics of Type I and Type II EN

Figures:

- Figure 1: EEG analysis demonstrating epileptic nystagmus (EN).

(a) EN in patient #1. 60s EEG page with left and right tempore mystagmus. Channel P7-01 indicates the seizure onset wit

The large positivity at F7 at points mystagmus. Channel P7-01 indicates the seizure onset with paroxysmal fast activity at A.
The large positivity at F7 at points B & C indicates the initial leftward eye deviation. D
indicates corrective rightward saccades. T
	-
	- The large positivity at F7 at points B & C indicates the initial leftward eye deviation. D
indicates corrective rightward saccades. TC for the first two and last two channels is at 1
(whereas that for P7-O1 is 0.1s.
Channe indicates corrective rightward saccades. TC for the first two and last two channels is at whereas that for P7-01 is 0.1s.
Channel F7-Cz with TC of 10s demonstrating the oculogram for patient #1. Horizontal
dotted green lin whereas that for P7-01 is 0.1s.
Channel F7-Cz with TC of 10s demonstrating the oculogram for patient #1. Horizontal
dotted green line represents the EEG baseline. Slow phases are mostly linear. Both slow and
fast phases cr Channel F7-Cz with TC of 10s d
dotted green line represents th
fast phases cross the midline.
EN in patient #2. 30s EEG page
nystagmus. Channel P8-O2 indi
The large positivity at F7 at poi
deviation is produced by a 'stai
 (c) EN in patient #2.30s EEG page with left and right temporal channels demonstrating the mystagmus. Channel P8-O2 indicates the seizure onset with paroxysmal fast activity at The large positivity at F7 at points B & C ind fast phases cross the midline.
EN in patient #2. 30s EEG page with left and right temporal channels demonstrating the
nystagmus. Channel P8-02 indicates the seizure onset with paroxysmal fast activity at A.
The large posit EN in patient #2. 30s EEG pag
nystagmus. Channel P8-02 inc
The large positivity at F7 at po
deviation is produced by a 'sta
2s whereas that for P8-02 is 0
Channel F7-Fp1 with TC of 2s
dotted green line represents t
exponen
	- mystagmus. Channel P8-02 indicates the seizure onset with paroxysmal fast activity at A.
The large positivity at F7 at points B & C indicates the overall leftward eye deviation. Thi
deviation is produced by a 'staircase' o The large positivity at F7 at points B & C indicates the overall leftward eye deviation. This
deviation is produced by a 'staircase' of leftward saccades. TC for the first four channels i:
2s whereas that for P8-O2 is 0.1s deviation is produced by a 'staircase' of leftward saccades. TC for the first four channels is
2s whereas that for P8-02 is 0.1s.
2s whereas that for P8-02 is 0.1s.
Channel F7-Fp1 with TC of 2s demonstrating the oculogram 25 whereas that for P8-02 is 0.1s.

	Channel F7-Fp1 with TC of 2s demonstrating the oculogram for patient #2. Horizontal

	dotted green line represents the EEG baseline. The prolonged slow phases show a negative

	exponential Channel F7-Fp1 with TC of 2s dem
dotted green line represents the E
exponential decay approaching th
lines. This is consistent with declin
arrows indicate progressively long
: Eye movement animation repres
ing with rightwa dotted green line represents the EEG baseline. The prolonged slow phases show a negare exponential decay approaching the baseline, as indicated by the two parallel dotted red lines. This is consistent with declining veloci exponential decay approaching the baseline, as indicated by the two parallel dotted red
lines. This is consistent with declining velocity as the eyes approach the midline. Black
arrows indicate progressively longer slow ph

Videos:

exponential decay approach the midline, Black
lines. This is consistent with declining velocity as the eyes approach the midline. Black
arrows indicate progressively longer slow phases.

: Eye movement animation representi lines. This is consistent with declining velocity as the eyes approach the midline. Black arrows in the set of the set of the set of the sing with rightward post-saccadic slow drifts without
attive of nystagmus seen in Figure 1c. The eye ske
attive of nystagmus seen in Figure 1c. The eye ske
alternating with ri

Video 1: Lye movement animation representing type I EN. Video demonstrates leftward saccades
alternating with rightward post-saccadic slow drifts without crossing the midline. This is
representative of nystagmus seen in Fi representative of nystagmus seen in Figure 1c. The eye sketch is created using BioRender
Video 2: Eye movement animation representing type II EN. Video demonstrates leftward
phases alternating with rightward saccades cross **Video 2:** Eye movement animation representing type II EN. Video demonstrates leftward slow
phases alternating with rightward saccades crossing the midline. This is representation of
nystagmus seen in Figure 1a. The eye sk |
|
| Video 2: Eye movement animation representing type II En. Video demonstrates leftward slow
phases alternating with rightward saccades crossing the midline. This is representation of
nystagmus seen in Figure 1a. The eye sket phases alternative with rightward same same same middle with rightward same middle. The middle m nystagmus seen in Figure 1a. The eye sketch is created using BioRender.com.

References:

]
| |
| |
| |

mystagmus. Neurology, 43(12), 2508–2514. https://doi.org/10.1212/wnl.43.12.2508.

[2] Tusa, R. J., Kaplan, P. W., Hain, T. C., & Naidu, S., 1990. Ipsiversive eye deviation and epile

nystagmus. Neurology, 40(4), 662–665. h nystagmus. Neurology, 12(12), 12) 1212 1212. https://mate.gy. 121212/mm. 12121212. [2]

Tusa, R. J., Kaplan, P. W., Hain, T. C., & Naidu, S., 1990. Ipsiversive eye deviation

nystagmus. Neurology, 40(4), 662–665. https://d Inystagmus. Neurology, 40(4), 662–665. https://doi.org/10.1212/wnl.40.4.662.

[3] Lüders, H.O., Noachtar, S., 2000. Atlas and classification of electroencephalography.

Saunders, Philadelphia.

[4] Leigh, R.J., Zee,D.S., 2 nystagmus. Neurology, 1002–1001m, perspectively, 1002–701m, 100002

Saunders, Philadelphia.

[4] Leigh, R.J., Zee, D.S., 2015. The neurology of eye movements. Oxford Univ

York.

[5] Lee, S. U., Suh, H. I., Choi, J. Y., Hu

[4] Leigh, R.J., Zee, D.S., 2015. The neurology of eye movements. Oxford University Press
York.
The neurology of eye movements. Oxford University Press
York.
The Lee, S. U., Suh, H. I., Choi, J. Y., Huh, K., Kim, H. J., & [4] Leigh, R.J., Zee,D
York.
[5] Lee, S. U., Suh, H
case report and systema
https://doi.org/10.1016
[6] Nicita, F., Papett
Epileptic nystagmus: de
findings. Journal of the r
https://doi.org/10.1016 [5] Lee, S. U., Suh, H. I., Choi, J. Y., Huh, K., Kim, H. J., & Kim, J. S., 2014. Epileptic nystagmus: A
case report and systematic review. Epilepsy & behavior case reports, 2, 156–160.
https://doi.org/10.1016/j.ebcr.2014. [5]
[5]
case r
https:
[6]
Epileµ
findin
https:
[7] case report and systematic review. Epilepsy & behavior case reports, 2, 156–160.

https://doi.org/10.1016/j.ebcr.2014.08.004

[6] Nicita, F., Papetti, L., Spalice, A., Ursitti, F., Massa, R., Properzi, E., & lannetti, P., https://doi.org/10.1016/j.ebcr.2014.08.004

[6] Nicita, F., Papetti, L., Spalice, A., Ursitti, F., Massa, R., Properzi, E., & lannett

Epileptic nystagmus: description of a pediatric case with EEG correlation and SPE

find Formulary, 1997)

(6) Nicita, F., Papetti, L., Spalice, A., Ursitt

Epileptic nystagmus: description of a pediatr

findings. Journal of the neurological sciences,

https://doi.org/10.1016/j.jns.2010.08.022.

[7] Bekdik, P Epileptic nystagmus: description of a pediatric case with EEG correlation and SPECT
findings. Journal of the neurological sciences, 298(1-2), 127–131.
https://doi.org/10.1016/j.jns.2010.08.022.
[7] Bekdik, P., Sener, U., A

Findings. Journal of the neurological sciences, 298(1-2), 127–131.

https://doi.org/10.1016/j.jns.2010.08.022.

[7] Bekdik, P., Sener, U., Asan, I. F., Ozcelik, M., & Zorlu, Y., 2006. Epileptic nystagr

disorders : interna

managery of the neurological sciences.
https://doi.org/10.1016/j.jns.2010.08.022.
[7] Bekdik, P., Sener, U., Asan, I. F., Ozcelik, M., & Zorlu, Y., 200
disorders : international epilepsy journal with videotape, 8(4), 30
[8 Electrical Bekdik, P., Sener, U., Asan, I. F., Ozcel
disorders : international epilepsy journal wi
[8] Harris, C. M., Boyd, S., Chong, K., Harl
infancy. Journal of the neurological sciences
510x(97)00102-0.
[9] Fotedar, N. disorders : international epilepsy journal with videotape, $8(4)$, 305–308..

[8] Harris, C. M., Boyd, S., Chong, K., Harkness, W., & Neville, B. G., 1997. Epileptic nystagmus in

infancy. Journal of the neurological scien disorders international epileptic disorders : international epileptic of the equilibrations of the neurological sciences, $151(1)$, $111-114$. https://do $510x(97)00102-0$.
[9] Fotedar, N., Gajera, P., Pyatka, N., Nasrall 1 Harry. Journal of the neurological sciences, 151(1), 111–114. https://doi.org/10.1016/s0022-
10x(97)00102-0.
19] Fotedar, N., Gajera, P., Pyatka, N., Nasralla, S., Kubota, T., Vaca, G. F., Shaikh, A. G., & Lüders,
1. O., 510x(97)00102-0.

[9] Fotedar, N., Gajera, P., Pyatka, N., Nasralla, S., Kubota, T., Vaca, G. F., Shaikh, A. G., & Lüde

H. O., 2022. A descriptive study of eye and head movements in versive seizures. Seizure, 98, 44-

htt [9] Fotedar, N.,
H. O., 2022. A describitions://doi.org/10
[10] García-Past
case study video-El
videotape, 4(1), 23
[11] Ma, Y., Wan
report. Internation

H. O., 2022. A descriptive study of eye and head movements in versive seizures. Seizure, 98, 44–50
https://doi.org/10.1016/j.seizure.2022.04.003.
[10] García-Pastor, A., López-Esteban, P., & Peraita-Adrados, R., 2002. Epil https://doi.org/10.1016/j.seizure.2022.04.003.

[10] García-Pastor, A., López-Esteban, P., & Peraita-Adrados, R., 2002. Epileptic nystagmus: a

case study video-EEG correlation. Epileptic disorders : international epilepsy [10] García-Pastor, A., López-Esteban, P., & Pe
case study video-EEG correlation. Epileptic disor
videotape, 4(1), 23–28.
[11] Ma, Y., Wang, J., Li, D., & Lang, S., 2015. Ty
report. International journal of clinical and e case study video-EEG correlation. Epileptic disorders : international epilepsy journal with
videotape, 4(1), 23–28.
[11] Ma, Y., Wang, J., Li, D., & Lang, S., 2015. Two types of isolated epileptic nystagmus: case
report. I

videotape, 4(1), 23–28.

[11] Ma, Y., Wang, J., Li, D., & Lang, S., 2015. Two types of isolated epileptic nystagmus: c

report. International journal of clinical and experimental medicine, 8(8), 13500–13507.

Fig. 3500–13 [11] Ma, Y., Wang, J., I
report. International journals of $\frac{1}{2}$, $\frac{1}{2}$
report. International journals of $\frac{1}{2}$ Figure 1.1, μ , μ , representational journal of clinical and experimental medicine, 8(8), 13500–13507.
The contract medicine, 8000–13507.
The contract medicine, 8000–13500–13507.

