

Burden and risk factors for Achilles tendon rupture in the military population from 2006 to 2015: A retrospective cohort study.

John J. Fraser, DPT, PhD, FACSM*¹

email: john.j.fraser8.mil@health.mil; phone: (757) 438-0390; ORCID: 0000-0001-9697-3795;
Twitter/Med-Mastodon: @NavyPT

Jennifer A. Zellers, DPT, PhD^{2,3}

email: jzellers@wustl.edu; phone: (314) 286-1436

Christopher K Sullivan, MD⁴

email: christopher.k.sullivan5.mil@health.mil; phone: (619) 532-8429

Cory F. Janney, MD⁴

email: cory.f.janney.mil@health.mil; phone: (619) 532-8429

1. Operational Readiness and Health Directorate, Naval Health Research Center, 140 Sylvester Road, San Diego, CA 92106-3521, USA

2. Program in Physical Therapy, Washington University School of Medicine in St. Louis, 4444 Forest Park Avenue, St. Louis, MO 63108, USA

3. Department of Orthopaedic Surgery, Washington University School of Medicine in St. Louis, 660 S. Euclid Avenue, St. Louis, MO 63110, USA

4. Department of Orthopedic Surgery, Naval Medical Center San Diego, 34800 Bob Wilson Drive, San Diego, CA 92134, USA

***Correspondence to:** John J. Fraser, Operational Readiness and Health Directorate, Naval Health Research Center, 140 Sylvester Road, San Diego, CA 92106-3521, USA. email: john.j.fraser8.mil@health.mil; Twitter: @NavyPT

Disclaimer: JJF, CKS, and CJF are military service members or employees of the U.S. Government. This work was prepared as part of their official duties. Title 17, U.S.C. §105 provides that copyright protection under this title is not available for any work of the U.S. Government. Title 17, U.S.C. §101 defines a U.S. Government work as work prepared by a military service member or employee of the U.S. Government as part of that person's official duties. The views expressed in this article are those of the authors and do not necessarily reflect the official policy or position of the Department of the Navy, Department of Defense, nor the U.S. Government. The study protocol was approved by the Naval Health Research Center Institutional Review Board in compliance with all applicable federal regulations governing the protection of human subjects. Research data were derived from approved Naval Health Research Center Institutional Review Board protocol, number NHRC.2020.0207-NHSR

Author Contributions: JJF, JAZ, and CKS contributed to the drafting of this manuscript. JJF contributed to the conception, study design, data acquisition, and analysis. All authors contributed to data interpretation, critical revision, and final approval of the study.

Conflict of Interest Statement: Dr. Fraser reports grants from Congressionally Directed Medical Research Programs and the Office of Naval Research, outside of the submitted work. In addition, Dr. Fraser has a patent pending for an Adaptive and Variable Stiffness Ankle Brace, U.S. Provisional Patent Application No. 63254,474.

1 **Burden and risk factors for Achilles tendon rupture in the military population from 2006**
2 **to 2015: A retrospective cohort study**

3

4 **ABSTRACT**

5 **Background:** Achilles tendon rupture (ATR) is a serious musculoskeletal injury that results in
6 substantial functional decline, especially in highly physically demanding occupations such as
7 service in the military

8 **Purpose:** The objective of this study was to evaluate the burden and associated factors of ATR
9 in US military service members.

10 **Study Design:** Descriptive Epidemiology Study

11 **Methods:** The Defense Medical Epidemiology Database was used to identify all diagnosed ATR
12 in military personnel from 2006 to 2015. Prevalence of ATR was calculated and compared by
13 year, service branch, and military rank. Unadjusted and adjusted assessments of injury risk were
14 calculated.

15 **Results:** Officers incurred 15 978 episodes at a prevalence of 7.43/1000 (male: 8.11/1000;
16 female: 3.89/1000). Among enlisted personnel, there were 59 242 episodes of ATR that occurred
17 at a prevalence of 6.23 episodes per 1000 (male enlisted: 6.49/1000; female enlisted: 4.48/1000).
18 Apart from enlisted aviation specialists (where there was no significant difference in risk
19 between men and women), both female officers and enlisted service members had significantly
20 lower risk of ATR compared with their male counterparts in all occupations (prevalence ratio
21 [PR]: 0.26-0.73). Aviation and service officers demonstrated significantly lower risk of ATR
22 episodes (PR: 0.87-0.91) and administration, operations, intelligence, and logistic officers
23 demonstrated increased risk (PR: 1.16-1.31) compared with ground and naval gunfire officers.
24 Among enlisted specialties, all but mechanized/armor and combat engineers had significantly

25 higher risk of ATR risk compared with infantry (PR: 1.14-2.13), with the highest risk observed
26 in the administration, intelligence, and communication fields.

27 **Conclusions:** ATR was ubiquitous in the US military, with multiple risk factors identified,
28 including male sex, older age, rank, military occupation, and service branch.

29 **Clinical Relevance:** Based on the burden of ATR in the US military observed in this study, these
30 findings highlight both the need for prophylactic interventions and identification of the
31 populations who can most greatly benefit from preventive screening and care.

32

33 **Key Terms:** leg injuries, military personnel, occupational injuries, sports medicine, public health

34

35 **What is known about the subject:** In an earlier study of the ATR burden in US military
36 members, 1441 ATR cases were identified between 1998 and 2001, occurring at a rate of 30.9
37 per 100 000 person-years. In a more recent study of care episodes for Achilles tendinopathy in
38 the military, the prevalence was 17.65 per 1000 in officers and 12.22 per 1000 in enlisted
39 members, with male sex, older age, senior rank, military occupation, and service branch found to
40 be salient associated factors. It is highly plausible that occupation may also be a salient factor for
41 ATR, given that tendinopathy may be a prodromal sign of future tendon failure.

42 **What this study adds to existing knowledge:** Due to the changes in operational requirements,
43 training, engagement in overseas contingency operations, and force composition that have
44 occurred over the past 22 years, this provides an updated assessment of burden and the
45 associated risk factors of ATR.

46

47

48 INTRODUCTION

49 The Achilles tendon, also known as the calcaneal tendon, has an important role in propulsion,
50 force attenuation, and energy return during walking, running, and jumping during function.
51 Achilles tendon rupture (ATR) is a serious musculoskeletal injury that results in substantial
52 functional decline, especially in highly physically demanding occupations such as service in the
53 military.¹ In the general public, ATR affects an estimated 32 906 individuals in the US per year,
54 at an overall rate of 2.1 per 100 000 person-years.² Some of the risk factors associated with ATR
55 include previous musculoskeletal injury, chronic Achilles tendon inflammation and
56 tendinopathy, regular participation in athletic activity, spring season, genetic factors, chronic
57 systemic conditions (diabetes, hyperparathyroidism, and renal failure), and medication use
58 (steroids, quinolones, and oral bisphosphonate).³ ATR management includes both conservative
59 and surgical approaches, with return to sport occurring in 80% of affected athletes⁴ and return to
60 duty in 90-100% of affected military members.^{1,5} Among military members, rehabilitation takes
61 an average of 8.2 months in nonoperative cases¹ and 5.4 to 6.7 months following surgery^{1,5}
62 before they are cleared for return to full, unrestricted duty.

63 In an earlier study of the ATR burden in US military members, 1441 ATR cases were
64 identified between 1998 and 2001, occurring at a rate of 30.9 per 100 000 person-years.⁶ The 15-
65 fold higher burden in the military may in part be due to the unique exposures incurred during
66 military service. In a more recent study of care episodes for Achilles tendinopathy in the
67 military, the prevalence was 17.65 per 1000 in officers and 12.22 per 1000 in enlisted members,
68 with male sex, older age, senior rank, military occupation, and service branch found to be salient
69 associated factors.⁷ It is highly plausible that occupation may also be a salient factor for ATR,
70 given that tendinopathy may be a prodromal sign of future tendon failure. Due to the changes in

71 operational requirements, training, engagement in overseas contingency operations, and force
72 composition that have occurred over the past 22 years, an updated assessment of burden and the
73 associated risk factors of ATR is warranted. Therefore, the purpose of this study was to
74 determine the burden of ATR in the military population with consideration of military
75 occupation and identify risk factors that may contribute to this condition.

76

77 **METHODS**

78 A population-based epidemiological retrospective cohort study of all service members in the US
79 Armed Forces was performed, assessing sex, military occupation, rank, branch, and year on ATR
80 prevalence from 2006 to 2015. The Defense Medical Epidemiology Database (DMED; Defense
81 Health Agency, Falls Church, VA, <https://bit.ly/DHADMED5>) was utilized to identify relevant
82 healthcare encounters. This database provides aggregated data for International Classification of
83 Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) codes and de-identified patient
84 characteristics, including sex, categories of military occupations, and branch of service for all
85 active duty and reserve military service members. The database is HIPAA compliant, does not
86 include any personally identifiable or personal health information, and has been used previously
87 for the epidemiological study of ankle-foot injury in the military.^{8,9} This study was approved as
88 non-human-subjects research by the Institutional Review Board [REDACTED]
89 [REDACTED]. The Strengthening the Reporting of Observational Studies in
90 Epidemiology (STROBE) guidelines¹⁰ were used to guide reporting.

91 The database was queried for the number of distinct care episodes for a primary diagnosis
92 of ATR (ICD-9-CM codes 727.67, nontraumatic rupture of the Achilles tendon, and 845.09,
93 other sprains and strains of ankle) from 2006 to 2015. Prevalence of ATR care episodes was

94 calculated for male and female military members, enlisted and officers, in each service branch
95 (Army, Navy, Marine Corps, and Air Force) and occupational category.^{8,9} Prevalence ratio (PR)
96 point estimates and 95% confidence intervals (CIs), risk difference point estimates, attributable
97 risk, number needed to harm, and chi-square statistics were calculated in the assessment of sex
98 and occupation category using Microsoft Excel for Mac 2016 (Microsoft Corp., Redmond, WA)
99 and a custom epidemiological spreadsheet.¹¹ In the unadjusted assessment of sex and occupation
100 on ATR risk, male service members, enlisted infantry, and ground/naval gunfire officer groups
101 were used as the reference categories.

102 Due to overdispersion of the data, a negative binomial model was employed over a
103 Poisson model for the adjusted assessment of female sex (male reference), branch (US Army
104 reference), officer rank (enlisted reference), and year (2006 reference) on the risk of ATR.¹²
105 Since there were excess zeros noted in the data, with many demographic categories not having
106 any reported outcomes of interest (especially among the smaller subpopulations), assessment of
107 convergence between the unadjusted and adjusted (hurdle) negative binomial regression models
108 were performed.⁷ In the hurdle negative binomial regression model, a logistic link is employed to
109 remove demographic categories that contributed no episodes of ATR from the final count model.
110 By doing so, excess zeros that contribute to skewed point estimates, standard errors, and
111 overdispersion are parsed in the truncated regression model.¹² Results of hurdle negative
112 binomial regression are reported using calculations of the predictors regressed on count data
113 (count model assessing the rate of the outcome) as well as a linked logistic regression (zero
114 model that assessed the probability within demographic categories of not having the outcome).
115 Both unadjusted and adjusted models are reported in the supplemental material. The regression
116 analyses were performed using the PSCL (version 1.5) and MASS (version 7.3-58.1) packages

117 on R (version 3.5.1, The R Foundation for Statistical Computing, Vienna, Austria). The level of
118 significance was $p < 0.05$ for all analyses. PR point estimates were considered statistically
119 significant if CIs did not cross the 1.00 threshold. Convergence of p values, effect sizes, and 95%
120 CIs were considered when evaluating significant findings.

121

122 **RESULTS**

123 **Supplemental tables 1 and 2** detail the counts and prevalence of ATR in officer and enlisted
124 personnel. In the study epoch, officers incurred 15 978 episodes of ATR at a prevalence of 7.43
125 per 1000 (male officers: 8.11 per 1000; female officers: 3.89 per 1000). Among enlisted
126 personnel, 59 242 ATR episodes occurred at a prevalence of 6.23 episodes per 1000 (male
127 enlisted: 6.49 per 1000; female enlisted: 4.48 per 1000). **Table 1** details the univariate factor of
128 sex within occupational specialty on ATR episodes. Apart from enlisted aviation specialists
129 (with no significant difference in risk was between men and women), both female officers and
130 enlisted service members had significantly lower risk of ATR compared with their male
131 counterparts in all occupations (PR: 0.26-0.73).

132

133
134
135
TABLE 1
Assessment of Risk of Achilles Tendon Rupture by Sex Within Occupation in Members of the
US Armed Forces, 2006-2015^a

Occupational Specialty	PR	95% CI	Risk Difference (per 1000 person- years)	AR (%)	p Value	NNH
Enlisted specialties						
Artillery/Gunnery	0.73	(0.55-0.97)	-1.5	36.8	0.03	-682
Aviation	0.92	(0.66-1.30)	-0.4	-8.1	0.65	-2641
Combat Engineers	0.26	(0.08-0.82)	-2.8	279.7	0.01	-357
Maintenance	0.67	(0.63-0.70)	-2.1	-50.1	<0.001	-488
Administration, Intelligence, & Communication	0.53	(0.51-0.55)	-4.1	90.0	<0.001	-243
Logistics	0.70	(0.65-0.74)	-2.1	-43.4	<0.001	-472
Maritime/Naval Specialties	0.65	(0.53-0.81)	-2.3	-53.4	<0.001	-444
Enlisted total	0.69	(0.67-0.71)	-2.0	-44.9	<0.001	-497
Officer specialties						
Ground/Naval Gunfire	0.73	(0.57-0.93)	-2.0	-37.2	0.01	-497
Aviation	0.41	(0.32-0.54)	-3.9	-141.2	<0.001	-260
Engineering & Maintenance	0.37	(0.30-0.45)	-4.9	-172.7	<0.001	-205
Administration	0.37	(0.32-0.43)	-7.2	-171.1	<0.001	-139
Operations & Intelligence	0.50	(0.43-0.59)	-4.6	-98.7	<0.001	-218
Logistics	0.39	(0.33-0.46)	-6.2	-155.4	<0.001	-162
Services	0.48	(0.44-0.53)	-4.2	-107.3	<0.001	-239
Officer total	0.48	(0.45-0.51)	-4.2	-108.6	<0.001	-237

136 ^aFemale service members referenced to male members. Bolded values indicate statistical significance. AR,
137 attributable risk; CI, confidence interval; NNH, number needed to harm; PR, prevalence ratio.

138
139
140 In the assessment of occupational exposure, aviation and service officers demonstrated
141 significantly lower risk of ATR episodes (PR: 0.87-0.91) and administration, operations,
142 intelligence, and logistic officers demonstrated increased risk (PR: 1.16-1.31) compared with
143 ground and naval gunfire officers (**Table 2**). There was no significant difference in PR between
144 engineering and maintenance officers and the reference group. Among enlisted specialties, all
145 but mechanized/armor and combat engineers had significantly higher ATR risk compared with
146 infantry (PR: 1.14-2.13), with the highest risk observed in the administration, intelligence, and
147 communication fields.

148
149
150
TABLE 2
Assessment of Risk of Achilles Tendon Rupture by Occupation in the US Armed Forces, 2006-
2015^a

Occupational Specialty	PR	95% CI	Risk Difference (per 1000 person- years)	AR (%)	<i>p</i> value	NNH
Enlisted specialties^b						
Special Operations Forces	1.14	(1.01-1.28)	0.5	12.0	0.03	2024
Mechanized/Armor	1.03	(0.93-1.14)	0.1	3.1	0.55	8635
Artillery/Gunnery	1.48	(1.40-1.57)	1.8	32.5	<0.001	571
Aviation	1.38	(1.26-1.51)	1.4	27.6	<0.001	724
Combat Engineers	1.03	(0.94-1.13)	0.1	3.1	0.48	8580
Maintenance	1.64	(1.59-1.70)	2.3	39.1	<0.001	428
Administration, Intelligence, & Communication	2.13	(2.05-2.20)	4.1	53.0	<0.001	244
Logistics	1.83	(1.76-1.90)	3.0	45.4	<0.001	331
Maritime/Naval Specialties	1.67	(1.54-1.80)	2.4	40.0	<0.001	412
Officer specialties^c						
Aviation	0.87	(0.82-0.92)	-1.0	-15.0	<0.001	-1041
Engineering & Maintenance	0.97	(0.91-1.03)	-0.2	-3.2	0.27	-4336
Administration	1.31	(1.23-1.39)	2.2	23.4	<0.001	446
Operations & Intelligence	1.16	(1.09-1.23)	1.2	13.7	<0.001	860
Logistics	1.22	(1.15-1.30)	1.6	18.1	<0.001	617
Services	0.91	(0.87-0.96)	-0.6	-9.5	<0.001	-1571

151 ^aBolded values indicate statistical significance. AR, attributable risk; CI, confidence interval; NNH, number
 152 needed to harm; PR, prevalence ratio.

153 ^bReferenced to enlisted infantry.

154 ^cReferenced to ground and naval gunfire officers.

155
 156

157 **Table 3** details the results of the hurdle negative binomial regression model (adjusted for
 158 zero inflation) assessing sex, age range, rank, branch, and year on ATR risk. When controlling
 159 for other factors, female sex and senior officers demonstrated significantly lower PR of ATR
 160 than the male and junior enlisted reference groups, respectively. Older age (>30 years) and
 161 service in the US Marine Corps and Air Force were salient associated factors for increased risk
 162 of ATR (PR: 1.17-2.50). **Supplemental table 3** compares the negative binomial models with and
 163 without adjustment for zero inflation. While the results converged between the models for most
 164 of the variables assessed, the hurdle model outperformed the standard negative binomial model.
 165

166

167

TABLE 3

168 Negative Binomial Regression (Adjusted for Zero Inflation) Assessing Sex, Age, Rank, Service
 169 Branch, and Year on Prevalence of Achilles Tendon Rupture in the US Armed Forces, 2006-
 170 2015^a

	PR	95% CI	
		LL	UL
Sex			
Female	0.71	0.65	0.79
Age, y			
20-29	0.92	0.75	1.14
30-39	1.81	1.47	2.23
40+	2.50	2.01	3.11
Rank			
Senior enlisted	0.97	0.85	1.12
Junior officer	0.92	0.79	1.06
Senior officer	0.76	0.65	0.89
Service branch			
Navy	1.05	0.92	1.19
Marine Corps	1.38	1.20	1.58
Air Force	1.17	1.03	1.33
Year			
2007	0.92	0.75	1.13
2008	0.94	0.76	1.15
2009	0.91	0.74	1.12
2010	0.82	0.66	1.01
2011	0.79	0.64	0.97
2012	0.88	0.71	1.08
2013	0.94	0.77	1.16
2014	0.98	0.79	1.20
2015	1.08	0.88	1.32
Negative log-likelihood ratio		2358 on 41 <i>df</i>	

171 ^aBolded values indicate statistical significance ($p < 0.05$). Reference groups: sex (male), age (18-19), rank
 172 (junior enlisted), service branch (US Army), year (2006). CI, confidence interval, LL, lower limit; PR, prevalence
 173 ratio; UL, upper limit.

174

175

176 **DISCUSSION**

177 This study found that ATRs were common in the US military and identified multiple salient
 178 associated risk factors, including male sex, older age, enlisted or junior officer rank, military
 179 occupation, and service in the US Marine Corps or Air Force. Among military occupations,
 180 administration, operations, intelligence, and logistics specialties had the greatest risk for ATR in
 181 both officer and enlisted personnel. Integrating our understanding from the literature on civilian

182 populations with the present study presents unique opportunities to interrogate specific aspects of
183 mechanisms underlying ATR risk, including the contribution of exposure to activities with high
184 Achilles tendon loads, sex, and age.

185 We found the burden of ATR more closely aligns with those previously described in the
186 civilian population,² contrasted with the high rates previously reported in the US military.⁶ The
187 annual burden was consistent throughout the study epoch, despite the changes in exposure and
188 tempo of military operations. This finding diverges from studies in the civilian population
189 describing a progressive increased burden of ATR beginning in the 1990s and continuing to the
190 present.¹³ It may be that an increasingly sedentary civilian population engaged in irregular bouts
191 of activity (“weekend warriors”) has led to increased burden of ATR over time compared with
192 the more consistently physically active military population.

193 Findings from this study underscore the importance of activity exposure on ATR risk.
194 Interestingly, this study supports the framework that both overloading and underloading with
195 bouts of high loads can be detrimental to Achilles tendon health. It may seem contradictory that
196 service in the US Marine Corps (routinely encountering high Achilles tendon loading activities)
197 and Air Force (routinely involved in low Achilles tendon loading activities) both had increased
198 risk of Achilles tendon rupture compared with the US Army. We posit that these two branches
199 represent different occupational risk exposures. Individuals in the US Marine Corps are routinely
200 engaged in high-load running and jumping tasks and may represent an “overload” mechanism
201 contribution to ATR risk. Alternatively, individuals in the Air Force are more routinely involved
202 in technical occupations with longer bouts of sitting interspersed with less-frequent bouts of high
203 Achilles loading activities. This group may represent a more traditional, “underload with
204 intermittent high loading” mechanism contributing to ATR risk. Similarly, there is a lower risk

205 of ATR in senior officers (when accounting for age), supporting the idea that individuals with
206 high administrative loads, generally associated with longer bouts of sedentary activities without
207 intermittent high loading bouts, have less occupational exposure to the high eccentric loads,
208 leading to ATR.

209 A higher burden of ATR in men than in women has been well established in the civilian
210 population, with injuries occurring 3-7 times more frequently in men.¹⁴ The mechanisms
211 underlying this sexual dimorphism is a point of ongoing study. Studies of preclinical models of
212 the Achilles tendon have found regulation of female hormones to be an important contributor to
213 tendon mechanical behavior in the context of maintaining homeostasis and healing.^{15,16}
214 Furthermore, women have been found to have greater passive dorsiflexion motion compared
215 with men, a function of triceps surae extensibility.¹⁷ Recent studies in human tendon have
216 identified the presence of estrogen receptors in tendon tissue and estrogen effects on tendon
217 biomechanical function, supporting the notion of sex hormone-mediated signaling in tendon.¹⁸⁻²⁰
218 Studies investigating sex-mediated differences in patient outcomes after ATR have found
219 conflicting results, though the propensity for male patients to incur this injury more than female
220 patients is highly consistent across studies.²¹⁻²³

221 The influence of sex on tendon homeostasis and healing in humans, though, is potentially
222 confounded by a host of factors outside of direct hormonal influence. For example, it is possible
223 that behavioral differences between sexes relating to engaging in recreational sport activity²⁴
224 could partially account for differences in ATR burden, given that rupture tends to be associated
225 with forceful eccentric contractions encountered during team sports like basketball and football.
226 The present study informs this conversation by demonstrating that burden of ATR is higher in
227 men than women when occupational exposure is comparable (for every occupational specialty

228 included in this study, except for enlisted aviation). However, the relative risk encountered by
229 women compared with men was lowest in specialties in which exposures are generally
230 considered to have lower physical demands and are more sedentary (e.g., officer specialties,
231 particularly administrative and logistics roles). The findings of this study add support to the
232 notion that there is a mechanism by which sex influences ATR risk, and this risk is also
233 influenced by repeated exposure to activities in which the Achilles tendon is likely to experience
234 higher loads.

235 Service members above the age of 30 were found to be at increased risk of ATR. This
236 finding aligns well with prior studies of ATR in civilian and elite athlete populations, where the
237 highest incidence occurs in individuals aged 30-50 years.^{13,14,25,26} Individuals in middle and older
238 age groups have also consistently been found to have less optimal functional outcomes compared
239 with younger individuals,^{21,23} though individuals over the age of 65 have been reported to have
240 greater improvements in self-reported postinjury outcomes compared with younger age groups.²²
241 Identifying mechanisms underlying the isolated factor of age on ATR risk is challenging because
242 it is often difficult to disentangle behavioral changes (increased sedentary behavior) and tendon
243 maturation (age-related mechanical and physiological changes).²⁷ That said, it does seem that
244 aging can have an effect on tendon cellular function and matrix composition, though regular
245 activity mitigates many of these changes.²⁷ Similarly, aging has inconsistently been associated
246 with reduced muscle cross-sectional area and altered Achilles tendon mechanical behavior, but
247 these characteristics also seem to be moderated by physical activity level.^{27,28} Reductions in
248 muscle cross-sectional area and Achilles tendon extensibility, findings typically associated with
249 aging, were not observed when participants were matched for height, weight, and activity level.²⁹

250 The findings from the present study further support that the risk of ATR does not likely reflect
251 aging alone but rather the combined effects of both age and exposure.

252 A substantial strength of this study was that using DMED allowed for assessment of
253 population-level differences in ATR across several strata, including sex, occupation, military
254 rank, branch of service, and year. This study also had some limitations. Using diagnostic codes
255 has inherent weaknesses, especially in military populations where many determinants can
256 influence care seeking.³⁰ The data used in this study only represent individuals who sought
257 treatment for their injuries and do not include individuals who self-managed their condition,
258 though ATR presents with substantial functional deficits that would be difficult to self-manage,
259 mitigating concerns regarding care-seeking behavior. This study also included each occurrence
260 of ATR as a separate care episode. Individuals with a history of ATR have been found to have
261 increased burden of rupture on the contralateral limb compared with the general population,³¹ so
262 it is possible that a single person may have contributed more than one episode of care in the
263 present study. Lastly, although personnel were categorized by their military occupation, a
264 detailed task analysis to determine the specific duties and exposures that may be associated with
265 these injuries was not possible.

266

267 **CONCLUSION**

268 ATR was ubiquitous in the US military, with multiple risk factors identified, including male sex,
269 older age, rank, military occupation, and service branch. Based on the burden of ATR in the US
270 military observed in this study, these findings highlight both the need for prophylactic
271 interventions and identification of the populations who can most greatly benefit from preventive
272 screening and care.

273

274 **REFERENCES**

- 275 1. Renninger CH, Kuhn K, Fellars T, Youngblood S, Bellamy J. Operative and Nonoperative
276 Management of Achilles Tendon Ruptures in Active Duty Military Population. *Foot Ankle*
277 *Int.* 2016;37(3):269-273. doi:10.1177/1071100715615322
- 278 2. Lemme NJ, Li NY, DeFroda SF, Kleiner J, Owens BD. Epidemiology of Achilles Tendon
279 Ruptures in the United States: Athletic and Nonathletic Injuries From 2012 to 2016. *Orthop*
280 *J Sports Med.* 2018;6(11):2325967118808238. doi:10.1177/2325967118808238
- 281 3. Xergia SA, Tsarbou C, Liveris NI, Hadjithoma M, Tzanetakou IP. Risk factors for Achilles
282 tendon rupture: an updated systematic review. *Phys Sportsmed.* Published online June 10,
283 2022:1-11. doi:10.1080/00913847.2022.2085505
- 284 4. Zellers JA, Carmont MR, Silbernagel KG. Return to Play Post Achilles Tendon Rupture: A
285 Systematic Review and Meta-Analysis of Rate and Measures of Return to Play. *Br J Sports*
286 *Med.* 2016;50(21):1325-1332. doi:10.1136/bjsports-2016-096106
- 287 5. Tropf J, Piscocoy AS, Lundy A, Nelson S, Eckel TT. No Difference in Return to Duty
288 Between Minimally Invasive, Percutaneous Achilles Tendon Repair and Nonoperative
289 Functional Rehabilitation in Active Duty Military Members. *Foot Ankle Orthop.*
290 2022;7(2):2473011421S00544. doi:10.1177/2473011421S00544
- 291 6. Lum G. Spontaneous Ruptures of the Achilles Tendon, US Armed Forces, 1998-2001. *Med*
292 *Surveill Mon Rep.* 2002;8.
- 293 7. Sullivan CK, Janney CF, Fraser JJ. Burden and risk factors for Achilles tendinopathy in the
294 military population from 2006 to 2015. A retrospective cohort study. *J Athl Train.* 2023;(In
295 Press).
- 296 8. MacGregor AJ, Fogleman SA, Dougherty AL, Ryans CP, Janney CF, Fraser JJ. Sex
297 Differences in the Incidence and Risk of Ankle–Foot Complex Stress Fractures Among U.S.
298 Military Personnel. *J Womens Health.* 2022;31(4):586-592. doi:10.1089/jwh.2021.0292
- 299 9. Fraser JJ, MacGregor AJ, Ryans CP, Dreyer MA, Gibboney MD, Rhon DI. Sex and
300 occupation are salient factors associated with lateral ankle sprain risk in military tactical
301 athletes. *J Sci Med Sport.* 2021;24(7):677-682. doi:10.1016/j.jsams.2021.02.016
- 302 10. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The
303 Strengthening the Reporting of Observational Studies in Epidemiology (STROBE)
304 Statement: Guidelines for reporting observational studies. *Int J Surg.* 2014;12(12):1495-
305 1499. doi:10.1016/j.ijsu.2014.07.013
- 306 11. LaMorte WW. Epidemiology/Biostatistics Tools [Excel workbook]. Published online 2006.
307 Accessed August 3, 2019. <http://bit.ly/Lamort3>
- 308 12. Feng CX. A comparison of zero-inflated and hurdle models for modeling zero-inflated count
309 data. *J Stat Distrib Appl.* 2021;8(1):8. doi:10.1186/s40488-021-00121-4

- 310 13. Sheth U, Wasserstein D, Jenkinson R, Moineddin R, Kreder H, Jaglal SB. The epidemiology
311 and trends in management of acute Achilles tendon ruptures in Ontario, Canada: a
312 population-based study of 27 607 patients. *Bone Jt J.* 2017;99-B(1):78-86.
313 doi:10.1302/0301-620X.99B1.BJJ-2016-0434.R1
- 314 14. Lantto I, Heikkinen J, Flinkkilä T, Ohtonen P, Leppilahti J. Epidemiology of Achilles tendon
315 ruptures: increasing incidence over a 33-year period. *Scand J Med Sci Sports.*
316 2015;25(1):e133-138. doi:10.1111/sms.12253
- 317 15. Fryhofer GW, Freedman BR, Hillin CD, et al. Postinjury biomechanics of Achilles tendon
318 vary by sex and hormone status. *J Appl Physiol.* 2016;121(5):1106-1114.
319 doi:10.1152/jappphysiol.00620.2016
- 320 16. Pardes AM, Freedman BR, Fryhofer GW, Salka NS, Bhatt PR, Soslowsky LJ. Males have
321 inferior Achilles Tendon Material Properties Compared to Females in a Rodent Model. *Ann*
322 *Biomed Eng.* 2016;44(10):2901-2910. doi:10.1007/s10439-016-1635-1
- 323 17. Riemann BL, DeMont RG, Ryu K, Lephart SM. The Effects of Sex, Joint Angle, and the
324 Gastrocnemius Muscle on Passive Ankle Joint Complex Stiffness. *J Athl Train.*
325 2001;36(4):369-375.
- 326 18. Bryant AL, Clark RA, Bartold S, et al. Effects of estrogen on the mechanical behavior of the
327 human Achilles tendon in vivo. *J Appl Physiol.* 2008;105(4):1035-1043.
328 doi:10.1152/jappphysiol.01281.2007
- 329 19. Leblanc DR, Schneider M, Angele P, Vollmer G, Docheva D. The effect of estrogen on
330 tendon and ligament metabolism and function. *J Steroid Biochem Mol Biol.* 2017;172:106-
331 116. doi:10.1016/j.jsbmb.2017.06.008
- 332 20. Astill BD, Katsma MS, Cauthon DJ, et al. Sex-based difference in Achilles peritendinous
333 levels of matrix metalloproteinases and growth factors after acute resistance exercise. *J Appl*
334 *Physiol.* 2017;122(2):361-367. doi:10.1152/jappphysiol.00878.2016
- 335 21. Carmont MR, Zellers JA, Brorsson A, Nilsson-Helander K, Karlsson J, Grävare Silbernagel
336 K. Age and tightness of repair are predictors of heel-rise height after Achilles tendon rupture.
337 *Orthop J Sports Med.* 2020;8(3):2325967120909556.
- 338 22. Cramer A, Jacobsen NC, Hansen MS, Sandholdt H, Hölmich P, Barfod KW. Outcome after
339 acute Achilles tendon rupture is not negatively affected by female sex and age over 65 years.
340 *Knee Surg Sports Traumatol Arthrosc.* 2020;28(12):3994-4002. doi:10.1007/s00167-020-
341 06003-7
- 342 23. Olsson N, Petzold M, Brorsson A, Karlsson J, Eriksson BI, Grävare Silbernagel K.
343 Predictors of clinical outcome after acute Achilles tendon ruptures. *Am J Sports Med.*
344 2014;42(6):1448-1455.
- 345 24. Rosenfeld CS. Sex-dependent differences in voluntary physical activity. *J Neurosci Res.*
346 2017;95(1-2):279-290. doi:10.1002/jnr.23896

- 347 25. Parekh SG, Wray WH, Brimmo O, Sennett BJ, Wapner KL. Epidemiology and Outcomes of
348 Achilles Tendon Ruptures in the National Football League. *Foot Ankle Spec.* 2009;2(6):283-
349 286. doi:10.1177/1938640009351138
- 350 26. Houshian S, Tscherning T, Riegels-Nielsen P. The epidemiology of achilles tendon rupture
351 in a Danish county. *Injury.* 1998;29(9):651-654. doi:10.1016/S0020-1383(98)00147-8
- 352 27. Magnusson DM, Eisenhart M, Gorman I, Kennedy VK, E. Davenport T. Adopting
353 Population Health Frameworks in Physical Therapist Practice, Research, and Education: The
354 Urgency of Now. *Phys Ther.* 2019;99(8):1039-1047. doi:10.1093/ptj/pzz048
- 355 28. Lazarus NR, Harridge SDR. Declining performance of master athletes: silhouettes of the
356 trajectory of healthy human ageing? *J Physiol.* 2017;595(9):2941-2948.
357 doi:10.1113/JP272443
- 358 29. Couppé C, Svensson RB, Grosset JF, et al. Life-long endurance running is associated with
359 reduced glycation and mechanical stress in connective tissue. *Age Dordr Neth.*
360 2014;36(4):9665. doi:10.1007/s11357-014-9665-9
- 361 30. Fraser JJ, Schmied E, Rosenthal MD, Davenport TE. Physical therapy as a force multiplier:
362 population health perspectives to address short-term readiness and long-term health of
363 military service members. *Cardiopulm Phys Ther J.* 2020;31(1):22-28.
364 doi:10.1097/CPT.0000000000000129
- 365 31. Arøen A, Helgø D, Granlund OG, Bahr R. Contralateral tendon rupture risk is increased in
366 individuals with a previous Achilles tendon rupture. *Scand J Med Sci Sports.* 2004;14(1):30-
367 33. doi:10.1111/j.1600-0838.2004.00344.x

368