1 Transactional sex and age-disparate sexual partnerships among adolescent girls and young

2 women in Tanzania

- 3 Katherine B. Rucinski^{1*}, Gaspar Mbita^{2,3}, Kaitlyn Atkins⁴, Esther Majani⁵, Albert Komba⁶,
- 4 Caterina Casalini⁶, Mary Drake⁶, Neema Makyao⁷, Anthony Galishi⁸, Yeronimo Mlawa⁹, Joyce
- 5 Wamoyi¹⁰, Sheree Schwartz⁴, Stefan Baral⁴, Kelly Curran¹¹
- 6
- 7 1 Department of International Health, Johns Hopkins School of Public Health, Baltimore, MD,
- 8 USA
- 9 2 Jhpiego, Monrovia, Liberia
- 10 3 Athena Institute, Vrije Universiteit, Amsterdam, The Netherlands
- 11 4 Department of Epidemiology, Johns Hopkins School of Public Health, Baltimore, MD, USA
- 12 5 Embassy of Switzerland in Tanzania, Dar es Salaam, Tanzania
- 13 6 Jhpiego, Dar es Salaam, Tanzania
- 14 7 Amref Health Africa, Dar es Salaam, Tanzania
- 15 8 National Malaria Control Program, Ministry of Health, Dodoma, Tanzania
- 16 9 USAID Tanzania, Dar es Salaam, Tanzania
- 17 10 National Institute of Medical Research, Mwanza, Tanzania
- 18 11 Jhpiego, Baltimore, MD
- 19
- 20 ^{*} Corresponding author: Katherine B. Rucinski
- 21 615 N. Wolfe Street, E5034
- 22 Baltimore, MD, USA
- 23 rucinski@jhu.edu

24 Abstract

25 Adolescent girls and young women (AGYW) continue to experience a high incidence of HIV in 26 southern and eastern Africa, even in the context of large-scale HIV prevention interventions. In Tanzania, 27 AGYW account for the largest proportion of new infections and have a higher risk of HIV acquisition 28 than males of comparable age. We used routinely collected data from the PEPFAR/USAID-funded Sauti 29 Project, a large combination HIV prevention program, to examine the relationship between transactional 30 sex and sex with older partners among AGYW in Tanzania (2015-2020). Out-of-school AGYW 15-24 31 years completed a vulnerability index and were tested for HIV. We estimated weighted prevalence ratios 32 (PR) and 95% confidence intervals (CI) for the associations of transactional sex (sex exchanged for 33 money/services/gifts outside of sex work) and sex with older partners (\geq 5-years older, \geq 10-years older) 34 with prevalent HIV. We assessed potential synergism between exposures, and subgroup analyses explored 35 associations among girls 15-19. 67,357 AGYW completed the vulnerability index and 14,873 had 36 captured HIV testing records. Median age was 20 years (IQR 18-22). Transactional sex and age-disparate 37 sex were common (35% and 28%, respectively); 13% of AGYW reported both behaviors. HIV prevalence 38 was associated with both transactional sex (PR:1.28; 95% CI 1.00-1.63) and age-disparate sex (PR:1.26, 39 95% CI 0.99-1.60). In common referent analysis, transactional sex remained strongly associated with 40 HIV, even in the absence of age-disparate sex (PR 1.41; 95% CI 1.02-1.94). Evidence of statistical 41 synergism was not present, suggesting both transactional sex and age-disparate sex operate through 42 similar pathways to increase HIV risk. Increased specificity within HIV prevention programs is needed to 43 better meet the sexual and reproductive health needs of AGYW at high risk of HIV in Tanzania, including 44 investment in tailored youth-friendly strategies for AGYW who have been marginalized from the current 45 HIV response.

46

48 Introduction

49 Over the last decade, substantial global investment and the expansion of critical prevention and 50 treatment programs have resulted in overall declines in HIV incidence in countries across sub-Saharan 51 Africa, including in Tanzania (1,2). Despite this progress, the burden of HIV remains high among key and 52 other vulnerable populations such as adolescent girls and young women (AGYW) ages 15–24 (3). In 53 2022, AGYW account for the highest proportion of new infections in Tanzania and remain at greater risk 54 of infection relative to their male counterparts and other age groups (1.4.5). 55 Structural, economic and social dynamics in eastern and southern Africa, along with increased 56 biological vulnerability, heighten young women's risk of acquiring HIV (6.7). During adolescence and 57 early adulthood, key transitions including physical, physiological, emotional, and social changes happen 58 rapidly, affecting relationship formation and sexual decision making (8-11). As young people navigate 59 these changes, gender norms that assign greater social and economic power to men create the material and 60 ideological conditions that make it difficult for young women to refuse sex, negotiate condom use, or 61 advocate for their own sexual interests (12–15). These dynamics may be heightened for young women 62 living in poverty, including AGYW that live in food-insecure households and are unable to readily meet 63 their basic needs from sources other than boyfriends or other sexual partners (16). 64 Sexual relationships with older male partners can increase the risk of HIV for AGYW in eastern 65 and southern Africa. These relationships, which often comprise a sexual relationship with a male partner 66 ten or more years older, can also lead to transmission of other sexually transmitted infections, early 67 pregnancy, and child marriage (17,18). In South Africa, increasing differences in partner age are 68 associated with risk of HIV infection, particularly for adolescents (19). Phylogenetic studies also indicate 69 that men 25-40 years are the main source of HIV transmission for AGYW ages 15-25 (20). Similar 70 findings have emerged in Kenya, Uganda, and Zimbabwe, with partner age differences as little as five

71 years predictive of elevated HIV risk (21,22).

72 Transactional sex, defined by the exchange of sex for money or material support, has also been 73 shown to be consistently associated with HIV acquisition (23-25). Transactional sex exists across 74 a continuum for AGYW, with exchanges occurring within established partnerships, casual encounters, or 75 within the context of more formalized sexual exchanges such as sex work (26,27). AGYW who report 76 transactional sex have also reported a high prevalence of emotional and sexual violence, substance use, 77 and condomless sex. When transactional sexual relationships occur with older males, these HIV risks can 78 be further amplified given a higher prevalence of infection (28–30). 79 Study of transactional sex with older partners among AGYW in eastern and southern Africa has 80 increased over the last five years, with inferences largely made using established research cohorts in 81 countries such as South Africa, Uganda, and Kenya (23,31,32). While substantial qualitative work has 82 been done to contextualize experiences of transactional sex in Tanzania, few empiric studies have 83 quantitatively examined the relationship between transactional sex and HIV in the context of age-84 disparate relationships (22,27,33,34). Here, we use routinely collected data from a large combination HIV 85 prevention program to better define the relationship between transactional sex and sex with older partners, 86 and their joint impact on the HIV burden among AGYW in Tanzania. 87

88 Methods

89 Study setting and program

Data included in this analysis are from the Sauti Project, a large community-based HIV
prevention program funded by the United States Agency for International Development (USAID) through
the President's Emergency Plan for AIDS Relief (PEPFAR). Implemented by Jhpiego in partnership with
EngenderHealth, Pact, and the Tanzania National Institute for Medical Research, Sauti provided a core
package of client-centered sexual and reproductive health services for key populations such as female sex
workers and men who have sex with men (35–39). Sauti also delivered critical HIV-prevention services
for vulnerable AGYW ages 15-24 within 14 regions of Tanzania between 2015 and 2020, and was also an

97	implementing partner of the DREAMS (Determined, Resilient, Empowered, AIDS-free, Mentored, and
98	Safe) Initiative (Figure 1) (34,40). Programming delivered through Sauti for AGYW included biomedical
99	services such as HIV testing and counseling, HIV case management, sexual risk assessments, family
100	planning counseling and provision of contraception, screening for sexually transmitted infections (STI)
101	and tuberculosis, nutritional assessments, and additional screening and referrals for substance use and
102	gender-based violence. Further programming included behavioral change sessions and social protection
103	interventions such as savings and loans clubs.
104	
105	Figure 1. Sauti Project regions in Tanzania, 2015-2020
106	

107 Recruitment, eligibility, and procedures

108 AGYW between the ages of 15 and 19 who were not in school full time (defined as not having 109 attended classes more than 10 days in the last three months when not on public holidays or school 110 vacation) and those aged 20-24 were recruited beginning in 2015. Venues included biomedical service-111 delivery points such as community-based HIV testing and counseling sites, workplaces, bars, guest 112 houses, salons, markets, other venues previously identified as "hotspots" by the program, or AGYW-113 identified DREAMS safe spaces including religious houses and local government offices. Trained peer 114 facilitators recruited AGYW from each site and assessed their eligibility for program services using a 115 brief tool. This vulnerability index was adapted using existing tools from Tanzania and other countries in 116 eastern and southern Africa (40). Methods to develop this index have been previously described (40). 117 Briefly, measures included basic demographics, indicators of sexual experience including partner number 118 and age, engagement in transactional sex, experiences of intimate partner violence, depression, and food 119 insecurity, among others. The full index is reported in the Supplemental Materials (S1 File). Further, 120 AGYW who accessed HIV testing and other biomedical services through Sauti also completed a health 121 screening and service tool (HSST). Typically, these HIV testing visits occurred after AGYW had engaged

in behavior change sessions, economic empowerment sessions or other Sauti biomedical services.
Information routinely collected through the HSST included basic demographics, screening for harmful
drinking of alcohol and substance use, STI screening and treatment, and each AGYW's HIV testing

125 results.

126

127 Study population

128 We assembled an analytic sample using data collected among AGYW through the vulnerability 129 index and merged these data with biomedical HIV testing records collected through the HSST using 130 participant-specific alphanumeric codes. Data were restricted to those AGYW that accessed services 131 through Sauti between 2015-2019, and to those AGYW that were eligible for inclusion based on the 132 established eligibility criteria as described. Notably, some AGYW who did not meet full eligibility 133 requirements, including those that were still in school, were also referred for Sauti services if they 134 expressed interest in engaging with the program; these AGYW were excluded from analyses. In some 135 cases, AGYW had duplicate or repeat vulnerability index records. For these AGYW, we used the earliest 136 record available in the database. Similarly, some AGYW had multiple HIV testing results reported over 137 multiple years. For these AGYW, we used the date of first HIV testing record under the assumption that 138 this visit occurred closet to initial program engagement. Young women ages 18-24 who identified as 139 female sex workers were also excluded from analyses, as they were ineligible to complete the AGYW 140 vulnerability index.

141

142 **Outcome and Exposures**

143The primary outcome was prevalent HIV infection (yes/no), defined as a positive HIV test result144documented through the Sauti HSST. Standard HIV testing comprised serial testing using the SD Bioline145HIV-1/2 3.0 (Standard Diagnostics Inc., Suwon, Korea) and Uni-Gold HIV-1/2 (Trinity Biotech,

146 Wicklow, Ireland) assays.

1	4	7
---	---	---

148	The two primary exposures included in this analysis were self-reported engagement in
149	transactional sex and sex with older partners, both assessed through the vulnerability index. Transactional
150	sex was defined as having ever engaged in any sex for money, services, or gifts (yes/no) (41). Age-
151	disparate sex was defined as having engaged in sex with a partner five or more years older;
152	intergenerational sex was defined as having ever engaged in sex with a partner ten or more years older
153	(yes/no) (42,43).
154	
155	Statistical Analysis
156	Demographics and characteristics of AGYW participants who completed the vulnerability index
157	were summarized using proportions for categorical variables and medians for continuous variables. Chi-
158	square tests were used to compare the differences in proportions between groups for categorical variables,
159	and Wilcoxon signed-rank sum tests were used for continuous data (alpha= 0.05).
160	We assessed differences in sample characteristics among participants who did and did not have a
161	linked HIV testing result confirmed through the HSST. Among participants with an HIV testing result, we
162	further assessed differences in history of transactional sex, comparing those who had engaged in
163	transactional sex with those who reported never having engaged in transactional sex. We used a modified
164	Poisson regression model with a robust variance estimator to estimate prevalence ratios (PR) and 95%
165	confidence intervals (CI) for the association between transactional sex and prevalent HIV infection.
166	Models were restricted to AGYW who were sexually active. A directed acyclic graph (DAG) was used to
167	identify a minimally sufficient set of confounders for adjustment. Covariates in the minimally sufficient
168	set included age, age-disparate sex/intergenerational sex, adult support, marital status, prior pregnancy,
169	food insecurity, year, and early sexual debut. Models were similarly developed to assess associations
170	between age-disparate sex and HIV, as well as intergenerational sex and HIV. Interaction was assessed
171	between transactional sex and age-disparate sex, and between transactional sex and intergenerational sex

using common referent analysis and through comparison of observed and expected joint effects (44).

173 Subgroup analyses assessed main associations among adolescent girls ages 15-19.

- 174
- 175 Weights

176 Given missing HIV testing outcomes among those AGYW that did not have a linked record 177 through the HSST, we derived non-response weights under the assumption that these data were missing at 178 random (MAR) to address the possibility of selection bias and to improve the internal validity of our 179 findings (45,46). Weights were created using the full database of 67,357 unique and eligible AGYW 180 participants who completed the vulnerability index. An indicator variable for having an observed versus a 181 missing HIV testing result was created and estimated as a function of a participant's measured covariates 182 including age, partner status, prior pregnancy, food insecurity, sexual debut, experiences of violence, 183 region, and survey version (year). Covariates were chosen based on preliminary analyses to identify 184 patterns of missing information in these data, and we used multiple imputation by chained equations 185 (MICE) to fill in missing data for all covariates (n=10 imputations) (47,48). We calculated and applied 186 weights to all effect estimates for each series of imputations. Weights were stabilized by the marginal 187 probability of having an observed HIV testing outcome and were truncated at the 5th and 95th percentiles 188 to further improve stability. Weighted effect estimates and standard errors were pooled to calculate final 189 PRs and 95% CIs, standardized to the full population of AGYW who completed the vulnerability index. 190 All analyses were conducted using SAS 9.4 (SAS Institute, Cary, NC).

191

192 Ethics

193 The National Institute for Medical Research and the Ministry of Health Community 194 Development, Gender, Elderly and Children of the United Republic of Tanzania provided ethical 195 clearance for primary data collection activities under Sauti. The vulnerability index was administered in 196 the context of routine service delivery, and all participants provided verbal informed consent. Parental

 Tanzania law. Minors under the age of 18 who were parents were considered emancipated, and thus did not require parental consent. Ethical approval for the use of de-identified routine data was provided by the National Institute for Medical Research (NIMR/HQ/R.8c/Vol.1/678) and the Johns Hopkins Institutional Review Board (IRB No. 00006673). RESULTS Sample characteristics A total of 67,357 unique AGYW participants completed the Sauti vulnerability index between 2015-2019 (Figure 2). Based on their age and schooling status, 65,185 were eligible for inclusion and retained in further analyses. Of these AGYW participants, 19,748 (30.4%) lived in Dar es Salaam, 11,240 (17.3%) in Mbeya, and 8,891 (13.7%) in Shinyanga; the remaining participants were from Arusha, Dodoma, Iringa, Kilimanjaro, Morogoro, Njombe, Songwe, and Tabora. Figure 2. Construction of analytic sample including 14,873 AGYW using program data from the Sauti Project, 2015-2019 A total of 14,873 (22.8%) AGYW were able to be linked to their HIV testing record through the HSST using a unique identifier. Of these 14,873 AGYW, the median age was 20 years (IQR 15, 24) (Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime experience of transactional sex and age-disparate sex were common (35.6% and 39.1%, respectively). A total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of 	197	consent was only sought for HIV testing if the participant was under the age of 18 in accordance with
 not require parental consent. Ethical approval for the use of de-identified routine data was provided by the National Institute for Medical Research (NIMR/HQ/R.8c/Vol.1/678) and the Johns Hopkins Institutional Review Board (IRB No. 00006673). RESULTS Sample characteristics A total of 67,357 unique AGYW participants completed the Sauti vulnerability index between 2015-2019 (Figure 2). Based on their age and schooling status, 65,185 were eligible for inclusion and retained in further analyses. Of these AGYW participants, 19,748 (30,4%) lived in Dar es Salaam, 11,240 (17.3%) in Mbeya, and 8,891 (13.7%) in Shinyanga; the remaining participants were from Arusha, Dodoma, Iringa, Kilimanjaro, Morogoro, Njombe, Songwe, and Tabora. Figure 2. Construction of analytic sample including 14,873 AGYW using program data from the Sauti Project, 2015-2019 A total of 14,873 (22.8%) AGYW were able to be linked to their HIV testing record through the HSST using a unique identifier. Of these 14,873 AGYW, the median age was 20 years (IQR 15, 24) (Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime experience of transactional sex and age-disparate sex were common (35,6% and 39.1%, respectively). A total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of 	198	Tanzania law. Minors under the age of 18 who were parents were considered emancipated, and thus did
 National Institute for Medical Research (NIMR/HQ/R.8c/Vol.1/678) and the Johns Hopkins Institutional Review Board (IRB No. 00006673). RESULTS Sample characteristics A total of 67,357 unique AGYW participants completed the Sauti vulnerability index between 2015-2019 (Figure 2). Based on their age and schooling status, 65,185 were eligible for inclusion and retained in further analyses. Of these AGYW participants, 19,748 (30.4%) lived in Dar es Salaam, 11,240 (17.3%) in Mbeya, and 8,891 (13.7%) in Shinyanga; the remaining participants were from Arusha, Dodoma, Iringa, Kilimanjaro, Morogoro, Njombe, Songwe, and Tabora. Figure 2. Construction of analytic sample including 14,873 AGYW using program data from the Sauti Project, 2015-2019 A total of 14,873 (22.8%) AGYW were able to be linked to their HIV testing record through the HSST using a unique identifier. Of these 14,873 AGYW, the median age was 20 years (IQR 15, 24) (Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime experience of transactional sex and age-disparate sex were common (35.6% and 39.1%, respectively). A 	199	not require parental consent. Ethical approval for the use of de-identified routine data was provided by the
 Review Board (IRB No. 00006673). RESULTS Sample characteristics A total of 67,357 unique AGYW participants completed the Sauti vulnerability index between 2015-2019 (Figure 2). Based on their age and schooling status, 65,185 were eligible for inclusion and retained in further analyses. Of these AGYW participants, 19,748 (30.4%) lived in Dar es Salaam, 11,240 (17.3%) in Mbeya, and 8,891 (13.7%) in Shinyanga; the remaining participants were from Arusha, Dodoma, Iringa, Kilimanjaro, Morogoro, Njombe, Songwe, and Tabora. Figure 2. Construction of analytic sample including 14,873 AGYW using program data from the Sauti Project, 2015-2019 A total of 14,873 (22.8%) AGYW were able to be linked to their HIV testing record through the HSST using a unique identifier. Of these 14,873 AGYW, the median age was 20 years (IQR 15, 24) (Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime experience of transactional sex and age-disparate sex were common (35.6% and 39.1%, respectively). A total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of 	200	National Institute for Medical Research (NIMR/HQ/R.8c/Vol.1/678) and the Johns Hopkins Institutional
202203 RESULTS 204 Sample characteristics 205A total of 67,357 unique AGYW participants completed the Sauti vulnerability index between2062015-2019 (Figure 2). Based on their age and schooling status, 65,185 were eligible for inclusion and207retained in further analyses. Of these AGYW participants, 19,748 (30.4%) lived in Dar es Salaam, 11,240208(17.3%) in Mbeya, and 8,891 (13.7%) in Shinyanga; the remaining participants were from Arusha,209Dodoma, Iringa, Kilimanjaro, Morogoro, Njombe, Songwe, and Tabora.210Figure 2. Construction of analytic sample including 14,873 AGYW using program data from the Sauti212Project, 2015-2019213A total of 14,873 (22.8%) AGYW were able to be linked to their HIV testing record through the215HSST using a unique identifier. Of these 14,873 AGYW, the median age was 20 years (IQR 15, 24)216(Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime217experience of transactional sex and age-disparate sex were common (35.6% and 39.1%, respectively). A218total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of	201	Review Board (IRB No. 00006673).
 RESULTS Sample characteristics A total of 67,357 unique AGYW participants completed the Sauti vulnerability index between 2015-2019 (Figure 2). Based on their age and schooling status, 65,185 were eligible for inclusion and retained in further analyses. Of these AGYW participants, 19,748 (30.4%) lived in Dar es Salaam, 11,240 (17.3%) in Mbeya, and 8,891 (13.7%) in Shinyanga; the remaining participants were from Arusha, Dodoma, Iringa, Kilimanjaro, Morogoro, Njombe, Songwe, and Tabora. Figure 2. Construction of analytic sample including 14,873 AGYW using program data from the Sauti Project, 2015-2019 A total of 14,873 (22.8%) AGYW were able to be linked to their HIV testing record through the HSST using a unique identifier. Of these 14,873 AGYW, the median age was 20 years (IQR 15, 24) (Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime experience of transactional sex and age-disparate sex were common (35.6% and 39.1%, respectively). A total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of transactional sex. 	202	
 Sample characteristics A total of 67,357 unique AGYW participants completed the Sauti vulnerability index between 2015-2019 (Figure 2). Based on their age and schooling status, 65,185 were eligible for inclusion and retained in further analyses. Of these AGYW participants, 19,748 (30.4%) lived in Dar es Salaam, 11,240 (17.3%) in Mbeya, and 8,891 (13.7%) in Shinyanga; the remaining participants were from Arusha, Dodoma, Iringa, Kilimanjaro, Morogoro, Njombe, Songwe, and Tabora. Figure 2. Construction of analytic sample including 14,873 AGYW using program data from the Sauti Project, 2015-2019 A total of 14,873 (22.8%) AGYW were able to be linked to their HIV testing record through the HSST using a unique identifier. Of these 14,873 AGYW, the median age was 20 years (IQR 15, 24) (Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime experience of transactional sex and age-disparate sex were common (35.6% and 39.1%, respectively). A total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of 	203	RESULTS
205A total of 67,357 unique AGYW participants completed the Sauti vulnerability index between2062015-2019 (Figure 2). Based on their age and schooling status, 65,185 were eligible for inclusion and207retained in further analyses. Of these AGYW participants, 19,748 (30.4%) lived in Dar es Salaam, 11,240208(17.3%) in Mbeya, and 8,891 (13.7%) in Shinyanga; the remaining participants were from Arusha,209Dodoma, Iringa, Kilimanjaro, Morogoro, Njombe, Songwe, and Tabora.210Figure 2. Construction of analytic sample including 14,873 AGYW using program data from the Sauti212Project, 2015-2019213A total of 14,873 (22.8%) AGYW were able to be linked to their HIV testing record through the215HSST using a unique identifier. Of these 14,873 AGYW, the median age was 20 years (IQR 15, 24)216(Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime217experience of transactional sex and age-disparate sex were common (35.6% and 39.1%, respectively). A218total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of219transactional sex.	204	Sample characteristics
 206 2015-2019 (Figure 2). Based on their age and schooling status, 65,185 were eligible for inclusion and retained in further analyses. Of these AGYW participants, 19,748 (30.4%) lived in Dar es Salaam, 11,240 (17.3%) in Mbeya, and 8,891 (13.7%) in Shinyanga; the remaining participants were from Arusha, Dodoma, Iringa, Kilimanjaro, Morogoro, Njombe, Songwe, and Tabora. 210 Figure 2. Construction of analytic sample including 14,873 AGYW using program data from the Sauti Project, 2015-2019 213 A total of 14,873 (22.8%) AGYW were able to be linked to their HIV testing record through the HSST using a unique identifier. Of these 14,873 AGYW, the median age was 20 years (IQR 15, 24) (Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime experience of transactional sex and age-disparate sex were common (35.6% and 39.1%, respectively). A total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of 	205	A total of 67,357 unique AGYW participants completed the Sauti vulnerability index between
 retained in further analyses. Of these AGYW participants, 19,748 (30.4%) lived in Dar es Salaam, 11,240 (17.3%) in Mbeya, and 8,891 (13.7%) in Shinyanga; the remaining participants were from Arusha, Dodoma, Iringa, Kilimanjaro, Morogoro, Njombe, Songwe, and Tabora. Figure 2. Construction of analytic sample including 14,873 AGYW using program data from the Sauti Project, 2015-2019 A total of 14,873 (22.8%) AGYW were able to be linked to their HIV testing record through the HSST using a unique identifier. Of these 14,873 AGYW, the median age was 20 years (IQR 15, 24) (Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime experience of transactional sex and age-disparate sex were common (35.6% and 39.1%, respectively). A total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of transactional sex. 	206	2015-2019 (Figure 2). Based on their age and schooling status, 65,185 were eligible for inclusion and
 (17.3%) in Mbeya, and 8,891 (13.7%) in Shinyanga; the remaining participants were from Arusha, Dodoma, Iringa, Kilimanjaro, Morogoro, Njombe, Songwe, and Tabora. Figure 2. Construction of analytic sample including 14,873 AGYW using program data from the Sauti Project, 2015-2019 A total of 14,873 (22.8%) AGYW were able to be linked to their HIV testing record through the HSST using a unique identifier. Of these 14,873 AGYW, the median age was 20 years (IQR 15, 24) (Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime experience of transactional sex and age-disparate sex were common (35.6% and 39.1%, respectively). A total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of transactional sex. 	207	retained in further analyses. Of these AGYW participants, 19,748 (30.4%) lived in Dar es Salaam, 11,240
 Dodoma, Iringa, Kilimanjaro, Morogoro, Njombe, Songwe, and Tabora. Figure 2. Construction of analytic sample including 14,873 AGYW using program data from the Sauti Project, 2015-2019 A total of 14,873 (22.8%) AGYW were able to be linked to their HIV testing record through the HSST using a unique identifier. Of these 14,873 AGYW, the median age was 20 years (IQR 15, 24) (Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime experience of transactional sex and age-disparate sex were common (35.6% and 39.1%, respectively). A total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of transactional sex. 	208	(17.3%) in Mbeya, and 8,891 (13.7%) in Shinyanga; the remaining participants were from Arusha,
 Figure 2. Construction of analytic sample including 14,873 AGYW using program data from the Sauti Project, 2015-2019 A total of 14,873 (22.8%) AGYW were able to be linked to their HIV testing record through the HSST using a unique identifier. Of these 14,873 AGYW, the median age was 20 years (IQR 15, 24) (Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime experience of transactional sex and age-disparate sex were common (35.6% and 39.1%, respectively). A total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of transactional sex. 	209	Dodoma, Iringa, Kilimanjaro, Morogoro, Njombe, Songwe, and Tabora.
 Figure 2. Construction of analytic sample including 14,873 AGYW using program data from the Sauti Project, 2015-2019 A total of 14,873 (22.8%) AGYW were able to be linked to their HIV testing record through the HSST using a unique identifier. Of these 14,873 AGYW, the median age was 20 years (IQR 15, 24) (Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime experience of transactional sex and age-disparate sex were common (35.6% and 39.1%, respectively). A total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of transactional sex. 	210	
 Project, 2015-2019 A total of 14,873 (22.8%) AGYW were able to be linked to their HIV testing record through the HSST using a unique identifier. Of these 14,873 AGYW, the median age was 20 years (IQR 15, 24) (Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime experience of transactional sex and age-disparate sex were common (35.6% and 39.1%, respectively). A total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of transactional sex. 	211	Figure 2. Construction of analytic sample including 14,873 AGYW using program data from the Sauti
213214215215216(Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime21721821821921921921921921921921921921121221321421521521621721821921921921921021121221321421521621721821921921921121221321421521521621721821921921921021121221321421521521621721821821921921021102122132142152152162172182182192192102110212213214215216217218218219 <t< td=""><td>212</td><td>Project, 2015-2019</td></t<>	212	Project, 2015-2019
214A total of 14,873 (22.8%) AGYW were able to be linked to their HIV testing record through the215HSST using a unique identifier. Of these 14,873 AGYW, the median age was 20 years (IQR 15, 24)216(Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime217experience of transactional sex and age-disparate sex were common (35.6% and 39.1%, respectively). A218total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of219transactional sex.	213	
 HSST using a unique identifier. Of these 14,873 AGYW, the median age was 20 years (IQR 15, 24) (Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime experience of transactional sex and age-disparate sex were common (35.6% and 39.1%, respectively). A total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of transactional sex. 	214	A total of 14,873 (22.8%) AGYW were able to be linked to their HIV testing record through the
 (Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime experience of transactional sex and age-disparate sex were common (35.6% and 39.1%, respectively). A total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of transactional sex. 	215	HSST using a unique identifier. Of these 14,873 AGYW, the median age was 20 years (IQR 15, 24)
 experience of transactional sex and age-disparate sex were common (35.6% and 39.1%, respectively). A total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of transactional sex. 	216	(Table 1). One-fifth (20.9%) were married. Almost half (40.7%) had previously been pregnant. Lifetime
 total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of transactional sex. 	217	experience of transactional sex and age-disparate sex were common (35.6% and 39.1%, respectively). A
219 transactional sex.	218	total of 278 (1.9%) AGYW were living with HIV. Characteristics differed significantly by experience of
	219	transactional sex.

221 **Table 1.** Characteristics of 14,873 AGYW accessing combination HIV prevention services in Tanzania

through the Sauti Project, 2015-2019^a

	Overall N=14,87	No transactional sex N=9,521 (65.4%)		Transactional sex N=5,047 (34.6%)			
	Median	IQR	Median	IQR	Median	IQR	p- value
Age, years	20.0	18.0-	20.0	18.0-	21.0	19.0-	<.001
		22.0		22.0		23.0	
	n	%	n	%	n	%	p- value
Married	3,063	20.9	1,916	20.5	1,096	22.0	<.001
Adult support ^b	7,949	55.0	4,381	47.1	3,482	70.2	<.001
Food insecurity ^c	4,302	29.0	2,081	22.3	2,179	44.4	<.001
Ever pregnant	6,035	40.7	3,369	35.4	2,587	51.3	<.001
Sexual debut <15 years	2,544	17.7	1,099	11.8	1,412	28.7	<.001
Sexual partner >5 years older	4,162	28.3	2,156	22.8	1,953	39.1	<.001
Sexually active, ever	12,730	86.4	7,509	79.0	5,006	99.9	<.001
Sexually active, last 12 months	10,895	76.1	6,230	67.0	4,522	92.9	<.001
Using modern contraception ^d	4,774	36.8	2,663	31.5	2,111	46.5	<.001
Condomless sex, last 12 months ^e	791	74.1	472	68.0	319	85.5	<.001
>1 partner, last 12 months	5,014	34.8	1,777	19.0	3,127	65.4	<.001
Sexual violence, ever	1,901	13.1	771	8.2	1,109	22.4	<.001
Sex with HIV+ partner, last 12 months	411	2.9	186	2.0	216	4.4	<.001
Prevalent HIV	278	1.9	150	1.6	125	2.5	.006

^a Missing. transactional sex: 305 (2.1%); married: 244 (1.6%); adult support: 427 (2.9%); food insecurity: 484 (3.3%); pregnancy: 25

224 (0.2%); sexual debut: 470 (3.2%); partner age: 153 (1.0%); ever sexually active: 547 (3.7%); sexually active in the last year: 139

225 (0.9%); contraception: 659 (4.8%); number of sex partners: 438 (2.9%); sexual violence: 390 (2.6%); sex with HIV+ partner: 438

226 (2.9%); HIV: 341 (2.3%)

^bHas an adult in the household or community who provides unconditional emotional and/ or financial support

228 °Could not afford to buy food or there was not enough food to eat at home

^dUse of modern contraception asked on versions 1 and 2 of the index

^e Used a condom never, almost never, or sometimes when having vaginal sex; asked on version 3 of the index

231

232 The remaining 50,312 participants did not have a matching HIV testing record in the database. Compared

to those for whom an HIV testing record was not available, AGYW with observed HIV testing outcomes

251 which is the first interval $(20.570, 5.25.070, p)$ of $(0.001, 0.001)$	to have experienced recent food insecurity	< 0.001), to hav	(20.9% vs. 23.6%;	were less likely to be married	234
-----------------------------------------------------------------------------	--------------------------------------------	-------------------	-------------------	--------------------------------	-----

- 235 (29.9% vs. 40.8%; p<0.001), to have ever experienced sexual violence (13.1% vs. 21.8%; p<0.001), and
- to have engaged in transactional sex (37.0% vs. 45.1% p<0.001). Characteristics among AGYW
- 237 participants with and without a linked HIV testing record are reported in S1 Table.
- 238

239 Transactional sex and HIV

Lifetime experience of transactional sex was associated with HIV prevalence in unadjusted (PR: 1.41;

241 95% CI 1.10, 1.80) and adjusted models (PR: 1.34; 95% CI 1.04, 1.74) (Table 2). Weighted estimates

similarly demonstrated a population-level association between transactional sex and HIV (PR: 1.28; 95%)

243 CI 1.00, 1.63).

244

Table 2. Unadjusted, adjusted, and weighted associations of A) transactional sex with HIV prevalence; B)

age-disparate sex^a with HIV prevalence; and C) intergenerational sex^b with HIV prevalence among

247 12,730 sexually active AGYW accessing combination HIV services in Tanzania through the Sauti Project,

248 2015-2019^{c,d}

		Unadjusted PR (95% CI)	Adjusted PR (95% CI)	Weighted PR (95% CI)
Α.	No transactional sex	1 (REF.)	1 (REF.)	1 (REF.)
	Transactional sex	1.41 (1.11, 1.80)	1.34 (1.04, 1.74)	1.28 (1.00, 1.63)
В.	No age-disparate sex	1 (REF.)	1 (REF.)	1 (REF.)
	Age-disparate sex	1.20 (0.94, 1.55)	1.13 (0.88, 1.46)	1.26 (0.99, 1.60)
C.	No intergenerational sex	1 (REF.)	1 (REF.)	1 (REF.)
	Intergenerational sex	1.09 (0.71, 1.67)	0.97 (0.64, 1.47)	1.11 (0.77, 1.60)

249 Abbreviations. PR: prevalence ratio, CI: confidence interval

²⁵⁰ ^a Defined as a sexual relationship with a male partner ≥5 years older than the participating AGYW.

251 ^b Defined as a sexual relationship with a male partner ≥10 years older than the participating AGYW.

252 • Among 12,730 AGYW who reported having ever engaged in sex, of whom 309 (2.4%) were missing measures of transactional sex

and/or partner age and were excluded from analysis.

^dA directed acyclic graph (DAG) was used to identify a minimally sufficient adjustment set of covariates, and potential confounders

255 were included based on prior literature. Confounders included in the adjustment set were age, adult support, marital status, prior

pregnancy, food insecurity, early sexual debut, and survey version. For model A, intergenerational sex was also included as a potential confounder.

258

259 Older partners and HIV

- 260 There was a moderate association between age-disparate sex and HIV in unadjusted analysis (PR: 1.20;
- 261 95% CI 0.94, 1.55) that was attenuated when adjusted for potential confounders (PR: 1.15; 95% CI 0.89,
- 262 1.50). In weighted analysis, age-disparate sex was associated with a 26% increase in HIV prevalence (PR:
- 263 1.26; 95% CI 0.99, 1.60). Intergenerational sex was not strongly associated with HIV prevalence in
- 264 unadjusted (PR: 1.09; 95% CI 0.71, 1.67), adjusted (PR: 1.00; 95% CI 0.65, 1.53), or weighted models
- 265 (PR: 1.11, 95% CI 0.77, 1.60).
- 266

267 Combined association of transactional sex and sex with older partners

- 268 In common referent analysis, AGYW who engaged in transactional sex in the absence of age-disparate
- 269 partners had 1.41 (95% CI 1.02, 1.94) times the prevalence of HIV compared to AGYW who reported
- 270 neither behavior (Table 3). Conversely, age-disparate sex was not associated with HIV prevalence in the
- absence of transactional sex (PR: 1.19, 95% CI 0.81, 1.67). The prevalence ratio for those who reported
- both transactional sex and age-disparate sex was 1.45 (95% CI 1.01, 2.08), which was less than the
- 273 expected prevalence ratio using both additive and multiplicative criteria.
- 274
- **Table 3.** Interaction between a) transactional sex and age-disparate sex^a and b) transactional sex and
- 276 intergenerational sex^b on HIV prevalence among 12,730 sexually active AGYW in Tanzania, 2015-2019^c

	No transactional sex	Transactional sex
A. Age-disparate sex strata	Adjusted PR (95% CI)	Adjusted PR (95% CI)
No age-disparate sex	1.00 (REF)	1.41 (1.02, 1.94)
Age-disparate sex	1.16 (0.81, 1.67)	1.45 (1.01, 2.08)
B. Intergenerational sex strata	Adjusted PR (95% CI)	Adjusted PR (95% CI)
No intergenerational sex	1.00 (REF)	1.30 (1.00, 1.70)

Intergenerational sex 0.78 (0.40, 1.55) 1.55 (0.90, 2.65) 277 Abbreviations. PR: prevalence ratio, CI: confidence interval 278 ^a Defined as a sexual relationship with a male partner \geq 5 years older than the participating AGYW. 279 ^b Defined as a sexual relationship with a male partner ≥ 10 years older than the participating AGYW. 280 ^c Confounders included in each adjustment set were age, adult support, marital status, prior pregnancy, food insecurity, early sexual 281 debut, and survey version. 282 283 Subgroup analysis among AGYW age 15-19 284 Transactional sex was also common among AGYW age ≤ 19 (Table 4). Transactional sex, age-disparate 285 sex, and intergenerational sex were not significantly associated with HIV prevalence among younger 286 AGYW (S2 Table). 287 288 Table 4. Characteristics of 5,816 adolescent girls ages 15-19 accessing combination HIV prevention

289 services in Tanzania through the Sauti Project, 2015-2019^a

	Overall N=5,816		No transa sex N=3,879	actional (68.2%)	Transact N=1,813	ional sex (31.9%)	
	Median	IQR	Median	IQR	Median	IQR	p- value
Age, years	18.0	17.0-	18.0	17.0-	18.0	17.0-	<.001
		19.0		19.0		19.0	
	n	%	n	%	n	%	p- value
Married	553	9.7	320	8.4	219	12.3	<.001
Adult support ^b	3,015	53.5	1,758	46.5	1,230	68.9	<.001
Food insecurity ^c	1,735	30.9	899	23.8	820	46.2	<.001
Ever pregnant	1,321	22.7	679	17.5	627	34.6	<.001
Sexual debut <15 years	1,310	23.2	594	15.6	702	39.5	<.001
Sexual partner >5 years older	1,282	22.3	648	16.8	617	34.4	<.001
Sexually active, ever	4,272	74.3	2,399	61.9	1,800	100.0	<.001

Sexually active, last 12 months	3,576	63.7	1,913	50.3	1,618	92.1	<.001
Using modern contraception ^d	1,363	26.8	706	20.6	657	39.6	<.001
Condomless sex, last 12	222	70.3	136	63.3	86	85.1	<.001
months ^e							
>1 partner, last 12 months	1,657	29.4	577	14.1	1,074	60.9	<.001
>2 partners, last 12 months	526	9.3	162	4.2	363	20.6	<.001
Sexual violence, ever	673	11.9	307	8.1	355	19.9	<.001
Sex with HIV+ partner, last 12	163	2.9	76	2.0	8	4.6	<.001
months							
Prevalent HIV	65	1.2	34	0.9	29	1.7	.002

^a Missing. Married: 103 (1.8%); adult support: 185 (3.2%); food insecurity: 202 (3.5%); pregnancy: 5 (0.1%); sexual debut: 180

291 (3.1%); partner age: 58 (1.0%); ever sexually active: 68 (1.2%); sexually active in the last year 204 (3.5%); number of sex partners:

292 174 (3.0%); sexual violence: 176 (3.0%); sex with HIV+ partner: 165 (2.8%); HIV: 191 (3.3%)

^bHas an adult in the household or community who provides unconditional emotional and/ or financial support

^cCould not afford to buy food or there was not enough food to eat at home

295 °Could not afford to buy food or there was not enough food to eat at home

^d Use of modern contraception asked on versions 1 and 2 of the index

^e Used a condom never, almost never, or sometimes when having vaginal sex; asked on version 3 of the index

298

299 **DISCUSSION**

300 In this study, we examined the relationship between HIV burden and transactional sex among over 14,000

301 AGYW accessing real-world HIV prevention services through the PEPFAR/USAID-funded Sauti Project

302 in Tanzania. Transactional exchanges of money, services, or gifts for sex were common among AGYW

303 enrolled in Sauti programing, including among girls as young as age 15. These exchanges occurred

304 among AGYW who were mostly unmarried, had previously been pregnant, and were receiving emotional

305 and financial support from older adults, including family and other community members. AGYW who

306 reported transactional sex had an elevated burden of HIV, and transactional sex was strongly associated

307 with HIV prevalence even in the absence of older partners. Given the multiple social and structural

308 vulnerabilities that can lead to transactional sexual relationships, including those with older partners, there

is need to identify HIV prevention approaches that are more responsive to AGYW who may necessitatetailored and targeted programmatic support early in adolescence.

311

312 The burden of HIV was not evenly distributed among AGYW in this analysis, with estimates of HIV 313 prevalence the highest among young women who reported both transactional sex and sexual relationships 314 with older partners. While intergenerational and transactional sex are often conceptualized in the literature 315 as compatible risk factors for HIV acquisition (23,24,29,49), few studies have explored the extent to 316 which asymmetry in partner age and transactional sex interact to increase HIV-associated vulnerability 317 (28). Consistent with our findings, one case-control study in South Africa found women who reported 318 both transactional sex and intergenerational partners had nearly twice the odds of HIV compared to 319 women who reported either experience alone (22). These findings contribute to a growing body of 320 literature that aim to further distinguish heterogeneities of risk among AGYW and more effectively guide 321 HIV prevention programming. Importantly, early findings from the DREAMS initiative have 322 demonstrated little to no improvement in reducing HIV incidence across multiple countries in southern 323 and eastern Africa (50). Our results suggest that aligning HIV prevention services with the needs of 324 AGYW who necessitate more targeted and intense approaches- such as young women who engage in 325 resource-driven transactional relationships with older partners- are needed to deliver meaningful 326 reductions in incidence for AGYW moving forward. Approaches may include economic empowerment 327 for broader social asset building (51), integrated family planning and maternal health services to support 328 the needs of adolescent mothers (52), as well as vocational training to facilitate employment and improve 329 income (53), among others.

330

Experiences of transactional sex are often predicated on structural vulnerabilities, gender norms, and power differentials that motivate AGYW to engage in resource-driven sexual relationships. The high burden of food insecurity, prior pregnancy, and sexual violence among AGYW in this study, particularly among those reporting transactional sex, reinforce the larger social and structural determinants of HIV

335 that underly sexual exchanges and increase risk of infection. For AGYW living in poverty, transactional 336 sex offers a means to provide food, clothing, shelter and other resources, and can establish pathways to 337 support and security. These more explicit transactional relationships may be even riskier for younger 338 AGYW, who may be further marginalized or restricted from accessing sexual and reproductive health 339 services due to their age (54). While transactional sex and sex with older partners were not strongly 340 associated with HIV prevalence among adolescents in subgroup analyses, an alarmingly high prevalence 341 of condomless sex among girls as young as 15 engaged in transactional sex indicates a potential high 342 incidence of infection in the context of these larger social and structural factors (7,49,55). 343 344 AGYW are broadly considered a priority population for HIV epidemic control, but not all AGYW have a 345 high risk of acquiring HIV. While young women >18 years who identified as female sex workers were 346 explicitly excluded from analyses, a high prevalence of condomless sex, violence, and food insecurity 347 among adolescent girls accessing Sauti services suggests some overlap with sex work. Notably, few 348 programs in southern and eastern Africa have evaluated the impact of tailored HIV programming for 349 young women who sell sex (56–58). In Zimbabwe, DREAMS programming that implemented 350 community mobilization and social protection interventions alongside dedicated HIV testing and PrEP for 351 young women who sell sex found increased uptake of clinical services; however, no effect on HIV 352 incidence was observed (59). Additional efforts in Zimbabwe, Mozambique, and Kenya have worked to 353 engage young women who sell sex, but more broadly this population remains under-represented in both 354 research and programs (58,60). This gap may be further perpetuated by issues of identity. Young women 355 who sell sex may not identify as sex workers until they are older (61, 62), but may be marginalized from 356 existing AGYW HIV prevention and treatment programs (16,63,64). They may also feel ostracized by 357 older sex workers, thus limiting the extent to which they engage in FSW HIV prevention and treatment 358 efforts such as drop-in centers. Thus, there is an urgent need to engage this population in both research 359 and program design efforts, and to grow the evidence base of effective interventions for young women 360 who engage in necessity-driven transactional sex.

361

362 A strength of this study was our ability to leverage a large database of observational data collected during 363 the delivery of routine services through the Sauti Project in Tanzania. In general, population-level 364 inferences made using programmatic data tend to be more representative than research data, which 365 generally are collected using narrow eligibility criteria that may threaten external validity (65). Well-366 controlled research studies may also exclude adolescents who are most at risk for poor sexual and 367 reproductive health outcomes such as those who live and work on the street, potentially limiting 368 generalizability of findings (66,67). Yet programmatic data remain a relatively under-utilized resource to 369 examine the effectiveness of HIV-related interventions and implementation strategies for key populations 370 (68–71). As programs continue to roll out biomedical prevention interventions for AGYW such as long-371 acting PrEP, program data offer an opportunity to keep pace and quickly tailor and adapt implementation 372 strategies, thus providing for a more nuanced and more efficient HIV response.

373

374 This study also had several limitations. Our use of program data necessitated the merging of multiple 375 databases. Rigorous quality assurance procedures were implemented throughout the duration of the 376 program; however, challenges with unique identifiers limited the number of AGYW who could be linked 377 across program records. We used non-response weights to try and improve internal validity under the 378 assumption these unlinked records were MAR given the observed patterns in which the "missingess" of 379 records occurred. Additionally, changes to the vulnerability index and question format during program 380 implementation precluded us from fully assessing key measures including frequency of condom use, 381 which may have provided additional insight into behavioral risks within transactional sexual partnerships. 382 Further, we were also limited by temporality and were unable to determine if transactional or 383 intergenerational sexual partnerships were more recent, and whether HIV was sexually or pediatrically 384 acquired. Nevertheless, these findings provide critical insights into multiple relationship dynamics among 385 AGYW in Tanzania. Given relatively few research data available for AGYW in the context of HIV

prevention and treatment uptake in Tanzania (72), these estimates fill an epidemiologic gap and caninform targets for intervention moving forward.

388

389 Conclusions

390 Declines in HIV incidence among AGYW over the last several years have been smaller than expected

391 given substantial investments in HIV programming for AGYW in eastern and southern Africa through

392 large combination HIV prevention programs such as DREAMS. This study contributes to a growing body

393 of evidence that HIV risks among AGYW are not homogenous, and that additional implementation

394 strategies are needed to link AGYW at high-risk of HIV to behavioral, structural and biomedical

395 prevention interventions such as PrEP. These efforts likely include increased specificity among AGYW

396 programs to better meet the sexual and reproductive health needs of AGYW at high risk of HIV, but also

397 investment in tailored youth-friendly strategies for AGYW who have been marginalized from the current

398 HIV response, including young women who engage in sexual relationships with older partners and those

399 who sell sex.

400

401 Acknowledgements

402 The authors wish to thank the participants that engaged with the Sauti Project, as well as the Sauti

403 program staff that worked diligently for many years to deliver critical health services for key and priority

404 populations across Tanzania.

405 **References**

406 1. Joint United Nations Programme on HIV/AIDS. UNAIDS Data 2021 [Internet]. Geneva, 407 Switzerland; [cited 2022 Jan 4]. Available from: 408 https://www.unaids.org/en/resources/documents/2021/2021 unaids data 409 2. The DREAMS core package of interventions: A comprehensive approach to preventing HIV 410 among adolescent girls and young women [Internet], [cited 2020 Jul 6]. Available from: 411 https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0208167 412 3. Adolescents-HIV-Eastern-Southern-Africa-2021.pdf [Internet]. [cited 2022 Jun 16]. Available from: https://www.unicef.org/esa/media/8791/file/Adolescents-HIV-Eastern-413 414 Southern-Africa-2021.pdf 415 4. PHIA Project [Internet]. [cited 2022 Jan 5]. Tanzania. Available from: 416 https://phia.icap.columbia.edu/countries/tanzania/ 417 5. Brown K. Status of HIV Epidemic Control Among Adolescent Girls and Young Women 418 Aged 15–24 Years — Seven African Countries, 2015–2017. MMWR Morb Mortal Wkly 419 Rep [Internet]. 2018 [cited 2020 Jul 5];67. Available from: 420 https://www.cdc.gov/mmwr/volumes/67/wr/mm6701a6.htm 421 6. Harrison A, Colvin CJ, Kuo C, Swartz A, Lurie M. Sustained High HIV Incidence in Young 422 Women in Southern Africa: Social, Behavioral, and Structural Factors and Emerging 423 Intervention Approaches. Curr HIV/AIDS Rep. 2015 Jun;12(2):207-15. 424 7. Comins CA, Rucinski KB, Baral S, Abebe SA, Mulu A, Schwartz SR. Vulnerability profiles 425 and prevalence of HIV and other sexually transmitted infections among adolescent girls and 426 young women in Ethiopia: A latent class analysis. PloS One. 2020;15(5):e0232598. 427 8. Delany-Moretlwe S, Cowan FM, Busza J, Bolton-Moore C, Kelley K, Fairlie L. Providing 428 comprehensive health services for young key populations: needs, barriers and gaps. J Int 429 AIDS Soc. 2015;18(2 Suppl 1):19833. 430 9. Pettifor AE, Measham DM, Rees HV, Padian NS. Sexual power and HIV risk, South Africa. 431 Emerg Infect Dis. 2004;10(11):1996. 432 10. Brook DW, Morojele NK, Zhang C, Brook JS. South African adolescents: pathways to risky 433 sexual behavior. AIDS Educ Prev. 2006;18(3):259-72. 434 11. Sanders RA. Adolescent psychosocial, social, and cognitive development. Pediatr Rev. 2013 435 Aug;34(8):354-8; quiz 358-9. 436 12. Jewkes RK, Dunkle K, Nduna M, Shai N. Intimate partner violence, relationship power 437 inequity, and incidence of HIV infection in young women in South Africa: a cohort study. 438 The Lancet. 2010 Jul 3;376(9734):41-8.

439 13. Harrison A, O'Sullivan LF, Hoffman S, Dolezal C, Morrell R. Gender role and relationship 440 norms among young adults in South Africa: measuring the context of masculinity and HIV 441 risk. J Urban Health Bull N Y Acad Med. 2006 Jul;83(4):709-22. 442 14. Gottert A, Barrington C, McNaughton-Reves HL, Maman S, MacPhail C, Lippman SA, et al. 443 Gender Norms, Gender Role Conflict/Stress and HIV Risk Behaviors Among Men in 444 Mpumalanga, South Africa. AIDS Behav. 2018 Jun 1;22(6):1858-69. 445 15. Palermo T, Chzhen Y, Balvin N, Kajula L, Palermo T, Groppo V, et al. Examining 446 determinants of gender attitudes: evidence among Tanzanian adolescents. BMC Womens 447 Health. 2020 Sep 10;20(1):195. 448 16. Rucinski KB, Schwartz SR, Mishra S, Phaswana-Mafuya N, Diouf D, Mothopeng T, et al. 449 High HIV Prevalence and Low HIV-Service Engagement Among Young Women Who Sell 450 Sex: A Pooled Analysis Across 9 Sub-Saharan African Countries. J Acquir Immune Defic 451 Syndr 1999. 2020 Oct 1;85(2):148-55. 452 17. Hokororo A, Kihunrwa A, Hoekstra P, Kalluvya SE, Changalucha JM, Fitzgerald DW, et al. 453 High prevalence of sexually transmitted infections in pregnant adolescent girls in Tanzania: a 454 multi-community cross-sectional study. Sex Transm Infect. 2015 Nov;91(7):473-8. 455 18. Erulkar A, Medhin G, Weissman E. The impact and cost of child marriage prevention in 456 rural Tanzania. Poverty Gend Youth [Internet]. 2017 Jan 1; Available from: 457 https://knowledgecommons.popcouncil.org/departments sbsr-pgy/536 458 19. Topazian HM, Stoner MCD, Edwards JK, Kahn K, Gómez-Olivé FX, Twine R, et al. 459 Variations in HIV Risk by Young Women's Age and Partner Age Disparity in Rural South 460 Africa (HPTN 068). J Acquir Immune Defic Syndr 1999. 2020 Apr 1;83(4):350-6. 20. de Oliveira T, Kharsany ABM, Gräf T, Cawood C, Khanyile D, Grobler A, et al. 461 Transmission networks and risk of HIV infection in KwaZulu-Natal, South Africa: a 462 463 community-wide phylogenetic study. Lancet HIV. 2017 Jan;4(1):e41-50. 464 21. Okiring J, Getahun M, Gutin SA, Lebu S, Lee J, Maeri I, et al. Sexual partnership 465 concurrency and age disparities associated with sexually transmitted infection and risk 466 behavior in rural communities in Kenya and Uganda. Int J Infect Dis. 2022 Jul 1;120:158-467 67. 468 22. McCloskey LA, Eloff I, Doran K. Determinants of intergenerational sexual relationships and 469 HIV risk among South African women outpatients in Gauteng. AIDS Care. 2021 470 May;33(5):654–62. 471 23. Kilburn K, Ranganathan M, Stoner MCD, Hughes JP, MacPhail C, Agyei Y, et al. 472 Transactional sex and incident HIV infection in a cohort of young women from rural South 473 Africa. AIDS Lond Engl. 2018;32(12):1669-77. 474 24. Jewkes R, Dunkle K, Nduna M, Shai NJ. Transactional sex and HIV incidence in a cohort of 475 young women in the stepping stones trial. J AIDS Clin Res. 2012;3(5). 20

- 476 25. Wamoyi J, Stobeanau K, Bobrova N, Abramsky T, Watts C. Transactional sex and risk for
 477 HIV infection in sub-Saharan Africa: a systematic review and meta-analysis. J Int AIDS Soc.
 478 2016;19(1):20992.
- 479 26. Stoebenau K, Heise L, Wamoyi J, Bobrova N. Revisiting the understanding of "transactional 480 sex" in sub-Saharan Africa: A review and synthesis of the literature. Soc Sci Med. 2016 Nov 481 1;168:186–97.
- 482 27. Wamoyi J, Heise L, Meiksin R, Kyegombe N, Nyato D, Buller AM. Is transactional sex
 483 exploitative? A social norms perspective, with implications for interventions with adolescent
 484 girls and young women in Tanzania. Fielding-Miller RK, editor. PLOS ONE. 2019 Apr
 485 2;14(4):e0214366.
- 28. Ranganathan M, Kilburn K, Stoner MCD, Hughes JP, MacPhail C, Gomez-Olive FX, et al.
 The Mediating Role of Partner Selection in the Association Between Transactional Sex and
 HIV Incidence Among Young Women. J Acquir Immune Defic Syndr 1999. 2020
 01;83(2):103–10.
- 490 29. Stoner MCD, Nguyen N, Kilburn K, Gómez-Olivé FX, Edwards JK, Selin A, et al. Age491 disparate partnerships and incident HIV infection in adolescent girls and young women in
 492 rural South Africa. AIDS Lond Engl. 2019 27;33(1):83–91.
- 30. Rucinski KB, Kilburn K, delong, Stephanie Marie, Gómez-Olivé FX, Kahn K, Twine R, et
 al. Developmental trajectories of transactional sex and age-disparate relationships during
 adolescence: An HPTN 068 analysis. 2019 Jul; Mexico City, Mexo.
- 496 31. Full article: Sexual health of adolescent girls and young women in Central Uganda:
 497 exploring perceived coercive aspects of transactional sex [Internet]. [cited 2022 Jul 15].
 498 Available from: https://www.tandfonline.com/doi/full/10.1080/26410397.2019.1700770
- 32. Becker ML, Bhattacharjee P, Blanchard JF, Cheuk E, Isac S, Musyoki HK, et al.
 Vulnerabilities at First Sex and Their Association With Lifetime Gender-Based Violence and
 HIV Prevalence Among Adolescent Girls and Young Women Engaged in Sex Work,
 Transactional Sex, and Casual Sex in Kenya. J Acquir Immune Defic Syndr 1999. 2018
 01;79(3):296–304.
- 33. Wamoyi J, Wight D, Plummer M, Mshana GH, Ross D. Transactional sex amongst young
 people in rural northern Tanzania: an ethnography of young women's motivations and
 negotiation. Reprod Health. 2010 Apr 29;7(1):2.
- 34. Gichane MW, Wamoyi J, Atkins K, Balvanz P, Maman S, Majani E, et al. The influence of
 cash transfers on engagement in transactional sex and partner choice among adolescent girls
 and young women in Northwest Tanzania. Cult Health Sex. 2020 Sep 16;1–15.
- 35. Mbita G, Mwanamsangu A, Plotkin M, Casalini C, Shao A, Lija G, et al. Consistent Condom
 Use and Dual Protection Among Female Sex Workers: Surveillance Findings from a LargeScale, Community-Based Combination HIV Prevention Program in Tanzania. AIDS Behav.
 2019 Aug 23;

- 36. Mbita G, Komba AN, Casalini C, Bazant E, Curran K, Christensen A, et al. Predictors of
 HIV Among 1 Million Clients in High-Risk Male Populations in Tanzania. AIDS Behav.
 2022 Apr 1;1–14.
- 37. Nyato D, Nnko S, Komba A, Kuringe E, Plotkin M, Mbita G, et al. Facilitators and barriers
 to linkage to HIV care and treatment among female sex workers in a community-based HIV
 prevention intervention in Tanzania: A qualitative study. PLOS ONE. 2019 Nov
 19;14(11):e0219032.
- 38. Kuringe E, Materu J, Nyato D, Majani E, Ngeni F, Shao A, et al. Prevalence and correlates
 of depression and anxiety symptoms among out-of-school adolescent girls and young women
 in Tanzania: A cross-sectional study. PLOS ONE. 2019 Aug 16;14(8):e0221053.
- 39. Wambura M, Drake M, Kuringe E, Majani E, Nyato D, Casalini C, et al. Cash Transfer to
 Adolescent Girls and Young Women to Reduce Sexual Risk Behavior (CARE): Protocol for
 a Cluster Randomized Controlled Trial. JMIR Res Protoc. 2019 Dec 20;8(12):e14696.
- 40. Han H, Yang F, Murray S, Mbita G, Bangser M, Rucinski K, et al. Characterizing a sexual
 health and HIV risk stratification scale for sexually active adolescent girls and young women
 (AGYW) in Tanzania. PloS One. 2021;16(3):e0248153.
- 530 41. Stoebenau K, Heise L, Wamoyi J, Bobrova N. Revisiting the understanding of "transactional 531 sex" in sub-Saharan Africa: A review and synthesis of the literature. Soc Sci Med. 2016 532 Nov;168:186–97.
- 42. Leclerc-Madlala S. Age-disparate and intergenerational sex in southern Africa: the dynamics
 of hypervulnerability. AIDS Lond Engl. 2008 Dec;22 Suppl 4:S17-25.
- 43. UNAIDS Terminology Guidelines 2015 [Internet]. [cited 2023 Jan 9]. Available from:
 https://www.unaids.org/en/resources/documents/2015/2015_terminology_guidelines
- 44. Knol MJ, VanderWeele TJ. Recommendations for presenting analyses of effect modification
 and interaction. Int J Epidemiol. 2012 Apr 1;41(2):514–20.
- 45. Howe CJ, Cole SR, Lau B, Napravnik S, Eron JJ. Selection Bias Due to Loss to Follow Up
 in Cohort Studies. Epidemiol Camb Mass. 2016 Jan;27(1):91–7.
- 541 46. Cole SR, Hernán MA. Constructing inverse probability weights for marginal structural
 542 models. Am J Epidemiol. 2008 Sep 15;168(6):656–64.
- 543 47. Stuart EA, Azur M, Frangakis C, Leaf P. Multiple Imputation With Large Data Sets: A Case
 544 Study of the Children's Mental Health Initiative. Am J Epidemiol. 2009 May 1;169(9):1133–
 545 9.
- 48. Azur MJ, Stuart EA, Frangakis C, Leaf PJ. Multiple imputation by chained equations: what
 is it and how does it work? Int J Methods Psychiatr Res. 2011 Feb 24;20(1):40–9.

548 49. Gichane MW, Rosenberg NE, Zimmer C, Pettifor AE, Maman S, Maseko B, et al. Individual
549 and Relationship-Level Correlates of Transactional Sex Among Adolescent Girls and Young
550 Women in Malawi: A Multilevel Analysis. AIDS Behav. 2022;26(3):822–32.

- 50. Birdthistle I, Kwaro D, Shahmanesh M, Baisley K, Khagayi S, Chimbindi N, et al.
 Evaluating the impact of DREAMS on HIV incidence among adolescent girls and young
 women: A population-based cohort study in Kenya and South Africa. PLoS Med. 2021
 Oct;18(10):e1003837.
- 555 51. Iwelunmor J, Nwaozuru U, Obiezu-Umeh C, Uzoaru F, Ehiri J, Curley J, et al. Is it time to
 556 RE-AIM? A systematic review of economic empowerment as HIV prevention intervention
 557 for adolescent girls and young women in sub-Saharan Africa using the RE-AIM framework.
 558 Implement Sci Commun. 2020 Jun 10;1(1):53.
- 559 52. Groves AK, Maman S, Stankard PH, Gebrekristos LT, Amon JJ, Moodley D. Addressing the
 560 unique needs of adolescent mothers in the fight against HIV. J Int AIDS Soc. 2018 Jun
 561 29;21(6):e25155.
- 562 53. Kasirye R, Laurenzi C, Nabulya A, Nakijoba B. Safe spaces, vocational training, and
 563 prevention programs protect young Ugandan women: findings from Uganda Youth
 564 Development Link's DREAMS initiative for rural communities. Vulnerable Child Youth
 565 Stud. 2023 Apr 3;18(2):195–206.
- 566 54. Kangaude G, Coast E, Fetters T. Adolescent sexual and reproductive health and universal
 567 health coverage: a comparative policy and legal analysis of Ethiopia, Malawi and Zambia.
 568 Sex Reprod Health Matters. 28(2):1832291.
- 569 55. Gichane MW, Moracco KE, Pettifor AE, Zimmer C, Maman S, Phanga T, et al.
 570 Socioeconomic Predictors of Transactional Sex in a Cohort of Adolescent Girls and Young
 571 Women in Malawi: A Longitudinal Analysis. AIDS Behav. 2020 Dec 1;24(12):3376–84.
- 56. Busza J, Mtetwa S, Mapfumo R, Hanisch D, Wong-Gruenwald R, Cowan F. Underage and
 underserved: reaching young women who sell sex in Zimbabwe. AIDS Care. 2016;28 Suppl
 2:14–20.
- 575 57. Napierala S, Chabata ST, Fearon E, Davey C, Hargreaves J, Busza J, et al. Engagement in
 576 HIV Care Among Young Female Sex Workers in Zimbabwe. J Acquir Immune Defic Syndr
 577 1999. 2018 01;79(3):358–66.
- 578 58. Chabata ST, Makandwa R, Hensen B, Mushati P, Chiyaka T, Musemburi S, et al. Strategies
 579 to Identify and Reach Young Women Who Sell Sex With HIV Prevention and Care Services:
 580 Lessons Learnt From the Implementation of DREAMS Services in Two Cities in Zimbabwe.
 581 JMIR Public Health Surveill. 2022 Jul 27;8(7):e32286.
- 582 59. Chabata ST, Hensen B, Chiyaka T, Mushati P, Musemburi S, Dirawo J, et al. The impact of
 583 the DREAMS partnership on HIV incidence among young women who sell sex in two
 584 Zimbabwean cities: results of a non-randomised study. BMJ Glob Health. 2021 Apr
 585 1;6(4):e003892.

- 60. Bhattacharjee P, Musau A, Manguro G, Ongwen P, Mutegi J, Kioko J, et al. HIV prevention
 programme with young women who sell sex in Mombasa, Kenya: learnings for scale-up. J
 Int AIDS Soc. 2022;25(8):e25969.
- 589 61. Hensen B, Chabata ST, Floyd S, Chiyaka T, Mushati P, Busza J, et al. HIV risk among
- 590 young women who sell sex by whether they identify as sex workers: analysis of
- respondent-driven sampling surveys, Zimbabwe, 2017. J Int AIDS Soc [Internet]. 2019 Dec
- 592 3 [cited 2020 May 9];22(12). Available from:
- 593 https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6887898/
- 62. Ranganathan M, MacPhail C, Pettifor A, Kahn K, Khoza N, Twine R, et al. Young women's
 perceptions of transactional sex and sexual agency: a qualitative study in the context of rural
 South Africa. BMC Public Health. 2017 Aug 22;17:666.
- 63. Milovanovic M, Jewkes R, Matuludi M, Dunkle K, Hlongwane K, Vanleeuw L, et al. Sex
 work and young women: a cross sectional study to understand the overlap of age and sex
 work as a central tenet to epidemic control in South Africa. AIDS Care. 2022 Apr 4;1–9.

600 64. Bowring AL, Ketende S, Rao A, Njindam IM, Decker MR, Lyons C, et al. Characterising
601 unmet HIV prevention and treatment needs among young female sex workers and young
602 men who have sex with men in Cameroon: a cross-sectional analysis. Lancet Child Adolesc
603 Health. 2019 Jul 1;3(7):482–91.

- 604 65. Degtiar I, Rose S. A Review of Generalizability and Transportability. Annu Rev Stat Its
 605 Appl. 2023;10(1):null.
- 606 66. HIV and adolescents: focus on young key populations Bekker 2015 Journal of the
 607 International AIDS Society Wiley Online Library [Internet]. [cited 2022 Jul 15]. Available
 608 from: https://onlinelibrary.wiley.com/doi/full/10.7448/IAS.18.2.20076
- 609 67. Cheuk E, Isac S, Musyoki H, Pickles M, Bhattacharjee P, Gichangi P, et al. Informing HIV
 610 Prevention Programs for Adolescent Girls and Young Women: A Modified Approach to
 611 Programmatic Mapping and Key Population Size Estimation. JMIR Public Health Surveill.
 612 2019 Apr 1;5(2):e11196.
- 613 68. Rucinski K, Masankha Banda L, Olawore O, Akolo C, Zakaliya A, Chilongozi D, et al. HIV
 614 Testing Approaches to Optimize Prevention and Treatment for Key and Priority Populations
 615 in Malawi. Open Forum Infect Dis. 2022 Apr 1;9(4):ofac038.
- 616 69. Olawore O, Astatke H, Lillie T, Persaud N, Lyons C, Kamali D, et al. Peer Recruitment
 617 Strategies for Female Sex Workers Not Engaged in HIV Prevention and Treatment Services
 618 in Côte d'Ivoire: Program Data Analysis. JMIR Public Health Surveill. 2020;6(4):e18000.
- 619 70. Ortblad KF, Mawandia S, Bakae O, Tau L, Grande M, Mogomotsi GP, et al. Using routine
 620 programmatic data to measure HIV incidence among pregnant women in Botswana. Popul
 621 Health Metr. 2022 Mar 4;20:10.

- 71. Rao A, Mhlophe H, Comins C, Young K, Mcingana M, Lesko C, et al. Persistence on oral
 pre-exposure prophylaxis (PrEP) among female sex workers in eThekwini, South Africa,
 2016-2020. PloS One. 2022;17(3):e0265434.
- 625 72. Low A, Gummerson E, Schwitters A, Bonifacio R, Teferi M, Mutenda N, et al. Food
 626 insecurity and the risk of HIV acquisition: findings from population-based surveys in six
 627 sub-Saharan African countries (2016-2017). BMJ Open. 2022 Jul 12;12(7):e058704.

628

629 Supporting Information

630 S1 File. SAUTI Adolescent Girls and Young Women Vulnerability Index

631

- 632 S1 Table. Characteristics of 65,185 AGYW accessing combination HIV prevention services
- 633 in Tanzania through the Sauti Project, 2015-2019

- 635 S2 Table. Unadjusted and adjusted associations of A) transactional sex with HIV
- 636 prevalence; B) age-disparate sex with HIV prevalence; and C) intergenerational sex with
- 637 HIV prevalence among 4,272 sexually active adolescent girls ages 15-19 accessing
- 638 combination HIV services in Tanzania through the Sauti Project, 2015-2019



Figure 1



Figure 2