Training-of-trainers program for community health workers involved in an innovative and community-based intervention among goldminers in the Guiana Shield: a quality and effectiveness evaluation.

- 1 Carboni Carlotta^{1*}, Jimeno Maroto Irene¹, Galindo Muriel¹, Plessis Lorraine¹, Lambert Yann¹,
- Bardon Teddy¹, Vreden Stephen², Suárez-Mutis Martha³, Miller Bordalo Jane⁴, Douine 2
- 3 Maylis¹, Sanna Alice¹
- ¹ Centre d'Investigation Clinique Antilles-Guyane (Inserm 1424), Département Recherche, 4
- 5 Innovation et Santé Publique, Centre Hospitalier de Cayenne, Cayenne, French Guiana, France
- 6 ² Foundation for the Advancement of Scientific Research in Suriname (SWOS), Paramaribo,
- 7 Suriname
- 8 ³ Laboratório de Doenças Parasitárias, Instituto Oswaldo Cruz, Fiocruz, Rio de Janeiro, Brazil
- 9 ⁴DPAC Fronteira, Oiapoque, Brazil
- * Correspondence: 10
- 11 Carboni Carlotta
- carbonicarlotta@gmail.com 12
- 14 Keywords: Training-of-trainers, Training Programs, Community Health Workers, Capacity-
- 15 building, Malaria, Goldminers, Amazon.
- 16 **Abstract**

- **Introduction:** An innovative and community-based intervention is implemented in the Guiana Shield 17
- 18 to eliminate malaria among people involved in artisanal and small-scale gold mining. The intervention
- 19 is possible in the field thanks to community health workers (CHWs) who are previously trained to
- 20 master all intervention's procedures, including training gold miners to treat themselves for malaria.
- This article describes thus a training-of-trainers (ToT) program for CHWs, as well as the method and 21
- results of its evaluation in terms of quality and effectiveness. Methods: A mixed-method case study 22
- 23
- was implemented in two steps and based on knowledge survey, satisfaction test, observations and semi-
- structured interviews. Quantitative and qualitative data were triangulated. Results: The project team's 24
- 25 previous experience and the Guiana Shield countries' commitment to the WHO E-2025 initiative were
- identified as levers for the quality of the ToT training, while the complexity of the project context was 26
- a challenge. Group dynamics and adaptations were found to be central elements of a high-quality 27
- training program. CHWs' satisfaction was elevated especially regarding training format and learning 28
- 29 results. Improvements on knowledge level demonstrated good effectiveness of the training.
- 30 Nevertheless, some difficulties persisted regarding certain tasks to be carried out by CHWs during the
- intervention procedure. Further on-the-job training permitted to address them, improving CHWs' 31
- practices in the field. **Discussion**: High-quality, effective, and appropriate training programs are 32
- required for effective and sustainable intervention involving CHWs profiles. Training design is a 33
- 34 crucial point to attend quality and effectiveness. ToT model has been shown to allow a high level of
- 35 satisfaction, good learning results and satisfactory implementation in the field. Initial and continuing

- 36 training is an indispensable continuum to sustain good practices in the field and CHWs' motivation.
- 37 Training evaluation permits to standardize methods and facilitate transferability of successful efforts.

1 Introduction

- 40 Although since 2000 the malaria burden considerably decreased in the Americas, it is still endemic in
- the Guiana Shield (1-3). In this area, the major malaria reservoir is represented by a mobile, and hard-41
- 42 to-reach population challenging the fight against malaria (4,5). This population is mainly represented
- 43 by Brazilians (95-98%) involved in artisanal and small-scale gold mining (ASGM) in the Amazonian
- 44 rainforest, where they live in a situation of vulnerability with reduced access to the health care system
- 45 (6,7).

38

- 46 Despite the complexity of reaching populations suffering from social inequality, strategies have been
- 47 developed to provide health care access (8). Indeed, at a global scale community-based interventions
- 48 play a fundamental role in improving population health, and increasing access to health care and
- 49 services (9,10). Worldwide, a range of effective interventions involving community health workers
- 50 (CHWs) exists, especially with the aim of fighting malaria (11,12). In malaria elimination programs,
- 51 CHWs are often involved in case management, malaria surveillance, diagnosis, treatment, prevention
- 52 and health promotion activities (13).
- 53 A complex, innovative and community-based intervention has been implemented in 2018-2020 as a
- 54 result of a scientific partnership established between Brazil, French Guiana and Suriname focusing on
- populations involved in ASGM in the Guiana Shield in situations of extreme isolation from health 55
- 56 services (14). The aim was the evaluation of a strategy that consisted in providing access to diagnosis
- 57 and treatment through the distribution of kits (called "malakits"), associated with a training to use it
- 58 correctly (15). The malakit was composed of self-testing kits with rapid diagnostic tests (RDTs) and
- 59 self-treatment kits in case of malaria This strategy was evaluated as effective and acceptable by the
- 60 community, and has been scaled up in Suriname (16–18).
- Started in 2023, another intervention's evaluation study Curema expands the foundations of the 61
- 62 Malakit project to also include the radical cure of *Plasmodium vivax* (19,20). Curema intervention
- 63 consists of two modules offered at inclusion sites located in Brazil and Suriname, in resting and
- 64 logistical places across the border in French Guiana, regularly frequented by people involved in ASGM
- in French Guiana: the "malakit" module and the "radical cure" module. The latter aims to prevent 65
- relapses and thus reduce the number of human hosts capable of transmitting the parasite; asymptomatic 66
- individuals at risk of carrying *P.vivax* hypnozoites are offered presumptive treatment with chloroquine 67
- 68 and aminoquinolines.
- 69 The intervention is made possible by CHWs, specifically recruited for the Curema project, and working
- in pairs on the field at the inclusion sites. As part of the Curema project, CHWs are responsible for 70
- 71 enrolling participants and providing health education to participants followed by individual training
- 72 during the inclusion process (Figure 1). Furthermore, they carry out collective outreach activities on
- 73 malaria in the community.
- 74 Before the launch of the project, the CHWs participated in a training program which enabled them to
- 75 take part in a bidirectional working space, to master health research procedures, to present information
- 76 effectively, to lead activities that reinforce participants' learning, and to collaborate on the design of

- 77 working tools. Moreover, they were supported during the gradual launch of the project in the field and
- 78 regularly supervised during the whole project implementation.
- 79 Despite the large integration of CHWs in public health interventions, little is known about best
- practices for training them. To ensure effective CHWs activities, it seems vital to provide a good 80
- 81 training and a continuous professional development through supervisions. Nevertheless, literature is
- often lacking evidence on the implementation and evaluation of training programs. A number of 82
- 83 theoretical frameworks have been developed for evaluating this process, but the introduction of
- 84 standardized evaluation methods in the context of training evaluations is increasingly necessary (21–
- 85 24).

90

- Therefore, it seems crucial to share findings and experiences in this domain with the scientific 86
- 87 community, in order to facilitate the transferability of successful efforts.
- This present article aims to illustrate the results of an evaluation of the quality and effectiveness of the 88
- 89 training program offered to CHWs involved in the Curema project.

2 Materials and methods

91 **Description of the Curema training program**

92 2.1.1 CHWs' recruitment

- 93 The recruitment process was conducted separately by local partners for Brazilian and Surinamese
- 94 inclusion sites with the support of the project's promotor, the Hospital of Cayenne, French Guiana.
- 95 Their profile is similar to CHWs involved in malaria intervention in other countries (25,26). CHWs'
- 96 recruitment criteria were: being over 18 years of age, belonging to the community involved in ASGM
- 97 or knowing it closely, speaking fluent Portuguese, being able to use information technology tools
- 98 (tablets, smartphones, etc.), living in or being prepared to move to one of the project locations for
- 99 inclusion sites. Moreover, a technical diploma as nurse-assistant was an asset for CHWs assigned to
- 100 Brazil.
- 101 The CHWs working at the Brazilian inclusion sites were employed by the NGO DPAC-Fronteira
- 102 (Desenvolvimento, Prevencao, Acompanhamento e Cooperacao de Fronteiras; Development,
- 103 Prevention, Support and Cooperation on Borders) and were only involved in activities linked to the
- 104 Curema project. DPAC-Fronteira selected CHWs before they underwent the training program. At the
- 105 Surinamese inclusion sites, the CHWs were employed by the Surinamese research foundation SWOS
- 106 for the Curema project (half-time) and by the Surinamese National Malaria Elimination Programme
- 107 (NMEP) for the second half-time as "malaria service deliverer" (MSD). This MSD role included in
- 108 malaria case management (diagnosis, treatment, notification, and investigation) as well as in
- 109 distributing mosquito nets, in remote areas related to ASGM (27). CHWs for Suriname inclusions sites
- 110 underwent a two-steps recruitment process, consequently a larger number of potential CHWs was
- 111 invited to attend the training program. Most CHWs invited to attend the Curema training program,
- 112 already functioned as MSD in the mining fields. It was made clear to all trainees, that participating in
- 113 the training was no guarantee for being selected to work in the Curema project. Final CHWs were
- 114 selected after the end of the initial training, just before the launch of the project in the field.
- 115 Prior to the Curema training program, CHWs undertook basic training sessions separately as a pre-
- 116 work. In Brazil, a basic one-day training on malaria was provided by Fiocruz and a training on NGO

- 117 work and community mediation was provided by DPAC. In Suriname, CHWs received initial and
- 118 refresher training on malaria prevention, diagnosis, and treatment techniques by NMEP.

119 2.1.2 Design and development of the training program

- 120 With the aim of making CHWs able to master all the procedures and tasks they need to carry out, the
- 121 Curema training program was designed using a training-of-trainers (ToT) model with two components:
- 122 an initial training and an on-the-job training (Figure 2).
- Adult learning principles were applied to design the training program focusing on practice-based 123
- 124 activities to be appropriate for CHWs' needs and to give them the opportunity to practice (28). To be
- 125 more effective, and to reach all of the audience, varied activities and methods were implemented
- 126 mobilizing cognitive, affective, psychomotor learning principles, and the visual, auditory and
- 127 kinesthetic (VAK) learning model (29–31). Moreover, the training program was tailored and centered
- 128 on CHWs. Therefore, the training program was conducted in Portuguese using learning tools targeting
- 129 on an audience with a low level of scholarity.
- All partners and organizations involved in the Curema project participated in the training program, 130
- 131 increasing contributions from all staff and from professionals with different experiences and expertise.
- 132 Trainers used a horizontal approach to make CHWs feel respected and to make the learning
- 133 environment safe and supportive. To be more responsive to the needs of the CHWs and the community
- 134 in the field, a bidirectional approach was used, and CHWs were asked to collaborate in the design of
- 135 operating tools.
- 136 The initial training component took place in Paramaribo, Suriname, jointly by the entire team of CHWs
- 137 both from Suriname and Brazil. Considering the type and number of topics to be addressed, the initial
- 138 training was developed for three-weeks to allow enough time to ensure an effective knowledge transfer,
- 139 an exhaustive practice of skills and new procedures, an appropriate adaptation of tools, and a good
- 140 team and capacity building.
- The on-the-job training component took place at each inclusion site and lasted two to three days. Its 141
- aim was to support CHWs while the project was gradually rolled out in the field and the first inclusions 142
- 143 carried out.
- 144 The components included on the ToT program, as Centers for Disease Control and Prevention
- 145 recommend, are shown in **Table 1** (32).

146 Evaluation of the quality and effectiveness of the training program

147 2.2.1 Framework

- 148 Evaluation was carried out for the quality and effectiveness of training.
- 149 Quality was defined by the International Standard Vocabulary as "all the characteristics of an entity
- that influence its ability to satisfy expressed and implicit needs" (33). In this context, authors referred 150
- 151 to "needs" as the outcomes to be achieved by CHWs through the training program.
- Effectiveness corresponded to the extent to which the training outcomes are achieved. According to 152
- 153 Kirkpatrick's theoretical framework, effectiveness is assessed at 4 levels: reaction, learning, behavior,
- 154 and system (34).

155 The elements to be evaluated were defined as follow (24):

Quality assessment

- O Context: external factors which can influence the training program, both as obstacles and levers and which cannot be controlled by the training deliverers.
- Input: internal conditions under which the training program takes place (eg. methods used by trainers, CHWs' prior knowledge, group dynamism, etc.).
- Process: the training delivery process, including the adaptation, dose delivered, CHWs reached.

Effectiveness assessment

- o Reaction: level of CHWs' satisfaction in terms of dose and content.
- o Learning: changes in level of knowledge, know-how and interpersonal skills.
- o Behavior: level of concordance of CHWs' practice with the reference (Curema protocol)
- o System: impact of CHWs' practice on community health education

2.2.2 Methods

156

157

158

159

160 161

162

163 164

165

166

167

168

169

- The design of the training evaluation consisted of a mixed-method case study, executed in two steps 170
- as shown in Figure 2. Firstly, during the initial training in Paramaribo developed in a classroom 171
- 172 environment, and secondly during the on-the-job training during the launch of the intervention on
- inclusion sites. First step is nurture by quantitative and qualitative methods, while second step 173
- 174 correspond to qualitative method. The quantitative component provided computable data on
- 175 satisfaction, knowledge improvement and practice, while the qualitative component enabled giving
- 176 context and an in-depth explanation of the quantitative results.
- Data were collected through knowledge surveys, a satisfaction test, observations, and semi-structured 177
- 178 interviews, and analyzed as follows (Table 2):
- 179 The analysis was conducted using a theoretical framework conceptualized for the evaluation of training
- 180 programs for the control of infectious diseases in cross-border contexts, by the Centre for Infectious
- Disease Control, National Institute for Public Health of Health of The Netherlands and The EU Healthy 181
- 182 Gateways Joint Action (24).

183 **2.2.3** Ethics

- Curema protocol was approved by the ethics committees of the countries involved in the project 184
- (Brazil: CONEP:5,507,241 and Suriname: CMWO05/22). The evaluation of quality and effectiveness 185
- 186 of CHWs' training program is part of the secondary objectives of the Curema protocol.
- According to the article R1121-1 of the French Public Health Code, the regulatory classification of this 187
- 188 evaluation is a "research not involving the human being" (outside the scope of the *Loi Jardé* or *RNIPH*)
- 189 because it "aims to evaluate the methods of practice of healthcare health professionals or teaching
- 190 practices in the field of health" (35). A collective oral information was reached out and participants'
- 191 non-opposition was collected. For semi-structured interviews, consent was requested.
- 192 The evaluation was conducted in accordance with the principles of the Declaration of Helsinki. Data
- 193 collected were recorded in accordance with French law and the European Union General Data
- 194 Protection Regulation. Data collected, field notes and transcribed recordings were anonymized and
- 195 stored on a secure server.

3 **Results**

196

201

209

210

- The initial training took place between February the 13th 2023 and March the 03rd 2023. On-the-job 197
- training stared with the launch of the intervention at the first inclusion site on March the 13th 2023 and 198
- lasted two to three days per site. The evaluation was carried out between February the 13th 2023 and 199
- April the 03rd 2023. 200

CHWs profile 3.1

- A total of 20 CHWs were invited for the training program, 14 from Suriname employer-organizations 202
- 203 and 6 from Brazil employer-organization. The characteristics of the trainees are summarized in Table
- 204 3. The majority were women over 40, with Portuguese as their mother tongue. Five CHWs had Spanish
- 205 as their mother tongue, but could proficiently speak and understand Portuguese. Less than a quarter
- 206 had previously worked in research. CHWs profiles were different in Brazil and in Suriname because
- 207 of different recruitment criteria. Of the 20 CHWs, 4 took part in individual semi-structured interviews
- 208 during Evaluation Step 2.

Results of the quality evaluation

3.2.1 Contextual levers and difficulties

- 211 Two elements were identified as levers for the training. The first is the political, economic and
- 212 intellectual environment in which the Curema project - and consequently the training program -
- 213 navigates. In fact, Suriname and French Guiana have committed to eliminate malaria from their
- 214 territories by 2025 (World Health Organization's E-2025 initiative) and Brazil by 2035 (1,36,37). There
- 215 is no doubt that this commitment ensured good conditions for the start of Curema intervention and its
- 216 training program. Second, previous lessons learned during Malakit project by our team in training the
- 217 same community in a similar context and partners experiences were precious to guide the design and
- 218 development of the ToT program, particularly regarding the methodology, the time, the appropriate
- 219 tools for the audience (38).
- On the other hand, the complexity of Curema project also influenced the training because of a large 220
- 221 number of interacting components that are sometimes difficult to align in order to make possible the
- 222 training program development (39). In fact, Curema is an international project which means several
- 223 organizational levels, country-specific features for implementation on the field, communication in
- 224 different foreign languages depending on the actors, logistical issues linked to differences in
- 225 bureaucracy and regulatory requirements between countries (40). Moreover, it was important to
- 226 consider that the number and level of articulations of the activities required by CHWs complexified
- 227 the training program design. Finally, differences on recruitment and employment frameworks made
- 228 CHWs group composition heterogeneous: previous baselines knowledge were dissimilar due to
- 229 previous experiences and to a pre-work organized separately for Brazil and Suriname.

230 3.2.2 The importance of group dynamics

- 231 Among the CHWs who attended the training program, 4 (20.0%) had already worked in the research
- 232 field and 10 (50%) had already collaborated with one of the three employer-organizations involved in
- 233 the Curema project. We observed that, for these CHWs, prior knowledge of certain procedures
- 234 increased mastery of the subject and reduced the time needed to learn. In addition, thanks to a
- 235 participatory approach to training and the inclusion of practical sessions of pair and group work in the
- 236 schedule, peer education dynamics took place. The attention in the choice of the classroom and the
- 237 setting is also decisive in the generation of these dynamics. Thus, these CHWs became a resource for

perpetuity.
It is made available under a CC-BY-NC 4.0 International license .

- 238 supporting training. However, certain disadvantages were also identified such as the transfer of some
- 239 erroneous automatisms already adopted in their practice. Feedback activities allowed to readjust
- 240 practices.
- 241 The observed interpersonal relationships among CHWs had a strong influence on group dynamics for
- 242 learning, active participation in training and engagement in the project. Because of the different
- 243 recruitment framework, at the beginning of the training program CHWs appeared to form two distinct
- 244 groups. Based on team building and adult-learning principles, more collaborative activities were
- 245 implemented in order to facilitate exchanges and integration. Some spontaneous team-building
- 246 initiatives were also pointed out.
- 247 "Afterwards, the group work was done, which [...] allowed people to feel more in tune with each other
- 248 and communication improved [...] which was fun and at the same time a joint learning process, it was
- 249 very good." (Interview verbatim, CHW from Suriname)

3.2.3 A flexible and evolutive process

- 251 The training was delivered in 105 hours over a period of three consecutive weeks (**Table 1**). The
- 252 training program was designed to be part of a continuous professional development approach centered
- 253 on CHWs. Thus, it was flexible and evolutive. Numerous adaptations were made to ensure constant
- 254 improvement of CHW knowledge and practice and to perfect training (schedule, activities, tools) and
- 255 project elements. Indeed, many of those elements had already been co-designed through community
- 256 involvement. The bidirectional working space during the training sessions enabled to adapt vocabulary
- 257 and content of the inclusion questionnaire, to add supplementary material in the facilitation manual, to
- 258 make a video for the informed consent procedure. As they reported during informal conversations and
- 259 semi-structured interviews, CHWs greatly appreciated the opportunity to be part of this evolution
- 260 process.

261

262

250

3.3 Results of the effectiveness evaluation

3.3.1 Reaction: participant's satisfaction

- 263 Overall, CHWs affirmed to be very satisfied with the training. The methodology used for the training
- 264 was particularly appreciated by CHWs with reference to activities using kinesthetic learning model,
- practice-based activities, group-work and bidirectional approach. All items (Table 2) explored on the 265
- 266 scale of 1-5 ranging from "not at all satisfied" to "completely satisfied", were ranked over 4.
- Satisfaction is shown to be directly linked to learning: 267
- 268 "It was fun, creative, and useful for memorizing, because you remember [what you learn using these
- 269 techniques], isn't it? We don't forget. The games were very good" (Interview verbatim, CHW from
- 270 Suriname).
- Indeed, the 61.1% of the 18 CHWs answering the satisfaction test declared had reached "optimum 271
- 272 mastery", the 27.8% reported "sufficient mastery but needed support during the launch in the field";
- 273 and only one person felt "insufficient mastery" asking for "more time for training".
- CHWs were observed to be satisfied also during on-the-job training. In fact, some of them stated 274
- 275 trainees indicated that the sessions provided the knowledge and practice needed to start the job in the
- 276 field.

3.3.2 Learning: evaluation of knowledge improvement

- 278 The evaluation found a knowledge improvement with a significant increase in the average scores of
- 279 the test between pre and post survey with a p<0.05 both for test scores at t_0 - t_1 and t_0 - t_2 (**Figure 3**). On
- 280 the contrary, differences in scores between tests at t₁ vs t₂ were not significant (p=0.26). The same
- 281 trend was shown for analyzing individual scores' increase for CHWs which underwent the knowledge
- 282 survey at t₀, t₁, and t₂. The bivariate analysis for age, sex, recruitment country, mother tongue and
- 283 previous collaboration did not show any differences in these categories, apart from a better basic
- 284 knowledge in the t_2 test among those with previous experience in research field (p<0.05).
- 285 Systematic inclusion process monitoring through the checklist pointed out that two-thirds of observed
- 286 inclusion simulations (n=16) contained inaccuracies in the step-by-step process. Nevertheless,
- 287 inaccuracies often concerned the same steps. The highest rate of inaccuracy was found for the
- 288 explanation of the patient information note for informed consent procedure that was inaccurate in
- 289 29.2% of the observed inclusion simulations, followed by the process of installing the Curema
- 290 smartphone application, which had an inaccuracy rate of 16.7%. These results are in line with the
- 291 results of the qualitative analyses based on informal conversations and interviews with CHWs who
- 292 also reported reasons why those two steps were difficult in their opinions. First, the information note,
- 293 using scientific terms and constructions, was dense of complex concepts and difficult to be understand
- 294 for people with poor level of scholarity.
- 295 "What really frightened us, I can say for myself and for several other people, was the informative note
- 296 [...] What really complicates things for me is the informative note." (Interview verbatim, CHW from
- 297 Suriname)

277

- 298 Second, the Curema smartphone application was not enough explored during the training program.
- 299 Indeed, at that time, some technical developments of the application were in progress. Thus, the final
- 300 version and procedure to use it were not integrally exposed.

301 3.3.3 Behavior: from the training to the field

- 302 During the observed launch at the inclusion sites, the level of concordance of CHWs' practices with
- 303 the Curema standard procedures was adequate even if, during on-the-job training, further support by
- 304 trainers was needed to guide CHWs towards autonomy. Moreover, CHWs expressed the need for more
- 305 time and fieldwork to master the step-by-step process and re-elaborate concepts for an effective
- 306 knowledge transfer to the community.
- 307 Constructive professional relationships, good team working, and a sense of belonging to a common
- 308 cause were also pointed out by CHWs as factors able to influence behavior in the field, leading to
- 309 engagement and proactive participation in the intervention.
- 310 "If you don't feel good in the team, this can have an influence on your motivation, for example, in the
- 311 way the work is run." (Interview verbatim, CHW from Suriname)

312 4 **Discussion**

- High-quality, effective, and appropriate training are required for effective and sustainable intervention 313
- 314 involving CHWs profiles. This involves a great deal of effort in development, implementation, and
- 315 evaluation. Moreover, this study highlights the importance of quality to achieve effectiveness of a
- 316 training program. ToT model has been shown to allow a high level of satisfaction, good learning results
- 317 and satisfactory implementation in the field.

perpetuity.

It is made available under a CC-BY-NC 4.0 International license.

Limits of our study 4.1

- 319 The evaluation was carried out by researchers previously involved in the design, development, and
- 320 implementation of the training. Despite this could lead to attribution bias, several data sources – both
- quantitative and qualitative were used for the evaluation, enabling data triangulation and increasing 321
- 322 internal validity.

318

329

- 323 The Kirkpatrick's theoretical framework used recommends to assess effectiveness at 4 levels: reaction,
- 324 learning, behavior, and system. The higher the impact in terms of effectiveness, the better the achieved
- 325 the final training outcome (24). Nevertheless, the assessment at the system level is more challenging
- 326 and we were unable to evaluate it. A further evaluation of learning transfer to the community at the
- 327 end of the intervention could be realized by the post-intervention cross-sectional survey and would be
- 328 very useful.

Training design: a crucial step 4.2

- 330 Developing a high-quality training program demands careful considerations of the learning methods
- 331 and modules to be designed. We designed the training program on a ToT model based on adult-learning
- 332 principles. Consequently, the training program contained ample interactive sessions including group
- 333 discussions, role plays, practical activities, and simulations. This methodology is considered more
- 334 effective in comparison to traditional teaching approaches, especially among CHWs with poor level of
- 335 scholarity (9,41). Despite complexities, we can affirm that the quality of the training was good and met
- 336 the needs for the implementation of the Curema intervention in the field. The significant increase in
- 337 pre-post scores obtained at the knowledge survey suggests the achievement of theorical concepts.
- 338 Moreover, the ample panel of activities helped increase the satisfaction level, in accordance with the
- 339 literature (24).
- 340 A crucial point to be discussed is adaptations. During the training program, several adaptations were
- 341 necessary to be adopted both for training program and for the intervention on the field. We could even
- 342 say that one of the aims of the training was to enable these adaptations to be made, thanks to the
- 343 bidirectional space set up with the CHWs. Despite the balance between adaptation and fidelity in
- 344 implementation science is still a topic of debate (42-44), several examples in literature described
- 345 helpful adaptations: Miller et al. adapted terminology of their intervention to reduce the possibility of
- 346 misinterpretation of their study population, while Holtrop et al. shown training program or recruitment
- 347 adaptations due to outside forces or made to improve feasibility or engagement (45,46). A review by 348
- Stirman et al. identified most common reasons intervention adaptation were to address language,
- 349 cultural differences, literacy, or situational constraints (47). In our experience, we can argue that
- 350 adaptive aptitude, embedded in a community-based approach, may have enabled our intervention to be
- 351 more appropriate, effective, and sustainable by fostering the bidirectional transfer of knowledge, the
- 352 empowerment of CHWs as a central actor of the intervention, and their engagement in the Curema
- 353 project.
- 354 To be appropriate, trainings of the CHWs should take place in their work environment to practice in a
- 355 real-life situation (48,49). Our quantitative and qualitative results suggested that, although many
- 356 simulations were carried out during the initial training, real practice in the field during on-the-job
- 357 training permitted CHWs to improve mastery of procedures and to appropriately perform participants'
- 358 inclusions and education activities regarding malaria. Similar results have been shown in previous
- 359 experiences among the same community and intervention patterns by Galindo et al. who concluded
- 360 that practical training in the field was the most appropriate learning method for CHWs (38). Besides,
- 361 a scoping review describing training, supervision and quality of care in community-based programs in

perpetuity.
It is made available under a CC-BY-NC 4.0 International license .

- 362 different contexts from the studied one, also concluded that training in the field and on-site have been
- 363 shown to improve CHWs' practices (50).
- Another important element to address is team-building, which has been found out as a crucial element 364
- for learning process and daily work on the field. During the training, the team-building activities helped 365
- to motivate CHWS and encouraged peer learning. Moreover, the sense of unity jointly with the 366
- 367 engagement towards the goal of eliminating malaria increased CHW's motivation in the daily working
- 368 context, particularly in regions where malaria is in the process of being eliminated. Similar results
- 369 showing how relationships with other cadres and actors could play a major role on sustain motivation
- 370 are reported in literature (51,52).

371 **Evaluation** 4.3

- 372 Nowadays, limited number of studies exists regarding quality and effectiveness evaluation of training
- programs, especially those regarding infectious diseases in cross-border areas and those addressed to 373
- 374 CHWs. Indeed, a recent review of the literature analyzed 62 articles on the evaluation of ToT, and
- 375 found only 5 in the field of infectious diseases and in cross-border contexts, with heterogeneous and
- 376 non-standardized methodologies (24). Reviewing articles on CHWs-based programs for cardiovascular
- 377 disease management in low-income and middle-income countries with the aim of evaluate training
- 378 program effectiveness, Abdel-All and al. found only 8 articles reporting training details on evaluation,
- 379 among 90 eligible for their systematic review (53).
- Scientists call for studies examining the effectiveness and characteristics of training programs for 380
- 381 CHWs to provide reliable evidence, to avoid replication of programming difficult, to ensure high-
- 382 quality of training programs, to make possible standardization on the evaluation (53–57). This study
- 383 shows as developing methodology for training evaluation is possible, feasible and necessary to ensure
- 384 advancements on this field. Evaluation of training programs must be integrated in all CHWs-based
- 385 health intervention.

386

4.4 Initial and continuing training: an indispensable continuum

- 387 ToT is a good predictor of the long-term sustainability of public health initiatives (23). This ToT
- 388 program was the starting point of a continuous professional development scheme; indeed we
- 389 considered continuing re-training as important as initial training. In fact, despite a widespread lack of
- 390 ongoing training or other forms of continuing professional development, several studies have found
- 391 the positive association between training and maintaining good standards of practices. (58–61). This is
- 392 shown to be linked: (i) to refresh of acquired skills and knowledge and (ii) to motivation. For instance,
- 393 Curtale et al. suggest that even few days additional training/supervision, if regularly provided, results
- 394
- in improved quality of service (62). Furthermore, access to training and supervision seems to be 395 associated with non-monetary incentive sustaining CHWs motivation and engagement (13,61).
- 396 Moreover, a continuing training centered on CHWs seems to be an asset to the strategy of continuous
- 397 professional development. Puchalski Ritchie et al. in their training course, established training contents
- 398 on needs they identified through a prior qualitative survey performed by CHWs (63). These
- 399 recommendations sustain the design we adopted.
- 400 Ongoing training, including various moments of evaluation and reflection, enable interventions to be
- 401 adjusted in time to ensure proper implementation. Constant efforts throughout the project are essential
- 402 to maintain quality while adapting to the inevitable changes in the context in which the intervention
- 403 evolves.

5 Figure legends Figure 1: CHWs tasks over Curema enrolling process and knowledge/skills associated with. Figure 2: Timeline and flowchart of the training design and training evaluation over Curema intervention phases. **Figure 3**: Evolution over time in the knowledge survey score for basic knowledge (at t₀, t₁, and t₂) among all CHWs (A), for advanced knowledge (at t₁, and t₂) among all CHWs (B), for basic knowledge (at t_0 , t_1 , and t_2) per CHWs (C), for advanced knowledge (at t_1 , and t_2) per CHWs (D).

404

405

406

407

408

409

410

6 Tables

Table 1: The ToT components included on the Curema training program.

ToT component	Pre-assessment	Pre-work	Training agenda	Facilitation manual	Modelling of skills and tasks
Aim	Identifying pre-training knowledge and skills of the CHW, to determine training needs	Providing CHWs with the knowledge and background needed before the training program	Providing some degree of control over what, when and how CHWs will learn during the training program.	Organizing tools and improving the capacity of CHWs to transfer knowledge	Enabling CHWs to form a clear idea of how to carry out a task by showing them how to perform a skill while describing each step
Initial Training	Pre-training knowledge survey	Topics: basic knowledge about malaria (diagnosis and treatment), epidemiology in the Guiana Shield and malaria control strategies Pre-work training sessions took place separately for the CHWs working on the inclusion sites located in Brazil or Suriname, due to different recruitment frameworks. Those sessions were provided respectively by two different organizations, due to legitimacy of respective countries	Provided on the first day of the training	Audio-visual tools (i.e. a step- by-step tutorial about how to perform a Malakit RDT) were uploaded to tablets Visual sheets for explanation support were put together in a CHW binder Games and teach-back cards were also provided as teaching tools	Following training units: Audio-visual tutorial for Malakit RDT and for G6PD test; Information Education and Communication (IEC) strategies and tools use; Inclusion process simulation role-play; Informative note and informed consent procedure role-play
On-the-job training	Post-initial-training knowledge survey Feedback of the trainers regarding the initial training program outcomes	Initial training	Not applicable	The same tools and CHW blinder with the addition of some updated sheets	Not applicable

ToT component	Adults learning principles	Skill practice	Feedback	Action planning	Planned follow-up support	
Aim	Diversifying activities based on VAK Model and Bloom's Taxonomy, enhancing experience, designing a problem-centered training program to reach CHWs and make the training effective	Providing the opportunity for the practice of selected training activities or contents.	Collecting CHWs' feedback regarding the learning process to be adjusted.	Taking CHWs through the process of creating or adjusting tools and practices for the intervention to succeed.	Strengthening the knowledge and skills of CHWs, so they can be applied effectively.	
Initial Training	Implementation of games, practical simulations, video making, and a variety of activities with movements needed Activities in small working groups composed of heterogeneous profiles for skills and personalities	Practice of: data collection, tablet use, G6PD test execution, Malakit RDT execution monitoring, participants and community training, health education, public speaking, inclusion process simulations, installation of Curema application for smartphone	Daily debriefing at the beginning of the day regarding previous content, doubts, difficulties. Use of a checklist to help on completing the whole inclusion process Satisfaction assessment and discussion group on the last day	Information Note video - for the signature of the ICF (Informed Consent Form) Feedback about Questionnaire	Continuous professional development through ongoing training for the intervention launch on the inclusion sites.	
On-the-job training	Directly linked to on-site work and problem-solving	Inclusion process, community outreach activities	Debriefing together at the end of inclusion process (either real process or simulations)	CHWs' initiatives encouraged (group of discussions, projection of promotional videos, marauding and outreach activities, etc)	Continuous professional development and targeted follow-up: (i) regular supervision visits (ii) remote support via a specific WhatsApp group	

Table 2: Methodology used to evaluate quality and effectiveness of the Curema training program.

Evaluation method	Collection Tool	Inclusion criteria	Temporal aspects	Analysis	Software
Knowledge survey	Basic and advanced multiple choice and true-false questions	All CHWs attending the training program who agreed to undergo the knowledge survey (N=18 at t ₀ ; N=16 at t ₁ ; N=15 at t ₂)	Evaluation step 1: Pre-training (t ₀), immediate post- training (t ₁) and 1- month follow-up (t ₂) questionnaires	Mann–Whitney–Wilcoxon U test to compare average scores; Wilcoxon signed ranks test to compare average scores only for CHWs answered the three tests; Bivariate analyse of the following variables: age, gender, recruitment country, mother tongue, previous collaboration, retained after training	Microsoft® Excel® for Microsoft 365 MSO (Version 2304 Build 16.0.16327.20200) Stata® (Version 14.2 Copyright 1985-2015 StataCorp LLC).
Observation	Observation guide to collect:	Evaluation step 1:	Evaluation step 1:	Inductive content analysis;	ATLAS.ti®
(direct and indirect)	Informal conversations with participants and trainers;	All CHWs attending the initial training who agreed to participate in the	During the initial training	Descriptive analysis for quantitative data	(Version 23.1.2.0, licence L-685-998).
	Participant observation;	observation (N=20)			Microsoft® Excel® for Microsoft 365 MSO (Version 2304 Build 16.0.16327.20200)
	Systematic inclusion process monitoring through a check-list;	Evaluation step 2:			
	Documentary research;	Convenience sampling depending on the	Evaluation step 2:		
	Discussion groups	inclusion site to be launched. CHWs assigned to the predetermined inclusion site who agreed to participate in the observation (N=4)	During on-the-job training		
Satisfaction Test	Single-choice and 5-point-Likert-	All CHWs attending the initial training	Evaluation step 1:	Descriptive analysis	Mentimeter® (Version 3.2.7, 26-08-2021)
	scale test (anonymous test) to explore opinion on methodology, relevance, clarity, trainers support, practice, facilitation manual, group dynamics and logistic	who accepted to fill in the satisfaction test (N=18)	At the end of the initial training		
Semi- structured interviews	CHWs interviews: an interview guide was developed based on preliminary results of the observations	Evaluation step 2:	Evaluation step 2:	Description analysis on the verbatim	ATLAS.ti®
		Convenience sampling depending on the inclusion site to be launched. CHWs	During on-the-job training	(Brazilian Portuguese) mainly deductive with inductive elements	(Version 23.1.2.0, licence L-685-998).
		assigned to the predetermined inclusion site who agreed to be interviewed (N=4)	(4h 38' 18")		

Table 3: Characteristics of CHWs who attended initial training program in Paramaribo and who participated to semi-structured interviews

CHWs characteristics		Involved in the initial training in Paramaribo (N=20)		Involved in semi-structured interviews (N=4)	
		Percentage	Number	Percentage	Number
		(%)	(n)	(%)	(n)
Sex M	lale	20.0	4	25.0	1
Age (years)					
	-40	20.0	4	25.0	1
	>40	80.0	16	75.0	3
Country of the employer-organization					
Bra	azil	30.0	6	0.0	0
Surina	ıme	70.0	14	100.0	4
Mother tongue		75.0	1.5	100.0	4
Portugue Span		75.0 25.0	15 5	100.0 0.0	4 0
Span	11511	23.0	3	0.0	U
Previous collaboration with the employer-organization	Yes	50.0	10	100.0	4
Previous work experience in research	Yes	20.0	4	0.0	0
	ies	20.0	4	0.0	U
Belonging to ASCM- related community					
	Yes	70.0	14	100.0	4
Retained after training	Yes	75.0	15	100.0	4

418

perpetuity.

It is made available under a CC-BY-NC 4.0 International license .

7 Conflict of Interest

- The authors declare that the research was conducted in the absence of any commercial or financial
- relationships that could be construed as a potential conflict of interest.

424 8 Author Contributions

- 425 CC, JMI, GM, MD and AS participated in the conceptualization. CC and JMI developed the
- 426 methodology. CC made formal analysis and drafted the manuscript. JMI, MD and AS reviewed
- initial draft. All the authors read, corrected, and approved the final manuscript.

428 9 Funding

421

- 429 Curema project is funded by European Funds FEDER-PCIA and Synergy no 8754, and by Regional
- 430 Health Agency of French Guiana.

431 10 Acknowledgments

- 432 Authors would like to thank all CHWs who participated to the training and the evaluation, and who are
- 433 currently make possible the implementation of Curema intervention in the field. Moreover, authors
- 434 thank Surinamese MoH and particularly the NMEP for their support to CHWs' training program in
- 435 Paramaribo.

436 11 References

- 437 1. World Health Organization. World malaria report 2022 [Internet]. Geneva: World Health
- 438 Organization; 2022. Available from: https://www.who.int/publications/i/item/9789240064898
- 439 2. Ministério da Saúde. Boletim Epidemiológico Especial Malária 2021 [Internet]. Secretaria de
- Vigilância em Saúde. 2021 [cited 2023 Sep 13]. Available from: https://www.gov.br/saude/pt-
- 441 br/centrais-de-
- conteudo/publicacoes/boletins/epidemiologicos/especiais/2021/boletim epidemiologico especial
- 443 malaria 2021.pdf/view
- 3. Musset L, Pelleau S, Girod R, Ardillon V, Carvalho L, Dusfour I, et al. Malaria on the Guiana
- Shield: a review of the situation in French Guiana. Mem Inst Oswaldo Cruz. 2014
- 446 Aug;109(5):525–33.
- 447 4. Douine M, Lambert Y, Musset L, Hiwat H, Blume LR, Marchesini P, et al. Malaria in Gold
- Miners in the Guianas and the Amazon: Current Knowledge and Challenges. Curr Trop Med
- 449 Rep. 2020 Jun 1;7(2):37–47.
- 5. Douine M, Musset L, Corlin F, Pelleau S, Pasquier J, Mutricy L, et al. Prevalence of Plasmodium
- spp. in illegal gold miners in French Guiana in 2015: a hidden but critical malaria reservoir.
- 452 Malar J. 2016 Jun 9;15:315.
- 453 6. Douine M, Mosnier E, Le Hingrat Q, Charpentier C, Corlin F, Hureau L, et al. Illegal gold
- 454 miners in French Guiana: a neglected population with poor health. BMC Public Health. 2017 Jul
- 455 17;18(1):23.


- 456 7. Le Tourneau FM. Chercheurs d'or - L'orpaillage clandestin en Guyane française [Internet]. CNR
- 457 editions. 2020 [cited 2023 Sep 12]. Available from:
- 458 https://www.cnrseditions.fr/catalogue/geographie-territoires/chercheurs-d-or/
- 8. Marmot M, Friel S, Bell R, Houweling TAJ, Taylor S, Commission on Social Determinants of 459
- 460 Health. Closing the gap in a generation: health equity through action on the social determinants
- 461 of health. Lancet Lond Engl. 2008 Nov 8;372(9650):1661-9.
- 462 Bhutta, Zulfiqar & Lassi, Zohra & Pariyo, George & Huicho, Luis. Global Experience of
- Community Health Workers for Delivery of Health Related Millennium Development Goals: A 463
- 464 Systematic Review, Country Case Studies, and Recommendations for Integration into National
- 465 Health Systems. WHOGlobal Health Workforce Alliance (GHWA) Geneva: World Health
- 466 Organization. 2010.
- 467 10. Rocha R, Soares RR. Evaluating the Impact of Community-Based Health Interventions: Evidence
- 468 from Brazil's Family Health Program. IZA DP. 2009.
- 469 11. Christopher JB, Le May A, Lewin S, Ross DA. Thirty years after Alma-Ata: a systematic review
- 470 of the impact of community health workers delivering curative interventions against malaria,
- 471 pneumonia and diarrhoea on child mortality and morbidity in sub-Saharan Africa. Hum Resour
- 472 Health. 2011 Oct 24;9:27.
- 473 12. Chanda P, Hamainza B, Moonga HB, Chalwe V, Pagnoni F. Community case management of
- 474 malaria using ACT and RDT in two districts in Zambia: achieving high adherence to test results
- 475 using community health workers. Malar J. 2011 Jun 9;10(1):158.
- 476 13. Sunguya BF, Mlunde LB, Ayer R, Jimba M. Towards eliminating malaria in high endemic
- 477 countries: the roles of community health workers and related cadres and their challenges in
- 478 integrated community case management for malaria: a systematic review. Malar J. 2017 Jan
- 479 3;16(1):10.
- 480 14. Galindo MS, Lambert Y, Mutricy L, Garancher L, Bordalo Miller J, Gomes JH, et al. Setting-up
- 481 a cross-border action-research project to control malaria in remote areas of the Amazon:
- 482 describing the birth and milestones of a complex international project (Malakit). Malar J. 2021
- 483 Dec;20(1):216.
- 484 15. Douine M, Sanna A, Galindo M, Musset L, Pommier de Santi V, Marchesini P, et al. Malakit: an
- 485 innovative pilot project to self-diagnose and self-treat malaria among illegal gold miners in the
- 486 Guiana Shield. Malar J. 2018 Apr 10;17(1):158.
- 487 16. Douine M, Lambert Y, Galindo MS, Mutricy L, Sanna A, Peterka C, et al. Self-diagnosis and
- 488 self-treatment of malaria in hard-to-reach and mobile populations of the Amazon: results of
- 489 Malakit, an international multicentric intervention research project. Lancet Reg Health – Am
- 490 [Internet]. 2021 Dec 1 [cited 2022 Nov 9];4. Available from:
- 491 https://www.thelancet.com/journals/lanam/article/PIIS2667-193X(21)00039-9/fulltext
- 492 17. Parent AA, Galindo MS, Lambert Y, Douine M. Combatting malaria disease among gold miners:
- 493 a qualitative research within the Malakit project. Health Promot Int. 2022 Aug 1;37(4):daac058.

- 494 18. Douine M, Cairo H, Galindo MS, Vreden S, Lambert Y, Adenis A, et al. From an interventional
- 495 study to a national scale-up: lessons learned from the Malakit strategy at the French Guiana-
- 496 Suriname border. Malar J. 2023 Aug 16;22(1):237.
- 497 19. Curema project - Malakit [Internet]. 2023 [cited 2023 Sep 25]. Available from:
- 498 https://www.malakit-project.org/curema-project/
- 499 20. Sanna A, Lambert Y, Musset L, Vreden S, Suárez-Mutis M, White M, et al. CUREMA: an
- 500 operational research project aiming malaria elimination among hard-to-reach mobile population
- in the Guiana Shield. [Internet]. 2021. Available from: 501
- 502 https://www.researchgate.net/publication/358646601 CUREMA an operational research proje
- 503 ct aiming malaria elimination among hard-to-
- 504 reach mobile population in the Guiana Shield
- 21. Kobel S, Wartha O, Wirt T, Dreyhaupt J, Lämmle C, Friedemann EM, et al. Design, 505
- 506 Implementation, and Study Protocol of a Kindergarten-Based Health Promotion Intervention.
- 507 BioMed Res Int. 2017;2017:1-9.
- 508 22. Lai AY, Stewart SM, Mui MW, Wan A, Yew C, Lam TH, et al. An Evaluation of a Train-the-
- 509 Trainer Workshop for Social Service Workers to Develop Community-Based Family
- Interventions. Front Public Health. 2017 Jun 30;5:141. 510
- 23. Mormina M, Pinder S. A conceptual framework for training of trainers (ToT) interventions in 511
- 512 global health. Glob Health. 2018 Dec;14(1):100.
- 513 24. De Rooij D, Belfroid E, Hadjichristodoulou C, Mouchtouri VA, Raab J, Timen A, et al.
- 514 Educating, training, and exercising for infectious disease control with emphasis on cross-border
- 515 settings: an integrative review. Glob Health. 2020 Dec;16(1):78.
- 516 25. de Sousa A, Tiedje KE, Recht J, Bjelic I, Hamer DH. Community case management of childhood
- illnesses: policy and implementation in Countdown to 2015 countries. Bull World Health Organ. 517
- 2012 Mar 1;90(3):183-90. 518
- 519 26. Olaniran A, Smith H, Unkels R, Bar-Zeev S, van den Broek N. Who is a community health
- 520 worker? - a systematic review of definitions. Glob Health Action. 2017;10(1):1272223.
- 521 27. Hiwat H, Martínez-López B, Cairo H, Hardjopawiro L, Boerleider A, Duarte EC, et al. Malaria
- 522 epidemiology in Suriname from 2000 to 2016: trends, opportunities and challenges for
- 523 elimination. Malar J. 2018 Nov 12;17(1):418.
- 524 28. Knowles MS. Andragogy: Adult Learning Theory in Perspective. Community Coll Rev. 1978
- 525 Jan;5(3):9–20.
- 526 29. Bloom, B. S. Taxonomy of educational objectives: The classification of educational goals. 1st ed.
- Longman Group.; 1956. 527
- 30. Krathwohl DR. A Revision of Bloom's Taxonomy: An Overview. Theory Pract. 2002 Nov 528
- 529 1;41(4):212-8.

- 530 31. Barbe WB, Swassing RH, Milone MN. Teaching through modality strengths: concepts and
- 531 practices. Columbus, Ohio: Zaner-Bloser; 1979. 117 p.
- 532 32. and N center for chronic disease prevention, health promotion, division of population health.
- CDC. Understanding the training of trainers model. Available from: 533
- 534 https://www.cdc.gov/healthyschools/professional development/documents/17 279600 Trainers
- 535 Model-FactSheet v3 508Final.pdf
- 33. 14:00-17:00. ISO. [cited 2023 Sep 12]. ISO 8402:1994. Available from: 536
- 537 https://www.iso.org/standard/20115.html
- 538 34. Kirkpatrick DL, Kirkpatrick JD. Evaluating training programs: the four levels. 3rd ed. San
- 539 Francisco, CA: Berrett-Koehler; 2006. 372 p.
- 35. Article R1121-1 Code de la santé publique Légifrance [Internet]. [cited 2023 Sep 12]. 540
- 541 Available from: https://www.legifrance.gouv.fr/codes/article_lc/LEGIARTI000043723460
- 542 36. Global Malaria Programme WHO. Zeroing in on malaria elimination: Final report of the E-2020
- 543 initiative [Internet]. Geneva: World Health Organization; 2021. Available from:
- 544 https://www.who.int/publications/i/item/9789240024359
- 545 37. Ministério da Saúde. Secretaria de Vigilância em Saúde. Departamento de Imunização e Doenças
- 546 Transmissíveis. Elimina Malária Brasil: Plano Nacional de Eliminação da Malária. Brasília:
- 547 Ministério da Saúde; 2022.
- 548 38. Galindo MS, Lambert Y, Mutricy L, Garancher L, Miller JB, Gomes JH, et al. Implementation of
- 549 a novel malaria management strategy based on self-testing and self-treatment in remote areas in
- 550 the Amazon (Malakit): confronting a-priori assumptions with reality. BMC Public Health. 2022
- 551 Apr 15;22(1):770.
- 552 39. Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M, et al. Developing and
- 553 evaluating complex interventions: the new Medical Research Council guidance. BMJ. 2008 Sep
- 554 29;337:a1655.
- 555 40. Sanna A, Suarez Mutis M, Lambert Y, Carvalho L, Cairo H, Cox H, et al. Cooperation for
- 556 malaria control and elimination in the Guiana Shield – Many hands make light work. 2023.
- 557 (under publication).
- 558 41. Bazeyo W, Bagonza J, Halage A, Okure G, Mugagga M, Musoke R, et al. Ebola a reality of
- 559 modern Public Health; need for Surveillance, Preparedness and Response Training for Health
- 560 Workers and other multidisciplinary teams: a case for Uganda. Pan Afr Med J. 2015 Apr
- 561 23;20:404.
- 562 42. Peters DH, Tran NT, Adam T. Implementation research in health: a practical guide. Alliance for
- 563 Health Policy and Systems Research, World Health Organization; 2013.
- 564 43. Carroll C, Patterson M, Wood S, Booth A, Rick J, Balain S. A conceptual framework for
- 565 implementation fidelity. Implement Sci. 2007 Nov 30;2(1):40.

- 566 44. von Thiele Schwarz U, Giannotta F, Neher M, Zetterlund J, Hasson H. Professionals'
- 567 management of the fidelity-adaptation dilemma in the use of evidence-based interventions—an
- 568 intervention study. Implement Sci Commun. 2021 Mar 16;2(1):31.
- 569 45. Miller CJ, Wiltsey-Stirman S, Baumann AA. Iterative Decision-making for Evaluation of
- 570 Adaptations (IDEA): A decision tree for balancing adaptation, fidelity, and intervention impact. J
- 571 Community Psychol. 2020 May;48(4):1163–77.
- 572 46. Holtrop JS, Gurfinkel D, Nederveld A, Phimphasone-Brady P, Hosokawa P, Rubinson C, et al.
- Methods for capturing and analyzing adaptations: implications for implementation research. 573
- 574 Implement Sci. 2022 Jul 29;17(1):51.
- 575 47. Stirman SW, Miller CJ, Toder K, Calloway A. Development of a framework and coding system
- 576 for modifications and adaptations of evidence-based interventions. Implement Sci. 2013 Jun
- 577 10;8(1):65.
- 48. World Health Organization (WHO). What do we know about community health workers? A 578
- 579 systematic review of existing reviews. 2020. (Human Resources for Health Observer Series).
- 580 49. Rega PP, Fink BN. Immersive simulation education: a novel approach to pandemic preparedness
- 581 and response. Public Health Nurs Boston Mass. 2014;31(2):167–74.
- 582 50. Bosch-Capblanch X, Marceau C. Training, supervision and quality of care in selected integrated
- 583 community case management (iCCM) programmes: A scoping review of programmatic evidence.
- 584 J Glob Health. 2014 Dec;4(2):020403.
- 585 51. Dambisya YM, Matinhure S. Policy and programmatic implications of task shifting in Uganda: a
- 586 case study. BMC Health Serv Res. 2012 Mar 12;12:61.
- 587 52. Peltzer K, Tabane C, Matseke G, Simbayi L. Lay counsellor-based risk reduction intervention
- 588 with HIV positive diagnosed patients at public HIV counselling and testing sites in Mpumalanga,
- 589 South Africa. Eval Program Plann. 2010 Nov;33(4):379-85.
- 590 53. Abdel-All M, Putica B, Praveen D, Abimbola S, Joshi R. Effectiveness of community health
- 591 worker training programmes for cardiovascular disease management in low-income and middle-
- 592 income countries: a systematic review. BMJ Open. 2017 Nov 3;7(11):e015529.
- 593 54. Adams LB, Richmond J, Watson SN, Cené CW, Urrutia R, Ataga O, et al. Community Health
- 594 Worker Training Curricula and Intervention Outcomes in African American and Latinx
- 595 Communities: A Systematic Review. Health Educ Behav Off Publ Soc Public Health Educ. 2021
- 596 Aug;48(4):516–31.
- 55. O'Brien MJ, Squires AP, Bixby RA, Larson SC. Role Development of Community Health 597
- 598 Workers. Am J Prev Med. 2009 Dec;37(6 Suppl 1):S262-9.
- 599 56. Lee CN, Matthew RA, Orpinas P. Design, implementation, and evaluation of community health
- 600 worker training programs in Latinx communities: A scoping review. J Community Psychol. 2023
- 601 Jan;51(1):382-405.

- 602 57. Schleiff MJ, Aitken I, Alam MA, Damtew ZA, Perry HB. Community health workers at the
- 603 dawn of a new era: 6. Recruitment, training, and continuing education. Health Res Policy Syst.
- 604 2021 Oct 12;19(3):113.
- 605 58. Takasugi T, Lee ACK. Why do community health workers volunteer? A qualitative study in 606 Kenya. Public Health. 2012 Oct; 126(10):839-45.
- 607 59. Kok MC, Muula AS. Motivation and job satisfaction of Health Surveillance Assistants in 608 Mwanza, Malawi: an explorative study. Malawi Med J. 2013 Mar;25(1):5–11.
- 609 60. Martinez R, Vivancos R, Visschers B, Namatovu L, Nyangoma E, Walley J. Training needs, 610 practices and barriers in the work of community reproductive health workers in Masindi district,
- 611 Uganda. Trop Doct. 2008 Apr;38(2):93-5.
- 612 61. Kok MC, Kane SS, Tulloch O, Ormel H, Theobald S, Dieleman M, et al. How does context
- 613 influence performance of community health workers in low- and middle-income countries?
- Evidence from the literature. Health Res Policy Syst. 2015 Mar 7;13(1):13. 614
- 62. Curtale F, Siwakoti B, Lagrosa C, LaRaja M, Guerra R. Improving skills and utilization of 615 community health volunteers in Nepal. Soc Sci Med 1982. 1995 Apr;40(8):1117–25. 616
- 617 63. Puchalski Ritchie LM, van Lettow M, Barnsley J, Chan AK, Schull MJ, Martiniuk ALC, et al.
- 618 Lay Health Workers experience of a tailored knowledge translation intervention to improve job
- 619 skills and knowledge: a qualitative study in Zomba district Malawi. BMC Med Educ. 2016 Feb
- 9;16(1):54. 620
- 621 **Data Availability Statement**
- 622 Data available under reasonable request to the corresponding author.



Launch of the Prior activities to the setting up of the Curema Curema intervention Curema intervention on going in the field intervention at the inclusion sites **QUALITY AND EFFECTIVENESS EVALUATION EVALUATION STEP 2 EVALUATION STEP 1** Pre-training Post-training Observation, Observation knowledge knowledge survey, In-depth interviews Satisfaction test survey ON-THE-JOB TRAINING **INITIAL TRAINING RETRAINING (DO-CHECK-ADJUST)** TRAINING PROGRAM **SUPERVISION CONTINUOUS PROFESSIONAL DEVELOPMENT**

