An efficient design for whole genome trio sequencing identifies key variants in rare neurological disorder cases

K. Ramsey², S. Kruglyak¹ M. Naymik², B. R. Lajoie¹, K. N. Wiseman¹, M. Sanchez-Castillo², S. Billings¹, W. Jepsen², M. Huentelman², V. Narayanan²; ¹Element Biosciences, San Diego, CA, ²TGen, Phoenix, AZ

Abstract

We sequenced nine trios in which the probands in an underserved population were affected by a rare and undiagnosed disorder with neurological features. Sequencing was performed with one trio per flowcell on a benchtop sequencing instrument, leveraging the design of sequencing the proband at twice the coverage of the parents. The reduced coverage in the parents led to sequencing efficiencies while retaining the benefits of trio sequencing: the ability to discover de novo variants and the ability to trace inheritance patterns of rare variants. Once the sequencing data was generated, our two teams used independent informatics pipelines for variant calling and interpretation. In five of the nine cases, both teams found a single SNV or small indel that was deemed causal pending clinical validation. In three of the nine cases, neither team had a significant finding. In the final case, an additional scan for large CNVs performed by one of the teams identified a de novo deletion and duplication in the proband which is the likely cause of the underlying disease. The results across cases with significant findings showed a variety of affected genes (CFAP52, DYNC1H1, FANCE, TCF4, and TOP3A), variant types, and inheritance patterns. All findings were clinically validated, after which the families were counseled about disease management, current research studies (i.e. gene therapy), and family planning. With six of nine families receiving findings, the study demonstrated an efficient and effective trio sequencing design strategy.

Introduction

Whole genome sequencing is a powerful tool for discovering causal variants in rare disease studies^{1–3}. In previous work, whole genome sequencing by avidity effectively identified causal variants in approximately half of the genomes sequenced in cases of rare eye disease⁴. The work primarily focused on single genomes and all findings involved inherited SNPs or small indels. In this study, we sequenced nine trios consisting of a proband affected by a rare, undiagnosed neurological disorder and both parents to enable identification of *de novo* variants, which are often the cause of genetic neurological disorders⁵. We chose to sequence the proband at a typical coverage level of approximately 35-40X and the parents at half that coverage (~15-20X). The design is efficient in cost and time because the three genomes can be sequenced in a single flowcell of the AVITI⁶ benchtop sequencing instrument. We hypothesized that we would be able to observe inheritance patterns and identify *de novo* variants even with the reduced coverage in the parents.

The families for the study were selected from an underserved population and enrolled at TGen's Center for Rare Childhood Disorders. The families had little to no access to previous genetic testing. Enrollment in this study provided a means of finding the underlying cause of their child's disorder and hope in ending the diagnostic odyssey. The probands' symptoms included global developmental delay, intellectual disability (ID), seizures, hypotonia, microcephaly, situs inversus, facial dysmorphism, thrombocytopenia, small stature, and cardiac malformations (see Table 1 for a complete list of phenotypes).

Results

Extracted DNA was used to make PCR-free sequencing libraries for the 27 samples (9 affected probands and their parents). The libraries were pooled by trio, with each proband having twice the concentration as the parents. Following whole genome sequencing and FASTQ generation, the data was independently analyzed by the teams at Element Biosciences and at TGen. We wanted to determine whether teams working independently and with different informatics pipelines would arrive at the same interpretations for each of the cases. One team used a combination of DNAScope (Sentieon, San Jose California) and Franklin (Genoox, Tel Aviv, Israel) for secondary and tertiary analysis. The other team leveraged Varseq (Golden Helix, Inc., Bozeman, MT) and Emedgene (Illumina, San Diego, CA) tools. Once interpretation was completed, final results were shared and discussed. In five of the cases, the two teams identified the same variant as causal. In 3 of the cases, neither team identified a causal variant. In the final case, one team identified a large CNV as the likely cause of the disorder, while the other team did not have CNV interpretation implemented within their pipeline. Following data review, both teams agreed on the six findings and samples were sent for clinical validation either via Sanger sequencing in a CLIA lab (for SNPs and small indels) or for CMA in a CLIA lab for the large CNV (GeneDx, Gaithersburg, MD). All variants were confirmed, at which point results could be shared with the treating clinicians and family counseling could begin. Table 1 summarizes the findings.

Table 1: Findings across all cases. Coordinates are provided with respect to the hg19 / GRCh37 reference.

ID	Gene	Variant	Inheritance	Phenotypes	TGen Recommendations		
		c.4580delinsCC,		Global developmental delay, ID, Seizures,	Referred family to support group, research,		
1	DYNC1H1	p.L1527Pfs*9	Het <i>de novo</i>	Cerebral hemorrhage	discussed recurrence risk		
					Referred family to the support group and		
		c.923-2 A>G,		Global developmental delay, Nonverbal,	registry at the Pitt Hopkins Research		
		splice acceptor		Hypotonia, Short stature, Happy	Foundation, gene therapy and therapeutic		
2	TCF4	variant	Het <i>de novo</i>	demeanor	drug research, discussed recurrence risk		
		Terminal deletion					
		1.45 Mb Xp22.33		Developmental delay, Microcephaly,	Provided resources from Unique,		
		and terminal		Hypotonia, Hypoplasia of corpus	recommendations of FISH studies in at least		
	ChrX	duplication 15.1		callosum, High pain tolerance, Seizures,	mother but also proband's siblings to		
	terminal	Mb Xq27.1 to		Hypothyroidism, Facial dysmorphism,	determine if mother has a pericentric		
a3	del/dup	Xq28	Het de novo	Cryptorchidism and Micropenis	inversion		
				Global developmental delay, Hearing			
		c.1162del	Homo	impairment, Dextrocardia, Transposition	Recommendations for creating social media		
4	CFAP52	p.S388Afs*21	recessive	of the great arteries, Situs inversus totalis	support group, family planning		
				Global developmental delay,			
				Microcephaly, Short stature, Decreased	Recommended sister chromatid exchange		
				body weight, Periventricular	studies, follow-up ECHO, discussion of		
		c.1685 G>A,	Homo	leukomalacia, Delayed gross motor	"mitochondrial cocktail" with her healthcare		
5	TOP3A	p.G562E	recessive	development	team, discussed recurrence risk		
				Thrombocytopenia, VSD, Single kidney,			
				Unilateral deafness, R Thumb	Referral to the Fanconi Anemia Research		
				abnormality, Short stature, Triangular	Fund and Family Support Group, discussion		
		c.1111 C>T,	Homo	face, Narrow nasal bridge, Learning	of hematopoietic cell transplant, cancer		
6	FANCE	p.R371W	recessive	delays	surveillance, family planning		
					Counseled mother about risk for HBOC,		
					recommendations for increased screening,		
					risk reducing surgery, testing of family		
					members; mother does not meet NCCN		
		c.2257 C>T,		Incidental Finding: Risk for Hereditary	criteria for testing based on personal and		
4	PALB2	p.R753*	Het maternal	Breast and Ovarian Cancer (HBOC)	family history		

In three cases (Families 1, 2, and 3), *de novo* variants were determined to be causal. One of the variants was an indel in the *DYNC1H1* gene and the other was a splice acceptor variant in *TCF4*. Figure 1 shows the IGV displays for the two calls and demonstrates how the trio design was leveraged to confirm that neither variant is inherited.

A	102.468,875 bp 102.468,880 bp 102.468,885 bp 102.468,890 bp 102.468,895 bp 102.468,900 bp 102.468,905 bp 102.468,910 bp 102.468,915 bp 102.468,920 bp 102.468,925 bp 102.468,930 bp 102.468,930 bp 102.468,940 bp 102.468,945 bp 102.468,945 bp 102.468,940 bp 102.468,945 bp 102.46
	ле отттттолло олго ото то то то то ло о то ло лтоло о то ло олго лто лто лто то то то то то то то то то лто то
52	
0_	Proband (Family 1)
26	
0_	Mother (Family 1)
20 0_	Father (Family 1)
в	
	2 C A O T C T O C O O C T O C C O C T T C C T C T
80 ⁻	Proband (Family 2)
35 ⁻ 0_	Mother (Family 2)
51 0_	
	Father (Family 2)

Figure 1: (A) De novo indel in DYNC1H1 gene. The proband is in the top display and the parents are below. The reference T is replaced by a CC, causing a heterozygous frameshift. The data from the parents shows no evidence of the variant, confirming that the mutation occurred de novo. (B) Splice altering de novo variant in TCF4. The top display in the proband confirms the heterozygous variant, while the maternal and paternal displays below show no evidence for the variant in the parents, confirming the de novo call.

In one of the cases (Family 3), a del/dup event in chromosome X (chrX) was discovered. WGS did not detect a chromosomal rearrangement or del/dup on chrX in the mother. Figure 2 shows the structural variants in chrX in the trio.

A									12:00					78,en
607	Proband (Fami	ily 3)	KA TERRETARI AND KA PENDARI KA PENDARI	10108-0010-0-000 8-08-0900-0-0-0		I NUMBER	III I	10.041 (-0.041-01) - 1.041 (000) 10.011 - 1.043 (01) - 1.041 (000) 10.011 - 1.043 (01) - 1.041 (01)		анинно наста нинно сова а 	10000-1 - 10000 - 10000 100	I I IIII CHANGON (IIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIII		
0	Mother (Family	v 3)	eliterationen en time.	ter anitality and	and a state of the second	a Mariana	nezotaten erantzi deriotzera	en förstalla som en konstanten som en so	-	i i i i i i i i i i i i i i i i i i i i		. I	-	an a
0	Father (Family	3)	auraniaturinyaaniyya	inter of the laget of the contract of the laget		nd and franciscon	nyados yoo ta'aa maaanaa ay ahaa ahaa	alay to a straight of the state	หน่งของแปลใจอากุกรับกับอย่าง	z z z	n na star star star star star star star sta	senso ann air an		erendenstelsensenstelseren.
B	Breakend::	chrX	:1,506	,467/ch	nrX:140	,707,484	1							
	Sample Proband Mother Father Total	Alt 28 0 0 28	Ref 129 135 83 347	Amb 195 95 103 393										
	Alternate All Proband (Family	ele / 3)												
						_				-		_		
	Mother (Family 3	3) 3)					No read	ls found						
							No read	ls found						

Figure 2: (A) Chromosome X coverage in the proband (top) followed by the mother and father. The proband has a deletion in the beginning of the chromosome marked in green. This is part of the PAR region so a copy number of 2 is expected in males, as shown in the father, but the proband has a stretch of copy number 1. The proband also has a large duplication at the end of the chromosome. (B) SVVIZ⁷ display of the junction between the duplication and an early part of the chromosome. The proband shows extensive read support for the junction, while the parents show no support. This leads to the conclusion that the event is de novo.

In three cases (Families 4, 5, and 6), the primary finding was a small homozygous variant in the proband (in genes *CFAP52, TOP3A*, and *FANCE*, respectively). Surprisingly, each allele was extremely rare in the population, yet the proband inherited a copy from each of the unrelated parents. Intuition would suggest that a different rare allele would be inherited from each parent. Table 2 provides the gnomAD allele frequencies and the relatedness coefficients⁸ for each case. The rare nature of these variants and the cooccurrence in a single family may imply the presence of a founder mutation or a 'mutation hotspot.'

Gene	Variant	gnomAD het	Parent relatedness		
		frequency	coefficient		
CFAP52	c.1162del p.S388Afs*21	0.00006572	2.4%		
ТОРЗА	c.1685 G>A, p.G562E	0	2.3%		
FANCE	c.1111 C>T, p.R371W	0.00003288	-2.0%		

Table 2: Allele frequencies and relatedness coefficients for the three homozygous variants. Surprisingly, each parent is a carrier for the same rare variant. Coordinates are provided with respect to the hg19 / GRCh37 reference.

Discussion

In this study, we used a novel trio sequencing design to solve six of nine rare neurological disorder cases. Each trio was sequenced on a single flowcell of a benchtop sequencer, obviating the need for batching large numbers of samples. The design of sequencing parents at half the typical depth was effective in identifying *de novo* variants. The design also decreased manual interpretation effort (a costly step of the entire process) by dramatically shortening the list of variants prioritized by the automated machine learning algorithms. This is because rare heterozygous variants inherited from a healthy parent could be ruled out.

Both families 1 and 2 were counseled of a less than 1% recurrence risk (due to germline mosaicism). Both families were informed of online social media groups for the disorders, and Family 2 was directed to the Pitt Hopkins Research Foundation (<u>https://pitthopkins.org/</u>), which has support for families, a registry, and a list of currently funded research, including gene therapy and repurposing of drug compounds. For Families 4, 5, and 6, additional counseling was provided regarding recurrence risk and family planning for autosomal recessive disorders. Two of the families were interested in having additional children and were counseled on the recurrence risk of 25%, as well as the availability of PGD testing, sperm or egg donation, and adoption.

In Family 3, WGS identified a del/dup of chromosome X in the proband. Similar del/dups of chromosome X have been seen in individuals after a meiotic recombination event involving a phenotypically normal mother with a pericentric inversion on the X chromosome ^{9,10}. However, a thorough analysis of both maternal and paternal WGS shows no indication of an inversion, suggesting the event is *de novo* in the proband. Confirmation studies using FISH would help support this finding, given other unexplained conditions in the pedigree (Figure 3).



Figure 3: Pedigree of family 3. We had suspected a pathogenic variant segregating in the pedigree because multiple individuals are affected. However, we found the del / dup to be de novo and the conditions in the pedigree are highly varied.

However, at this time, it has not been possible to coordinate additional testing in the mother or siblings. The duplicated region contains more than 130 genes, including the *MECP2* gene. Duplications of 0.3 Mb and larger in *MECP2* are associated with *MECP2* duplication syndrome, causing early-onset hypotonia, ID, autistic features, seizures, spasticity, and lack of speech (Van Esch, 2008). The deleted region of the X chromosome contains ten genes, including *SHOX* and *CSF2RA*. The phenotype of *SHOX* deficiency disorders includes short stature, limb anomalies, including mesomelia and Madelung deformity¹¹. Variants in the *CSF2RA* gene have been associated with pulmonary alveolar proteinosis ¹². The proband's phenotype overlaps with individuals with *MECP2* duplication syndrome as well as other individuals with similar chromosome X del/dup ^{9,10,13}.

The proband in Family 5 was found to have a homozygous variant in *TOP3A* associated with a Bloom syndrome-like disorder ¹⁴. Unlike Bloom syndrome, individuals with *TOP3A* variants have shown evidence of mitochondrial dysfunction and dilated cardiomyopathy (sometimes fatal in childhood). Our patient had a normal echocardiogram at birth and is no longer being followed by cardiology. We recommended that since our patient was now older, she receive a follow-up echocardiogram. Based on the nature of the disorder, we also suggested the patient discuss a trial of "mitochondrial cocktail" (supplements including vitamins, Coenzyme Q10, and L-carnitine) with her healthcare providers.

The proband for Family 6 presented to our center with a recent diagnosis of thrombocytopenia and a history of short stature, cardiac anomaly, a single kidney, unilateral deafness, and a smaller and inflexible right thumb. The geneticist suspected Fanconi anemia (FA) or Russell Silver syndrome. Genetic testing was crucial in confirming the diagnosis as well as identifying the mode of inheritance—FA is associated with 21 genes and has autosomal recessive, dominant, and X-linked modes of inheritance ¹⁵. Beyond the traditional suggested cancer screening for FA, there is additional cancer surveillance recommended for individuals with mutations in genes associated with hereditary breast and ovarian cancer (HBOC), making a genetic diagnosis crucial for management. The proband was found to have a mutation in *FANCE*, a gene that accounts for only 3% of FA. The family was counseled about treatments to improve blood counts and the use of hematopoietic stem cell transplantation. The family was also referred to the Fanconi Anemia Research Fund for support, registry information, and current research and clinical trials (https://www.fanconi.org/).

In Family 4, a pathogenic variant in the PALB2 gene was found in both the proband and his mother. They had consented to receiving incidental findings. The family was provided counseling for an increased risk of HBOC and the mother was encouraged to notify her health care providers to establish increased screening and discuss surgical recommendations. The proband's mother has no personal or family history of cancer and would not have met NCCN criteria for testing¹⁶. This case demonstrates the importance of universal screening for HBOC.

Conclusion

We implemented an efficient trio sequencing design for the study of rare disorders and used it to find key variants in 6 of 9 cases. The findings included multiple variant types, including SNPs, indels, and structural variants. In three solved cases, the variants in the proband arose de novo, whereas in the other three cases, they were inherited. In addition to the primary findings, there was an incidental finding: a SNP conferring cancer risk. Although the two teams performing interpretation used different informatics tools for secondary and tertiary analysis, the findings matched closely. The only differences were encountered when one team used methods (e.g. CNV detection) that were not implemented by the other team. However, whenever each team had a finding for a case, the finding matched. All findings were clinically confirmed, and results were then returned. While having a diagnosis was already a positive result for the families in ending the diagnostic odyssey and suggesting support groups for the identified disorders, in some cases additional actions pertaining to family planning or clinical management were taken. For example, in the case of *de novo* variants, the families were told that recurrence risk in subsequent pregnancies would be minimal. By contrast, the risk in the cases where both parents were determined to be carriers for recessive disorders, the recurrence risk is 25%. Furthermore, certain findings suggested additional tests (e.g. ECHO cardiogram in the TOP3A case) or evaluation of potential gene therapy trials (e.g. the TCF4 case). We believe that the trio sequencing design described in this article could easily be scaled to more cases and effectively used in the study of rare genetic disorders.

Methods

Patient recruitment

Five male and four female probands, in the age range from 1-20 years, were either self-referrals to the Center for Rare Childhood Disorders or were referred by their neurologist or geneticist. Written informed consent for sequencing, data usage, and publication of clinical details was obtained from participants, legally authorized representatives and/or their guardians if under the age of 18 for all of the family members. The study protocol and consent procedure were approved by the WGC Institutional Review Board (WGC IRB; study number 20120789).

Whole blood was obtained for all participants except for one proband, where a buccal swab was obtained.

Library Preparation

Libraries were prepared PCR-free using 500ng of input gDNA using the KAPA HyperPrep kit. The libraries were sheared on Covaris ME220 to ~350bp average fragment size. KAPA UDI adapters were ligated and the libraries were size-selected with a dual-sided SPRI (0.48X/0.64X). All PCR-free libraries were circularized with the Element Adept Library Compatibility Kit v1.1 (cat# 830-00007). Following qPCR, the libraries were normalized to 1nM and pooled by volume as follows: proband 50%, mother 25%, and father 25%.

Sequencing

Pooled libraries were shipped to FYR Diagnostics (Missoula, MT) for sequencing with 2x150 bp reads and a trio per flowcell on an Element Biosciences AVITI sequencer. Table 3 summarizes the coverage and quality score statistics for each trio. FASTQ files were shared with the Element Biosciences team and the TGen team.

Sample Identifier	Data generated (GB)	%Q30
F1_1	130	95.3
F1_2	62	95.7
F1_3	53	95.6
F2_1	166	94.2
F2_2	70	95
F2_3	78	94.7
F3_1	164	94.7
F3_2	78	94.7
F3_3	87	95
F4_1	154	95.3
F4_2	87	95.2
F4_3	79	94.9
F5_1	160	95.4
F5_2	74	95.1
F5_3	83	95.9
F6_1	159	94
F6_2	84	95.5
F6_3	83	95.2
F7_1	113	92.3
F7_2	61	93
F7_3	75	94.3
F8_1	164	94.6
F8_2	79	94.6
F8_3	87	94.6
F9_1	171	94.6
F9_2	82	94.1
F9_3	78	94.3

Table 3: Summary of data collected and quality for each of the 27 samples in the study. Number following F is the deidentified family ID and the underscore 1, 2, 3, refers to proband, mother, and father, respectively.

Element Biosciences Analysis

BWA-MEM was used to align the FASTQ files to the hg38 / GRCh38 human reference. Sentieon DNA scope was used for variant calling in each individual. VCF files and phenotypes were uploaded into the Franklin by Genoox interpretation software. Each prioritized variant was manually reviewed for quality, inheritance mode, and phenotypic fit. In five of the cases, a single variant was identified. For all cases, we also reviewed a coverage depth by chromosome plot. In one of the cases, we noticed that chromosome X had slightly higher coverage in the proband (Figure 4).



Figure 4: Normalized coverage by chromosome plot for the pedigree. The small increase in coverage in chromosome X of the proband relative to the father led to the hypothesis that a large CNV might be present.

A review in IGV revealed a large duplication. At this point, we uploaded the FASTQ files into the Franklin software and repeated analysis of all cases because Franklin analysis from FASTQ includes structural variant calling. The five prior findings were reproduced, but there were two additional findings. In the case where we had observed the duplication, several structural variants were found, including a deletion near the beginning of chromosome X and the duplication at the end of chromosome X. To determine that the del dup originated *de novo*, we entered the breakpoints into SVVIZ⁷ and observed clear evidence in the proband but no evidence in either parent. For three of the cases, there were no significant findings.

TGen Analysis

Reads were aligned to the human genome (hg19/GRCh37) using the Burrows-Wheeler Aligner (BWA mem v.0.7.8). PCR duplicates were identified using Picard MarkDuplicates v1.79. Base quality recalibration and indel realignment were performed using the Genome Analysis Toolkit (GATK v3.5-1). Variants were jointly called with HaplotypeCaller and recalibrated with GATK. Quality controls were conducted using FASTQC v0.11.5. Called variants were annotated with SnpEff v3.0a against Ensembl GRCh37.66 and filtered against dbSNP137 1000 Genomes Project (minor allele frequency < 0.05), SnpEff Impact: High + Moderate, GATK quality score > 40, and known genes. Final annotation and reports were generated with Varseq (Golden Helix, Inc., Bozeman, MT) and Emedgene (Illumina, San Diego, CA). Relatedness coefficients used in Table 2 were part of the Emedgene analysis output.

Review of interpretation results

Once both teams completed interpretation of the cases, we held a conference call to discuss the results. For five of the cases with findings, both teams identified the same variant. For three of the cases, neither team had a compelling finding. In the case of the del dup on chromosome X, only one team ran a structural variant caller and saw the event. However, during the meeting,

the teams reviewed the findings in IGV, noted the genes in the affected region, and agreed that this was likely causal, pending clinical validation.

Clinical validation

DNA samples were sent for clinical validation in a CLIA lab either via Sanger for SNPs and small indels or for chromosomal microarray in a CLIA lab for the large CNVs (GeneDx, Gaithersburg, MD).

References

- 1. NIHR BioResource—Rare Disease, Next Generation Children Project, French CE, et al. Whole genome sequencing reveals that genetic conditions are frequent in intensively ill children. *Intensive Care Med*. 2019;45(5):627-636. doi:10.1007/s00134-019-05552-x
- Sanford EF, Clark MM, Farnaes L, et al. Rapid Whole Genome Sequencing Has Clinical Utility in Children in the PICU*: *Pediatr Crit Care Med*. 2019;20(11):1007-1020. doi:10.1097/PCC.00000000002056
- 3. Turro E, Astle WJ, Megy K, et al. Whole-genome sequencing of patients with rare diseases in a national health system. *Nature*. 2020;583(7814):96-102. doi:10.1038/s41586-020-2434-2
- 4. Pooja Biswas, Adda Villanueva, Benjamin J. Krajacich, et al. Avidity sequencing of whole genomes from retinal degeneration pedigrees identifies causal variants. *medRxiv*. Published online January 1, 2022:2022.12.27.22283803. doi:10.1101/2022.12.27.22283803
- 5. Brunet T, Jech R, Brugger M, et al. *De novo* variants in neurodevelopmental disorders—experiences from a tertiary care center. *Clin Genet*. 2021;100(1):14-28. doi:10.1111/cge.13946
- 6. Arslan S, Garcia FJ, Guo M, et al. *Sequencing by Avidity Enables High Accuracy with Low Reagent Consumption*. Genomics; 2022. doi:10.1101/2022.11.03.514117
- 7. Spies N, Zook JM, Salit M, Sidow A. svviz: a read viewer for validating structural variants. *Bioinformatics*. 2015;31(24):3994-3996. doi:10.1093/bioinformatics/btv478
- Pedersen BS, Quinlan AR. Who's Who? Detecting and Resolving Sample Anomalies in Human DNA Sequencing Studies with Peddy. *Am J Hum Genet*. 2017;100(3):406-413. doi:10.1016/j.ajhg.2017.01.017
- 9. Kokalj-Vokac N, Marcun-Varda N, Zagorac A, et al. Subterminal deletion/duplication event in an affected male due to maternal X chromosome pericentric inversion. *Eur J Pediatr*. Published online August 12, 2004. doi:10.1007/s00431-004-1519-5
- Mohandas T, Geller RL, Yen PH, et al. Cytogenetic and molecular studies on a recombinant human X chromosome: implications for the spreading of X chromosome inactivation. *Proc Natl Acad Sci*. 1987;84(14):4954-4958. doi:10.1073/pnas.84.14.4954

- Binder G, Rappold GA. SHOX Deficiency Disorders. In: Adam MP, Mirzaa GM, Pagon RA, et al., eds. GeneReviews[®]. University of Washington, Seattle; 1993. Accessed September 28, 2023. http://www.ncbi.nlm.nih.gov/books/NBK1215/
- 12. Al-Haidary AS, Alotaibi W, Alhaider SA, Al-Saleh S. A newly identified novel variant in the CSF2RA gene in a child with pulmonary alveolar proteinosis: a case report. *J Med Case Reports*. 2017;11(1):122. doi:10.1186/s13256-017-1285-4
- 13. Magini P, Poscente M, Ferrari S, et al. Cytogenetic and molecular characterization of a recombinant X chromosome in a family with a severe neurologic phenotype and macular degeneration. *Mol Cytogenet*. 2015;8(1):58. doi:10.1186/s13039-015-0164-1
- 14. Martin CA, Sarlós K, Logan CV, et al. Mutations in TOP3A Cause a Bloom Syndrome-like Disorder. *Am J Hum Genet*. 2018;103(2):221-231. doi:10.1016/j.ajhg.2018.07.001
- Mehta PA, Ebens C. Fanconi Anemia. In: Adam MP, Mirzaa GM, Pagon RA, et al., eds. *GeneReviews®*. University of Washington, Seattle; 1993. Accessed September 28, 2023. http://www.ncbi.nlm.nih.gov/books/NBK1401/
- Bevers TB, Niell BL, Baker JL, et al. NCCN Guidelines[®] Insights: Breast Cancer Screening and Diagnosis, Version 1.2023: Featured Updates to the NCCN Guidelines. *J Natl Compr Canc Netw*. 2023;21(9):900-909. doi:10.6004/jnccn.2023.0046