It is made available under a [CC-BY 4.0 International license](http://creativecommons.org/licenses/by/4.0/) .

¹⁵ **Abstract**

 Objective: To investigate and analyze the effects of the infarct volume on cytokines and immune status in patients with acute ischemic stroke. **Methods:** Patients with acute ischemic stroke that presented within 72 h of onset from October 2017 to October 2019 were enrolled. Patients with severe cerebral infarction (large-area cerebral infarction) (n=34) were enrolled and categorized as group A; Additionally, 33 patients with non- large-area cerebral infarction with matching baseline characteristics (sex and age) to group A were included in group B. We measured IL-2, IL-4, IL-6, IL-10, IL-17A, TNF- α, and IFN-γ levels in serum using a cytometric bead array.In addition,we compared 24 the absolute value of lymphocytes $(LYM#)$, lymphocyte percentage $(LYM\%)$, NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Key words

 Acute cerebral infarction, immunity, inflammation, cytokines, inflammatory cells,infarct volume

1. **Introduction**

 According to the all-cause mortality statistics of the World Health Organization (WHO), stroke is the second leading cause of death globally, accounting for 87%[1], and its high disability rate significantly increases the social burden. Because immunity is crucial in cerebral infarction and, in turn, is regulated by it, the immune mechanism has become a target that may benefit clinical treatment and has attracted increasing interest from researchers.

 Notably, immunity is crucial in the occurrence, progression, and outcome of ischemic stroke. The basic pathology of atherosclerotic cerebral infarction is

 atherosclerosis. Various immune cells, including lymphocytes and mononuclear macrophages, are involved in the formation of atherosclerotic plaques and the rupture of unstable plaques[2,3], an important cause of atherosclerotic cerebral infarction[4]. Immunity contributes to the pathogenesis of cerebral infarction and affects prognosis[5–8]. After local cerebral infarction, astrocytes and microglia, which are involved in the release of cytokines and chemokines[9], are rapidly activated, attract immune cells from the peripheral circulation to the ischemic area, and participate in inflammatory damage to the brain tissue[10]. In addition, in the recovery from chronic cerebral infarction, the immune status affects the brain tissue microenvironment, regulates neurogenesis[11,12] and angiogenesis[13], and affects the autoimmune response to brain tissue antigens after stroke, mediating the long-term prognosis[5,6]. Notably, as a stress event, ischemic stroke regulates systemic immunity through the sympathetic nervous system and the hypothalamus-pituitary-adrenal axis, which is an important mechanism of post-stroke immunosuppression, leading to increase mortality of patients with acute cerebral infarction [7,8,16-19]. We know that the size of different infarcts also affects the treatment and prognosis of acute ischemic stroke,(PMED: 28665171)In this study, we investigate and analyze the effects of the infarct volume on cytokines and immune status in patients with acute ischemic stroke.This enables the clinicians to make informed decisions in selecting immunotherapy approaches and tailor rehabilitation strategies to optimize patient outcomes by considering different infarct sizes.

2. **Materials and methods**

2.1 Research objects and samples

Inclusion criteria:

96 BD FACS Calibur flow cytometer (BD Company, USA); low-temperature high-speed centrifuge (Heraeus Sepaeeh, Germany); low-temperature refrigerator (Haier

 Company, Qingdao, China); BD™ Cytometric Bead Array (CBA) human Th1/Th2/Th17 kit (BD Company, USA); BD™ Cytometric Bead Array (CBA) Human Soluble Protein Master Buffer kit, (BD Company, USA).

2.3 Determination of cytokines in serum samples by CBA

method

 The BD™ Cytometric Bead Array (CBA) human Th1/Th2/Th17 kit was used to determine the cytokine (IL-2, IL-6, IL-4, IL-10, IL-17A, IFN-γ, TNF-α) levels in each serum sample according to the manufacturer's instructions.

2.4 Experimental group

 A total of 67 patients presenting with an onset time of 72 h were included in this study;patients with large-area cerebral infarction were included in the large-area cerebral infarction group and categorized as group A. A total of 34 patients, including patients with anterior circulation infarction and posterior circulation infarction, were included. For anterior circulation infarction, the maximum diameter of the infarct was >3 cm (25 participants), and regarding posterior circulation infarction, the maximum infarction cross-section was >1/3 of the brainstem (9 participants).

 Additionally, 33 patients with non-large-area cerebral infarction with matching sex and age with Group A were included in the non-large-area cerebral infarction group and categorized as Group B, including patients with anterior circulation infarction and posterior circulation infarction. For anterior circulation infarctions, the largest infarct diameter was <3 cm (27 participants), and for posterior circulation infarctions, the largest brainstem infarct area was <1/3 of the cross-sectional area (6 participants).

2.5 Statistical analysis and graph

Statistical analysis was performed using IBM SPSS Statistics version 25.0 (IBM

 Corp., Armonk, N.Y., USA), and GraphPad Prism version 7 (GraphPad Software, Inc., La Jolla, CA, USA) was used for graphing. The counted data were presented as the number of cases (n) and percentages (%). Measurement data with normal distribution 125 are presented as $\bar{x} \pm s$ and measurement data that are not normally distributed are presented as the median and quartile M (Q25–Q75). For counting data, the Chi-square 127 test $(\kappa 2 \text{ test})$ was used to compare groups. For the comparison between the two groups of measurement data, the t-test was used for normally distributed data, and the nonparametric rank-sum test (Mann–Whitney U test) was used for non-normal distribution. For the comparison between the three groups of measurement data, a one- way analysis of variance (ANOVA) was used for the normal distribution, and the variance was uniform. The LSD-t test was used for the pairwise comparison of the mean between the groups.

134 The groups showed statistically significant differences, with a P-value of ≤ 0.05 indicating statistical significance; for pairwise comparison of the mean between the groups, the statistical difference was based on the unadjusted P-value <0.017 and the adjusted P-value <0.05.

3. Results

 We compared the expression levels of various cytokines, including IL-2, IL-4, IL- 6, IL-10, and IL17A, in the large-area infarction group (group A) and the non-large- area infarction group (group B) based on the same baseline level. Furthermore, we compared the absolute lymphocyte value (LYM#), lymphocyte ratio lymphocyte percentage (LYM%), neutrophil/lymphocyte ratio (NLR), IFN-γ/IL-4 ratio, TNF-α/IL-4 ratio, and TNF-α/IL-10 ratio between groups A and B.

 The results indicated the following: (1) The cytokine levels in group A were significantly higher than in group B, as shown in Table 2 and Fig 1. These differences

 were considered statistically significant; P< 0.05. (2) Group A exhibited significantly lower LYM# and LYM% than group B, and the NLR was significantly higher. These findings are presented in Table 3 and Fig 2 and showed statistically significant 150 differences. (3) The TNF- α /IL-4 ratio in group A was significantly lower than in group B, which showed a statistically significant difference. The TNF-α/IL-10 ratio and IFN- γ /IL-4 ratio in group A demonstrated a decreasing trend but showed no statistically significant differences (Table 4 and Fig 3).

- 154
-

155 **Table 1. Baseline data between patients in group A and group B.**

Baseline	Group A	Group B	P-value
Sex, Male $(n, %)$	19 (55.88%)	24 (72.7%)	0.151
Female (n, %)	$15(44.12\%)$	$9(27.3\%)$	
Hypertension $(n, %)$	23 (67.65%)	26 (78.79%)	0.304
Diabetes $(n, %)$	$10(29.42\%)$	$10(30.30\%)$	0.936
Smoking history (n, %)	14(41.17%)	13 (39.39%)	0.882
Age (years)	$63.5(57-69)$	$62.5(56.25-69)$	0.768
$LDL-C$ (mmol/L)	$3.06(2.72 - 4.34)$	$2.9(2.29-3.86)$	0.075
Hcy (µmol/L)	$14.45(10.93-19.65)$	$15.3(13.4 - 20.8)$	0.166
Fasting blood	$5.84(5.23 - 7.93)$	$5.71(4.96 - 6.61)$	0.349
glucose (mmol/L)			
NIHSS score	$13(11-16.25)$	$4(3-8)$	0.000

156

157

158 **Table 2. Serum cytokine levels of patients of group A and group B.**

It is made available under a [CC-BY 4.0 International license](http://creativecommons.org/licenses/by/4.0/) .

Cytokine level	Group A	Group B	U-value	P-value
IL-6 (pg/mL)	$32.35(9.69 - 56.60)$	$9.81(3.09 - 28.60)$	320.00	0.003
IL-4 (pg/mL)	$4.02(3.45 - 4.76)$	$0.58(0.31-0.80)$	27.00	0.000
IL-2 (pg/mL)	$1.96(1.75-2.19)$	$0.55(0.32 - 0.98)$	84.50	0.000
IL-17A (pg/mL)	$8.15(6.30-9.66)$	$3.25(1.37 - 5.08)$	142.00	0.000
IFN- γ (pg/mL)	$2.58(2.30-2.99)$	$0.61(0.22 - 0.95)$	13.00	0.000
$TNFa$ (pg/mL)	$2.63(2.23-3.15)$	$1.71(0.80 - 2.82)$	284.00	0.001
IL-10 (pg/mL)	$3.43(2.66-4.52)$	$1.31(0.89 - 2.46)$	169.00	0.000

¹⁵⁹ Mann–Whitney U test was used. The cytokine levels in group A were significantly

- 161
- 162
-

163 **Fig 1. Serum cytokine levels of groups A and group B patients.**

- 164 ** P<0.01.
- 165
-

166 **Table 3. LYM#, LYM%, and NLR between group A and group B.**

167 Mann–Whitney U test was used. The LYM# and LYM% of the large-area group (group

168 A) were significantly lower than those of the non-large-area group (group B), while the

169 NLR of group A was significantly higher than that of group B.

170

171

¹⁶⁰ higher than in group B.

It is made available under a [CC-BY 4.0 International license](http://creativecommons.org/licenses/by/4.0/) .

Fig 2. LYM#, LYM%, NLR of group A and group B.

$$
173 \quad * \quad P < 0.01
$$

Table 4. Indicators of immune status of patients in group A and group B.

Mann–Whitney U test was used. The TNF-α/IL-4 ratio in group A was significantly

 lower than in group B; the TNF-α/IL-10 and IFN-γ/IL-4 ratios in group A showed a downward trend.

- **Fig 3. Immune status of patients in group A and group B.**
- 181 $** P< 0.01$.

4. Discussion

 Ischemic stroke is a common cerebrovascular disease; it causes irreversible brain tissue damage. The complex pathological mechanism includes excitatory glutamate toxicity, calcium overload, oxidative stress, and inflammatory damage.

 Inflammatory reaction is crucial in brain tissue injury following infarction. The immune state regulates local pathophysiological changes and significantly affects the outcome of ischemic tissues. The proinflammatory state aggravates local injury; however, the anti-inflammatory state protects the local brain tissue, promotes the removal of necrotic cells, and prevents further damage. Inflammation of the central nervous system and peripheral systemic immunity are closely linked and reflect the

 delicate and ingenious integrity of the human body. For example, the blood-brain barrier (BBB) is compromised after cerebral infarction and peripheral immune cells invade the brain tissue. The Th cells that invade the central nervous system secrete several cytokines and interact with microglia to initiate microglial polarization [14,15]; this affects the degree and outcome of the local brain tissue damage.

 The relationship between the nervous and immune systems is such that the immune system affects the pathophysiological changes of cerebral infarction and is, in turn, affected by cerebral infarction events, such as immunosuppression after stroke. The underlying mechanism is not fully understood.Post-stroke immunosuppression is a self-protective mechanism for suppressing central inflammation; however, it alters peripheral immune status and further affects the prognosis in patients with acute cerebral infarction[7–8].

 Considering the role of immune factors in the course of cerebral infarction, investigating the immune mechanisms has increasingly attracted the interest of researchers. The body's immunity is affected by several factors, and some related factors can be controlled artificially; therefore, the immune mechanism may be a potential clinical treatment target. In this study, we evaluated the levels of various 210 cytokines, including IL-2, IL-6, TNF- α , IFN- γ , IL-4, IL-10, and IL-17A in patients presenting with acute cerebral infarction within 72 h of onset, and calculated the INF- γ /IL-4, TNF- α /IL-4, and TNF- α /IL-10 ratios. The IFN- γ /IL-4 ratio indicates the balance between the subpopulations of helper T cells, reflecting the relative function between the Th1 and Th2 subpopulations. IFN-γ is an essential proinflammatory cytokine secreted by Th1-type cells, while IL-4 is a relatively specific cytokine of Th2-type cells and exerts an anti-inflammatory effect. The initial Th cells, Th0 cells, are regulated by several factors and differentiated into subpopulations. The most significant factor is the

 type and level of cytokines and the balance between each cytokine[20]. Therefore, the IFN-γ/IL-4 ratio is of great significance. Furthermore, the mononuclear macrophage system is divided into M1- and M2-type responses and is regulated by the Th1/Th2 immune status. Proinflammatory M1-type macrophages are polarized by binding to lipopolysaccharide (LPS) or Th1-type cytokines (such as IFN-γ), producing proinflammatory cytokines including IL-1β, IL- 23, TNF-α; anti-inflammatory M2 macrophages are polarized by Th2-related cytokines, such as IL-4 and IL-13, and secrete anti-inflammatory cytokines such as IL-10 and TGF-β[21], promoting tissue healing and inflammation resolution[22]. Therefore, the M1/M2 type immune response 227 can be evaluated by the TNF- α /IL-4 ratio and the TNF- α /IL-10 ratio.

 In this study, by collecting laboratory test results, imaging data, and medical history, we discovered that the levels of various inflammatory factors and immune status in patients with acute cerebral infarction were affected by the infarct size.

 The infarct size significantly affects the immune system. While several studies have focused on the effect of infarct size on adaptive immune function impairment, we explored its effects on the immune status and the balance between proinflammatory and anti-inflammatory responses. Fluctuations in various cytokine levels in the peripheral circulation can reflect changes in peripheral immunity following stroke. Our results 236 showed that the levels of cytokines (IFN- γ , TNF- α , IL-2, IL-4, IL-6, IL10, IL-17A) were significantly higher in the group with large-area cerebral infarction than those in non-large-area cerebral infarction group; however, the extent of increase varied for each cytokine. Furthermore, compared to the baseline-matched non-large cerebral infarction group, the absolute value and percentage of lymphocytes decreased significantly in patients with large cerebral infarctions; however, the NLR was significantly higher in the large-area cerebral infarction group. Neutrophils are crucial in innate immunity, and

 the NLR reflects the degree of immune disorder. A higher NLR indicated a greater degree of adaptive immunity impairment. Our findings suggest that patients with large- area cerebral infarctions have more impaired adaptive immune functions, and the relative levels of cytokines affect the immune status (adaptive immunity). Additionally, 247 we calculated the IFN- γ /IL-4, TNF- α /IL-4, and TNF- α /IL-10 ratios. The data showed that the TNF-α/IL-4 ratio was significantly decreased in the group with large-area cerebral infarction, indicating a more anti-inflammatory balance. Notably, the IFN- γ /IL-4 and TNF- α /IL-10 ratios were decreased, suggesting a shift towards an anti-inflammatory response in patients with severe infarction.

 The immune response is involved throughout cerebral infarction and plays an important role in the process of disease transformation with a great degree of variation. Therefore, it is a potential target for improving the prognosis and outcomes of neurological function. However, at present, there is no systematic research on the effects of the infarct volume on cytokines in patients with acute ischemic stroke.and there is a lack of a deeper understanding of the impact of the immune status on clinical adverse events. Cerebral infarction is a complex disease; several conditions induce it, and the pathological mechanism after its onset is complicated.

 This study collected first-hand clinical information on patients and systematically analyzed that severity of infarction may affect the serum cytokine levels and immune status of patients with acute cerebral infarction, and explored how it regulates immunity. This study provides valuable clinical data for related fields and enables the clinicians to make informed decisions in selecting immunotherapy approaches and tailor rehabilitation strategies to optimize patient outcomes by considering different infarct sizes.

It is made available under a [CC-BY 4.0 International license](http://creativecommons.org/licenses/by/4.0/) .

268

269

Acknowledgment

 We are highly thankful to Bethune First Hospital of Jilin University, Professor Ms. Wang, and Professor Ms.Ma for guiding us to finalize this article.Thanks to Xingqi Su, Lingmin Zhao,Jiulin You,LiangShu Feng,LiChong Zhu,Ying Chen,YuLei Hao,XinYu Wang ,JiaChun Feng,the above authors for their joint efforts.

Funding

None.

Data availability

 The data that support the findings of this study are available from the corresponding author upon reasonable request.

ethics statements

 The Ethics Committee of the First Hospital of Jilin University reviewed and approved the study, ensuring that it adhered to ethical principles and guidelines for human subject research. The approval number AF-IRB-032-02 indicates that the study has undergone thorough ethical scrutiny and met the necessary requirements.

 The participants were provided with detailed information about the study, including its purpose, procedures, potential risks and benefits, confidentiality measures, and their rights as participants. They were given ample time to ask questions and clarify any concerns before voluntarily signing the consent form.

 The written informed consent form documented the participant's agreement to participate in the study and their understanding of the information provided. It also stated that they could withdraw from the study at any time without penalty.

 To ensure confidentiality, the consent forms were securely stored and accessible only to the research team. Any identifying information was anonymized or coded to protect the participants' privacy.The research team will continue to prioritize the welfare and rights of the participants throughout the study, maintaining ethical standards and promptly addressing any concerns that may arise.

It is made available under a [CC-BY 4.0 International license](http://creativecommons.org/licenses/by/4.0/) .

- 303 Atients with acute ischemic stroke that presented within 72 h of onset from October 2017 to
- 304 October 2019 were enrolled.

perpetuity. preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in medRxiv preprint doi: [https://doi.org/10.1101/2023.10.16.23297078;](https://doi.org/10.1101/2023.10.16.23297078) this version posted October 16, 2023. The copyright holder for this

It is made available under a [CC-BY 4.0 International license](http://creativecommons.org/licenses/by/4.0/) .

References

- 1.Pan J, Li X, Peng Y. Remote ischemic conditioning for acute ischemic stroke: dawn in the darkness. Reviews in the Neurosciences. 2016;27.
- 2.Fernandez DM, Rahman AH, Fernandez NF, Chudnovskiy A, Giannarelli C. Single-cell immune landscape of human atherosclerotic plaques. Nature Medicine. 2019;25(10).
- 3.Pakzad B, Rajae E, Shahrabi S, Mansournezhad S, Saki N. T-Cell Molecular Modulation Responses in Atherosclerosis Anergy. Laboratory Medicine. 2020;(16).
-
- 4.Yi X, Zhu L, Sui G, Li J, Bao S. Inflammation and Endothelial Function Relevant Genetic Polymorphisms and Carotid Plaque in Chinese Population. Journal of Atherosclerosis and Thrombosis. 2020;27(9).
- 5.Korn T, Kallies A. T cell responses in the central nervous system. Nature Reviews Immunology. 2017;17(3):179-94. doi: 10.1038/nri.2016.144.
- 6.Becker KJ, Kalil AJ, Tanzi P, Zierath DK, Savos AV, Gee JM, et al. Autoimmune responses to the brain after stroke are associated with worse outcome. Stroke. 2011;42(10):2763-9. Epub
- 20110728. doi: 10.1161/strokeaha.111.619593.
- 7.Ren H, Liu X, Wang L, Gao Y. Lymphocyte-to-Monocyte Ratio: A Novel Predictor of the Prognosis of Acute Ischemic Stroke. J Stroke Cerebrovasc Dis. 2017;26(11):2595- 602.Epub20170826.doi:10.1016/j.jstrokecerebrovasdis.2017.06.019.
- 8.Nam KW, Kim TJ, Lee JS, Kwon HM, Lee YS, Ko SB, et al. High Neutrophil-to-Lymphocyte Ratio Predicts Stroke-Associated Pneumonia. Stroke. 2018;49(8):1886-92. doi: 10.1161/strokeaha.118.021228.
- 9.Wang Y, Zhang JH, Sheng J, Shao A. Immunoreactive Cells After Cerebral Ischemia. Front Immunol.2019;10:2781.Epub 20191126. doi:10.3389/fimmu.2019.02781.
- 10.Jian Z, Liu R, Zhu X, Smerin D, Zhong Y, Gu L, et al. The Involvement and Therapy Target of Immune Cells After Ischemic Stroke. Front Immunol. 2019;10:2167. Epub 20190911. doi: 10.3389/fimmu.2019.02167.
- 11.Saino O, Taguchi A, Nakagomi T, Nakano-Doi A, Kashiwamura S, Doe N, et al. Immunodeficiency reduces neural stem/progenitor cell apoptosis and enhances neurogenesis in the cerebral cortex after stroke. J Neurosci Res. 2010;88(11):2385-97. doi: 10.1002/jnr.22410.
- 12.Butovsky O, Ziv Y, Schwartz A, Landa G, Talpalar AE, Pluchino S, et al. Microglia activated by IL-4 or IFN-gamma differentially induce neurogenesis and oligodendrogenesis from adult stem/progenitor cells. Mol Cell Neurosci. 2006;31(1):149-60. Epub 20051116. doi:
- 10.1016/j.mcn.2005.10.006.
- 13.Albini A, Marchisone C, Del Grosso F, Benelli R, Masiello L, Tacchetti C, et al. Inhibition of angiogenesis and vascular tumor growth by interferon-producing cells: A gene therapy approach.Am J Pathol.2000;156(4):1381-93.doi: 10.1016/s0002-9440(10)65007-9.
- 14.Wang S, Zhang H, Xu Y. Crosstalk between microglia and T cells contributes to brain damage and recovery after ischemic stroke. Neurol Res. 2016;38(6):495-503. Epub 20160531. doi: 10.1080/01616412.2016.1188473.
- 15.Cramer JV, Benakis C, Liesz A. T cells in the post-ischemic brain: Troopers or paramedics?J Neuroimmunol. 2019;326:33-7. Epub 20181113. doi: 10.1016/j.jneuroim.2018.11.006.
- 16.Jickling GC, Liu D, Ander BP, Stamova B, Zhan X, Sharp FR. Targeting neutrophils in ischemic
- stroke: translational insights from experimental studies. J Cereb Blood Flow Metab.

2015;35(6):888-901. Epub 20150325. doi: 10.1038/jcbfm.2015.45.

- 17.Maestroni GJ. Sympathetic nervous system influence on the innate immune response. Ann N Y
- Acad Sci. 2006;1069:195-207. doi: 10.1196/annals.1351.017.
- 18.Yuan M, Han B, Xia Y, Liu Y, Wang C, Zhang C. Augmentation of peripheral lymphocyte-
- derived cholinergic activity in patients with acute ischemic stroke. BMC Neurol. 2019;19(1):236. Epub 20191015. doi: 10.1186/s12883-019-1481-5.
-
- 19.Prass K, Meisel C, Höflich C, Braun J, Halle E, Wolf T, et al. Stroke-induced immunodeficiency
- promotes spontaneous bacterial infections and is mediated by sympathetic activation reversal by
- poststroke T helper cell type 1-like immunostimulation. J Exp Med. 2003;198(5):725-36. Epub 20030825. doi: 10.1084/jem.20021098.
- 20.Sekiya T, Yoshimura A. In Vitro Th Differentiation Protocol. Methods Mol Biol. 2016;1344:183-91. doi: 10.1007/978-1-4939-2966-5_10.
- 21.Shapouri-Moghaddam A, Mohammadian S, Vazini H, Taghadosi M, Esmaeili SA, Mardani F,
- et al. Macrophage plasticity, polarization, and function in health and disease. J Cell Physiol.
- 2018;233(9):6425-40. Epub 20180301. doi: 10.1002/jcp.26429.
- 22.Tschoe C, Bushnell CD, Duncan PW, Alexander-Miller MA, Wolfe SQ. Neuroinflammation
- after Intracerebral Hemorrhage and Potential Therapeutic Targets. J Stroke. 2020;22(1):29-46. Epub
- 20200131. doi: 10.5853/jos.2019.02236.
-
-
-
-

It is made available under a CC-BY 4.0 International license. perpetuity. medRxiv preprint doi: [https://doi.org/10.1101/2023.10.16.23297078;](https://doi.org/10.1101/2023.10.16.23297078) this version posted October 16, 2023. The copyright holder for this
preprint (which was not certified by peer review) is the author/funder, who has granted

It is made available under a CC-BY 4.0 International license. perpetuity. medRxiv preprint doi: [https://doi.org/10.1101/2023.10.16.23297078;](https://doi.org/10.1101/2023.10.16.23297078) this version posted October 16, 2023. The copyright holder for this
preprint (which was not certified by peer review) is the author/funder, who has granted

Fig 1

medRxiv preprint doi: [https://doi.org/10.1101/2023.10.16.23297078;](https://doi.org/10.1101/2023.10.16.23297078) this version posted October 16, 2023. The copyright holder for this
preprint (which was not certified by peer review) is the author/funder, who has granted

medRxiv preprint doi: [https://doi.org/10.1101/2023.10.16.23297078;](https://doi.org/10.1101/2023.10.16.23297078) this version posted October 16, 2023. The copyright holder for this
preprint (which was not certified by peer review) is the author/funder, who has granted

