

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24

Cognitive Impairment and Noncommunicable Diseases in Egypt's Aging Population: Insights and Implications from the pilot of the Longitudinal Study of Egyptian Healthy Aging “AL-SEHA”

Sara A. Moustafa¹

Reem Deif¹

Nada Gaballah²

Mohamed Salama^{1,3}

¹Institute of Global Health and Human Ecology, The American University in Cairo

²Department of Computer Sciences, School of Sciences and Engineering, The American University in Cairo

³Global Brain Health Institute (GBHI), Trinity College Dublin (TCD), Dublin, Ireland

KEY MESSAGES

- The increasing prevalence of cognitive impairment among the aging population creates a need for a comprehensive understanding of its implications on the individuals' quality of life
- A significant association between cognitive impairment and non-communicable diseases exists and further research can help in the strategic allocation of resources for designing the most effective interventions

25

26

27

28 **Cognitive Impairment and Noncommunicable Diseases in Egypt's Aging Population: Insights and** 29 **Implications from the Pilot of the Longitudinal Study of Egyptian Healthy Aging “AL-SEHA”**

30

31 **Abstract**

32 As the global population ages, the prevalence of cognitive impairment among older individuals has been
33 steadily rising. Egypt, like many countries, is grappling with the challenges posed by an aging
34 demographic. Cognitive impairment not only affects the quality of life of older adults but also imposes
35 significant burdens on healthcare systems and societies as a whole. This paper presents findings from the
36 pilot phase of the Longitudinal Study of Egyptian Healthy Aging, known as "AL-SEHA," shedding light
37 on the intricate relationship between cognitive impairment and noncommunicable diseases (NCDs) in
38 Egypt's aging population.

39

40 **Introduction**

41 The concept of healthy aging, as advocated by the World Health Organization, emphasizes nurturing
42 functional ability in older age for an enriched quality of life. This approach integrates intrinsic capacity
43 (IC) and environmental factors, transitioning elder care from disease-centered to proactive personalized
44 interventions, notably in hospitalized older adults¹. The innovative notion of healthy aging encapsulates
45 the process of nurturing and sustaining functional ability in older age for an enriched quality of life. This
46 perspective underscores the essence of functional ability, encompassing health-related attributes that
47 empower older individuals to pursue valued endeavors. It intertwines an individual's (IC), a composite of

48 physical and mental capabilities, with environmental factors, thereby enhancing the potential for a higher
49 quality of life while reducing societal burdens. Traditional elder care approaches, centered on specific
50 disease markers, are transitioning towards a longitudinal, proactive model driven by personalized
51 interventions to amplify intrinsic capacity and functional ability, particularly notable in the context of
52 hospitalized older adults ^{1,2}.

53

54 **Assessing Intrinsic Capacity and Predicting Health Outcomes**

55

56 The assessment of IC's indicators predicts health outcomes, particularly functional capability. Domains
57 such as locomotion, vitality, cognition, psychological, and sensory elements are pivotal for maintaining
58 IC in older adults ³. Beard's study ⁴ corroborates this, identifying five subfactors within the English
59 Longitudinal Study on Ageing (ELSA) that forecast future functioning. However, a universally accepted
60 IC index is yet to emerge, warranting further exploration. The significance of assessing IC becomes
61 evident in its predictive value for health outcomes, specifically concerning functional capability. The
62 intricate nature of IC indicators raises inquiries about the most pertinent markers for comprehensive
63 evaluation of an individual's holistic physical and mental condition. Recent Chinese Longitudinal study
64 CHARLS further substantiates this structural validity through the ELSA approach ⁵, collectively
65 accentuating the predictive potential of IC decline and its domains within aging populations. However, a
66 universally acknowledged IC index for clinical or research purposes is yet to be established, necessitating
67 further exploration and quantification of IC concepts in varied contexts.

68

69 **Cognitive Impairment's Impact on Quality of Life**

70 Cognitive impairment's escalating prevalence, especially with age, necessitates a comprehensive
71 exploration of its implications for individuals' quality of life ^{6,7}. The escalation in cognitive decline,
72 particularly with increasing age and life expectancy, mirrors the mounting challenges facing developed
73 and developing nations. The resultant surge in dementia not only inflicts substantial economic and social
74 burdens on patients and families but also warrants a comprehensive exploration of its prevalence across
75 genders and residential areas. Discrepancies within the existing literature concerning the gender and
76 residential area-based prevalence of cognitive impairment highlight the need for a cohesive understanding
77 of these dimensions ^{6,7}.

78

79 **Socioeconomic Factors and Cognitive Impairment Risk**

80 Socioeconomic factors intricately interplay with cognitive health, with lower education, job insecurity,
81 and lifestyle choices influencing susceptibility to cognitive impairments⁹. Notably, individuals with
82 lower educational attainment, diminished IQ, and tenuous job security exhibit higher susceptibility to
83 cognitive impairments. Furthermore, lifestyle choices play a pivotal role, with tobacco and alcohol usage
84 surfacing as significant predictors of cognitive decline. These connections emphasize the multifaceted
85 nature of cognitive health, woven intricately with socioeconomic dimensions and behavioral patterns.

86

87

88 **Noncommunicable Diseases (NCDs) and Cognitive Decline**

89 NCDs, like diabetes and hypertension, impact cognitive faculties, revealing the interconnectedness of
90 physical and cognitive well-being (World Health Organization, 2016)¹⁰. NCDs, a formidable presence
91 within the global health landscape, assume a central role in shaping morbidity and mortality patterns.
92 Their impact resonates through reduced problem-solving capabilities, compromised verbal proficiency,
93 and waning numerical comprehension, underscoring the intricate interplay between physical and
94 cognitive well-being¹⁰.

95

96 **Cognitive Impairment and NCDs in Egypt: Gaps in Literature**

97 As people age, they are more likely to experience chronic non-communicable conditions. Based on the
98 recommendations of WHO, the National Institute of Aging (NIA), and the Centers for Disease Control
99 and Prevention

100 (CDC), fighting non-communicable diseases is broadly agreed to be the key to further health gains at an
101 older age and to making health and social policies sustainable. NCDs, malnutrition, and depression are
102 common in our elderly population, particularly among women and the under-educated¹¹.

103 Aging is often accompanied by cognitive function, memory, and mental health declines. Research on
104 effective interventions to prevent or delay these declines and improve the quality of life of older adults is
105 critical¹². With a prevalence of depression, a sense of isolation, and declined cognitive functions in the
106 Egyptian elderly^{12,13}. In older people, cognitive impairment is a strong predictor of functional disability
107 and the need for care. Mild cognitive impairment increases the risk of developing dementia, and available
108 evidence suggests that a five-year delay in the age of onset would reduce dementia prevalence by half¹⁴

109 .^{15,16}. There is limited data on mental health and cognitive functions in the Arab world, and specifically in
110 Egypt where data are specific to certain areas or specific clinical populations rather than those derived
111 from large nationwide studies. With this age group expected to grow by 2050, the consequences of
112 dementia will grow as well.

113 Existing research often focuses on cognitive impairment within NCDs, leaving a gap in understanding
114 cognitive health among the elderly holistically. Extant investigations into cognitive impairment have
115 predominantly focused on individuals afflicted by NCDs, imparting a limited scope to understanding the
116 broader landscape of cognitive health among the elderly. Prior research has, in various instances,
117 examined the intersection of cognitive issues and NCDs, potentially excluding a more holistic
118 comprehension of cognitive health among the elderly population. Addressing this void, our study
119 endeavors to illuminate the prevalence of cognitive impairment within Egypt's older demographic,
120 specifically those grappling with chronic diseases.

121

122 **Investigating Cognitive Impairment Among Egypt's Older Population**

123

124 **Methods**

125 Preliminary results of the pilot phase of the Egyptian Longitudinal Study for Aging (AL-SEHA), an HRS
126 sister study adopting the SHARE questionnaire presented in Table 3, offer insights into Egypt's aging
127 demographic. The pilot study includes respondents aged 50 and above, with a sample size of 299
128 participants. The results reflect a diversity of conditions, with varying percentages of cognitive
129 impairment across diseases. Responding to this need, an Egyptian aging study modeled after the Health
130 and Retirement Survey (HRS) network studies is being launched. Our findings are based on the AL-
131 SEHA questionnaire, which adapted the SHARE survey, targeting noninstitutionalized individuals aged
132 50 and above¹⁷. This encompassing longitudinal survey included all those above 50 within chosen
133 households.

134 The studies involving human participants were reviewed and approved by IRB of the American
135 University in Cairo (IRB-AUC). The patients/participants provided their written informed consent to
136 participate in this study. The current work has been approved by the IRB of the American University in
137 Cairo. IRBAUC_ Case# 2021-2022-029

138

139 **Research design**

140 The study used a cross-sectional design, assessing both dependent and independent variables
141 simultaneously. Findings were based on a questionnaire from the Longitudinal Study of Egyptian Healthy
142 Aging (AL-SEHA), adapted from the European Survey of Health, Ageing, and Retirement (SHARE).
143 Interviews included individuals aged 50 and above, as well as eligible partners living in the same
144 household. Some questionnaire modules were not given to all household respondents, and interview order
145 within couples was random.

146 **Inclusion criteria**

147 The study included individuals who could communicate and lived independently at home, in rented
148 accommodation, hostels, or retirement homes. Those with severe psychiatric issues were excluded. The
149 sample size was 299 participants, selected through a non-probability convenience sampling technique
150 based on the eligibility criteria.

151 **Fieldwork**

152 Investigators from five universities were trained to collect data using an Arabic-translated SHARE form.
153 A sample of eligible elderly individuals was chosen, and home visits were conducted to interview 15
154 participants per investigator. Data collection took about a month and was uploaded to the American
155 University in Cairo's Social Research Centre's internet server.

156 **Demographic factors**

157 AL-SEHA data set contains detailed data on demographics. Summary statistics of all demographic
158 variables used in the analyses can be found in Table 1.

159 **Table 1. Demographics**

	N	%
Gender		
Male	128	42.81
Female	171	57.19
Educational Attainment		
No education	48	16.05

Primary education	31	10.37
Preparatory education	18	6.02
High school	48	16.05
Medium high	18	6.02
Bachelor	112	37.46
Post grad	24	8.03
Age		
50-60	151	50.5
60-70	88	29.43
70-80	50	16.72
80-90	8	2.68
Above 90	2	0.67
Marital Status		
Unmarried	5	1.67
Engaged	0	0
Married	190	63.55
Separated	3	1
Divorced	13	4.35
Widow	88	29.43

160

161 **Cognitive function**

162 The data were extracted from the SPSS (Version 25) dataset for statistical analysis. Bivariate analyses
163 included chi-square tests. As appropriate, data were compared using independent t-tests, presented in
164 Table 2.

165

166

167

168

169 **Table 2. Cognitive functions**

	Impaired (n=118)		Unimpaired (n=181)		Chi Squared Test	p-value
Age	N	%	N	%		
50-60	52	44.07	99	54.7		
60-70	39	33.05	49	27.07		
70+	27	22.88	33	18.23	3.23	0.2
Mean (SD)	62.338(8.8087)		60.635(8.0505)			
Gender						
Male	44	37.29	84	46.41		
Female	74	62.71	97	53.59	2.07	0.15
Education						
No education	29	24.58	19	10.5		
Basic/intermediate	52	44.07	63	34.81		
University	37	31.36	99	54.7	18.968	0.0001
Marital Status						
Single	1	0.85	4	2.21		
Married	69	58.47	121	66.85		
Separated	3	2.54	0	0		
Divorced	3	2.54	10	5.52		

Widow	42	35.59	46	25.41	10.159	0.0378
-------	----	-------	----	-------	--------	---------------

170

171

172

173 **Dependent variable**

174 Cognitive performance was assessed using a composite measure based on three tests: immediate recall,
175 delayed recall, and verbal fluency. The immediate and delayed recall tasks evaluated short-term verbal
176 learning and memory. Verbal fluency measured executive function and language ability. The composite
177 score ranged from 0 to 1, where 0 to 0.5 indicated impairment and 0.5 to 1 indicated unimpaired cognitive
178 function.

179

180 **Covariates and moderators**

181 Cognitive performance in later life is influenced by various factors such as age, education, gender, and
182 marital status. To account for these variables, they were included as covariates in the analyses. Gender
183 was represented as a dummy variable with 1 for men and 2 for women. Age was also represented as a
184 dummy variable with 1 for ages between 50 and 60, 2 for ages between 60 and 70, and 3 for ages over 70.
185 Similarly, education was represented as a dummy variable with 1 for no education, 2 for basic to
186 intermediate education, and 3 for university or postgraduate studies.

187 **Analyses of data**

188 To begin, descriptive statistics were used to compute The means and standard deviations of continuous
189 variables, as well as the percentages and frequencies of categorical variables.

190 Bivariate analyses were performed in the second stage to examine the relationship between cognitive
191 health and the study variables, using independent t-tests for factors.

192 **Cognitive performance**

193 The researchers assessed cognitive functioning using three tasks: verbal fluency, immediate word recall,
194 and delayed word recall. Verbal fluency evaluated executive function by asking participants to name as
195 many animals as possible in one minute. Memory function was assessed through the Ten-Word Delayed
196 Recall Test, where participants recalled ten words immediately and after five minutes. These measures

197 allowed researchers to evaluate executive function and memory constructs, differentiating between age-
 198 related memory changes and significant impairments. The assessment was based on widely-used
 199 neuropsychological tasks for memory performance evaluation.

200

201

202 **Table 3. Non-communicable diseases and cognitive functions preliminary results**

Disease Name		Diseases	No disease	Percentage	Disease	Percentage%	
Cardiovascular Disease: (Heart disease, hypertension, cholesterol and angina)	Age group	50-60	492	81.46	112	18.54	
		60-70	244	69.32	108	30.68	
		70 and above	150	62.50	90	37.50	
	Gender	Male	371	72.46	141	27.54	
		Female	515	75.29	169	24.71	
	Education	No education	150	78.13	42	21.88	
		Basic/intermediate	335	72.83	125	27.17	
		University/post grad	401	73.71	143	26.29	
	Cognition	Unimpaired	563	77.76	161	22.24	
		Impaired	323	68.43	149	31.57	
	Neuro/psychiatric Disease: (Parkinson's disease, Insomnia, Fall, Fear of Falling,	Age group	50-60	834	92.15	71	7.85
			60-70	463	87.69	65	12.31
70 and above			297	82.50	63	17.50	
Gender		Male	695	90.49	73	9.51	

Dizziness, Brain Clots)		Female	899	87.71	126	12.29	
	Education	No education	254	88.50	33	11.50	
		Basic/intermediate	621	90.00	69	10.00	
		University/post grad	719	88.11	97	11.89	
	Cognition	Unimpaired	1002	92.35	83	7.65	
		Impaired	593	83.88	114	16.12	
Metabolic Disease: (Diabetes)	Age group	50-60	127	84.11	24	15.89	
		60-70	55	62.50	33	37.50	
		70 and above	28	46.67	32	53.33	
	Gender	Male	91	71.09	37	28.91	
		Female	119	69.59	52	30.41	
	Education	No education	34	70.83	14	29.17	
		Basic/intermediate	77	66.96	38	33.04	
		University/post grad	99	72.79	37	27.21	
	Cognition	Unimpaired	137	75.69	44	24.31	
		Impaired	73	61.86	45	38.14	
	Abdominal and Gastrointestinal Disease: (Stomach Ulcer, Urine	Age group	50-60	407	89.85	46	10.15
			60-70	215	81.44	49	18.56
70 and above			148	82.22	32	17.78	
Gender		Male	339	88.28	45	11.72	

Incontinence, Bowel Problems)		Female	431	84.02	82	15.98	
	Education	No education	127	88.19	17	11.81	
		Basic/intermediate	308	89.28	37	10.72	
		University/post grad	335	82.11	73	17.89	
	Cognition	Unimpaired	463	85.27	80	14.73	
		Impaired	307	86.72	47	13.28	
Chest Disease: (Asthma, Lung Disease, Hard Breathing, Cough)	Age group	50-60	565	93.54	39	6.46	
		60-70	314	89.20	38	10.80	
		70 and above	222	92.50	18	7.50	
	Gender	Male	467	91.21	45	8.79	
		Female	634	92.69	50	7.31	
	Education	No education	185	96.35	7	3.65	
		Basic/intermediate	428	93.04	32	6.96	
		University/post grad	488	89.71	56	10.29	
	Cognition	Unimpaired	689	95.17	35	4.83	
		Impaired	412	87.29	60	12.71	
	Cancer	Age group	50-60	149	98.68	2	1.32
			60-70	85	96.59	3	3.41
70 and above			58	96.67	2	3.33	
Gender		Male	124	96.88	4	3.13	

		Female	168	98.25	3	1.75
	Education	No education	47	97.92	1	2.08
		Basic/intermediate	113	98.26	2	1.74
		University/post grad	132	97.06	4	2.94
	Cognition	Unimpaired	176	97.24	5	2.76
		Impaired	116	98.31	2	1.69
Musculoskeletal Disease: (Osteoporosis, Arthritis, Hip Fracture, Back/Joint Pain)	Age group	50-60	480	79.47	124	20.53
		60-70	260	76.25	81	23.75
		70 and above	161	67.08	79	32.92
	Gender	Male	427	83.40	85	16.60
		Female	474	69.40	209	30.60
	Education	No education	131	68.23	61	31.77
		Basic/intermediate	365	79.52	94	20.48
		University/post grad	405	74.45	139	25.55
	Cognition	Unimpaired	554	76.52	170	23.48
		Impaired	347	73.67	124	26.33

203

204 **Prevalence of Noncommunicable Diseases (NCDs)**

205 Our study focuses on cardiovascular, abdominal, chest, neuro/psychiatric, metabolic, and musculoskeletal
 206 diseases and cancer's impact on cognitive impairment presented in Table 3.. The prevalence of these
 207 diseases varies across age groups and genders, reflecting the intricate web of interactions shaping Egypt's
 208 aging population.

209 **Neuro/Psychiatric Diseases and Cognitive Well-being**

210 Cognitive impairment reveals the interconnectedness of mental well-being and cognitive faculties.
211 Conditions like insomnia show statistically significant associations with cognitive impairment (p-value:
212 0.0169). The intricate interplay between cognitive well-being and mental health becomes evident when
213 exploring associations between neuro/psychiatric diseases and cognitive impairment.

214

215 **Metabolic Diseases and Their Impact on Cognition**

216 Metabolic diseases, epitomized by diabetes, demonstrate their nuanced relationship with cognitive health
217 through statistically significant associations with cognitive impairment (p-value: 0.0137). These findings
218 accentuate the urgency of interventions that holistically address metabolic and cognitive well-being,
219 thereby potentially ameliorating the progression of cognitive decline.

220

221 **Respiratory Health and Cognitive Implications**

222 Associations between chest diseases and cognitive impairment underline the intricate connection between
223 respiratory health and cognitive well-being. Conditions like hard breathing and cough show statistically
224 significant associations with cognitive impairment (p-values: 0.000076, 0.0032). The intricate correlation
225 between respiratory health and cognitive well-being materializes in associations between chest diseases
226 and cognitive impairment.

227

228 **Musculoskeletal Health and Functional Independence**

229 The intricate dance between physical well-being and cognitive health manifests as associations between
230 musculoskeletal diseases and cognitive impairment. Ailments like falls, fear of falls, and dizziness, often
231 rooted in musculoskeletal health, demonstrate statistically significant associations with cognitive
232 impairment, underscoring the profound interaction between the physical and cognitive domains (p-values:
233 0.00108, 0.001089, 0.0089).

234

235 **Cancer and Its Nuanced Prevalence**

236 Cancer prevalence in the older Egyptian population was notably low. Specifically, among those aged 50-
237 60 years, only 1.32% had cancer, with 98.68% without the disease. In the 60-70 age group, 3.41% were

238 affected by cancer, while 96.59% were not. Similarly, among individuals aged 70 and above, cancer was
239 found in 3.33%, with 96.67% unaffected. However, Larger and more diverse samples are needed to draw
240 more definitive conclusions about cancer rates in the Egyptian older population. Acknowledging these
241 limitations is essential for a balanced perspective on the findings, and further research is required for
242 more accurate insights into the cancer burden among the older population in Egypt.

243

244 **Discussion**

245 **Implications for Egypt's Aging Population**

246 Findings hold significance for Egypt's aging demographic, informing policies and interventions that cater
247 to older adults' unique needs. The implications of these findings extend far beyond statistical associations.
248 They reverberate through policy decisions, healthcare interventions, and societal strategies that
249 acknowledge the diverse needs and challenges faced by Egypt's aging demographic. These implications
250 underscore the imperatives of tailored interventions that holistically address the intricate interplay
251 between physical and cognitive well-being.

252 The policy implications derived from the AL-SEHA pilot study data present a comprehensive framework
253 for the development of effective strategies aimed at addressing noncommunicable diseases (NCDs) and
254 cognitive health concerns within Egypt's elderly population. To address these issues, it is imperative to
255 craft and implement targeted programs with a strong focus on the prevention and management of NCDs,
256 with particular attention to the promotion of cognitive health and overall well-being. Additionally, the
257 evaluation of the necessity for disease prevention initiatives, including adult vaccination campaigns,
258 awareness programs addressing tobacco use, and initiatives that encourage physical activity among older
259 adults, is crucial. Furthermore, enhancing the geriatric care landscape in Egypt is essential, involving the
260 incorporation of specialized care pathways that cater to the unique needs of the elderly. This also
261 necessitates a revision of the medical education curriculum to include geriatric medicine, emphasizing the
262 diagnosis, treatment, and management of multiple coexisting health conditions prevalent in the aging
263 population. Expanding the capacity and accessibility of geriatric care facilities is vital to ensure
264 comprehensive and effective support for older individuals. Furthermore, the integration of geriatric
265 healthcare services across primary, secondary, and tertiary healthcare systems, including community-
266 based health initiatives, is essential to provide holistic care. Cross-regional analyses should be conducted
267 to identify best practices in areas such as long-term care, home-based support, legal protections, and
268 healthcare expenditure, customized to the specific context of Egypt. Additionally, the formulation of
269 national strategies for dementia prevention, targeting both the general population and vulnerable

270 subgroups, is paramount. This should be complemented by the establishment of governmental initiatives
271 at both national and local levels to effectively address the multifaceted challenges of dementia care and
272 provide support for patients and caregivers in Egypt. These policy recommendations collectively form a
273 robust foundation for addressing the pressing healthcare needs of Egypt's elderly population in the context
274 of NCDs and cognitive health.

275

276 **Healthcare Planning and Resource Allocation**

277 Understanding associations between cognitive impairment and diseases aids in strategic resource
278 allocation for timely interventions. The recognition of associations between cognitive impairment and
279 diverse diseases augments the strategic blueprint for resource allocation within the Egyptian healthcare
280 system. Informed by these associations, policymakers can channel resources towards timely interventions
281 that ameliorate cognitive decline, enhancing not only individual well-being but also alleviating societal
282 burdens.

283

284 **Catalyzing Research and Collaboration**

285 Associations between cognitive impairment and diseases foster interdisciplinary research and
286 collaboration for holistic strategies. The implications of cognitive impairment's associations with diseases
287 extend beyond individual interventions, catalyzing collaborative research across disciplines in Egypt and
288 the MENA region. This interdisciplinary endeavor propels the creation of comprehensive strategies that
289 transcend traditional healthcare silos, forging innovative pathways to address the intricate web of
290 cognitive decline.

291

292 **Conclusion: Enabling a Brighter Aging Future for Egypt**

293 Evidence-based policies and interventions pave the way for a healthier and more inclusive aging
294 demographic. In conclusion, the synthesis of evidence-based policies and interventions emerges as the
295 cornerstone of facilitating a brighter, healthier aging trajectory for Egypt. The intricate interplay between
296 cognitive impairment and noncommunicable diseases unravels novel avenues for policymaking and
297 interventions, steering the nation toward a future that prioritizes the well-being and dignity of its aging
298 population.

299

300 **Limitations of the Study and Suggestions for Future Research in the Egyptian Context**

301 Acknowledging limitations, future research can employ longitudinal designs and objective measures to
302 enhance understanding. Admitting the inherent limitations of this study, future research endeavors within
303 the Egyptian context can harness the power of longitudinal designs and objective measurements. These
304 methodological refinements promise to illuminate deeper insights into the dynamic interplay between
305 cognitive impairment and diseases, thereby enriching the foundation for comprehensive interventions.

306 In conclusion, this paper delved into the intricate web connecting cognitive impairment and
307 noncommunicable diseases within Egypt's aging population. By investigating associations between
308 cognitive decline and diverse diseases, we illuminated a tapestry of interactions that transcend traditional
309 healthcare boundaries. The implications of these findings resonate through evidence-based policies,
310 cultural considerations, interdisciplinary collaboration, and most importantly, empowered aging
311 experiences. These insights paves the way for Egypt towards a future characterized by enhanced well-
312 being, dignity, and inclusivity for its elderly citizens.

313

314

315 **Correspondence to: Sara A. Moustafa**

316 **Full name: Sara Ayman Moustafa**

317 **Mailing address: The American University in Cairo**

318 **Email: saraayman@Aucegypt.edu**

319

320 **Acknowledgment**

321

322 The present work has been supported by the German Academic Exchange Services (DAAD), through the
323 funding program: Higher Education Dialogue with the Muslim World, project (Ageing in the East
324 Mediterranean Region: EMage). This work benefitted from an incentive grant from the World Health
325 Organization (WHO) as part of the WHO Global Noncommunicable Disease (NCD) Platform Young
326 Researchers Programme (Grant reg. № 2022/1256080).

327 The authors want to thank Dr. Tea Collins and Ms. Daria Berlina for their support and feedback on the
328 final draft.

329

330 **Patient involvement**

331 No patients were involved.

332

333 **Conflicts of Interest**

334 We have no interests to declare.

335

336

337

338 **References**

- 339 1. Chen X, Smith J, Strauss J, Wang Y, Zhao Y. China Health and Retirement Longitudinal Study
340 (CHARLS). In: Pachana N, eds. Encyclopedia of Geropsychology. Springer; 2015.
341 https://doi.org/10.1007/978-981-287-080-3_42-1
- 342 2. Cesari M, Araujo de Carvalho I, Amuthavalli Thiyagarajan J, et al. Evidence for the Domains
343 Supporting the Construct of Intrinsic Capacity. *J Gerontol A Biol Sci Med Sci*.
344 2018;73(12):1653-1660. doi:10.1093/gerona/gly011
- 345 3. Beard JR, Si Y, Liu Z, Chenoweth L, Hanewald K. Intrinsic Capacity: Validation of a New WHO
346 Concept for Healthy Aging in a Longitudinal Chinese Study. *J Gerontol A Biol Sci Med Sci*.
347 2022;77(1):94-100. doi:10.1093/gerona/glab226
- 348 4. Sharma D, Mazta S, Parashar A. Prevalence of cognitive impairment and related factors among
349 the elderly: a population-based study. *J Dr. NTR Univ. Heal. Sci*. 2013;2:171.
350 <https://doi.org/10.4103/2277-8632.117182>
- 351 5. Grover S, Nehra R, Malhotra R, Kate N. Positive aspects of caregiving experience among
352 caregivers of patients with dementia. *East Asian Arch. Psychiatry*. 2017;27:71-78.
- 353 6. Patel R, Singh U. Prevalence study of cognitive impairment and its associated sociodemographic
354 variables using mini-mental status examination among elderly population residing in field

- 355 practice areas of a medical college. *Indian J. Community Med.* 2018;43:113-116.
356 https://doi.org/10.4103/ijcm.IJCM_102_17
- 357 7. Abdel-Hady D, El-Gilany A-H. Tobacco use and its associated factors among older people: a
358 community-based study in Egypt. *East Mediterr Health J.* 2020;26(1):68–74.
359 <https://doi.org/10.26719/2020.26.1.68>
- 360 8. Egyptian Environmental Affairs Agency (EEAA). Egypt second national communication under
361 the united nations framework convention on climate change. Cairo: Egyptian Environmental
362 Affairs Agency (EEAA); 2010
- 363 9. El-Sherbiny NA, Younis A, Masoud M. A comprehensive assessment of the physical, nutritional,
364 and psychological health status of the elderly populace in the Fayoum Governorate (Egypt). *Arch*
365 *Gerontol Geriatr.* 2016;66:119-126. doi:10.1016/j.archger.2016.05.004
- 366 10. Okasha A. History of mental health in the Arab world. In: Okasha A, Maj M, eds. *Images in*
367 *Psychiatry: An Arab Perspective.* World Psychiatric Association; 2001:1-20.
- 368 11. Endrawes G, O'Brien L, Wilkes L. Mental illness and Egyptian families. *Int J Ment Health Nurs.*
369 2007;16:178-187. doi:10.1111/j.1447-0349.2007.00456.x
- 370 12. Aly HY, Hamed AF, Mohammed NA. Depression among the elderly population in Sohag
371 governorate. *Saudi Med J.* 2018;39(2):185–190. <https://doi.org/10.15537/smj.2018.2.21353>
- 372 13. Alsayyad E, et al. Gender difference in health issues and cognitive functions among an Egyptian
373 normal elderly population. *Egypt J Neuropsychiatr Neurosurg.* 2020;56(1). Available at:
374 <https://doi.org/10.1186/s41983-020-0146-z>
- 375 14. Odejimi O, Tadros G, Sabry N. A systematic review of the prevalence of mental and
376 neurocognitive disorders amongst older adults' populace in Egypt. *Middle East Curr Psychiatry.*
377 2020;27:47. <https://doi.org/10.1186/s43045-020-00055-8>
- 378 15. Hughes JC. Whole sight: clinical and research priorities in old age psychiatry. *Lancet Psychiatry.*
379 2017;4(2):94-95. doi:10.1016/S2215-0366(16)30353-4
- 380 16. Youssef W. Application of total scores of CERAD neuropsychological battery in detecting MCI
381 and dementia in a sample of Egyptian elderly. *J Posit School Psychol.* 2022;6(2):1535–1544.
- 382 17. Börsch-Supan A, Hank K, Jürges H. A new comprehensive and international view on ageing:
383 Introducing the 'Survey of Health, Ageing and Retirement in Europe'. *Eur J Ageing.* 2005;2:245-
384 253. doi:10.1007/s10433-005-0014-9
- 385 18. Cheke LG, Clayton NS. Do different tests of episodic memory produce consistent results in
386 human adults? *Learn Mem.* 2013;20(9):491-498. doi:10.1101/lm.030502.113