

1 **Changes in emergency department utilization in vulnerable populations after COVID-19**
2 **shelter in place orders**

3

4 Philip R. Wang BS¹, Akhil Anand MD^{1,2}, James Bena MS³, Shannon Morrison MS³, and Jeremy
5 Weleff DO⁴

6

- 7 1. Cleveland Clinic Lerner College of Medicine at Case Western Reserve University, EC-10
8 Cleveland Clinic, Cleveland, OH, USA
9 2. Department of Psychiatry and Psychology, Center for Behavioral Health, Neurological
10 Institute, Cleveland Clinic, Cleveland, OH, USA
11 3. Department of Quantitative Health Sciences, Cleveland Clinic, Cleveland, Ohio
12 4. Department of Psychiatry, Yale University School of Medicine, New Haven, Connecticut,
13 USA

14

15

16 **Corresponding author:**

17 Jeremy Weleff, DO
18 Yale University School of Medicine, Department of Psychiatry
19 300 George St, New Haven, CT, 06511
20 Phone/fax: 517-803-5675
21 jeremy.weleff@yale.edu

22

23 **Short title:** COVID-19's effect on patients with MH/SUD/SDOH

24 **Keywords:** social determinants of health, mental health, substance use disorders, emergency
25 department utilization, Z-codes

26 Word Count: 1462

27 Prior presentations: none

28 **Funding sources/disclosures:** This research did not receive any specific grant from funding
29 agencies in the public, commercial, or not-for-profit sectors. All authors have no conflicts of
30 interest, financial or otherwise, to declare related to this manuscript.

31 **Credit author statement:** Concept and design: PW, AA, JW. Acquisition of the data: JB, SM,
32 AA, JW. Analysis and interpretation of the data: PW, AA, JB, SM, JW. Drafting of the
33 manuscript: PW. Critical revision of the manuscript for important intellectual content: AA, JW,
34 JB, SM. Statistical expertise: JB, SM, JW. Acquisition of funding: None

35

36

37 **Abstract:**

38 **Purpose:** To compare emergency department (ED) utilization and admission rates for patients
39 with a history of mental health (MH), substance use disorder (SUD) and social determinants of
40 health (SDOH) before and after implementing COVID-19's shelter-in-place (SIP) orders.

41 **Methods:** This was a retrospective, multicenter study leveraging electronic medical record data
42 from 20 EDs across a large Midwest integrated healthcare system from 5/2/2019 to 12/31/2019
43 (pre-SIP) and from 5/2/2020 to 12/31/2020 (post-SIP). Diagnoses were documented in the
44 patient's medical records. Poisson and logistic regression models were used to evaluate ED
45 utilization and admission rate changes.

46 **Results:** 871,020 total ED encounters from 487,028 unique patients were captured. 2,572
47 (0.53%) patients had a documented Z code for SDOH. Patients with previously diagnosed MH or
48 SUDs were more likely to seek ED care after the SIP orders were implemented (RR: 1.20, 95%
49 CI: 1.18 – 1.22, $p < 0.001$), as were patients with SDOH (RR: 2.37, 95% CI: 2.19 – 2.55,
50 $p < 0.001$). Patients with both previously diagnosed MH or SUD and a documented SDOH had
51 even higher ED utilization (RR: 3.31, 95% CI: 2.83 – 3.88, $p < 0.001$) than those with either
52 condition alone. Patients with MH and SUDs (OR: 0.89, 95% CI: 0.86 – 0.92, $p < 0.001$) or
53 SDOH (OR: 0.67, 95% CI: 0.54, 0.83, $p < 0.001$) were less likely to be admitted post-SIP orders
54 while patients with a history of diseases of physiologic systems were more likely to be admitted.

55 **Conclusions:** Vulnerable populations with a history of MH, SUD, and SDOH experienced
56 increased ED utilization but a lower rate of hospital admissions after the implementation of SIP
57 orders. The findings highlight the importance of addressing these needs to mitigate the impact of
58 public health crises on these populations.

59

60 **Keywords:** social determinants of health, mental health, substance use disorders, emergency
61 department utilization, Z-codes

62 **Abbreviations:**

- 63 • Mental health – MH
- 64 • Substance use disorder - SUD
- 65 • Social determinant of health – SDOH
- 66 • Emergency department – ED
- 67 • Shelter in place – SIP

68

69 **Word Count (not including abstract):** 1496

70 **No. of text pages:** 4

71 **No. of figures/tables:** 2

72

73 **Background**

74 Patients with mental health (MH) and substance use disorders (SUD) face significant challenges
75 in accessing healthcare equitably.¹ In addition, social determinants of health (SDOH) profoundly
76 impact healthcare utilization, delivery, quality of life, and mortality. Identifying these patients for
77 targeted interventions is crucial²; however, inconsistent and non-standardized documentation of
78 SDOH across healthcare systems hinders public health surveillance efforts of this population. To
79 address this issue, the Centers for Medicare and Medicaid Services and other groups have
80 proposed using Z-Codes to standardize the documentation of social needs, but the utilization of
81 these codes remains limited.^{3,4}

82 The COVID-19 pandemic has exacerbated health disparities and disproportionately affected
83 individuals with MH, SUD, and SDOH.⁵ While research has shown that shelter-in-place (SIP)
84 orders have led to increased ED visits for mental health and substance use reasons⁶⁻⁸, few studies
85 have examined the impact on ED utilization and admissions in individuals with a prior history of
86 MH or SUD. Furthermore, the effects of the COVID-19 pandemic on ED utilization and
87 admissions by patients with SDOH have had been poorly studied thus far. The primary aim of
88 this study is to investigate the changes in ED utilization and hospital admission rates among
89 vulnerable populations with a history of MH, SUD, and SDOH before and after SIP orders
90 across a large integrated healthcare system. A secondary aim was to characterize the frequency
91 of utilization of Z-codes to identify patients with SDOH in the ED.

93 **Methods**

94 *Patient population*

95 Patients who visited any of 20 EDs ranging from large hospital-associated EDs to freestanding
96 EDs across a large Midwest integrated healthcare system Ohio from 5/2/2019-12/31/2019
97 (designated as the “pre-SIP order” time period) and 5/2/2020-12/31/2020 (the “post-SIP order”
98 time period) were identified. Demographics and diagnoses were extracted from the electronic
99 medical record (EMR). MH and SUDs were identified using the International Classification of
100 Diseases 10th Revision (ICD-10) codes; similarly, SDOH were identified in patients’ medical
101 histories using respective Z-Codes in the ICD-10. Historical use of these codes means that these
102 diagnoses and SDOH were documented in the patient's chart at any time, and not solely reliant
103 on ED encounter documentation. This study was approved by Cleveland Clinic IRB.

104 *Statistical analysis*

105
106 Poisson regression models were used to calculate absolute risk ratios (RR) to compare the total
107 number of cases between the pre-and post-SIP periods. Logistic regression models were used to
108 evaluate changes in admission rates between the two periods, and odds ratios (OR) were
109 calculated.

111 **Results**

112 *Patient demographics*

113 871,020 total ED encounters from 487,028 unique patients were captured; 473,449 visits (54%)
114 occurred in 2019, and 397,571 (46%) were in 2020. Our cohort was mostly White (65.4%),
115 female (53.8%), privately insured (68.9%), and had a mean age of 46.0 ± 24.0 . 2,572 (0.53%)
116 patients had a documented Z code for SDOH. The most coded SDOH in ED encounters was
117 problems related to housing and economic circumstances, followed by other problems related to
118 the primary support group and other psychosocial circumstances (Supplemental Table 1).
119 Compared to patients without SDOH, patients with a coded SDOH were more likely to be Black,
120 younger, identify as male, and were more likely to be on government or self-pay insurance
121 (Supplemental Table 2).

122 *ED utilization rates before and after shelter-in-place orders*

123 Compared to before the implementation of SIP orders, patients with previously diagnosed MH or
124 SUDs were more likely to seek ED care after the SIP orders were put in place (RR: 1.20, 95%
125 CI: 1.18 – 1.22, $p < 0.001$) (Table 1). Among patients with SDOH, those with a history of
126 problems related to the primary support group (RR: 2.89, 95% CI: 2.54 – 3.29) or housing and
127 economic circumstances (RR: 2.50, 95% CI: 2.23 – 2.80) were more likely to present to the ED
128 after SIP orders. Patients with both previously diagnosed MH or SUD and a documented SDOH
129 (RR: 3.31, 95% CI: 2.83 – 3.88, $p < 0.001$) had even higher ED utilization than those with either
130 condition alone.

131 *ED admission rates before and after shelter-in-place orders*

132 Compared to pre-SIP, patients with a previously diagnosed MH or SUD (OR: 0.89, 95% CI: 0.86
133 – 0.92, $p < 0.001$) or any documented SDOH (OR: 0.67, 95% CI: 0.54, 0.83, $p < 0.001$) were less
134 likely to be admitted to the hospital after presenting to the ED post-SIP orders (Table 2). Of the
135 SDOH diagnosis categories, only patients with problems related to their primary support group
136 and those with problems related to other psychosocial circumstances were significantly less
137 likely to be admitted. The difference in admission rate pre- and post-SIP orders in patients with
138 both MH/SUD and SDOH did not reach significance (OR: 0.65, 95% CI: 0.42, 1.02, $p = 0.061$).

139 **4. Discussion**

140 *The effect of shelter-in-place orders on vulnerable patient populations*

141 The COVID-19 pandemic has underscored the importance of addressing social determinants of
142 health, which are critical in shaping an individual's health status and access to care. Our analysis
143 of ED presentation and admission rates pre- and post-SIP orders found that patients with
144 previously diagnosed MH or SUDs and those with a documented SDOH were more likely to
145 present to the ED post-SIP. Prior studies that analyzed MH and substance use related ED visits
146 demonstrate increased visit rates for these conditions as a primary diagnosis or chief complaint.^{6–}
147 ⁸ This study extends previous work by demonstrating that SIP orders significantly impacted
148 vulnerable patients with a history of MH, SUDs and SDOH.

149 Moreover, our analysis revealed that patients with co-occurring MH/SUD and SDOH had the
150 highest increase in ED visits following SIP orders. These patients may have faced increased
151 difficulties in accessing care and maintaining their overall health due to the closure or reduction

152 of community-based resources⁹ and the exacerbation of existing socioeconomic inequities during
153 the pandemic.⁵ Individuals with multiple disadvantages are at a significantly higher risk of
154 premature mortality from avoidable causes than those with a single disadvantage¹⁰, highlighting
155 the urgent need for tailored interventions that address the complex interplay of social, economic,
156 and mental health-related factors contributing to poor health outcomes among these populations.
157 Furthermore, patients with MH/SUDs or SDOH were less likely to be admitted to the hospital
158 after presenting to the ED. While our study was not designed to examine the reasons behind
159 these disparities, possible explanations include inadequate inpatient resources for patients with
160 MH/SUD and SDOH or healthcare professional biases, as well as structural changes of
161 healthcare resources being shunted to emergent medical care beds.

162 *Low documentation of social determinants of health in patients visiting the ED*

163 Our study found that only 0.53% of ED patients had a documented Z-Code for a SDOH before
164 and during early COVID, which is lower than other rates previously reported.^{3,4,11} Molina et al.
165 estimated that 1.21% of ED visits have a coded SDOH; however, our paper analyzed SDOH
166 coding at the patient level rather than the visit level, which may explain the differences
167 observed.¹² The usage of Z-Codes to identify patients with SDOH has repeatedly been shown to
168 be underutilized and our results likely underestimate the number of patients with critical SDOH
169 seeking care at EDs.¹³ Future studies could consider using natural language processing models to
170 analyze provider notes to identify patients with SDOH that do not have a documented Z-code.
171 Identifying patients with SDOH is crucial because they are more likely to experience adverse
172 health outcomes and increased ED utilization.^{14,15} Coding SDOH is currently not reimbursable,
173 which is likely reflected in its low adoption rate; however, documentation of SDOH can
174 substantiate patient complexity and reveal trends in healthcare utilization. For instance, if
175 hospital systems accurately track SDOH, they can identify Z-Codes associated with higher
176 readmission rates and allocate appropriate resources to improve patient care and outcomes. In the
177 future, Z-Codes could play a role in determining payment rates and risk adjustment, but
178 consistent documentation of SDOH is necessary.¹⁶

179 *Limitations*

180 Our results should be interpreted in the context of their limitations. First, the retrospective design
181 of our study limits our ability to draw causal conclusions. Second, our data comes from one large
182 integrated healthcare system, which may differ from other healthcare systems or geographic
183 regions. Nonetheless, our study includes a large and diverse patient population from different
184 levels of care within the state. Third, our findings may be influenced by institution-specific
185 documentation practices that could affect SDOH coding rates and limit the generalizability of
186 our findings. Finally, our analysis relied on patients' past medical history documentation in the
187 EMR, which may not reflect their current diagnoses at the time of their ED visit. However, given
188 the time constraints of the ED, we considered capturing diagnoses from past medical history the
189 best way to represent our sample's health status.

190 *Conclusion*

191 Our study highlights the impact of the COVID-19 pandemic on vulnerable patient populations
192 with a history of MH, SUDs, and SDOH. Our findings demonstrate that these patients were more
193 likely to present to the ED but less likely to be admitted following SIP orders, indicating a need
194 for tailored interventions that address the complex interplay of socioeconomic and mental health-
195 related factors contributing to poor health outcomes among these populations. The findings
196 highlight the need for greater standardization and consistency in documenting SDOH and
197 addressing these factors in healthcare delivery.

198 **Credit author statement: Philip Wang:** Writing – original draft preparation, methodology,
199 conceptualization **Akhil Anand:** conceptualization, methodology, writing – review & editing,
200 supervision **James Bena:** software, formal analysis, data curation **Shannon Morrison:** software,
201 formal analysis, data curation **Jeremy Weleff:** conceptualization, methodology, writing – review
202 & editing, supervision

203

204 **References**

205

- 206 1. Ann Priester M, Browne T, Iachini A, Clone S, DeHart D, Seay KD. Treatment Access
207 Barriers and Disparities Among Individuals with Co-occurring Mental Health and Substance
208 Use Disorders: An Integrative Literature Review. *J Subst Abuse Treat.* 2016;61:47-59.
209 doi:10.1016/j.jsat.2015.09.006
- 210 2. Moulin A, Evans EJ, Xing G, Melnikow J. Substance Use, Homelessness, Mental Illness and
211 Medicaid Coverage: A Set-up for High Emergency Department Utilization. *West J Emerg*
212 *Med.* 2018;19(6):902-906. doi:10.5811/westjem.2018.9.38954
- 213 3. Truong HP, Luke AA, Hammond G, Wadhwa RK, Reidhead M, Joynt Maddox KE.
214 Utilization of Social Determinants of Health ICD-10 Z-Codes Among Hospitalized Patients in
215 the United States, 2016–2017. *Med Care.* 2020;58(12):1037-1043.
216 doi:10.1097/MLR.0000000000001418
- 217 4. McCormack LA, Madlock-Brown C. Social Determinant of Health Documentation Trends
218 and Their Association with Emergency Department Admissions. *AMIA Annu Symp Proc*
219 *AMIA Symp.* 2020;2020:823-832.
- 220 5. Perry BL, Aronson B, Pescosolido BA. Pandemic precarity: COVID-19 is exposing and
221 exacerbating inequalities in the American heartland. *Proc Natl Acad Sci.*
222 2021;118(8):e2020685118. doi:10.1073/pnas.2020685118
- 223 6. Anderson KN, Radhakrishnan L, Lane RI, et al. **Changes and Inequities in Adult Mental**
224 **Health–Related Emergency Department Visits During the COVID-19 Pandemic in the**
225 **US.** *JAMA Psychiatry.* 2022;79(5):475. doi:10.1001/jamapsychiatry.2022.0164

- 226 7. Holland KM, Jones C, Vivolo-Kantor AM, et al. Trends in US Emergency Department Visits
227 for Mental Health, Overdose, and Violence Outcomes Before and During the COVID-19
228 Pandemic. *JAMA Psychiatry*. 2021;78(4):372. doi:10.1001/jamapsychiatry.2020.4402
- 229 8. Smalley CM, Malone DA, Meldon SW, et al. The impact of COVID-19 on suicidal ideation
230 and alcohol presentations to emergency departments in a large healthcare system. *Am J Emerg
231 Med*. 2021;41:237-238. doi:10.1016/j.ajem.2020.05.093
- 232 9. Zangani C, Ostinelli EG, Smith KA, et al. Impact of the COVID-19 Pandemic on the Global
233 Delivery of Mental Health Services and Telemental Health: Systematic Review. *JMIR Ment
234 Health*. 2022;9(8):e38600. doi:10.2196/38600
- 235 10. Tweed EJ, Leyland AH, Morrison D, Katikireddi SV. Premature mortality in people affected
236 by co-occurring homelessness, justice involvement, opioid dependence, and psychosis: a
237 retrospective cohort study using linked administrative data. *Lancet Public Health*.
238 2022;7(9):e733-e743. doi:10.1016/S2468-2667(22)00159-1
- 239 11. Wray CM, Tang J, López L, Hoggatt K, Keyhani S. Association of Social Determinants of
240 Health and Their Cumulative Impact on Hospitalization Among a National Sample of
241 Community-Dwelling US Adults. *J Gen Intern Med*. 2022;37(8):1935-1942.
242 doi:10.1007/s11606-021-07067-y
- 243 12. Molina MF, Pantell MS, Gottlieb LM. Social Risk Factor Documentation in Emergency
244 Departments. *Ann Emerg Med*. 2023;81(1):38-46. doi:10.1016/j.annemergmed.2022.07.027
- 245 13. Wallace AS, Luther BL, Sisler SM, Wong B, Guo JW. Integrating social determinants of
246 health screening and referral during routine emergency department care: evaluation of reach
247 and implementation challenges. *Implement Sci Commun*. 2021;2(1):114. doi:10.1186/s43058-
248 021-00212-y
- 249 14. Davis CI, Montgomery AE, Dichter ME, Taylor LD, Blosnich JR. Social determinants and
250 emergency department utilization: Findings from the Veterans Health Administration. *Am J
251 Emerg Med*. 2020;38(9):1904-1909. doi:10.1016/j.ajem.2020.05.078
- 252 15. Braveman PA, Egerter SA, Mockenhaupt RE. Broadening the FocusThe Need to Address the
253 Social Determinants of Health. *Am J Prev Med*. 2011;40(1):S4-S18.
254 doi:10.1016/j.amepre.2010.10.002
- 255 16. American Medical Association. Social determinants of health and medical coding: What to
256 know. Accessed August 25, 2022. [https://www.ama-assn.org/practice-management/cpt/social-](https://www.ama-assn.org/practice-management/cpt/social-determinants-health-and-medical-coding-what-know)
257 [determinants-health-and-medical-coding-what-know](https://www.ama-assn.org/practice-management/cpt/social-determinants-health-and-medical-coding-what-know).

258

259

260

Table 1. ED visits of categories of historical diagnoses before and after shelter-in-place orders

Measure	Before shelter in place (5/2/2019 – 12/31/2019)	After shelter in place (5/2/2020 – 12/31/2020)	Relative Change P-value	Absolute Change Relative Risk	Absolute Change p-value
	N (%)	N (%)			
<i>Overall MH, SUD, and SDOH diagnoses</i>					
Mental Health Disorders	19,394 (4.1)	23,100 (5.8)	<0.001	1.19 (1.17, 1.21)	<0.001
Substance Use Disorders	15,420 (3.3)	19,444 (4.9)	<0.001	1.26 (1.23,1.29)	<0.001
Mental Health or Substance Use	32,320 (6.8)	38,711 (9.7)	<0.001	1.20 (1.18,1.22)	<0.001
Social Determinants of Health	955 (0.20)	2,260 (0.57)	<0.001	2.37 (2.19,2.55)	<0.001
Behavior Health or Substance Use Disorders and Social Determinants of Health	200 (0.04)	662 (0.17)	<0.001	3.31 (2.83,3.88)	<0.001
<i>Mental Behavioral & Neurodevelopmental Disorders</i>					
Mental disorders due to known physiological conditions	2,238 (0.47)	2,745 (0.69)	<0.001	1.23 (1.16,1.30)	<0.001
Schizophrenia, schizotypal, delusional, and other non-mood psychotic disorders	3,175 (0.67)	4,464 (1.1)	<0.001	1.41 (1.34,1.47)	<0.001
Mood [affective] disorders	6,983 (1.5)	7,242 (1.8)	<0.001	1.04 (1.00,1.07)	0.030
Anxiety, dissociative, stress-related, somatoform and other non-psychotic mental disorders	7,515 (1.6)	9,735 (2.4)	<0.001	1.30 (1.26,1.33)	<0.001
Behavioral syndromes associated with physiological disturbances and physical factors	291 (0.06)	304 (0.08)	0.008	1.04 (0.89,1.23)	0.59
Disorders of adult personality and behavior	411 (0.09)	581 (0.15)	<0.001	1.41 (1.25,1.60)	<0.001

Table 1. ED visits of categories of historical diagnoses before and after shelter-in-place orders

Measure	Before shelter in place (5/2/2019 – 12/31/2019)	After shelter in place (5/2/2020 – 12/31/2020)	Relative Change P-value	Absolute Change Relative Risk	Absolute Change p-value
	N (%)	N (%)			
Intellectual disabilities	153 (0.03)	207 (0.05)	<0.001	1.35 (1.10,1.67)	0.005
Pervasive and specific developmental disorders	233 (0.05)	402 (0.10)	<0.001	1.73 (1.47,2.03)	<0.001
Behavioral and emotional disorders with onset usually occurring in childhood	552 (0.12)	768 (0.19)	<0.001	1.39 (1.25,1.55)	<0.001
<i>Substance Use Disorders</i>					
Alcohol related disorders	7,247 (1.5)	8,341 (2.1)	<0.001	1.15 (1.12,1.19)	<0.001
Opioid related disorders	1,074 (0.23)	1,239 (0.31)	<0.001	1.15 (1.06,1.25)	<0.001
Cannabis related disorders	1,263 (0.27)	2,362 (0.59)	<0.001	1.87 (1.75,2.00)	<0.001
Sedative, hypnotic, or anxiolytic related disorders	175 (0.04)	217 (0.05)	<0.001	1.24 (1.02,1.51)	0.034
Cocaine related disorders	1,243 (0.26)	1,267 (0.32)	<0.001	1.02 (0.94,1.10)	0.63
Other stimulant related disorders	416 (0.09)	561 (0.14)	<0.001	1.35 (1.19,1.53)	<0.001
Hallucinogen related disorders	317 (0.07)	330 (0.08)	0.006	1.04 (0.89,1.21)	0.61
Nicotine dependence	2,940 (0.62)	5,219 (1.3)	<0.001	1.78 (1.70,1.86)	<0.001
Other psychoactive substance related disorders	2,788 (0.59)	3,074 (0.77)	<0.001	1.10 (1.05,1.16)	<0.001
<i>Social Determinants of Health</i>					
Problems related to education and literacy	9 (0.00)	6 (0.00)	0.66	0.67 (0.24,1.87)	0.44
Problems related to employment and unemployment	15 (0.00)	24 (0.01)	0.046	1.60 (0.84,3.05)	0.15
Problems related to housing and economic circumstances	420 (0.09)	1,051 (0.26)	<0.001	2.50 (2.23,2.80)	<0.001
Problems related to social environment	60 (0.01)	121 (0.03)	<0.001	2.02 (1.48,2.75)	<0.001

Table 1. ED visits of categories of historical diagnoses before and after shelter-in-place orders

Measure	Before shelter in place (5/2/2019 – 12/31/2019)	After shelter in place (5/2/2020 – 12/31/2020)	Relative Change P-value	Absolute Change Relative Risk	Absolute Change p-value
	N (%)	N (%)			
Other problems related to primary support group, including family circumstances	310 (0.07)	897 (0.23)	<0.001	2.89 (2.54,3.29)	<0.001
Problems related to other psychosocial circumstances	112 (0.02)	157 (0.04)	<0.001	1.40 (1.10,1.79)	0.006
<i>Disease Systems</i>					
Diseases of the Nervous System	20,831 (4.4)	21,560 (5.4)	<0.001	1.03 (1.02,1.05)	<0.001
Diseases of the Circulatory System	50,527 (10.7)	59,105 (14.8)	<0.001	1.17 (1.16,1.18)	<0.001
Diseases of the Respiratory System	63,045 (13.3)	44,616 (11.2)	<0.001	0.71 (0.70,0.72)	<0.001
Diseases of the Digestive System	42,809 (9.0)	42,061 (10.6)	<0.001	0.98 (0.97,1.00)	0.010
Diseases of the Skin	22,499 (4.7)	19,665 (4.9)	<0.001	0.87 (0.86,0.89)	<0.001
Diseases of the Musculoskeletal System	57,844 (12.2)	63,968 (16.1)	<0.001	1.11 (1.09,1.12)	<0.001
Diseases of the Genitourinary System	57,574 (12.1)	58,011 (14.6)	<0.001	1.01 (1.00,1.02)	0.20
<i>Suicide/Self-harm</i>					
Suicidal ideation	2,140 (0.45)	2,366 (0.59)	<0.001	1.11 (1.04,1.17)	<0.001
Suicide attempt and self-inflicted harm	4,153 (0.88)	4,070 (1.02)	<0.001	0.98 (0.94,1.02)	0.36
Opioid specific overdose	473 (0.10)	329 (0.08)	0.009	0.70 (0.60,0.80)	<0.001
Transgender Status	11 (0.00)	35 (0.01)	<0.001	3.18 (1.62,6.26)	<0.001
Severe Presentation	6,133 (1.3)	6,277 (1.6)	<0.001	1.02 (0.99,1.06)	0.20

Table 2. Admission rates of categories of historical diagnoses before and after shelter-in-place orders

Measure	Before shelter in place (5/2/2019 – 12/31/2019)		After shelter in place (5/2/2020 – 12/31/2020)		Odds Ratio	p-value
	Encounters (N)	Admissions N (%)	Encounters N	Admissions N (%)		
<i>Overall MH, SUD, and SDOH diagnoses</i>						
Mental Health Disorders	19,374	7,244 (37.4)	23,087	8,140 (35.3)	0.91 (0.88, 0.95)	<0.001
Substance Use Disorders	15,398	5,653 (36.7)	19,425	6,453 (33.2)	0.86 (0.82,0.90)	<0.001
Mental Health or Substance Use Disorders	32,282	11,851 (36.7)	38,680	13,174 (34.1)	0.89 (0.86,0.92)	<0.001
Social Determinants of Health	955	155 (16.2)	2,259	260 (11.5)	0.67 (0.54,0.83)	<0.001
Behavior Health or Substance Use and Social and Economic Disadvantage	200	32 (16.0)	662	73 (11.0)	0.65 (0.42,1.02)	0.061
<i>Mental Behavioral & Neurodevelopmental Disorders</i>						
Mental disorders due to known physiological conditions	2,235	1,288 (57.6)	2,745	1,610 (58.7)	1.04 (0.93,1.17)	0.47
Schizophrenia, schizotypal, delusional, and other non-mood psychotic disorders	3,171	1,152 (36.3)	4,462	1,481 (33.2)	0.87 (0.79,0.96)	0.004
Mood [affective] disorders	6,972	3,064 (43.9)	7,236	3,212 (44.4)	1.02 (0.95,1.09)	0.60
Anxiety, dissociative, stress-related, somatoform and other non-psychotic mental disorders	7,511	2,207 (29.4)	9,726	2,501 (25.7)	0.83 (0.78,0.89)	<0.001
Behavioral syndromes associated with physiological disturbances and physical factors	291	193 (66.3)	304	185 (60.9)	0.79 (0.56,1.10)	0.17
Disorders of adult personality and behavior	411	139 (33.8)	580	172 (29.7)	0.82 (0.63,1.08)	0.16
Intellectual disabilities	153	74 (48.4)	207	82 (39.6)	0.70 (0.46,1.07)	0.098
Pervasive and specific developmental disorders	233	98 (42.1)	402	121 (30.1)	0.59 (0.42,0.83)	0.002
Behavioral and emotional disorders with onset usually occurring in childhood	552	202 (36.6)	768	207 (27.0)	0.64 (0.51,0.81)	<0.001
<i>Substance Use Disorders</i>						

Table 2. Admission rates of categories of historical diagnoses before and after shelter-in-place orders

Measure	Before shelter in place (5/2/2019 – 12/31/2019)		After shelter in place (5/2/2020 – 12/31/2020)		Odds Ratio	p-value
	Encounters (N)	Admissions N (%)	Encounters N	Admissions N (%)		
Alcohol related disorders	7,237	2,630 (36.3)	8,336	3,098 (37.2)	1.04 (0.97,1.11)	0.29
Opioid related disorders	1,074	428 (39.9)	1,238	461 (37.2)	0.90 (0.76,1.06)	0.20
Cannabis related disorders	1,261	319 (25.3)	2,361	587 (24.9)	0.98 (0.83,1.14)	0.77
Sedative, hypnotic, or anxiolytic related disorders	175	85 (48.6)	217	95 (43.8)	0.82 (0.55,1.23)	0.34
Cocaine related disorders	1,238	462 (37.3)	1,267	507 (40.0)	1.12 (0.95,1.32)	0.17
Other stimulant related disorders	416	97 (23.3)	558	138 (24.7)	1.08 (0.80,1.46)	0.61
Hallucinogen related disorders	317	30 (9.5)	330	43 (13.0)	1.43 (0.87,2.35)	0.15
Nicotine dependence	2,933	1,612 (55.0)	5,212	1,925 (36.9)	0.48 (0.44,0.53)	<0.001
Other psychoactive substance related disorders	2,785	855 (30.7)	3,071	942 (30.7)	1.00 (0.89,1.12)	0.98
<i>Social Determinants of Health</i>						
Problems related to education and literacy	9	5 (55.6)	6	2 (33.3)	0.40 (0.05,3.42)	0.40
Problems related to employment and unemployment	15	1 (6.7)	24	3 (12.5)	2.00 (0.19,21.23)	0.57
Problems related to housing and economic circumstances	420	52 (12.4)	1,051	119 (11.3)	0.90 (0.64,1.28)	0.57
Problems related to social environment	60	36 (60.0)	120	69 (57.5)	0.90 (0.48,1.69)	0.75
Other problems related to primary support group, including family circumstances	310	18 (5.8)	897	25 (2.8)	0.47 (0.25,0.86)	0.016
Problems related to other psychosocial circumstances	112	46 (41.1)	157	46 (29.3)	0.59 (0.36,0.99)	0.046
<i>Disease Systems</i>						
Diseases of the Nervous System	20,798	11,024 (53.0)	21,531	11,529 (53.5)	1.02 (0.98,1.06)	0.26
Diseases of the Circulatory System	50,424	32,519 (64.5)	59,003	34,179 (57.9)	0.76 (0.74,0.78)	<0.001
Diseases of the Respiratory System	63,004	19,574 (31.1)	44,572	17,801 (39.9)	1.48 (1.44,1.51)	<0.001
Diseases of the Digestive System	42,747	16,633 (38.9)	42,017	17,063 (40.6)	1.07 (1.04,1.10)	<0.001

Table 2. Admission rates of categories of historical diagnoses before and after shelter-in-place orders

Measure	Before shelter in place (5/2/2019 – 12/31/2019)		After shelter in place (5/2/2020 – 12/31/2020)		Odds Ratio	p-value
	Encounters (N)	Admissions N (%)	Encounters N	Admissions N (%)		
Diseases of the Skin	22,494	5,238 (23.3)	19,655	5,292 (26.9)	1.21 (1.16,1.27)	<0.001
Diseases of the Musculoskeletal System	57,817	9,110 (15.8)	63,946	9,818 (15.4)	0.97 (0.94,1.00)	0.053
Diseases of the Genitourinary System	57,529	23,281 (40.5)	57,952	24,565 (42.4)	1.08 (1.06,1.11)	<0.001
<i>Suicide/Self-harm</i>						
Suicidal ideation	2,139	582 (27.2)	2,366	824 (34.8)	1.43 (1.26,1.62)	<0.001
Suicide attempt and self-inflicted harm	4,150	908 (21.9)	4,065	852 (21.0)	0.95 (0.85,1.05)	0.31
Opioid specific overdose	471	47 (10.0)	329	38 (11.6)	1.18 (0.75,1.85)	0.48
Transgender Status	11	5 (45.5)	34	13 (38.2)	0.74 (0.19,2.93)	0.67
Severe Presentation	6,129	1,417 (23.1)	6,272	1,612 (25.7)	1.15 (1.06,1.25)	<0.001

Supplemental Table 1: Counts and frequencies of patients with historical diagnoses of interest

Diagnosis	Total (N=487,028)
Overall MH, SUD, and SDOH diagnoses	
Mental Health Disorders	49,742 (10.2)
Substance Use Disorders	24,549 (5.0)
Social Determinants of Health,	2,572 (0.53)
Mental Behavioral & Neurodevelopmental Disorders	
Mental disorders due to known physiological conditions,	4,134 (0.85)
Schizophrenia, schizotypal, delusional, and other non-mood psychotic disorders,	4,212 (0.86)
Mood [effective] disorders,	11,084 (2.3)
Anxiety, dissociative, stress-related, somatoform and other non-psychotic mental disorders,	14,208 (2.9)
Behavioral syndromes associated with physiological disturbances and physical factors,	541 (0.11)
Disorders of adult personality and behavior,	702 (0.14)
Intellectual disabilities,	250 (0.05)
Pervasive and specific developmental disorders,	481 (0.10)
Behavioral and emotional disorders with onset usually occurring in childhood,	1,152 (0.24)
Substance Use Disorders	
Alcohol related disorders,	10,052 (2.1)
Opioid related disorders,	1,897 (0.39)
Cannabis related disorders,	3,277 (0.67)
Sedative, hypnotic, or anxiolytic related disorders,	379 (0.08)
Other stimulant related disorders,	873 (0.18)
Hallucinogen related disorders,	474 (0.10)
Nicotine dependence,	7,699 (1.6)
Other psychoactive substance related disorders,	4,678 (0.96)
Social Determinants of Health	
Problems related to education and literacy,	15 (0.00)
Problems related to employment and unemployment,	39 (0.01)
Occupational exposure to risk factors,	45 (0.01)
Problems related to housing and economic circumstances,	802 (0.16)
Problems related to social environment,	177 (0.04)
Problems related to upbringing,	12 (0.00)
Other problems related to primary support group, including family circumstances	1,157 (0.24)
Problems related to certain psychosocial circumstances,	4 (0.00)
Problems related to other psychosocial circumstances,	263 (0.05)
Disease Systems	
Neoplasms,	13,424 (2.8)
Diseases of the blood and blood-forming organs and certain disorders involving immune mechanism,	31,867 (6.5)
Endocrine, nutritional and metabolic diseases,	82,877 (17.0)
Diseases of the Nervous System,	35,789 (7.3)
Diseases of the eye and adnexa,	14,523 (3.0)

Supplemental Table 1: Counts and frequencies of patients with historical diagnoses of interest

Diagnosis	Total (N=487,028)
Diseases of the ear and mastoid process,	14,830 (3.0)
Diseases of the Circulatory System,	80,081 (16.4)
Diseases of the Respiratory System,	85,219 (17.5)
Diseases of the Digestive System,	68,523 (14.1)
Diseases of the Skin and subcutaneous tissue,	35,984 (7.4)
Diseases of the musculoskeletal system and connective tissue,	98,638 (20.3)
Diseases of the genitourinary system,	84,660 (17.4)
Pregnancy, childbirth and the puerperium,	676 (0.14)
Certain conditions originating in the perinatal period,	1,002 (0.21)
Congenital malformations, deformations and chromosomal abnormalities,	2,155 (0.44)
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified,	267,956 (55.0)
Injury, poisoning and certain other consequences of external causes,	155,944 (32.0)
Codes for special purposes,	11,244 (2.3)
External causes of morbidity,	21,455 (4.4)
Factors influencing health status and contact with health services,	101,058 (20.7)
<i>Suicide/Self-harm</i>	
Suicidal ideation,	3,956 (0.81)
Suicide attempt and self-inflicted harm,	7,879 (1.6)
Suicide death by underlying cause of death,	48 (0.01)
Suicide deaths by method,	48 (0.01)
Overdoses,	0 (0.00)
Opioid specific overdose,	732 (0.15)
Transgender Status,	46 (0.01)
Severe Presentation,	11,244 (2.3)
Behavior Health/Substance Abuse and Social and Economic Disadvantage,	525 (0.11)

*Data not available for all subjects. Missing values: Other problems related to primary support group, including family circumstances = 485,871. Statistics presented as N (column %).

Supplemental Table 2. Demographics of patients with and without social determinants of health

Factor	Total (N=487,028)	No Coded Social Determinant of Health (N=484,456)	Coded Determinants of Health (N=2,572)	p-value
Patient Race, *				<0.001 ^c

Supplemental Table 2. Demographics of patients with and without social determinants of health

Factor	Total (N=487,028)	No Coded Social Determinant of Health (N=484,456)	Coded Determinants of Health (N=2,572)	p-value
AMERICAN INDIAN/ALASKA NATIVE	611 (0.13)	609 (0.13)	2 (0.08)	
ASIAN	4,549 (0.97)	4,524 (0.97)	25 (1.04)	
BLACK	132,895 (28.3)	131,905 (28.3)	990 (41.3)	
MULTIRACIAL/MULTICULTURAL	22,721 (4.8)	22,485 (4.8)	236 (9.8)	
OTHER	1,374 (0.29)	1,369 (0.29)	5 (0.21)	
WHITE	306,850 (65.4)	305,708 (65.5)	1,142 (47.6)	
Patient Gender, *				<0.001^c
FEMALE	262,222 (53.8)	261,028 (53.9)	1,194 (46.4)	
MALE	224,754 (46.2)	223,376 (46.1)	1,378 (53.6)	
NONBINARY	18 (0.00)	18 (0.00)	0 (0.00)	
Ethnic Group Name, *				<0.001^c
African American	132,494 (27.4)	131,505 (27.3)	989 (38.8)	
Asian/Pacific Islander	4,597 (0.95)	4,571 (0.95)	26 (1.02)	
Caucasian	291,401 (60.3)	290,287 (60.4)	1,114 (43.7)	
Hispanic	36,939 (7.6)	36,717 (7.6)	222 (8.7)	
Multi-Racial	12,031 (2.5)	11,895 (2.5)	136 (5.3)	
Native American	609 (0.13)	607 (0.13)	2 (0.08)	
Other	5,465 (1.1)	5,403 (1.1)	62 (2.4)	
Hispanic Race, *				0.027^c
Hispanic	36,939 (7.8)	36,717 (7.8)	222 (8.9)	
Not Hispanic	438,901 (92.2)	436,641 (92.2)	2,260 (91.1)	
Current Age, mean ± sd*	46.0 ± 24.0	46.1 ± 23.9	28.7 ± 26.0	<0.001^a
Age at Admission, mean ± sd*	44.9 ± 24.0	45.0 ± 23.9	27.6 ± 25.8	<0.001^a
ED Disposition - Admit/Transfer, *	111,010 (22.8)	110,491 (22.8)	519 (20.2)	0.001^c
Insurance,				<0.001^c
Government (Medicare/Medicaid) + Self-Pay	151,708 (31.1)	150,620 (31.1)	1,088 (42.3)	
Private + Other inc. CCHS	335,320 (68.9)	333,836 (68.9)	1,484 (57.7)	

*Data not available for all subjects. Missing values: Patient Race = 18,028; Patient Gender = 34; Ethnic Group Name = 3,492; Hispanic Race = 11,188; Current Age = 78; Age at Admission = 78; ED Disposition - Admit/Transfer = 599.

Statistics presented as Mean ± SD, N (column %).

p-values: a=Satterthwaite t-test, c=Pearson's chi-square test.