1	Full title: High prevalence of asymptomatic and sub-patent <i>Plasmodium falciparum</i> infections in
2	Urban Bouaké, Côte d'Ivoire, but no hrp2 deletions: Implications for Malaria Control
3	
4	Short title: The hidden burden of malaria infections in urban Côte d'Ivoire
5	
6	Author names: André Barembaye Sagna ^{1,2,3} , Yilekal Gebre ² , Claudia Abigail Vera-Arias ² ,
7	Dipomin François Traoré ^{3,4} , Bertin N'cho Tchekoi ³ , Serge Brice Assi ⁵ , Amnan Alphonsine
8	Koffi ³ , Christophe Rogier ^{6,7} , Franck Remoue ¹ , Cristian Koepfli ²
9	
10	Affiliations
11	1) MIVEGEC, University of Montpellier, IRD, CNRS, Montpellier, France
12	2) Department of Biological Sciences and Eck Institute for Global Health, University of Notre,
13	Dame, Notre Dame, IN, USA
14	3) Institut Pierre Richet, Institut National de la Santé Publique, Bouaké, Côte d'Ivoire
15	4) Université Alassane Ouattara, Bouaké, Côte d'Ivoire
16	5) Programme National de Lutte Contre le Paludisme en Côte d'Ivoire, Ministère de la Santé et de
17	l'Hygiène Publique, Abidjan, Côte d'Ivoire.
18	6) Primum vitare, Paris, France
19	7) Institut Pasteur de Madagascar, Antananarivo, Madagascar
20	
21	Corresponding authors:
22	André B. SAGNA, andre.sagna@ird.fr and Cristian Koepfli, ckoepfli@nd.edu

23 Abstract:

Asymptomatic *Plasmodium falciparum* infections are common in endemic settings, vet 24 diagnosing these infections remains challenging because they are often below the limit of 25 detection of conventional light microscopy (LM) or rapid diagnostic tests (RDTs). Deletions of 26 the hrp2/3 gene, encoding the protein detected by the most sensitive class of RDTs, present a 27 further threat. In this study, the prevalence of asymptomatic and sub-patent P. falciparum 28 29 infections was characterized in the rainy season in the city of Bouaké, Central Côte d'Ivoire. A cross-sectional survey was conducted in nine neighborhoods of Bouaké, Côte d'Ivoire, in 30 2016. Matched LM, RDTs, and varATS gPCR, were used to determine the prevalence of P. 31 32 falciparum infections and to compare the performance of the different diagnostic methods. P. falciparum hrp2/3 deletions were typed by digital PCR. Among 2313 individuals, malaria 33 prevalence was 10.8% by microscopy, 13.0% by RDT, 35.1% by qPCR, and 37.3% (863/2313) 34 when all three methods were combined. 96.4% (832/863) of individuals with P. falciparum 35 infections did not report a recent fever. 57.1% (493/863) of infections were sub-patent. The 36 prevalence was highly heterogeneous across the neighborhoods, ranging from 15% to 55.2%, and 37 differed among age groups (<5 years: 17.7%, 5-14 years: 42.7%, \geq 15 years: 35.9%). Compared 38 to varATS qPCR, LM and RDT had a low sensitivity of 24.3% and 34.2%, respectively, although 39 40 both methods were highly specific (>96%). Sub-microscopic malaria infections were more prevalent in \geq 15-year-olds (69.9%, 349/499) and in the neighborhood of Dar es Salam (75.9%, 41 44/58). No https://deletions.were observed, and two samples carried *https*/deletion/wild-type mixed 42 infections. The high prevalence of asymptomatic and sub-patent infections prompts the 43 implementation of strategies targeting these parasite reservoirs to achieve reductions in malaria 44 burden in this high-transmission city of Côte d'Ivoire. 45

46 Introduction

The overall number of malaria infections is undeniably much higher than the estimated 241 47 million clinical cases (1) in 2020, owing to the large proportion of asymptomatic infections 48 among people living in endemic areas (2). Asymptomatic (or subclinical) malaria infection refers 49 to malarial parasitemia of any density in the absence of fever or other acute symptoms in 50 individuals who have not received recent antimalarial treatments (3). While some asymptomatic 51 52 infections may have parasitemia levels that are detectable by light microscopy (LM) or rapid diagnostic tests (RDTs), others can only be detected by molecular methods and are termed sub-53 patent infections (4). Asymptomatic and sub-patent individuals remain untreated in countries 54 55 where control focuses on clinical malaria cases and serve as reservoirs for onward transmission (5.6). Ouantifying the extent of submicroscopic infections and the underlying true prevalence of 56 parasites in different settings is central to understanding the dynamics of malaria transmission 57 and adapting control measures accordingly. 58 The asymptomatic reservoir presents a major challenge for control, with up to 95% of 59 transmission stemming from asymptomatic carriers (7–9). Numerous approaches are currently 60 being trialed or implemented to shrink the asymptomatic reservoir, such as mass screening and 61 treatment (10,11), reactive case detection (12), focal mass drug administration (13), or seasonal 62 63 malaria chemoprophylaxis (14). The success of control strategies for conducting active screening for infections relies heavily on 64 the sensitivity of the diagnostic test used. These programs typically use RDTs, the only 65

66 diagnostic method that can readily be used away from health centers. Yet, the sensitivity of RDTs

67 is limited. Compared to highly sensitive PCR, RDTs often miss more than half of all infections

(15). The most sensitive RDTs for *P. falciparum* detect the HRP2 and HRP3 proteins. Deletions 68 of the hrp2/3 genes result in false-negative RDTs, even if parasite density is high. Initially 69 reported in Peru (16), the frequency of hrp2/3 is high in the Horn of Africa (17–20), and deletions 70 were also reported from Ghana (21.22). Surveillance of hrp 2/3 is needed to decide whether 71 HRP2-based RDTs can be used for *P. falciparum* diagnosis. 72 The epidemiology of urban malaria in Côte d'Ivoire is not well documented, with few studies 73 74 conducted in the southern (23-25) and northern parts of the country (26). These studies found a high prevalence of infection by microscopy, ranging from 12% in the north to more than 60% in 75 the south. This difference in infection prevalence could mainly be explained by climatic factors. 76 77 The climate is tropical (Sub-Sudanese) in the north with two main seasons, and equatorial in the south with two rainy seasons and two dry seasons (27). Other factors such as age groups (5-14 78 years), traveling to rural areas, and agriculture practices have also been found to be risk factors 79 for malaria infections in urban areas (23-26). 80 Bouaké, in central Côte d'Ivoire, is the second-most populous city in the country, with more than 81

750,000 inhabitants in 2015. Like many cities in sub-Saharan Africa, urban agriculture is 82 widespread within and around the city to increase food security. This agricultural activity has 83 been shown to significantly impact malaria transmission with the creation of numerous larval 84 85 habitats and vector proliferation (28-30). Bouaké is a highly endemic area with >160,000 malaria cases recorded each year (annual incidence >200 per 1000 inhabitants) and more than 100 deaths 86 (31). Bouaké suffered from an armed conflict and a sociopolitical crisis from 2000 to 2011, 87 leading to large population moves, environmental modifications, and interruption in the 88 implementation and/or maintenance of malaria control strategies for the whole period (32,33). 89

90 Following the crisis, the population has reinvested in the city and resettled in formerly abandoned

- 91 neighborhoods. Only a few epidemiological studies have been conducted in the city of Bouaké,
- 92 and the real burden of malaria is underestimated, with no published data available on the
- 93 prevalence of asymptomatic and submicroscopic malaria infections. Its characterization will help
- 94 the national malaria control program (NMCP) adapt control measures for a more efficient use of
- 95 limited resources.
- 96 The objectives of this study were to measure the burden of asymptomatic and sub-patent *P*.
- 97 *falciparum* infections by LM, RDT, and qPCR in the highly endemic city of Bouaké and to
- 98 investigate potential *P. falciparum hrp2* and/or *hrp3* deletion.

99 Materials and Methods

100 Ethical approval and source of funding

101 Funding was provided by the French Initiative 5% and Expertise France, Grant N°12INI210. This

study followed the ethical principles recommended by the Edinburgh revision of the Declaration

103 of Helsinki and was approved by the Ethics Committee of the Ministry of Health of Côte d'Ivoire

104 (June 2015; No. 029/MSLS/CNER-dkn) and the University of Notre Dame Institutional Review

105 Board (18-08-4803). Before collecting samples, all participants over the age of 18 provided

106 written informed consent, as did the parent/guardian of each participant under the age of 18.

107 Study area and demographic data collection

108 In Côte d'Ivoire, malaria transmission occurs all year round, with a peak during the rainy season.

109 The annual incidence rates vary between 100 and >500 cases per 1000 inhabitants, with more

- than 80% of the whole Ivoirian population living in areas with an annual incidence rate \geq 300 per
- 111 1000 inhabitants (31). Despite a national coverage of long-lasting insecticidal nets (LLINs)

112	estimated at 95% since 2015	the number of	of malaria-related	cases has	been increa	sing since	e 2016
		, the mannoor (Ji manana ionatoa	cubeb mub	occur mercu	Shing Shines	2 2010

- from 3,754,504 to 7,295,068 cases in 2021 (34). The number of malaria-related deaths has
- decreased during the same period, from 3,340 to 1,276 deaths.
- 115 The survey was conducted in three health districts of the city of Bouaké: Northwestern-Bouaké,
- 116 Northeastern-Bouaké, and Southern-Bouaké. All health districts had a high coverage rate (>80%)
- 117 of standard pyrethroid-based LLINs. This area has year-round malaria transmission, with a peak
- 118 during the wet season (April–October). P. falciparum is the dominant species identified in more

than 95% of malaria cases (35). The Anopheles gambiae species complex is the dominant vector

120 species, with *An. funestus* and *An. nili* also present. The local vector populations are highly

resistant to almost all classes of insecticides used for vector control (36,37).

122 A cross-sectional study was performed from August 1st to August 27th, 2016, in nine

neighborhoods of the city (three neighborhoods per health district): Dar-es-Salam,

124 Djezoukouamekro, and N'gattakro in Northwestern-Bouaké; Sokoura, Belle-ville, and Attienkro

125 in Northeastern-Bouaké; Kennedy, Air-France, and N'gouattanoukro in Southern-Bouaké. This

126 study was part of a large operational research project named PALEVALUT

127 (<u>http://www.mesamalaria.org/mesa-track/palevalut</u>), which aimed to evaluate the effectiveness of

128 malaria control resources currently deployed and identify the barriers to their effectiveness in five

129 different sub-Saharan countries (Benin, Cameroon, Côte d'Ivoire, Niger, and Madagascar). In

130 each selected neighborhood, 50 households were randomly selected, and all members who were

131 at home were invited to participate in the study. After enrollment, all households were geo-

132 located using GPS (Garmin, Switzerland) instruments with up to 5 m of accuracy. A socio-

demographic questionnaire was completed with the head of the family to collect information

about the sanitary conditions of the household and household characteristics (house type and 134 construction materials, bathroom type, access to public services, etc.). Information on age, sex, 135 history of fever, LLIN use, and knowledge of malaria was recorded for each participant. 136 Case definitions 137 An asymptomatic malaria infection was defined as a *P. falciparum* positive test result in an 138 afebrile individual (temperature $<37.5^{\circ}$ C) at the time of sampling with no history of fever in the 139 140 previous 48 hours. A clinical malaria case was defined as a positive test result with a history of fever or a temperature >37.5 °C. Sub-patent infections were defined as infections that were only 141 positive by *var*ATS qPCR. 142 **Blood sample collection** 143 During the house visit, blood samples were collected from each study participant using a 1 mL 144 insulin syringe (Luer-Lok[™], BD, Franklin Lakes, NJ, USA) to prepare thick and thin blood 145 smears (both thick and thin smears were collected on the same slide), one RDT, and two spots on 146 Whatman qualitative filter paper N°3. The remaining blood was poured into a 1.5-mL 147 microcentrifuge tube. At the laboratory, microcentrifuge tubes were centrifuged, and plasma was 148 removed and stored at -20°C. The red cell pellets were also stored at -20°C until DNA extraction. 149 All samples of each study participant were labeled with a unique code and stored accordingly 150

151 until use.

152 **RDT-based diagnosis and treatment**

The Pf/HRP2-based RDT (Standard Diagnostics Inc.; Yongin, Republic of Korea) recommended
by the NMCP of Côte d'Ivoire was used in this study. This RDT is based on histidine-rich protein
2 (HRP2) detection and is specific to *P. falciparum* (Pf). All febrile individuals that tested

- 156 positive for RDT were treated with anti-malarials according to the national drug policy. Those
- 157 found positive by microscopy or qPCR were not treated because microscopy examination and
- 158 qPCR assays were not performed at the point of collection.
- 159 Microscopic examination
- 160 Giemsa-stained peripheral blood smears were examined under a compound microscope
- 161 (Olympus CX21 LED) at 1000× magnification to detect malaria parasites by trained technicians.
- 162 Ring, trophozoite, or gametocyte stages were identified and assessed for parasite density
- 163 (parasites/µl blood) against 200 white blood cells (WBCs), and densities were calculated
- 164 considering an average of 8000 WBC/µl.
- 165 All positive slides and 10% of randomly chosen negative blood slides were cross-examined by
- 166 independent laboratory personnel as part of a standard protocol to monitor examination quality.
- 167 Microscopists who examined the participants' blood slides were not aware of the RDT results.

168 DNA extraction and qPCR detection of P. falciparum infections

- 169 DNA was extracted from 100 µL of blood using the Genomic DNA Extraction Kit (Macherey-
- 170 Nagel, Düren, Germany) and eluted in 100 μ L of elution buffer. 4 μ L of extracted DNA was
- 171 screened for *P. falciparum* using ultrasensitive qPCR that amplifies a conserved region of the var
- gene acidic terminal sequence (*var*ATS) according to a previously published protocol (38). The
- 173 qPCR results were converted to parasites/ μ L using an external standard curve of ten-fold serial
- dilutions (5-steps) of 3D7 *P. falciparum* parasites quantified by droplet digital PCR (ddPCR)
- 175 (39,40).

176 P. falciparum hrp2/3 typing by digital PCR

Samples positive for *P. falciparum* at a density of approximately >5 parasites/ μ L by *var*ATS qPCR were typed for *hrp2* and *hrp3* deletion by dPCR (41). To this end, the assay was successfully transferred from the BioRad QX 200 droplet digital PCR system to the Qiagen QIAcuity dPCR system. Assay conditions were maintained as previously published (41). In this assay, *hrp2* or *hrp3* and a control gene (*serine-tRNA ligase*) were quantified in a single tube with very high specificity, thus providing highly accurate data on deletion status. The assay amplified parts of *hrp2* exon 2 that encode for the antigen detected by the RDT.

184 Data analysis

Field data were collected and stored using the Open Data Kit toolkit. Electronic data was 185 exported to MS Excel for analysis. A data summary was done using descriptive statistics. 186 Malaria prevalence was calculated for each diagnostic test (light microscopy and RDT) and 187 compared to the gold standard of *var*ATS qPCR using McNemar's or Cochran's test. The 188 comparison of malaria prevalence by age group, gender, neighborhood, symptoms, and LLIN use 189 was done using the Pearson χ^2 test or Fisher's exact test, as appropriate. Differences in parasite 190 densities between age groups were determined by ANOVA's Dunn's multiple comparisons test. 191 The limits of detection (LoD) of light microscopy (LM) and RDT, defined as the lowest parasite 192 density that would be detected with 95% probability, were determined by logistic regression 193 194 based on *var*ATS qPCR parasite density. Univariate and multivariate logistic regression models were used to investigate the association between sub-patent malaria infections and risk factors. 195 Variables significant at a p-value of 0.30 in the univariate logistic regression were selected for the 196 multivariate logistic regression analysis model. For all statistical tests, a 5% level of significance 197 was used. Odds ratios with 95% confidence intervals (CI) were reported. All statistical analyses 198

- 199 were performed using Graph Pad Prism 9.5 (Graph Pad Software, San Diego, CA, USA) and
- 200 MedCalc 20.218 (MedCalc Software Ltd., Ostend, Belgium).
- 201 **Results**
- 202 Study population demography
- 203 Matched LM, RDT, and varATS qPCR data were available for 2313 individuals (Figure 1). The
- 204 median age was 18 years (interquartile range: 8-33). Females represented 57.2% of the study
- 205 population. The majority of individuals (60.4%) reported using an LLIN within the last 15 days,
- which varied between neighborhoods (Table 1).

207 Table 1: Social demographic and clinical characteristics of the study population: Bouaké,

208 Côte d'Ivoire, 2016

Characteristics	Participants (N=2313)
Age	
<5 years	345 (14.9%)
5-14 years	646 (27.9%)
≥15 years	1322 (57.2%)
Median age [interquartile range]	18.0 [8.0 - 33.0]
Female sex	1324 (57.2%)
Neighborhood	
Attienkro	310 (13.4%)
Belle-ville	315 (13.6%)
Sokoura	320 (13.8%)
Dar es Salam	257 (11.1%)
Djézoukouamékro	192 (08.3%)
Ngattakro	220 (09.5%)
Kennedy	173 (07.5%)
Air-France	204 (08.8%)
Ngouatanoukro	322 (13.9%)
Bed net use (last two weeks, n=2090)	
Every night	1263 (60.4%)
Sometimes or never	827 (39.6%)
Clinical characteristics	
Cases of fever	66 (2.8%)
Positive (with RDT)	300 (13.0%)
Positive (with microscopy)	249 (10.8%)
Positive (with varATS qPCR)	812 (35.1%)
Total positive	863 (37.3%)
Total clinical malaria (with fever)	31/66 (47.0%)
Total asymptomatic cases (without fever)	832/863 (96.4%)
Total sub-patent infections	493/863 (57.1%)
Symptomatic, sub-patent infections (with fever)	9/493 (1.8%)
Asymptomatic, sub-patent infections (without fever)	484/493 (98.2%)
Symptomatic, sub-patent infections (with fever) among all positives	9/863 (1.0%)
Asymptomatic, sub-patent infections (without fever) among all positives	484/863 (56.1%)

209 Symptomatic, asymptomatic and sub-patent P. falciparum infections

Among the full study population, 37.3% (863/2313) of participants were diagnosed positive by
LM, RDT, and qPCR combined (Table 1). Figure 2 shows the distribution of *P. falciparum*

212 infections according to the three diagnostic methods used. The prevalence of *P. falciparum*

infections varied significantly according to the diagnostic method and was 10.8% (249/2313) by

214 LM, 13% (300/2313) by RDT, and 35.1% (812/2313) by qPCR (Cochran's Q= 823.9, DF= 2, p<

215 0.001). Of all *P. falciparum* infections, 96.4% (832/863) were classified as asymptomatic and

216 57.1% (493/863) as sub-patent (Table 1, Figure 2). Sixty-six (66) persons reported fever or a

history of fever within 48 hours, of which 31 (47%) were positive by LM, RDT, and qPCR

218 combined. The number of clinical malaria cases was variable depending on the diagnostic

219 method and was 17 by LM, 20 by RDT, and 27 by qPCR. Nine clinical malaria cases were

220 missed by both LM and RDT (Figure 2).

221 *P. falciparum* infection prevalence differed significantly among age groups, sex, neighborhoods,

and symptoms (Table 2). The prevalence of infections peaked in 5–14-year-olds (irrespective of

diagnostic method, all p < 0.0001). Although malaria prevalence tended to be lower in individuals

 ≥ 15 years compared to ≤ 5 years by LM and RDT, it was 2-fold higher in individuals ≥ 15 years

compared to <5 years by qPCR. Prevalence was higher in males compared to females by all three

diagnostic methods (all p < 0.05; **Table 2**). The prevalence was highest in Attienkro, health

district of Northwestern Bouaké (28.7% by LM, 19.0% by RDT, and 55.2% by qPCR). The

lowest prevalence by LM (1.6%) and qPCR (15.0%) was recorded in Sokoura, health district of

229 Northwestern Bouaké, and by RDT (5.1%) in Dar es Salam, health district of Northeastern

230 Bouaké (Table 2). Aside from the neighborhood of Attienkro, health district of Northeastern

Bouaké, where prevalence was higher for LM (28.7%) compared to RDT (19.0%, p< 0.001), all

- neighborhoods displayed an equal or higher prevalence by RDT as compared to LM. qPCR
- 233 detected higher numbers of infections than LM or RDT in all neighborhoods (all p < 0.001).
- 234 The prevalence of *P. falciparum* infections was significantly higher in individuals who reported
- 235 fever or a history of fever within 48 hours compared to afebrile individuals by LM and RDT (all
- 236 p < 0.0001), but not by qPCR (p = 0.316).
- 237 The overall parasite density expressed as geometric mean parasite density by qPCR was 40
- parasites/µL (95% CI: 33.7-47.5, Figure 3A). Individuals who were positive by all three tests had
- a significantly higher parasite density (856.8 parasites/ μ L; 95% CI: 707.3–1,038) than those who
- 240 were positive by *var*ATS qPCR only (13.7 parasites/ μ L; 95% CI: 11.4–16.3) (p< 0.0001) (Figure
- **3A**). Parasite density varied significantly by age group and was lower in \geq 15-year-olds (27.2)
- 242 parasites/µL; 95% CI: 22.2-33.5) compared to <5-year-olds (73.5 parasites/µL; 95% CI: 36.9-
- 243 146.2; p=0.004) and 5-14-year-olds (76.5 parasites/µL; 95% CI: 55.9-104.7; p< 0.001, Figure
- 244 **3B**).

Characteristics		N	Tested positive by LM	p-value	Tested positive by RDT	p-value	Tested positive by qPCR	p-value
	<5	345	30 (8.7%)	<0.0001	42 (12.2%)	<0.0001	61 (17.7%)	<0.0001
Age group	5-14	646	118 (18.3%)		153 (23.7%)		276 (42.7%)	
	≥15	1322	101 (7.6%)		105 (7.9%)		475 (35.9%)	
Candar	Male	989	123 (12.4%)	0.026	148 (15.0%)	0.014	376 (38.0%)	0.012
Gender	Female	1324	126 (9.5%)	- 0.026	152 (11.5%)	- 0.014	436 (32.9%)	
	Attienkro	310	89 (28.7%)		59 (19.0%)		171 (55.2%)	<0.0001
	Belle-ville	315	29 (9.2%)	<0.0001	39 (12.4%)	<0.0001	103 (32.7%)	
	Sokoura	320	5 (1.6%)		16 (8.4%)		48 (15.0%)	
	Dar es Salam	257	9 (3.5%)		13 (5.1%)		58 (22.6%)	
Neighborhood	Djézoukouamékro	192	27 (14.1%)		36 (18.8%)		94 (49.0%)	
	Ngattakro	220	16 (7.3%)		17 (7.7%)		65 (29.5%)	
	Kennedy	173	13 (7.5%)		26 (15.0%)		69 (39.9%)	
	Air-France	204	22 (10.8%)		36 (17.6%)		81 (39.7%)	
	Ngouatanoukro	322	39 (12.1%)		58 (18.0%)		123 (38.2%)	
	Febrile	66	17 (25.8%)	- <0.0001	20 (30.3%)	- <0.0001	27 (40.9%)	- 0.316
Symptoms	Afebrile	2247	232 (10.3%)		280 (12.5%)		785 (34.9%)	
Bed net use (last	Every night	1263	147 (11.6%)	0.279	166 (13.1%)	- 0.854	460 (36.4%)	- 0.727
two weeks)	Sometimes/never	827	86 (10.4%)	0.378	111 (13.4%)		295 (35.7%)	

245 Table 2: Prevalence of *P. falciparum* infections by microscopy, RDT and *var*ATS qPCR (N=2313).

247 Specificity and sensitivity of microscopy and RDT

- 248 Compared to the gold standard qPCR, the sensitivity of RDT and LM was low at 34.24% (95%
- 249 CI: 30.97–37.61) and 24.34% (95% CI: 21.47–27.49), respectively (Table 3). However, both
- 250 methods were highly specific: 98.53% (95% CI: 97.79–99.08) for RDT, and 96.60% (95% CI:
- 251 95.56–97.46) for LM. The measurement of agreement (Kappa) was high between RDT and
- 252 microscopy (60.6%, p< 0.001).
- 253 Table 3: Test performance of the different diagnostic assays used for evaluating *P. falciparum*
- 254 infection

		R	DT	LM		
		Positive	Negative	Positive	Negative	
V ATC DOD	Positive	278	534	198	614	
varA18 qPCR	Negative	22	1479	51	1450	
Sensitivity (95% CI)		34.24% (30.97	- 37.61)	24.38% (21.47 - 27.49)		
Specificity (95% CI)		98.53% (97.79 – 99.08)		96.60% (95.56 - 97.46)		
PPV % (95% CI)		92.66 (89.20 -	95.08)	79.51 (74.28 - 83.92)		
NPV % (95% CI)		73.48 (72.49 - 74.44)		70.26 (69.41 - 71.09)		
AUC (95% CI)		75.97 (74.17 –	77.69)	71.25 (69.36 - 73.09)		

255 RDT= Rapid Diagnostic Test; LM= Light Microscopy; PPV= Positive Predictive Value; NPV=

- 256 *Negative Predictive Value; AUC= Area Under the Curve*
- 257 The limit of detection (LoD), which is defined here as the lowest parasite density that could be
- detected with 95% probability, was 1,117 (95% CI: 968–1331) parasites/µL for LM and 972
- 259 (95% CI: 834–1173) parasites/µL for RDT. The LoD of the *var*ATS qPCR method was
- previously determined at 0.29 (95% CI: 0.14-2.30) parasites/ μ L (42).
- 261

262 Risk factors for sub-patent infections

- Age group, the neighborhood of residency, and absence of fever within 48 hours and at the time
- of sampling were associated with an increased risk of sub-patent malaria infection by univariate
- analysis (Table 4). All risk factors were tested in the univariate analysis, and only those with
- 266 $p \le 0.30$, such as age group, gender, neighborhood of residency, and symptoms, were selected and
- included in the multivariate analysis. Only age \geq 15 years (aOR: 1.09; 95% CI: 1.06 1.13; p<
- 268 0.0001), the neighborhood of residency, specifically Dar es Salam (aOR: 3.97; 95% CI: 1.91 -
- 269 8.27; p< 0.001), and absence of symptoms (aOR: 2.83; 95% CI: 1.22 6.51; p= 0.015), were
- significantly associated with the prevalence of sub-patent infections in multivariable logistic
- 271 regression analysis. The gender and bed net use did not correlate with sub-patent malaria
- infection prevalence (all p > 0.05) (**Table 4**).

Risk factor	Category	Malaria positive (n=863)	Sub-patent malaria (%)	Univariate cOR [95% CI]	p-value	Multivariate aOR [95% CI]	p-value
	<5	75	27 (36)	1		1	
Age group	5-14	289	117 (40.5)	1.21 [0.71-2.05]	0.479	1.0 [0.90-1.12]	0.922
	≥15	499	349 (69.9)	4.13 [2.49-6.88]	<0.0001	1.09 [1.06-1.13]	<0.0001
Candar	Male	402	218 (54.2)	1	- 0.108	1	0.204
Gender	Female	461	275 (59.7)	1.25 [0.95-1.64]		1.17 [0.87-1.55]	0.294
	Attienkro	202	95 (47.0)	1		1	
	Belle-ville	105	63 (60.0)	1.69 [1.05-2.73]	0.032	1.19 [1.06-1.33]	0.002
	Sokoura	50	31 (62.0)	1.84 [0.97-3.47]	0.060	1.27 [1.06-1.52]	0.007
	Dar es Salam	58	44 (75.9)	3.54 [1.83-6.86]	<0.001	3.97 [1.91-8.27]	<0.001
Neighborhood	Djézoukouamékro	96	56 (58.3)	1.58 [0.97-2.58]	0.069	1.28 [0.98-1.68]	0.074
	Ngattakro	65	44 (67.7	2.36 [1.31-4.25]	0.004	1.41 [1.13-1.76]	0.002
	Kennedy	72	46 (63.9)	2.00 [1.14-3.47]	0.015	1.15 [1.05-1.26]	0.004
	Air-France	86	47 (54.7)	1.36 [0.82-2.25]	0.237	1.10 [1.02-1.19]	0.012
	Ngouatanoukro	129	67 (51.9)	1.22 [0.78-1.89]	0.384	1.06 [1.00-1.12]	0.037
S-matom s	Febrile	31	9 (29.0)	1	0.003	1	0.015
Symptoms	Afebrile	832	484 (58.2)	3.34 [1.5-7.3]		2.83 [1.22-6.51]	
Bed net use (last	Every night	489	277 (56.6)	1	0.870	1	- 0.430
two weeks)	Sometimes/never	311	178 (57.2)	1.02 [0.77-1.37]	0.870	1.13 [0.84-1.53]	

273 Table 4: Prevalence and risk factors for sub-patent malaria infections in malaria infected individuals (N=863)

274 cOR: Crude odds ratio; aOR: Adjusted odds ratio; CI: Confidence interval;

275 *P. falciparum hrp2/3 deletion typing*

A total of 344 *P. falciparum*-positive samples were successfully typed for *hrp2* and *hrp3* 276 deletion. No hrp2 deletions were observed. Two samples carried hrp3 deletion/wild-type mixed 277 infections. The first was from an 8-year-old asymptomatic male living in the neighborhood of Air 278 France (health district of Southern Bouaké) who was negative by RDT but positive by LM and 279 gPCR, with a parasite density of 149 parasites/uL. The second was from a 9-year-old 280 281 asymptomatic female from the neighborhood of Ngattakro (health district of Northeastern Bouaké) and was also positive by LM and qPCR only, with a parasite density of 190 282 parasites/µL. 283

284 Discussion

In 2016, in Bouaké, approximately a third of apparently healthy city dwellers had an

asymptomatic malaria infection during the rainy (high transmission) season. More than half of

287 malaria infections were detectable only by highly sensitive PCR (sub-patent infections).

288 Microscopy and rapid diagnostic tests detected fewer than 40% of qPCR-positive infections.

289 While prevalence was high, it was lower than in 2014, when a study conducted in three

290 neighborhoods reported a malaria prevalence of around 80% by microscopy (35). This sharp fall

in malaria prevalence between 2014 and 2016 could be attributed to intensive malaria control

efforts by the NMCP after the end of the crisis that occurred from 2000 to 2011. Two rounds of

large-scale distribution of LLINs were done in 2011 and 2014, leading to a coverage and usage

rate of 95% and 68%, respectively (43). In addition, the wide availability of RDTs and

artemisinin-combination therapy (ACTs) for testing and treating symptomatic individuals helped

296 decrease the burden of malaria in Bouaké.

The prevalence of malaria infection was highly heterogeneous across the neighborhoods, ranging 297 from 15% to more than 55%. Likewise, the prevalence of sub-patent infections also varied across 298 neighborhoods. The ecological environment and Anopheles density variability between 299 neighborhoods could be considerable factors that explain this variation. Bouaké is characterized 300 by the presence of lowlands used for agricultural activities, depending on the neighborhood (44). 301 This agricultural activity has been shown to have a significant impact on vector diversity and 302 303 density (30) and consequently on malaria transmission (28,29), which is highly heterogeneous across neighborhoods. 304 The prevalence of malaria infection was higher in school-age children (5-14 years) compared to 305

306 younger ones, in line with previous studies (26,45). The high number of malaria infections detected in school-aged children raises concern because they serve as a source of onward parasite 307 transmission (46,47), as this age group has been shown to be the least likely to benefit from 308 universal malaria interventions such as bed nets and access to prompt diagnosis and treatment 309 (48,49). Implementation of strategies that specifically target this age group could help further 310 decrease malaria transmission in this highly endemic area and improve the health of 311 schoolchildren (50–52). The proportion of sub-patent infections was higher in individuals above 312 15 years of age, probably because of greater immunity in older children and adults induced by 313

314 cumulative exposure to the parasite (53).

The sensitivity of LM and RDT is low in areas where a substantial number of asymptomatic and low-density infections are present (54). In this study, the sensitivity of LM and RDTs was 24% and 34%, respectively, as compared to the *var*ATS qPCR. Highly sensitive RDTs (55) or loopmediated isothermal amplification (LAMP) (56) could be promising options to improve the

diagnosis of asymptomatic and sub-patent infections in areas where they are common. In
Northern Côte d'Ivoire, for example, LAMP has been shown to be a relevant field diagnostic
alternative to standard RDTs for identifying asymptomatic infections with a sensibility and
specificity of 93.3% (95% CI 85.7–100) and 95.4% (95% CI 92.2–100), respectively (56).

For the first time in Côte d'Ivoire, parasites were typed for *hrp2/3* deletions. No *hrp2* deletions were detected in the study population, so the use of Pf/*hrp2*-based RDTs for the diagnosis of *P*. *falciparum* is still recommended. Two samples carried an *hrp3*/wild-type mixed infection. *Hrp3* deletion is not a threat for HRP2-based RDTs as long as *hrp2* is present (57). Future monitoring of *hrp2/hrp3* deletion is warranted in samples that are positive by microscopy or LDH-based RDT, but negative by HRP2-based RDT.

330 Conclusion

The prevalence of *P. falciparum* infection was underestimated when microscopy and RDT were 331 used for diagnosis because a substantial proportion of asymptomatic and sub-patent malaria 332 carriers were present in the population living in the city of Bouaké and could only be diagnosed 333 using the ultra-sensitive PCR diagnostic method (varATS qPCR). The prevalence of malaria 334 infection was highly heterogeneous across the city, and older children and adults constitute the 335 336 reservoirs of the parasite. These hidden parasite reservoirs are a major challenge for the malaria control program in the city. Various strategies, such as intermittent preventive treatment, periodic 337 mass parasite screening and treatment, or mass drug administration, could be suggested to further 338 reduce the burden of malaria in this city. But before that, more follow-up cohort studies will be 339 needed to identify the role of asymptomatic cases in the dynamics of malaria transmission, its 340

- 341 seasonal variability, and its contribution to the incidence of malaria in the region, and to provide
- 342 support in planning an effective and appropriate malaria elimination strategy to implement.

343 **Competing interests**

344 The authors declare that they have no competing interests.

345 Acknowledgements

- 346 We gratefully acknowledge the populations of 9 neighborhoods (DAR, DJZ, NGA, SOK, BLV,
- 347 ATK, KEN, AIF, and NGO) of Bouaké, especially householders, housewives, guardians of
- 348 children, and all the technical staff involved in blood sample collection in the field, for their kind
- 349 support and collaboration. We would also like to thank Mahamadou Barro, PhD student in
- 350 mathematics, for his invaluable help in analyzing the data.

351 Authors' contributions

- 352 ABS, CK and FR conceived the study, ABS, DFT, SBA, TBN conducted field work and recorded
- demographic data; YG and CAVA conducted the lab work, ABS analyzed the data and wrote the
- 354 manuscript. AAK, CR and CK revised the manuscript. All authors read and approved the final
- 355 manuscript.

356 **References**

- World Health Organization. World malaria report 2021: Tracking progress against malaria.
 Geneva, Switzerland: World Health Organization; 2021.
- Bousema T, Okell L, Felger I, Drakeley C. Asymptomatic malaria infections: detectability, transmissibility and public health relevance. Nat Rev Microbiol. 2014 Dec;12(12):833–40.
- Lindblade KA, Steinhardt L, Samuels A, Kachur SP, Slutsker L. The silent threat:
 asymptomatic parasitemia and malaria transmission. Expert Review of Anti-infective
 Therapy. 2013 Jun 1;11(6):623–39.

- 4. Okell LC, Ghani AC, Lvons E, Drakelev CJ. Submicroscopic Infection in Plasmodium 364 falciparum-Endemic Populations: A Systematic Review and Meta-Analysis. The Journal of 365 Infectious Diseases. 2009 Nov 15;200(10):1509-17. 366 367 5. Schneider P, Bousema JT, Gouagna LC, Otieno S, van de Vegte-Bolmer M, Omar SA, et al. Submicroscopic Plasmodium falciparum gametocyte densities frequently result in mosquito 368 infection. Am J Trop Med Hyg. 2007 Mar;76(3):470-4. 369 6. Coleman RE, Kumpitak C, Ponlawat A, Maneechai N, Phunkitchar V, Rachapaew N, et al. 370 371 Infectivity of Asymptomatic Plasmodium-Infected Human Populations to Anopheles dirus Mosquitoes in Western Thailand. Journal of Medical Entomology. 2004 Mar 1;41(2):201-8. 372 7. Gonçalves BP, Kapulu MC, Sawa P, Guelbéogo WM, Tiono AB, Grignard L, et al. 373 Examining the human infectious reservoir for Plasmodium falciparum malaria in areas of 374 differing transmission intensity. Nat Commun. 2017 Oct 26;8(1):1133. 375 Ouédraogo AL, Gonçalves BP, Gnémé A, Wenger EA, Guelbeogo MW, Ouédraogo A, et al. 376 8. Dynamics of the Human Infectious Reservoir for Malaria Determined by Mosquito Feeding 377 378 Assays and Ultrasensitive Malaria Diagnosis in Burkina Faso. J Infect Dis. 2016 Jan 1;213(1):90-9. 379 9. Sumner KM, Freedman E, Abel L, Obala A, Pence BW, Wesolowski A, et al. Genotyping 380 cognate Plasmodium falciparum in humans and mosquitoes to estimate onward transmission 381 382 of asymptomatic infections. Nat Commun. 2021 Feb 10;12(1):909. 10. Conner RO, Dieye Y, Hainsworth M, Tall A, Cissé B, Faye F, et al. Correction to: Mass 383 testing and treatment for malaria followed by weekly fever screening, testing and treatment 384 in Northern Senegal: feasibility, cost and impact. Malar J. 2020 Nov 30;19(1):443. 385 11. Kim S, Luande VN, Rocklöv J, Carlton JM, Tozan Y. A Systematic Review of the Evidence 386 on the Effectiveness and Cost-Effectiveness of Mass Screen-and-Treat Interventions for 387 Malaria Control. Am J Trop Med Hyg. 2021 Sep 7;105(6):1722-31. 388 12. Stuck L, Fakih BS, Al-Mafazy AWH, Hofmann NE, Holzschuh A, Grossenbacher B, et al. 389 390 Malaria infection prevalence and sensitivity of reactive case detection in Zanzibar. Int J Infect Dis. 2020 Aug;97:337-46. 391 13. Hsiang MS, Ntuku H, Roberts KW, Dufour MSK, Whittemore B, Tambo M, et al. 392 Effectiveness of reactive focal mass drug administration and reactive focal vector control to 393
- Effectiveness of reactive focal mass drug administration and reactive focal vector control to
 reduce malaria transmission in the low malaria-endemic setting of Namibia: a cluster randomised controlled, open-label, two-by-two factorial design trial. Lancet. 2020 Apr
 25;395(10233):1361–73.

- 14. Cairns M, Ceesay SJ, Sagara I, Zongo I, Kessely H, Gamougam K, et al. Effectiveness of
 seasonal malaria chemoprevention (SMC) treatments when SMC is implemented at scale:
 Case-control studies in 5 countries. PLoS Med. 2021 Sep;18(9):e1003727.
- 400 15. Yimam Y, Mohebali M, Afshar MJA. Comparison of diagnostic performance between
 401 conventional and ultrasensitive rapid diagnostic tests for diagnosis of malaria: A systematic
 402 review and meta-analysis. PLOS ONE. 2022 févr;17(2):e0263770.
- 403 16. Gamboa D, Ho MF, Bendezu J, Torres K, Chiodini PL, Barnwell JW, et al. A large
 404 proportion of P. falciparum isolates in the Amazon region of Peru lack pfhrp2 and pfhrp3:
 405 implications for malaria rapid diagnostic tests. PLoS One. 2010 Jan 25;5(1):e8091.
- 406 17. Golassa L, Messele A, Amambua-Ngwa A, Swedberg G. High prevalence and extended
 407 deletions in Plasmodium falciparum hrp2/3 genomic loci in Ethiopia. PLoS One.
 408 2020;15(11):e0241807.
- 18. Boush MA, Djibrine MA, Mussa A, Talib M, Maki A, Mohammed A, et al. Plasmodium
 falciparum isolate with histidine-rich protein 2 gene deletion from Nyala City, Western
 Sudan. Sci Rep. 2020 Jul 30;10(1):12822.
- Iriart X, Menard S, Chauvin P, Mohamed HS, Charpentier E, Mohamed MA, et al.
 Misdiagnosis of imported falciparum malaria from African areas due to an increased
 prevalence of pfhrp2/pfhrp3 gene deletion: the Djibouti case. Emerg Microbes Infect. 2020
 Dec;9(1):1984–7.
- 20. Menegon M, L'Episcopia M, Nurahmed AM, Talha AA, Nour BYM, Severini C.
 Identification of Plasmodium falciparum isolates lacking histidine-rich protein 2 and 3 in Eritrea. Infect Genet Evol. 2017 Nov;55:131–4.
- Amoah LE, Abuaku B, Bukari AH, Dickson D, Amoako EO, Asumah G, et al. Contribution
 of P. falciparum parasites with Pfhrp 2 gene deletions to false negative PfHRP 2 based
 malaria RDT results in Ghana: A nationwide study of symptomatic malaria patients. PLoS
 One. 2020;15(9):e0238749.
- 423 22. Bredu DG, Ahadzi GK, Dickson D, Peprah NY, Asamoah A, Asumah GA, et al. Nationwide
 424 Surveillance of Pfhrp2 Exon 2 Diversity in Plasmodium falciparum Circulating in
 425 Symptomatic Malaria Patients Living in Ghana. Am J Trop Med Hyg. 2022 May
 426 2;106(6):1660–6.
- 427 23. Assi SB, Ouattara L, Adja AM, Bla KB, N'Cho TB, Lingué KN, et al. Epidemiology of
 428 malaria in urban area (Adzopé city, Côte d'Ivoire). Am-Euras J Sci Res. 2010;5(2):94–100.
- 429 24. Bassa FK, Ouattara M, Silué KD, Adiossan LG, Baikoro N, Koné S, et al. Epidemiology of
 430 malaria in the Taabo health and demographic surveillance system, south-central Côte
 431 d'Ivoire. Malar J. 2016 Jan 6;15(1):9.

432 25. Wang SJ, Lengeler C, Smith TA, Vounatsou P, Cissé G, Tanner M. Rapid Urban Malaria
433 Appraisal (RUMA) III: epidemiology of urban malaria in the municipality of Yopougon
434 (Abidjan). Malar J. 2006 Apr 4;5:29.

- 435 26. Gbalégba CGN, Ba H, Silué KD, Ba O, Tia E, Chouaibou M, et al. Distribution of
 436 Plasmodium spp. infection in asymptomatic carriers in perennial and low seasonal malaria
 437 transmission settings in West Africa. Infect Dis Poverty. 2018 Apr 25;7:39.
- 438 27. Kouakou KE, Moussa H, Kouassi AM, Goula BTA, Savane I. Redefinition of homogeneous
 439 climatic zones in Cote d'Ivoire in a context of climate change. IJSER. 2017;8(11):453–62.
- 28. Dossou-Yovo J, Doannio JM, Diarrassouba S, Chauvancy G. [The impact of rice fields on malaria transmission in the city of Bouaké, Côte d'Ivoire]. Bull Soc Pathol Exot.
 1998;91(4):327–33.
- 29. Dossou-Yovo J, Ouattara A, Doannio JM, Diarrassouba S, Chauvancy G. [Malaria surveys in a humid savannah region in Côte d'Ivoire]. Med Trop (Mars). 1998;58(1):51–6.
- 30. Adja AM, Zoh DD, Sagna AB, Kpan DMS, Guindo-Coulibaly N, Yapi A, et al. Diversity of
 Anopheles Gambiae s.l., Giles (Diptera: Culicidae) Larval Habitats in Urban Areas and
 Malaria Transmission in Bouaké, Côte d'Ivoire. Vector Borne Zoonotic Dis. 2021
 Aug;21(8):593–601.
- 449 31. Ministère de la Santé et de l'Hygiène Publique. Rapport Annuel sur la situation Sanitaire
 450 (RASS) 2016. Abidjan, Côte d'Ivoire. 2017;354.

32. Betsi NA, Koudou BG, Cissé G, Tschannen AB, Pignol AM, Ouattara Y, et al. Effect of an
armed conflict on human resources and health systems in Côte d'Ivoire: prevention of and
care for people with HIV/AIDS. AIDS Care. 2006 May;18(4):356–65.

- 454 33. Adja AM, Yobo MC. Study of malaria transmission risk factors in a time of military-political
 455 crisis in Bouake urban area (Ivory Coast). European Scientific Journal, ESJ [Internet]. 2015
 456 Jan 30 [cited 2022 Nov 3];11(3). Available from:
 457 https://eujournal.org/index.php/esj/article/view/4986
- 458 34. World Health Organization. World malaria report 2022. Geneva, Switzerland: World Health
 459 Organization; 2022.
- 460 35. Assi SB, Ouattara L, Tchiekoi BN, Toure OA, Remoue F, Zika KD, et al. Cross sectional
 461 study on prevalence of malaria case and malaria infection in children under 15 years after the
 462 military crisis in Bouake, Cte dIvoire. JPVB. 2022 Jul 31;14(1):8–17.
- 36. Camara S, Koffi AA, Ahoua Alou LP, Koffi K, Kabran JPK, Koné A, et al. Mapping
 insecticide resistance in Anopheles gambiae (s.l.) from Côte d'Ivoire. Parasit Vectors. 2018
 Jan 8;11(1):19.

37. Zoh DD, Ahoua Alou LP, Toure M, Pennetier C, Camara S, Traore DF, et al. The current
insecticide resistance status of Anopheles gambiae (s.l.) (Culicidae) in rural and urban areas
of Bouaké, Côte d'Ivoire. Parasit Vectors. 2018 Mar 2;11:118.

- 38. Hofmann N, Mwingira F, Shekalaghe S, Robinson LJ, Mueller I, Felger I. Ultra-sensitive
 detection of Plasmodium falciparum by amplification of multi-copy subtelomeric targets.
 PLoS Med. 2015 Mar;12(3):e1001788.
- 472 39. Koepfli C, Nguitragool W, Hofmann NE, Robinson LJ, Ome-Kaius M, Sattabongkot J, et al.
 473 Sensitive and accurate quantification of human malaria parasites using droplet digital PCR
 474 (ddPCR). Sci Rep. 2016 Dec 16;6:39183.
- 40. Oduma CO, Ogolla S, Atieli H, Ondigo BN, Lee MC, Githeko AK, et al. Increased
 investment in gametocytes in asymptomatic Plasmodium falciparum infections in the wet
 season. BMC Infect Dis. 2021 Jan 9;21(1):44.
- 478 41. Vera-Arias CA, Holzschuh A, Oduma CO, Badu K, Abdul-Hakim M, Yukich J, et al. High479 throughput Plasmodium falciparum hrp2 and hrp3 gene deletion typing by digital PCR to
 480 monitor malaria rapid diagnostic test efficacy. Elife. 2022 Jun 28;11:e72083.
- 481 42. Holzschuh A, Koepfli C. Tenfold difference in DNA recovery rate: systematic comparison of
 482 whole blood vs. dried blood spot sample collection for malaria molecular surveillance. Malar
 483 J. 2022 Mar 15;21:88.
- 484 43. Ministère de la Santé et de l'Hygiène Publique. Plan national de développement sanitaire
 2016–2020. Abidjan, Côte d'Ivoire. 2016;88.
- 486 44. Dahoui MMC, Adou KA, Coulibaly B, Niamien KL, Koné A, Cornelie S, et al.
 487 Entomological drivers of uneven malaria transmission in urban lowland areas in Bouaké,
 488 Côte d'Ivoire. Malaria Journal. 2023 Jan 31;22(1):34.
- 489 45. Mwesigwa J, Achan J, Di Tanna GL, Affara M, Jawara M, Worwui A, et al. Residual malaria
 490 transmission dynamics varies across The Gambia despite high coverage of control
 491 interventions. PLoS One. 2017;12(11):e0187059.
- 492 46. Walldorf JA, Cohee LM, Coalson JE, Bauleni A, Nkanaunena K, Kapito-Tembo A, et al.
 493 School-Age Children Are a Reservoir of Malaria Infection in Malawi. PLoS One.
 494 2015;10(7):e0134061.
- 495 47. Coalson JE, Cohee LM, Buchwald AG, Nyambalo A, Kubale J, Seydel KB, et al. Simulation
 496 models predict that school-age children are responsible for most human-to-mosquito
 497 Plasmodium falciparum transmission in southern Malawi. Malar J. 2018 Apr 3;17(1):147.

48. Olapeju B, Choiriyyah I, Lynch M, Acosta A, Blaufuss S, Filemyr E, et al. Age and gender trends in insecticide-treated net use in sub-Saharan Africa: a multi-country analysis. Malaria Journal. 2018 Nov 14;17(1):423.

- 49. Coalson JE, Cohee LM, Walldorf JA, Bauleni A, Mathanga DP, Taylor TE, et al. Challenges
 in Treatment for Fever among School-Age Children and Adults in Malawi. Am J Trop Med
 Hyg. 2019 Feb;100(2):287–95.
- 504 50. Cohee LM, Valim C, Coalson JE, Nyambalo A, Chilombe M, Ngwira A, et al. School-based
 505 screening and treatment may reduce P. falciparum transmission. Sci Rep. 2021 Mar
 506 25;11(1):6905.
- 507 51. Staedke SG, Maiteki-Sebuguzi C, Rehman AM, Kigozi SP, Gonahasa S, Okiring J, et al.
 508 Assessment of community-level effects of intermittent preventive treatment for malaria in
 509 schoolchildren in Jinja, Uganda (START-IPT trial): a cluster-randomised trial. The Lancet
 510 Global Health. 2018 Jun 1;6(6):e668–79.
- 511 52. Tada MS, Ferreira R de GM, Katsuragawa TH, Martha RCD, Costa JDAN, Albrecht L, et al.
 512 Asymptomatic infection with Plasmodium falciparum and Plasmodium vivax in the Brazilian
 513 Amazon Basin: to treat or not to treat? Mem Inst Oswaldo Cruz. 2012 Aug;107(5):621–9.
- 53. Carneiro I, Roca-Feltrer A, Griffin JT, Smith L, Tanner M, Schellenberg JA, et al. Agepatterns of malaria vary with severity, transmission intensity and seasonality in sub-Saharan
 Africa: a systematic review and pooled analysis. PLoS One. 2010 Feb 1;5(2):e8988.
- 54. Lo E, Zhou G, Oo W, Afrane Y, Githeko A, Yan G. Low parasitemia in submicroscopic
 infections significantly impacts malaria diagnostic sensitivity in the highlands of Western
 Kenya. PLoS One. 2015;10(3):e0121763.
- 55. Niyukuri D, Sinzinkayo D, Troth EV, Oduma CO, Barengayabo M, Ndereyimana M, et al.
 Performance of highly sensitive and conventional rapid diagnostic tests for clinical and
 subclinical Plasmodium falciparum infections, and hrp2/3 deletion status in Burundi. PLOS
 Global Public Health. 2022 juil;2(7):e0000828.
- 56. Benié EMA, Silué KD, Ding XC, Yeo I, Assamoi JB, Tuo K, et al. Accuracy of a rapid
 diagnosis test, microscopy and loop-mediated isothermal amplification in the detection of
 asymptomatic Plasmodium infections in Korhogo, Northern Côte d'Ivoire. Malar J. 2022 Apr
 2;21(1):111.
- 57. Kong A, Wilson SA, Ah Y, Nace D, Rogier E, Aidoo M. HRP2 and HRP3 cross-reactivity
 and implications for HRP2-based RDT use in regions with Plasmodium falciparum hrp2 gene
 deletions. Malar J. 2021 Apr 29;20(1):207.

532 Figure legends

533	Figure 1: Study workflow diagram. Data for analysis consists of participants for whom resul	ts
534	from all three tests were available.	

535

536 Figure 2: Distribution of *P. falciparum* infections by the three diagnostic methods.

537

- 538 Figure 3: P. falciparum density as determined by varATS qPCR. Panel A shows P.
- *falciparum* density in the whole study population and in samples tested positive (+) or negative

540 (-) by microscopy and/or RDT. Panel B shows P. falciparum density by age group. Dots

- represent the parasite density of each participant. Red bars indicate geometric means with 95%
- 542 confidence intervals. P-values from Dunn's test with Bonferroni adjustment for multiple pairwise
- 543 comparisons are given.



Figure



Figure



Figure