It is made available under a CC-BY-NC-ND 4.0 International license .

Association Between Major Depressive Disorder and the Development of

Cerebral Small Vessel Disease: A Mendelian Randomization Study

Ting Lu^{1†}, Lijun Luo^{2†}, Jie Yang^{2†}, Yueying Li³, Daiyi Chen¹, Haiyang Sun¹, Huijuan Liao¹, Wen

Zhao¹, Zhixuan Ren¹, Yan Xu¹, Shiyao Yu¹, Xiao Cheng^{1,4,5,6}*, Jingbo Sun^{1,4,5,6}*

¹The Second School of Clinical Medicine, Guangzhou University of Chinese Medicine,

Guangzhou 510405, China

²Department of Neurology, The First Hospital of Wuhan, Wuhan 430022, China

³The Affiliated Hospital of Shaanxi University of Chinese Medicine, Xianyang 712000,

China

⁴Department of Neurology, Guangdong Provincial Hospital of Traditional Chinese Medicine, Guangzhou 510120, China

⁵State Key Laboratory of Dampness Syndrome of Chinese Medicine, The Second Affiliated

Hospital of Guangzhou University of Chinese Medicine, Guangzhou 510120, China

⁶Guangdong Provincial Key Laboratory of Research on Emergency in TCM, Guangzhou,

510120 China

[†]These three authors contributed equally to this work.

*Corresponding authors:

Jingbo Sun (ORCID: 0000-0001-6609-5274)

Department of Neurology, Guangdong Provincial Hospital of Traditional Chinese Medicine

Guangzhou 510120, China

Telephone: +86-136-4262-7661

Email address: gdszyysjb@gzucm.edu.cn

Xiao Cheng (ORCID: 0000-0002-8264-9713)

Department of Neurology, Guangdong Provincial Hospital of Traditional Chinese Medicine

Guangzhou 510120, China

It is made available under a CC-BY-NC-ND 4.0 International license .

Email address: chengxiaolucky@126.com

Abstract

Background: Although observational studies indicate a complex, bidirectional association between major depressive disorder (MDD) and cerebral small vessel disease (CSVD), the results are frequently inconsistent. This study investigated the potential correlation of MDD with both CSVD clinical outcomes and imaging markers, utilizing a bidirectional Mendelian randomization (MR) study design.

Methods: Instrumental variables for both MDD and CSVD were extracted from the latest or most extensive genome-wide association study (GWAS) data available for each phenotype. Clinical outcomes and imaging markers of CSVD were defined using several parameters. The inverse variance weighting (IVW) method with additional sensitivity and heterogeneity analyses was used. Furthermore, a separate GWAS for depression was used to validate our significant findings.

Results: In the forward MR analyses, the genetically predicted risk of MDD was positively associated with two CSVD phenotypes showing microscopic white matter (WM) damage: mean diffusivity (IVW OR = 2.191, 95 % CI 1.226 to 3.917, p = 0.008) and WM-enlarged perivascular space (OR = 1.053 95 % CI 1.006 to 1.101, p = 0.026). The use of an independent database for depression yielded no significant risk of depression associated with these two CSVD traits. Furthermore, reverse MR analyses showed no evidence of reverse causality between MDD and an altered CSVD risk.

Conclusions: This study utilizing MR imaging findings supports a substantial causal association between MDD and CSVD-related indicators of impaired WM microstructure. It is necessary to exercise caution when extending these results to individuals with depression.

Keywords: major depressive disorder; depression; cerebral small vessel disease; Mendelian randomization; microscopic white matter damage

Introduction

Major depressive disorder (MDD) is a global public health concern with a substantial socio-economic burden. Almost 20 % of individuals experience at least one episode of MDD during their lifetime (Kupfer, Frank, & Phillips, 2012). Cerebral small vessel disease (CSVD) comprises several syndromes affecting the small vessels of the brain. The most acute manifestations of CSVD are intracerebral hemorrhage (ICH) and small vessel ischemic stroke (SVS) (Markus & de Leeuw, 2023), and the principal neuroimaging markers include lacunar infarcts, white matter hyperintensities (WMH), brain microbleeds (BMBs), enlarged perivascular space (PVS), and cerebral atrophy (Chen et al., 2019). Interestingly, three decades of accumulated research evidence suggests a complex bidirectional relationship between MDD and CSVD, with WMH being the most extensively researched neuroimaging indicator.

WMH correlates with the occurrence of depressive symptoms (de Groot et al., 2000; Desmarais et al., 2021). Compared to the normal control group, patients with MDD exhibited greater severity of deep WMH signals (Brown, Lewine, Hudgins, & Risch, 1992). WMH severity is a crucial predictor of future depression risk in patients with CSVD (Park et al., 2015; Jaroonpipatkul et al.,

It is made available under a CC-BY-NC-ND 4.0 International license .

2022; Ottavi, Pepper, Bateman, Fiorentino, & Brodtmann, 2023; F. Zhang, Ping, Jin, Hou, & Song, 2023). Specifically, those with extensive WMH at baseline are more likely to develop major depressive symptoms (Qiu et al., 2017). Increased mean diffusivity (MD) and decreased fractional anisotropy (FA) are two key indicators of assessing WM microstructural damage (Liao et al., 2013; Nobuhara et al., 2006; Taylor et al., 2004). Lower FA values in adolescents (Vulser et al., 2018) and higher MD values (Özel et al., 2023) are correlated with greater individual risk and occurrence of depressive symptoms, respectively.

Both lacunar infarcts (Direk et al., 2016; Özel et al., 2023) and BMBs (Direk et al., 2016; Xu et al., 2017) are associated with the occurrence of depressive symptoms and disorders. CSVD severity correlates with the diagnosis of depression before and after ICH (Castello et al., 2022). In patients with MDD, a significant correlation was observed between the number of traumatic events and overall PVS volume (Ranti et al., 2022). However, these observational studies may be subject to reverse causation and residual confounding factors, leading to conflicting results (Ahmed et al., 2022; Mewton et al., 2019). Therefore, the causative link between MDD and CSVD remains to be elucidated.

Mendelian randomization (MR) represents a potent, up-and-coming method that employs genetic variance as an instrumental variable (IV) to evaluate the causal impact of an exposure on a given outcome (Emdin, Khera, & Kathiresan, 2017; Sekula, Del Greco M, Pattaro, & Köttgen, 2016). Herein, we conducted a bidirectional two-sample MR analysis to comprehensively evaluate the relationship between MDD and clinical outcomes associated with CSVD (ICH or SVS, lobar ICH or SVS, and non-lobar ICH or SVS) and radiological markers of CSVD (WMH, FA, MD, BMBs, and PVS). In addition, we used a separate depression genome-wide association study (GWAS) to validate our significant findings.

Materials and methods

Study design

This study was conducted in accordance with the STROBE-MR checklist (Supplement file 1: Table S1) (Skrivankova et al., 2021). We described our rationale, study design, and procedures in Figure 1. All data used in this study were publicly available and therefore did not require ethics approval.

Data sources

Details of the GWAS data sets are listed in Table 1. For MDD, we used the latest and largest GWAS summary statistics from the UK Biobank (UKB), which analyzed 170,756 MDD cases and 329,443 controls, and the Psychiatric Genomics Consortium (PGC) GWAS datasets, which analyzed 8,483,301 variants (Howard et al., 2019). The definition of MDD used by the UKB was based on self-reported symptoms, treatment, or electronic records (Howard et al., 2018). The definition of MDD used by the PGC was based on structured diagnostic interviews (Wray et al., 2018). To analyze depression, data were obtained from the FinnGen Study Release 9 of the

It is made available under a CC-BY-NC-ND 4.0 International license .

GWAS for depression, F5_DEPRESSIO (Kurki et al., 2023), which included 43,280 patients and 329,192 controls. Depression was defined according to the International Classification of Diseases.

The CSVD clinical outcomes were obtained from a meta-analysis of ICH GWAS results by location and SVS GWAS results from the MEGASTROKE study, which comprised "all location ICH or SVS," "lobar ICH or SVS," and "non-lobar ICH or SVS." The study sample included 6,255 ICH or SVS cases and 233,058 controls of European ancestry (Chung et al., 2019).

The seven radiological markers of CSVD included WMH, FA, MD, BMBs, WM-PVS, basal ganglia (BG)-PVS, and hippocampal (HIP)-PVS.

WMH represents pathological changes in MRI scan results, whereas FA and MD demonstrate greater sensitivity to the disruption of normal function and structure than measuring pathology in isolation. GWAS data for three WM markers, WMH (N = 18,381), FA (N = 17,663), and MD (N = 17,467), were obtained from the UKB, Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE), and WMH-Stroke multi-ethnic studies (Persyn et al., 2020). The WMH volume was measured using T1, T2, and fluid-attenuated inversion recovery sequences. FA and MD data were obtained from diffusion tensor imaging measurements of 48 brain regions.

The BMBs GWAS data were obtained from 11 population-based cohorts and 3 stroke cohort studies with case-control or case-only designs (Knol et al., 2020). BMBs were defined as small, low-signal lesions identifiable on a susceptibility-weighted imaging or T2-weighted gradient echo sequence. The study sample included 3,556 case and 22,306 control individuals, of whom 94 % were of European ethnicity.

The PVS data in European populations were obtained from a meta-analysis of 18 cohorts, including 9,317 case and 29,281 control individuals for WM-PVS, 8,950 case and 29,953 control individuals for BG-PVS, and 9,163 case and 29,708 control individuals for HIP-PVS (Duperron et al., 2023). PVSs were defined as fluid-filled spaces with the same signal as the cerebrospinal fluid and no high signal rim on T2-weighted or FLAIR sequences. The PVS burden was quantitatively defined by different scales, including the visual semiquantitative rating scale, among others.

Selection of instruments

The filtering of instrumental variables (IVs) must fulfill three core assumptions (Figure 1). In the forward MR analysis, SNPs satisfying a genome-wide significance level of p < 5E-08 were initially selected (Zhang et al., 2023). For fewer than three remaining SNPs, a relaxed threshold of p < 1E-05 was used. Relaxing the statistical threshold for IVs has been employed in several MR studies in the psychiatric domain (Lee et al., 2023; Li et al., 2023; Pan et al., 2023). To exclude genetic variants with strong correlations, the clumping process was performed using the 1000 Genomes Project European reference panel (R2 < 0.001 and a window size of 10,000 kb). The SNPs and corresponding statistics were extracted from the CSVD GWAS dataset, deleting SNPs with minor allele frequency (MAF) < 0.01. Further, the MDD and CSVD data were harmonized by removing all palindromic SNPs. To satisfy the independence assumption, SNPs from the

It is made available under a CC-BY-NC-ND 4.0 International license .

PhenoScannerV2 database that were significantly associated with confounding variables (Kamat et al., 2019) were collected. SNPs directly associated with CSVD were also excluded. We used the same rigorous criteria for selecting IVs in the reverse MR analysis. The variance explained (R^2) by each SNP was calculated using the formula: $R^2 = 2 \times MAF \times (1-MAF) \times beta^2$ (Si et al., 2021). The F-statistic was utilized to assess the strength of the genetic instrument. SNPs with F-statistic values < 10 (implying weak IVs) were removed from subsequent analyses (Burgess & Thompson, 2011).

Statistical analysis

The primary analytic method used in this survey was the inverse variance weighting (IVW) method, which is known for its superior detection power (Burgess, Butterworth, & Thompson, 2013). Four other methods, namely weighted median, simple mode, weighted mode, and MR–Egger, were used for complementary analyses. For heterogeneity analysis, we used Cochran's Q test (Greco M, Minelli, Sheehan, & Thompson, 2015) and leave-one-out analysis (Wu et al., 2021) to assess the level of heterogeneity among SNPs. The possible presence of pleiotropic effect was evaluated by the intercept of the MR–Egger regression test (Bowden, Davey Smith, & Burgess, 2015) and MR pleiotropy residual sum and outlier (MR-PRESSO) test (Verbanck, Chen, Neale, & Do, 2018). We used the same statistical analysis in the reverse MR analysis. We calculated odds ratios (ORs) to measure the causal effect between MDD and CSVD phenotypes. Considering the partial sample overlap between MDD and WMH in the UKB, we conducted a bias and type I error analysis to evaluate the potential bias risk attributable to sample overlap. Given the exploratory nature of this study, we did not perform correction of multiple comparisons.

The statistical software R (version 4.1.0, R Foundation for Statistical Computing, Vienna, Austria; https://www.R-project.org) was used to perform all statistical analyses and visualize the results obtained. The software packages were TwoSampleMR, MR-PRESSO, and Forestplot.

Results

Genetic instrument strength and sample overlap

For the 10 genetically related MDD-CSVD pairs, we only performed an MR analysis on pairs containing at least two qualifying IVs. In the forward MR analysis, all 10 pairs had SNPs available for analysis. In the reverse MR, two phenotypes, BG-PVS and HIP-PVS, had no IV remaining after the removal of the weak instrument (even when a loose threshold of p < 1E-05 was used); hence, they were not included in the MR analysis. The range of SNPs utilized as genetic instruments varied from 3 to 41, accounting for 0.16–23.63 % of the phenotypic variance, with F-statistics > 10 (range 20.92–1350.80) for all phenotypes, indicating the good strength of the IVs (Supplement file 1: Table S2). Details of all the SNPs included in the MR analysis are provided in the supplementary file (Supplement file 1: Table S3). The risk of bias due to sample overlap in the assessment was minimal (< 0.01, regardless of the overlap proportion).

Causal relationship between MDD and CSVD

It is made available under a CC-BY-NC-ND 4.0 International license .

First, to comprehensively assess the effect of MDD on CSVD clinical outcomes and radiological markers, we used the IVW approach for the main analysis. Notably, as shown in the forest plot (Figure 2), we found a significant effect of MDD on two radiological markers of CSVD: MD and WM-PVS.

The IVW approach demonstrated that a genetically determined higher risk of MDD was associated with increased MD (OR = 2.191, 95 % CI 1.226 to 3.917, p = 0.008). MR–Egger (OR = 97.532, 95 % CI 6.110 to 1556.906, p = 0.002) and weighted median (OR = 2.274, 95 % CI 1.117 to 4.631, p = 0.024) results were consistent with the IVW results (Figure 2). The MR–Egger intercept indicated the absence of pleiotropy (intercept = -0.116, p = 0.009) (Table 2), which may have caused the MR–Egger method to produce an OR with a wide range of results. Cochrane's Q test showed no heterogeneity (Cochrane Q = 53.499, p = 0.061). The leave-one-out method showed no significant change in the IVW causal estimates after deleting any variant, suggesting the robustness of the results (Supplemental file 2: Figure S1). In addition, the MR-PRESSO test detected no significant outliers (p = 0.051).

A genetically determined higher risk of MDD was also associated with an increased risk of WM-PVS (OR = $1.053\ 95\$ % CI 1.006 to 1.101, p = 0.026) (Figure 2). The other four methods (MR–Egger, weighted median, simple mode, weighted mode) did not yield significant results (p > 0.05) (Supplemental file 1: Table S2). The sensitivity and heterogeneity analyses passed the test (all p > 0.05) (Table 2). The leave-one-out analyses indicated that the observed associations were stable (Supplemental 2: Figure S1).

For the other phenotypes, there was no evidence of an effect of MDD on the risk of clinical CSVD outcomes (all p > 0.05). There was also no evidence of an effect of MDD on the risk of WMH, FA, BMBs, BG-PVS, and HIP-PVS (Figure 2).

Depression data from the FinnGen database revealed no association between depression and the risk of MD (OR =1.084, 95 % CI 0.557 to 2.110, p = 0.812) and WM-PVS (OR = 0.872, 95 % CI 0.722 to 1.052, p = 0.151) by the IVW method (Supplemental 1: Table S4).

Causal relationship between CSVD and MDD

In reverse MR, we examined the effect of eight phenotypes with qualified IVs on the risk of MDD. There was no evidence that ICH or SVS, lobar ICH or SVS, non-lobar ICH or SVS, WMH, FA, MD, BMBs, and WM-PVS affected the genetically predicted risk of MDD (Figure 2). Because there were only three SNPs in WM-PVS, the MR PRESSO Global test was not performed. No horizontal pleiotropy or heterogeneity (all p > 0.05) was found for the remaining phenotypes except for the BMBs (Cochrane Q = 30.089, p = 0.037, MR PRESSO Global test p = 0.046), suggesting that the results are reliable (Table 2).

Discussion

It is made available under a CC-BY-NC-ND 4.0 International license .

To our knowledge, this is the first MR study to comprehensively investigate the causal association of MDD with CSVD clinical outcomes (ICH or SVS) and radiological markers (WMH, MD, FA, BMBs, and PVS). In forward MR, we found that the genetically predicted risk of MDD was positively associated with increased MD and the presence of WM-PVS. However, when the FinnGen dataset on depression was used, no genetically predicted risk of depression was found to be associated with these two CSVD traits (MD, WM-PVS), thus validating the study results. Additionally, reverse MR did not identify any causal association between CSVD risk and MDD. Since genetic variants are assigned to the offspring during meiosis, MR analyses can reflect lifelong exposure, reducing interference from reverse causation and confounding factors in comparison to observational studies. The aforementioned findings indicate a correlation between MDD and CSVD, implying that the genetically predicted risk of MDD could result in weakened microstructural integrity of cerebral WM. Consequently, curative measures or prevention of MDD may prove to be beneficial in averting future occurrences of CSVD.

Numerous epidemiological studies have investigated the intricate reciprocal association between MDD (or depression) and cerebral WM lesions (Gu et al., 2022; Pasi et al., 2016; Zhang et al., 2021). Patients with CSVD exhibiting depressive symptoms demonstrate a notable decrease in network efficiency and WM integrity compared to their counterparts without such symptoms (Lu et al., 2022; van Uden et al., 2015). The current MR study evaluated the mutually influential association of MDD with the WM macrostructure (WMH volume) and microstructure (FA and MD). Surprisingly, the analyses showed no evidence of a genetic association between MDD and WMH. This outcome is comparable to the results of two recent cohort studies: Clancy et al. discovered that heightened WMH was not linked to depression (2023), whereas Ali et al. illustrated that the load of WMH was not associated with depression 1 year after a stroke (2023). Another study, which implemented a machine learning algorithm, demonstrated that the severity of depression over time was unaffected by WMH volume (Ahmed et al., 2022).

Furthermore, the genetically predicted risk of MDD was associated with increased MD but not with FA. As WMH volume indicates the accumulation of WM damage over time, microstructural measurements may have greater sensitivity, implying that damage to microstructural integrity precedes WMH. The disparity between the two microscopic indicators could be attributed to MD being linked to blood-brain barrier (BBB) leakage, which occurs earlier in WMH progression and may be more receptive to identifying minor injuries. Conversely, FA seems to be affected more directly by myelin changes and may identify more severe injuries (Cubon, Putukian, Boyer, & Dettwiler, 2011; O'Dwyer et al., 2011; Song et al., 2003).

This study demonstrated that the MDD correlated positively with another form of WM microstructure damage: the WM-PVS. A possible explanation relates to perturbations in the central nervous system microenvironment in patients with MDD, including impaired BBB permeability (Yarlagadda, Alfson, & Clayton, 2009), leading to the recruitment and accumulation of blood immune cells and immune mediators around blood vessels (Winkler et al., 2021), resulting in PVS enlargement (Wuerfel et al., 2008).

A prior MR study explored the causal link between MDD and ischemic stroke, specifically SVS, and found that MDD was associated with a higher risk of SVS (Cai et al., 2019). We used a

It is made available under a CC-BY-NC-ND 4.0 International license .

cross-phenotypic GWAS (Chung et al., 2019) and found no effect of genetically predicted risk of MDD on CSVD clinical phenotypes (ICH or SVS, lobar ICH or SVS, non-lobar ICH or SVS). One possible explanation for this discrepancy in results is that Cai et al.'s study used a looser genome-wide significance threshold (p < 1E-06) to allow the inclusion of more SNPs that contributed to MDD risk. In contrast, we used a more stringent threshold of p < 5E-08 when examining the three pairwise associations between MDD and CSVD clinical phenotypes. Furthermore, a recent MR in 2023 examined the risk association between MDD and ICH and found no causal relationship (Wang et al., 2022). Because we used larger and more comprehensive GWAS datasets for both MDD and CSVD clinical phenotypes, our findings may provide more reliable evidence of an association between the two.

Our study had some limitations. First, the positive results regarding the causal relationship between MDD and CSVD were not replicated with the FinnGen depression data. This suggests an association between MDD rather than depression and increased susceptibility to CSVD; therefore caution is needed in extending the results to individuals with depression who do not meet clinical diagnostic criteria for MDD.

Second, the association between CSVD and late-life depression (LLD), based on the "vascular depression hypothesis," has been extensively studied previously (Aizenstein et al., 2016; Alexopoulos et al., 1997). However, in the current study, the lack of high-quality GWAS data on LLD limited more precise analyses of the association between depressive subtypes and CSVD. Future studies are expected to explore the complex bidirectional relationship between CSVD and depression subtypes using more extensive GWAS data.

Third, women have a significantly higher prevalence of depression (Kessler, McGonagle, Swartz, Blazer, & Nelson, 1993; Kuehner, 2017). An observational study found a significant association of depressive symptoms with periventricular WMH, deep WMH, and the number of asymptomatic infarcts in women but not in men (Wendell et al., 2010). Although the original GWAS dataset used was adjusted for sex, sex-stratified data were lacking. Further future studies that stratify differential effects by sex would be valuable.

Finally, despite the various sensitivity analyses, the possibility of uncontrolled pleiotropy or heterogeneity cannot be excluded. More large-scale prospective studies are needed in the future to verify the causal relationship between MDD and CSVD.

Conclusion

In conclusion, the association between the risk of MDD and CSVD may have far-reaching physiological value and clinical implications, as identified in this MR study. The implication of this potential association involves the proactive identification of individuals at risk for the development and progression of CSVD, leading to the implementation of prevention strategies and improved clinical outcomes.

It is made available under a CC-BY-NC-ND 4.0 International license .

Availability of data and materials

All data generated or analyzed in this study are available in this published article and its supplementary information files.

Financial support

This work was supported by the National Key Research and Development Program of China (grant number 2019YFC1708601); the Specific Fund of State Key Laboratory of Dampness Syndrome of Chinese Medicine (grant number SZ2021ZZ14); the Guangdong Provincial Key Laboratory of Research on Emergency in traditional Chinese medicine (TCM) (grant numbers 2017B030314176, 2018-75, and 2019-140); and the Wuhan Municipal Health Commission (grant numbers WZ21M01 and WX20C35).

Ethical standards

Not applicable.

Author contributions

TL, XC, and JBS conceptualized the study design. TL, LJL, JY, YYL, DYC, HYS, and HJL analyzed the data. WZ, ZXR, YX, and SYY verified the accuracy of the data. TL, LJL, and JY were major contributors to the writing of the manuscript. All authors read and approved the final version of the manuscript.

Competing interests

The authors declare that they have no competing interests.

It is made available under a CC-BY-NC-ND 4.0 International license .

References

- Ahmed, R., Ryan, C., Christman, S., Elson, D., Bermudez, C., Landman, B. A., ... Taylor, W. D. (2022). Structural MRI-Based Measures of Accelerated Brain Aging do not Moderate the Acute Antidepressant Response in Late-Life Depression. *The American Journal of Geriatric Psychiatry* : Official Journal of the American Association for Geriatric Psychiatry, 30(9), 1015–1025. https://doi.org/10.1016/j.jagp.2021.11.011
- Aizenstein, H. J., Baskys, A., Boldrini, M., Butters, M. A., Diniz, B. S., Jaiswal, M. K., ... Tene, O. (2016, November). Vascular depression consensus report - a critical update. *BMC Medicine*, Vol. 14, p. 161. https://doi.org/10.1186/s12916-016-0720-5
- Alexopoulos, G. S., Meyers, B. S., Young, R. C., Campbell, S., Silbersweig, D., & Charlson, M. (1997). "Vascular depression" hypothesis. Archives of General Psychiatry, 54(10), 915–922. https://doi.org/10.1001/archpsyc.1997.01830220033006
- Ali, H. F., Fast, L., Khalil, A., Siebert, E., Liman, T., Endres, M., ... Kufner, A. (2023). White matter hyperintensities are an independent predictor of cognitive decline 3 years following first-ever stroke-results from the PROSCIS-B study. *Journal of Neurology*, 270(3), 1637–1646. https://doi.org/10.1007/s00415-022-11481-5
- Bowden, J., Davey Smith, G., & Burgess, S. (2015). Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *International Journal of Epidemiology*, 44(2), 512–525. https://doi.org/10.1093/ije/dyv080
- Brown, F. W., Lewine, R. J., Hudgins, P. A., & Risch, S. C. (1992). White matter hyperintensity signals in psychiatric and nonpsychiatric subjects. *The American Journal of Psychiatry*, 149(5), 620–625. https://doi.org/10.1176/ajp.149.5.620
- Burgess, S., Butterworth, A., & Thompson, S. G. (2013). Mendelian randomization analysis with multiple genetic variants using summarized data. *Genetic Epidemiology*, 37(7), 658–665. https://doi.org/10.1002/gepi.21758
- Burgess, S., & Thompson, S. G. (2011). Avoiding bias from weak instruments in Mendelian randomization studies. *International Journal of Epidemiology*, 40(3), 755–764. https://doi.org/10.1093/ije/dyr036
- Cai, H., Cai, B., Zhang, H., Sun, W., Wang, Y., Zhou, S., ... Liang, J. (2019). Major depression and small vessel stroke: a Mendelian randomization analysis. *Journal of Neurology*, 266(11), 2859–2866. https://doi.org/10.1007/s00415-019-09511-w
- Castello, J. P., Pasi, M., Kubiszewski, P., Abramson, J. R., Charidimou, A., Kourkoulis, C., ... Biffi, A. (2022). Cerebral Small Vessel Disease and Depression Among Intracerebral Hemorrhage Survivors. *Stroke*, 53(2), 523–531. https://doi.org/10.1161/STROKEAHA.121.035488
- Chen, X., Wang, J., Shan, Y., Cai, W., Liu, S., Hu, M., ... Lu, Z. (2019). Cerebral small vessel disease: neuroimaging markers and clinical implication. *Journal of Neurology*, 266(10), 2347–2362. https://doi.org/10.1007/s00415-018-9077-3
- Chung, J., Marini, S., Pera, J., Norrving, B., Jimenez-Conde, J., Roquer, J., ... Anderson, C. D. (2019). Genome-wide association study of cerebral small vessel disease reveals established and novel

It is made available under a CC-BY-NC-ND 4.0 International license .

loci. Brain : A Journal of Neurology, 142(10), 3176–3189. https://doi.org/10.1093/brain/awz233

- Clancy, U., Radakovic, R., Doubal, F., Hernández, M. D. C. V., Maniega, S. M., Taylor, A. M., ... Wardlaw, J. M. (2023). Are neuropsychiatric symptoms a marker of small vessel disease progression in older adults? Evidence from the Lothian Birth Cohort 1936. *International Journal of Geriatric Psychiatry*, 38(1), e5855. https://doi.org/10.1002/gps.5855
- Cubon, V. A., Putukian, M., Boyer, C., & Dettwiler, A. (2011). A diffusion tensor imaging study on the white matter skeleton in individuals with sports-related concussion. *Journal of Neurotrauma*, 28(2), 189–201. https://doi.org/10.1089/neu.2010.1430
- de Groot, J. C., de Leeuw, F. E., Oudkerk, M., Hofman, A., Jolles, J., & Breteler, M. M. (2000). Cerebral white matter lesions and depressive symptoms in elderly adults. *Archives of General Psychiatry*, 57(11), 1071–1076. https://doi.org/10.1001/archpsyc.57.11.1071
- Desmarais, P., Gao, A. F., Lanctôt, K., Rogaeva, E., Ramirez, J., Herrmann, N., ... Masellis, M. (2021). White matter hyperintensities in autopsy-confirmed frontotemporal lobar degeneration and Alzheimer's disease. *Alzheimer's Research & Therapy*, *13*(1), 129. https://doi.org/10.1186/s13195-021-00869-6
- Direk, N., Perez, H. S., Akoudad, S., Verhaaren, B. F. J., Niessen, W. J., Hofman, A., ... Tiemeier, H. (2016). Markers of cerebral small vessel disease and severity of depression in the general population. *Psychiatry Research. Neuroimaging*, 253, 1–6. https://doi.org/10.1016/j.pscychresns.2016.05.002
- Duperron, M.-G., Knol, M. J., Le Grand, Q., Evans, T. E., Mishra, A., Tsuchida, A., ... Debette, S. (2023). Genomics of perivascular space burden unravels early mechanisms of cerebral small vessel disease. *Nature Medicine*, 29(4), 950–962. https://doi.org/10.1038/s41591-023-02268-w
- Emdin, C. A., Khera, A. V, & Kathiresan, S. (2017). Mendelian Randomization. *JAMA*, *318*(19), 1925–1926. https://doi.org/10.1001/jama.2017.17219
- Greco M, F. Del, Minelli, C., Sheehan, N. A., & Thompson, J. R. (2015). Detecting pleiotropy in Mendelian randomisation studies with summary data and a continuous outcome. *Statistics in Medicine*, 34(21), 2926–2940. https://doi.org/10.1002/sim.6522
- Gu, Y., Zhao, P., Feng, W., Xia, X., Tian, X., Yan, Y., ... Li, X. (2022). Structural brain network measures in elderly patients with cerebral small vessel disease and depressive symptoms. *BMC Geriatrics*, 22(1), 568. https://doi.org/10.1186/s12877-022-03245-7
- Howard, D. M., Adams, M. J., Clarke, T.-K., Hafferty, J. D., Gibson, J., Shirali, M., ... McIntosh, A. M. (2019). Genome-wide meta-analysis of depression identifies 102 independent variants and highlights the importance of the prefrontal brain regions. *Nature Neuroscience*, 22(3), 343–352. https://doi.org/10.1038/s41593-018-0326-7
- Howard, D. M., Adams, M. J., Shirali, M., Clarke, T.-K., Marioni, R. E., Davies, G., ... McIntosh, A. M. (2018). Genome-wide association study of depression phenotypes in UK Biobank identifies variants in excitatory synaptic pathways. *Nature Communications*, 9(1), 1470. https://doi.org/10.1038/s41467-018-03819-3
- Jaroonpipatkul, C., Onwanna, J., Tunvirachaisakul, C., Jittapiromsak, N., Rakvongthai, Y., Chutinet, A., ... Maes, M. (2022). Depressive symptoms due to stroke are strongly predicted by the volume

It is made available under a CC-BY-NC-ND 4.0 International license .

and location of the cerebral infarction, white matter hyperintensities, hypertension, and age: A precision nomothetic psychiatry analysis. *Journal of Affective Disorders*, *309*, 141–150. https://doi.org/10.1016/j.jad.2022.04.041

- Kamat, M. A., Blackshaw, J. A., Young, R., Surendran, P., Burgess, S., Danesh, J., ... Staley, J. R. (2019). PhenoScanner V2: an expanded tool for searching human genotype-phenotype associations. *Bioinformatics (Oxford, England)*, 35(22), 4851–4853. https://doi.org/10.1093/bioinformatics/btz469
- Kessler, R. C., McGonagle, K. A., Swartz, M., Blazer, D. G., & Nelson, C. B. (1993). Sex and depression in the National Comorbidity Survey. I: Lifetime prevalence, chronicity and recurrence. *Journal of Affective Disorders*, 29(2–3), 85–96. https://doi.org/10.1016/0165-0327(93)90026-g
- Knol, M. J., Lu, D., Traylor, M., Adams, H. H. H., Romero, J. R. J., Smith, A. V, … Markus, H. S. (2020). Association of common genetic variants with brain microbleeds: A genome-wide association study. *Neurology*, 95(24), e3331–e3343. https://doi.org/10.1212/WNL.000000000010852
- Kuehner, C. (2017). Why is depression more common among women than among men? *The Lancet*. *Psychiatry*, 4(2), 146–158. https://doi.org/10.1016/S2215-0366(16)30263-2
- Kupfer, D. J., Frank, E., & Phillips, M. L. (2012). Major depressive disorder: new clinical, neurobiological, and treatment perspectives. *Lancet (London, England)*, 379(9820), 1045–1055. https://doi.org/10.1016/S0140-6736(11)60602-8
- Kurki, M. I., Karjalainen, J., Palta, P., Sipilä, T. P., Kristiansson, K., Donner, K. M., ... Palotie, A. (2023). FinnGen provides genetic insights from a well-phenotyped isolated population. *Nature*, *613*(7944), 508–518. https://doi.org/10.1038/s41586-022-05473-8
- Lee, K.-J., Kim, H., Lee, S. J., Duperron, M.-G., Debette, S., Bae, H.-J., & Sung, J. (2023). Causal Effect of the 25-Hydroxyvitamin D Concentration on Cerebral Small Vessel Disease: A Mendelian Randomization Study. *Stroke*, 54(9), 2338–2346. https://doi.org/10.1161/STROKEAHA.123.042980
- Li, J., Li, J., Shen, L., Wang, H., Zheng, T., Hui, Y., & Li, X. (2023). Investigating the causal association of postpartum depression with cerebrovascular diseases and cognitive impairment: a Mendelian randomization study. *Frontiers in Psychiatry*, 14, 1196055. https://doi.org/10.3389/fpsyt.2023.1196055
- Liao, Y., Huang, X., Wu, Q., Yang, C., Kuang, W., Du, M., ... Gong, Q. (2013). Is depression a disconnection syndrome? Meta-analysis of diffusion tensor imaging studies in patients with MDD. *Journal of Psychiatry & Neuroscience* □: JPN, 38(1), 49–56. https://doi.org/10.1503/jpn.110180
- Lu, Y., Li, Y., Feng, Q., Shen, R., Zhu, H., Zhou, H., & Zhao, Z. (2022). Rich-Club Analysis of the Structural Brain Network in Cases with Cerebral Small Vessel Disease and Depression Symptoms. *Cerebrovascular Diseases (Basel, Switzerland)*, 51(1), 92–101. https://doi.org/10.1159/000517243
- Markus, H. S., & de Leeuw, F. E. (2023). Cerebral small vessel disease: Recent advances and future directions. *International Journal of Stroke* : *Official Journal of the International Stroke*

It is made available under a CC-BY-NC-ND 4.0 International license .

Society, 18(1), 4-14. https://doi.org/10.1177/17474930221144911

- Mewton, L., Reppermund, S., Crawford, J., Bunce, D., Wen, W., & Sachdev, P. (2019). Cross-sectional and prospective inter-relationships between depressive symptoms, vascular disease and cognition in older adults. *Psychological Medicine*, 49(13), 2168–2176. https://doi.org/10.1017/S0033291718002994
- Nobuhara, K., Okugawa, G., Sugimoto, T., Minami, T., Tamagaki, C., Takase, K., ... Kinoshita, T. (2006). Frontal white matter anisotropy and symptom severity of late-life depression: a magnetic resonance diffusion tensor imaging study. *Journal of Neurology, Neurosurgery, and Psychiatry*, 77(1), 120–122. https://doi.org/10.1136/jnnp.2004.055129
- O'Dwyer, L., Lamberton, F., Bokde, A. L. W., Ewers, M., Faluyi, Y. O., Tanner, C., ... Hampel, H. (2011). Multiple indices of diffusion identifies white matter damage in mild cognitive impairment and Alzheimer's disease. *PloS One*, 6(6), e21745. https://doi.org/10.1371/journal.pone.0021745
- Ottavi, T. P., Pepper, E., Bateman, G., Fiorentino, M., & Brodtmann, A. (2023). Consensus statement for the management of incidentally found brain white matter hyperintensities in general medical practice. *The Medical Journal of Australia*, 219(6), 278–284. https://doi.org/10.5694/mja2.52079
- Özel, F., Hilal, S., de Feijter, M., van der Velpen, I., Direk, N., Ikram, M. A., ... Luik, A. I. (2023). Associations of neuroimaging markers with depressive symptoms over time in middle-aged and elderly persons. *Psychological Medicine*, 53(10), 4355–4363. https://doi.org/10.1017/S003329172200112X
- Pan, M., Roe, J. M., Nudel, R., Schork, A. J., Iakunchykova, O., Fjell, A. M., ... Wang, Y. (2023). Circulating S100B levels at birth and risk of six major neuropsychiatric or neurological disorders: a two-sample Mendelian Randomization Study. *Translational Psychiatry*, 13(1), 174. https://doi.org/10.1038/s41398-023-02478-3
- Park, J. H., Lee, S. B., Lee, J. J., Yoon, J. C., Han, J. W., Kim, T. H., ... Kim, K. W. (2015). Epidemiology of MRI-defined vascular depression: A longitudinal, community-based study in Korean elders. *Journal of Affective Disorders*, 180, 200–206. https://doi.org/10.1016/j.jad.2015.04.008
- Pasi, M., Poggesi, A., Salvadori, E., Diciotti, S., Ciolli, L., Del Bene, A., ... Pantoni, L. (2016). White matter microstructural damage and depressive symptoms in patients with mild cognitive impairment and cerebral small vessel disease: the VMCI-Tuscany Study. *International Journal of Geriatric Psychiatry*, 31(6), 611–618. https://doi.org/10.1002/gps.4368
- Persyn, E., Hanscombe, K. B., Howson, J. M. M., Lewis, C. M., Traylor, M., & Markus, H. S. (2020). Genome-wide association study of MRI markers of cerebral small vessel disease in 42,310 participants. *Nature Communications*, 11(1), 2175. https://doi.org/10.1038/s41467-020-15932-3
- Qiu, W. Q., Himali, J. J., Wolf, P. A., DeCarli, D. C., Beiser, A., & Au, R. (2017). Effects of white matter integrity and brain volumes on late life depression in the Framingham Heart Study. *International Journal of Geriatric Psychiatry*, 32(2), 214–221. https://doi.org/10.1002/gps.4469
- Ranti, D. L., Warburton, A. J., Rutland, J. W., Dullea, J. T., Markowitz, M., Smith, D. A., ... Morris, L. S. (2022). Perivascular spaces as a marker of psychological trauma in depression: A 7-Tesla

It is made available under a CC-BY-NC-ND 4.0 International license .

MRI study. Brain and Behavior, 12(7), e32598. https://doi.org/10.1002/brb3.2598

- Sekula, P., Del Greco M, F., Pattaro, C., & Köttgen, A. (2016). Mendelian Randomization as an Approach to Assess Causality Using Observational Data. *Journal of the American Society of Nephrology* : JASN, 27(11), 3253–3265. https://doi.org/10.1681/ASN.2016010098
- Si, S., Li, J., Tewara, M. A., Li, H., Liu, X., Li, Y., ... Xue, F. (2021). Identifying causality, genetic correlation, priority and pathways of large-scale complex exposures of breast and ovarian cancers. *British Journal of Cancer*, *125*(11), 1570–1581. https://doi.org/10.1038/s41416-021-01576-7
- Skrivankova, V. W., Richmond, R. C., Woolf, B. A. R., Yarmolinsky, J., Davies, N. M., Swanson, S. A., ... Richards, J. B. (2021). Strengthening the Reporting of Observational Studies in Epidemiology Using Mendelian Randomization: The STROBE-MR Statement. *JAMA*, 326(16), 1614–1621. https://doi.org/10.1001/jama.2021.18236
- Song, S.-K., Sun, S.-W., Ju, W.-K., Lin, S.-J., Cross, A. H., & Neufeld, A. H. (2003). Diffusion tensor imaging detects and differentiates axon and myelin degeneration in mouse optic nerve after retinal ischemia. *NeuroImage*, 20(3), 1714–1722. https://doi.org/10.1016/j.neuroimage.2003.07.005
- Taylor, W. D., MacFall, J. R., Payne, M. E., McQuoid, D. R., Provenzale, J. M., Steffens, D. C., & Krishnan, K. R. R. (2004). Late-life depression and microstructural abnormalities in dorsolateral prefrontal cortex white matter. *The American Journal of Psychiatry*, 161(7), 1293–1296. https://doi.org/10.1176/appi.ajp.161.7.1293
- van Uden, I. W. M., Tuladhar, A. M., de Laat, K. F., van Norden, A. G. W., Norris, D. G., van Dijk, E. J., ... de Leeuw, F.-E. (2015). White matter integrity and depressive symptoms in cerebral small vessel disease: The RUN DMC study. *The American Journal of Geriatric Psychiatry* 2: Official Journal of the American Association for Geriatric Psychiatry, 23(5), 525–535. https://doi.org/10.1016/j.jagp.2014.07.002
- Verbanck, M., Chen, C.-Y., Neale, B., & Do, R. (2018). Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases. *Nature Genetics*, 50(5), 693–698. https://doi.org/10.1038/s41588-018-0099-7
- Vulser, H., Paillère Martinot, M.-L., Artiges, E., Miranda, R., Penttilä, J., Grimmer, Y., ... Lemaitre, H. (2018). Early Variations in White Matter Microstructure and Depression Outcome in Adolescents With Subthreshold Depression. *The American Journal of Psychiatry*, 175(12), 1255–1264. https://doi.org/10.1176/appi.ajp.2018.17070825
- Wang, Q., Qi, Y., Li, Y., Yan, Z., Wang, X., Ma, Q., ... Zhang, H. (2022). Psychiatric traits and intracerebral hemorrhage: A Mendelian randomization study. *Frontiers in Psychiatry*, 13, 1049432. https://doi.org/10.3389/fpsyt.2022.1049432
- Wendell, C. R., Hosey, M. M., Lefkowitz, D. M., Katzel, L. I., Siegel, E. L., Rosenberger, W. F., & Waldstein, S. R. (2010). Depressive symptoms are associated with subclinical cerebrovascular disease among healthy older women, not men. *The American Journal of Geriatric Psychiatry* : Official Journal of the American Association for Geriatric Psychiatry, 18(10), 940–947. https://doi.org/10.1097/JGP.0b013e3181d57a2b

Winkler, B., Funke, D., Benmimoun, B., Spéder, P., Rey, S., Logan, M. A., & Klämbt, C. (2021). Brain

It is made available under a CC-BY-NC-ND 4.0 International license .

inflammation triggers macrophage invasion across the blood-brain barrier in Drosophila during pupal stages. *Science Advances*, 7(44), eabh0050. https://doi.org/10.1126/sciadv.abh0050

- Wray, N. R., Ripke, S., Mattheisen, M., Trzaskowski, M., Byrne, E. M., Abdellaoui, A., ... Sullivan, P. F. (2018). Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depression. *Nature Genetics*, 50(5), 668–681. https://doi.org/10.1038/s41588-018-0090-3
- Wu, P.-F., Zhang, X.-H., Zhou, P., Yin, R., Zhou, X.-T., & Zhang, W. (2021). Growth Differentiation Factor 15 Is Associated With Alzheimer's Disease Risk. *Frontiers in Genetics*, 12, 700371. https://doi.org/10.3389/fgene.2021.700371
- Wuerfel, J., Haertle, M., Waiczies, H., Tysiak, E., Bechmann, I., Wernecke, K. D., ... Paul, F. (2008). Perivascular spaces--MRI marker of inflammatory activity in the brain? *Brain* □: A Journal of Neurology, 131(Pt 9), 2332–2340. https://doi.org/10.1093/brain/awn171
- Xu, X., Chan, Q. L., Hilal, S., Goh, W. K., Ikram, M. K., Wong, T. Y., ... Venketasubramanian, N. (2017). Cerebral microbleeds and neuropsychiatric symptoms in an elderly Asian cohort. *Journal* of Neurology, Neurosurgery, and Psychiatry, 88(1), 7–11. https://doi.org/10.1136/jnnp-2016-313271
- Yarlagadda, A., Alfson, E., & Clayton, A. H. (2009). The blood brain barrier and the role of cytokines in neuropsychiatry. *Psychiatry (Edgmont (Pa.* □: *Township))*, 6(11), 18–22.
- Zhang, F., Ping, Y., Jin, X., Hou, X., & Song, J. (2023). White matter hyperintensities and post-stroke depression: A systematic review and meta-analysis. *Journal of Affective Disorders*, 320, 370–380. https://doi.org/10.1016/j.jad.2022.09.166
- Zhang, H., Yao, Y., Zhong, X., Meng, F., Hemminki, K., Qiu, J., & Shu, X. (2023). Association between intake of the n-3 polyunsaturated fatty acid docosahexaenoic acid (n-3 PUFA DHA) and reduced risk of ovarian cancer: A systematic Mendelian Randomization study. *Clinical Nutrition (Edinburgh, Scotland)*, 42(8), 1379–1388. https://doi.org/10.1016/j.clnu.2023.06.028
- Zhang, R., Yu, W., Wu, X., Jiaerken, Y., Wang, S., Hong, H., ... Huang, P. (2021). Disentangling the pathologies linking white matter hyperintensity and geriatric depressive symptoms in subjects with different degrees of vascular impairment. *Journal of Affective Disorders*, 282, 1005–1010. https://doi.org/10.1016/j.jad.2020.12.171

It is made available under a CC-BY-NC-ND 4.0 International license .

	Ethnicity	Ethnicity Sample size Adjusted		URL (data download)	Pubmed ID
Exposure					
MDD	European	170 756 cases and 329 443 controls	Sex, age, genotyping array, and the first 8 PCs	https://datashare.ed.ac.uk/handl e/10283/3203	30718901
Depression	European	43 280 cases and 329 192 controls	Sex, age, genotyping batch, and 10 PCs	https://r9.risteys.finngen.fi/pheno code/F5_DEPRESSIO	36653562
Outcome					
ICH or SVS	European	6 255 cases and 233 058 controls	Age, sex, and first 4 genetic PCs	https://cd.hugeamp.org/downloa ds.html	31430377
Lobar ICH or SVS	European	5 208 cases and 233 058 controls	Age, sex, and first 4 genetic PCs	https://cd.hugeamp.org/downloa ds.html	31430377
Non-lobar ICH or SVS	European	5 468 cases and 233 058 controls	Age, sex, and first 4 genetic PCs	https://cd.hugeamp.org/downloa ds.html	31430377
WMH	European	18 381 individuals	Age at MRI, sex, genotyping array, the UK Biobank assessment center, the first 10 PCs, and MRI head motion indicators	https://cd.hugeamp.org/downloa ds.html	32358547
FA	European	17 663 individuals	Age at MRI, sex, genotyping array, the UK Biobank assessment center, the first 10 PCs, and MRI head motion indicators	https://cd.hugeamp.org/downloa ds.html	32358547
MD	European	17 467 individuals	Age at MRI, sex, genotyping array, the UK Biobank assessment center, the first 10 PCs, and MRI head motion indicators	https://cd.hugeamp.org/downloa ds.html	32358547

Table 1. Characteristics of the genome-wide association study used in this study.

_

BMBs	Multiethnic (mainly European, 94%)	3 556 cases and 22 306 controls	Age, sex, PCs, family, and relations (if applicable)	https://cd.hugeamp.org/downloa ds.html	32913026
WM-PVS	European	9 317 cases and 29 281 controls	Age, sex, intracranial volume, PCs, and study site	https://www.ebi.ac.uk/gwas/studi es/GCST90244151	37069360
BG-PVS	European	8 950 cases and 29 953 controls	Age, sex, intracranial volume, PCs, and study site	https://www.ebi.ac.uk/gwas/studi es/GCST90244153	37069360
HIP-PVS	European	9 163 cases and 29 708 controls	Age, sex, intracranial volume, PCs, and study site	https://www.ebi.ac.uk/gwas/studi es/GCST90244155	37069360

Abbreviations: GWAS, genome-wide association study, MDD, major depressive disorder; ICH, intracerebral hemorrhage; SVS, small-vessel ischemic stroke; WMH, white matter hyperintensities; FA, fractional anisotropy; MD, mean diffusivity; BMBs, brain microbleeds; WM, white matter; PVS, perivascular space; BG, basal ganglia; HP, hippocampal; PCs, principal components

It is made available under a CC-BY-NC-ND 4.0 International license

	Outcome	- CND	Cochra	Cochrane's Q		egression	MR PRESSO Global test	
Exposure	Outcome	nonp	Q value	P value	intercept	P value	P value	
MDD	ICH or SVS	36	25.949	0.867	0.027	0.174	0.852	
MDD	Lobar ICH or SVS	37	36.115	0.463	0.035	0.048	0.437	
MDD	Non-lobar ICH or SVS	36	29.395	0.735	0.028	0.033	0.740	
MDD	WMH	40	53.176	0.065	-0.029	0.003	0.068	
MDD	FA	40	88.348	< 0.001	0.099	0.085	0.237	
MDD	MD	40	53.499	0.061	-0.116	0.009	0.051	
MDD	BMBs	40	34.709	0.666	0.044	0.123	0.681	
MDD	WM-PVS	41	47.365	0.197	0.001	0.844	0.210	
MDD	BG-PVS	41	61.910	0.015	0.005	0.239	0.018	
MDD	HIP-PVS	41	47.971	0.181	-0.004	0.264	0.183	
ICH or SVS	MDD	14	23.763	0.033	0.005	0.394	0.132	
Lobar ICH or SVS	MDD	13	20.619	0.056	-0.003	0.640	0.060	
Non-lobar ICH or SVS	MDD	28	32.248	0.223	-0.002	0.621	0.213	
WMH	MDD	6	3.600	0.608	-0.009	0.288	0.651	
FA	MDD	5	4.693	0.320	-0.009	0.281	0.408	
MD	MDD	4	0.203	0.977	-0.001	0.948	0.980	
BMBs	MDD	19	30.089	0.037	0.003	0.562	0.046	
WM-PVS	MDD	3	0.396	0.820	-0.002	0.859	NA	
BG-PVS	MDD	0	NA	NA	NA	NA	NA	
HIP-PVS	MDD	0	NA	NA	NA	NA	NA	

Table 2. Results of the pleiotropy and heterogeneity test results for major depressive disorder (MDD) and cerebral small vessel disease (CSVD).

Abbreviations: nSNP, number of SNP; MDD, major depressive disorder; ICH, intracerebral hemorrhage; SVS, small-vessel ischemic stroke; WMH, white matter hyperintensities; FA, fractional anisotropy; MD, mean diffusivity; BMBs, brain microbleeds; WM, white matter; PVS, perivascular space; BG, basal ganglia; HIP, hippocampal; NA, not applicable.

It is made available under a CC-BY-NC-ND 4.0 International license

It is made available under a CC-BY-NC-ND 4.0 International license

Figure Captions

Figure 1. Diagrammatic overview of this MR study. The GVs selected as IVs met three core assumptions. (1) the relevance assumption: IVs should be strongly related to the exposure; (2) the independence assumption: the IVs should not be associated with other confounding factors; (3) the exclusion restriction assumption: the effect of the IVs on outcome should be exclusively through exposure. Abbreviations MDD, major depressive disorder; CSVD, cerebral small vessel disease; ICH, intracerebral hemorrhage; SVS, small-vessel ischemic stroke; WMH, white matter hyperintensities; FA, fractional anisotropy; MD, mean diffusivity; BMBs, brain microbleeds; WM, white matter; PVS, perivascular space; BG, basal ganglia; HP, hippocampal; nSNP, number of SNP; GVs, genetic variants; IVs, instrumental variables.

Figure 2. Bidirectional causal relationships between MDD and CSVD using the IVW approach. A The foward MR analyses examined the effect of MDD on CSVD clinical outcomes (n=3) and CSVD radiological markers (n=7). B The reverse MR analyses examined the effect of CSVD clinical outcomes (n=3) and CSVD radiological markers (n=4) on MDD. BG-PVS and HIP-PVS did not have a qualified IV available for analysis, even with a relaxed threshold (P=1E-05), and were therefore not included in the reverse MR analysis. Abbreviations nSNP, number of SNP; OR, odds ratio; CI, confidence interval; CSVD, cerebral small vessel disease; ICH, intracerebral hemorrhage; SVS, small-vessel ischemic stroke; WMH, white matter hyperintensities; FA, fractional anisotropy; MD, mean diffusivity; BMBs, brain microbleeds; WM, white matter; PVS, perivascular space; BG, basal ganglia; HIP, hippocampal.

It is made available under a CC-BY-NC-ND 4.0 International license .

Figure 1. Diagrammatic overview of this MR study. The GVs selected as IVs met three core assumptions. (1) the relevance assumption: IVs should be strongly related to the exposure; (2) the independence assumption: the IVs should not be associated with other confounding factors; (3) the exclusion restriction assumption: the effect of the IVs on outcome should be exclusively through exposure. Abbreviations MDD, major depressive disorder; CSVD, cerebral small vessel disease; ICH, intracerebral hemorrhage; SVS, small-vessel ischemic stroke; WMH, white matter hyperintensities; FA, fractional anisotropy; MD, mean diffusivity; BMBs, brain microbleeds; WM, white matter; PVS, perivascular space; BG, basal ganglia; HP, hippocampal; nSNP, number of SNP; GVs, genetic variants; IVs,

instrumental variables.

It is made available under a CC-BY-NC-ND 4.0 International license .



It is made available under a CC-BY-NC-ND 4.0 International license .

Figure 2. Bidirectional causal relationships between MDD and CSVD using the IVW approach.

A The foward MR analyses examined the effect of MDD on CSVD clinical outcomes (n=3) and CSVD radiological

markers (n=7). B The reverse MR analyses examined the effect of CSVD clinical outcomes (n=3) and CSVD

radiological markers (n=4) on MDD. BG-PVS and HIP-PVS did not have a qualified IV available for analysis, even with

a relaxed threshold (P=1E-05), and were therefore not included in the reverse MR analysis. Abbreviations nSNP,

number of SNP; OR, odds ratio; CI, confidence interval; CSVD, cerebral small vessel disease; ICH, intracerebral

hemorrhage; SVS, small-vessel ischemic stroke; WMH, white matter hyperintensities; FA, fractional anisotropy; MD,

mean diffusivity; BMBs, brain microbleeds; WM, white matter; PVS, perivascular space; BG, basal ganglia; HIP,

hippocampal.

It is made available under a CC-BY-NC-ND 4.0 International license .

