It is made available under a CC-BY 4.0 International license .

| 1 | |
|--|---|
| 2 | |
| 3 | |
| 4 | EMS injury cause codes more accurate than |
| 5 | emergency department visit ICD-10-CM codes for |
| 6 | firearm injury intent in North Carolina |
| 7 | |
| 8 9 10 11 12 13 14 15 16 17 18 19 | Nicole L. Snyder ¹ , Amy Ising ^{1*} , Anna E. Waller ¹ ¹ Carolina Center for Health Informatics in the Department of Emergency Medicine, School of Medicine, University of North Carolina at Chapel Hill. |
| 20 21 22 23 24 | *Corresponding author E-mail: amy_ising@med.unc.edu (AI) |

It is made available under a CC-BY 4.0 International license .

25 Abstract

26 Background

The timeliness, accuracy, and completeness of data for firearm injury surveillance is crucial for public health surveillance efforts and informing injury prevention measures. While emergency department (ED) visit data can provide near real-time information on firearms injuries, there are concerns surrounding the accuracy of intent coding in these data. We examined whether emergency medical service (EMS) data provide more accurate firearm injury intent coding in comparison to ED data.

33 Methods

We applied a firearm injury definition to EMS encounter data in NC's statewide syndromic surveillance system (NC DETECT), from January 1, 2021, through December 31, 2022. Each record was manually reviewed to determine shooter and intent, and the corresponding manual classifications were compared to the injury cause codes entered in the EMS data and to ED visit records, where linkage was possible.

39 **Results**

40 We identified 9557 EMS encounters from January 1, 2021, through December 31, 2022 meeting 41 our firearm injury definition. After removing false positives and duplicates, 8584 records were 42 available for manual injury classification. Overall, our analysis demonstrated that manual and 43 EMS injury cause code classification were comparable. However, for the 3401 EMS encounters 44 that could be linked to an ED visit record, only 18.3% (n = 355) of the 1945 assaults and 22.2% (n = 38) of the 171 intentional self-harm suicide encounters we identified in the EMS records were 45 46 identified as assault firearm injures in the ED visit data. This demonstrates a marked difference 47 in the intent coding between the two data sources.

48 **Conclusions**

It is made available under a CC-BY 4.0 International license .

- 49 This study illustrates both the value of examining EMS encounters for firearm injury intent, and
- 50 the challenges of accurate intent coding in the ED setting. The results also suggest the importance
- 51 of developing better guidance around intent coding for firearm injuries in the ED.

It is made available under a CC-BY 4.0 International license .

59 Introduction

60 As firearm injury continues to be a significant cause of morbidity and mortality in the United States 61 (US), public health researchers continue to seek improvements to the timeliness, accuracy, and 62 completeness of firearm injury surveillance [1]. In North Carolina (NC), firearm injury surveillance 63 relies primarily on the NC Violent Death Reporting System (NC-VDRS), hospital administrative 64 data, and syndromic surveillance emergency department visit data [2]. While the syndromic 65 surveillance emergency department (ED) visit data provides near real time information on nonfatal firearm injury in NC, there are concerns about accuracy when subtyping firearm injuries by intent 66 67 (assault, unintentional, intentional self-harm). For example, while some studies estimate 68 unintentional firearm injury to represent 16% or less of all nonfatal firearm injury, approximately 69 60% of ED firearm visits in NC are identified as unintentional [3, 4]. Several other studies that 70 have documented these coding intent discrepancies and the potential overclassification of 71 nonfatal firearm injury as unintentional, relied on urban, Level I Trauma center data from a small 72 number of hospitals [5, 6]. For comparison, we aimed to quantify any coding intent discrepancies 73 using data from a statewide system in a predominantly rural state. To achieve this, we compared 74 firearm injury intent classifications as identified in statewide emergency medical services (EMS) 75 encounter data to the firearm injury intent classifications identified in syndromic surveillance ED 76 visit data, for those patients transported to the ED by EMS. We specifically chose to examine 77 EMS data because EMS narratives provide rich contextual details on the entire EMS encounter 78 and can, therefore, provide insights into firearm injury intent.

79 Methods

We conducted this evaluation in several steps. First, we applied a firearm injury definition to EMS encounter data for encounters sent to NC DETECT, NC's statewide syndromic surveillance system, from January 1, 2021 through December 31, 2022. We then manually reviewed each

83 record and documented shooter and intent. Third, we compared manual intent classifications to

84 those documented in the EMS data based solely on injury cause code. Finally, we linked the EMS

85 records to corresponding ED visit records, where possible, to compare our manually assigned

86 intent classifications to the intent coding in the ED visit data. The UNC IRB determined this study

to be exempt from full review.

88

89 EMS firearm V1 definition

90 The EMS firearm definition we used for this analysis is outlined in Table 1. It searches the

91 complaint for gunshot wound related injuries OR the injury cause field for firearm-related ICD-10-

- 92 CM codes. The definition also searches for gunshot wound related injuries in the narrative if the
- 93 dispatch complaint is Stab/Gunshot Wound/Penetrating Trauma. This initial definition was
- 94 designed to maximize sensitivity, while excluding EMS encounters that are not for active
- 95 responses based on EMS patient disposition.

96 97

Table 1. EMS Firearm V1 Definition.

| Complaint Terms | GSW or gunshot or "gun shot" or "shot myself" or "shot self*" or firearm or "I got shot" or "been shot" or "shot gun" or shotgun |
|--|---|
| Injury cause codes | W34, W34.0, W34.09, W34.00, W34.00XA, W33.02, W33.03, W33.09, W33.01, W32.0, W32, W34.1, W33.0, W33, X94.0XXA, X95.9, X95.9XXA, Y22, Y22.XXXA, Y23.1, X72.XXXA, X74.9, Y35.0, Y35.02, Y35.023, Y35.03, Y35.092, Y35.093, Y35.003, Y23.3, Y22.0, Y24.8, Y23.8, Y23, Y23.0, Y23.0XXA, Y38.4X, Y24.9, Y24.9XXA, Y23.9 |
| Dispatch complaint AND narrative | dispatch complaint is Stab/Gunshot Wound/Penetrating Trauma AND narrative keywords of GSW or gun shot or gunshot |
| Patient disposition exclusions | Assist, Agency; Assist, Public, Assist; Unit, Canceled (Prior to Arrival At Scene); Canceled on Scene (No Patient Contact); Canceled on Scene (No Patient Found); Standby-No Services or Support Provided; Standby- Public Safety, Fire, or EMS Operational Support Provided; Transport Non-Patient, Organs, etc. |

98

99 EMS encounter manual review

100 With approval from the NC Office of Emergency Medical Services, we downloaded all EMS 101 encounters from January 1, 2021 through December 31, 2022 meeting the firearm v1 definition 102 from NC's statewide syndromic surveillance system NC DETECT on February 1, 2023 with an 103 update to incorporate additional records on May 1, 2023[7]. The authors did not have access to 104 any information that could identify individual participants during or after data collection. For each 105 EMS encounter, we manually documented the shooter (self, other, unknown) and intent 106 (intentional, unintentional, undetermined) in a spreadsheet, which we also collapsed into the 107 categories used in injury intent coding in the International Classification of Diseases: assault 108 (other - intentional), self-inflicted (self - intentional), unintentional (other, unknown, and self) and 109 undetermined (other, unknown, and self). We also indicated false positives. We then added notes, 110 as needed, to document our rationale for any of these classifications. EMS encounters were 111 reviewed by only one person, but all challenging EMS encounters were reviewed by three 112 reviewers to achieve consensus on coding. We included both non-fatal and fatal firearm injuries 113 in our review. Additional decision points are listed below:

- Intent was assigned using complaint, narrative, and injury cause code fields. Sometimes
 these fields provided conflicting information. When that occurred, the reviewer used group
 consensus to determine intent.
- If injury cause was coded as undetermined, a more precise intent may have been assigned
 based on the narrative description.
- When the narrative described an incident as a drive-by shooting or shot by someone else
 while in a car with no documented/undetermined injury cause, intent was classified as
 assault.
- If a patient was transported to an ED by EMS, stabilized, and then transferred by EMS to
 another ED, both EMS encounters were coded.

It is made available under a CC-BY 4.0 International license .

- For deaths at the scene with inconclusive intent in the EMS record, death certificate data
- 125 were consulted to determine intent, if a linkage could be identified based on patient age,
- sex, race, county of residence and date of death.
- 127

128 False positives

- 129 We excluded any encounters identified by the firearm v.1 definition that described injuries caused
- 130 by bb guns, pellet guns, nail guns, air guns, or firearms used as a blunt weapon, e.g., "pistol whip."
- 131 We also excluded subsequent encounters for previous firearm injury and near misses.
- 132

133 EMS-specific descriptive analyses

134 After we completed the manual review and removed false positives, we conducted a descriptive 135 analysis of EMS firearm encounters. We calculated counts and crude rates statewide and by EMS 136 county by sex, race, ethnicity, and age group. We compared the intent as identified in the injury 137 cause field to the intent assigned manually during the review process. We classified EMS 138 encounters by injury cause code into assault, intentional self-harm/suicide, unintentional, 139 undetermined, and a final category of non-firearm injury cause codes. If the EMS encounter 140 record included two conflicting firearm injury cause codes, we classified the encounter based on 141 the first code listed.

142

143 EMS-ED linkage

The EMS encounters from January 1, 2021 through December 31, 2022 meeting the EMS firearm v1 definition were linked, when possible, to the corresponding ED visit record and provided to the authors for analysis on October 19, 2023. The linkage used an existing hierarchical deterministic process to match records based on drop-off time by EMS, arrival time in the ED, EMS destination

It is made available under a CC-BY 4.0 International license .

148 name, EMS destination type, patient date of birth, sex and race/ethnicity, ED facility name, and 149 ED transport mode [8]. The linkage rules use the same variables across multiple steps with 150 different matching criteria to maximize the number of linkages while minimizing false linkages. 151 Several updates were made to the linkage process as part of this analysis to increase the 152 likelihood of a successful linkage; we updated the EMS encounter destination type required for 153 linkage to include missing and *Freestanding Emergency Department* destination types as well as 154 Hospital-Emergency Department. We also allowed records in the linkage process with missing 155 race and updated the free text list of hospital names used by EMS agencies in the Destination 156 Name field to include additional aliases and misspellings, increasing that list from 3213 to 5739 157 entries.

158

159 EMS-ED linkage descriptive analyses

We determined the number and percentage of linked EMS records to corresponding ED visits and compared the EMS manually assigned intent to the intent in the ED visit data as identified by ED firearm injury intent definitions developed by the Centers for Disease Control and Prevention (CDC) and implemented in NC DETECT [9, 10].

164

165 **Results and discussion**

166

We identified 9557 EMS encounters from January 1, 2021, through December 31, 2022 meeting
our firearm injury definition. Of these, 969 (10%) were identified as false positives and four records
were duplicates, leaving 8584 EMS encounters for this analysis.

170

171 False positive analysis

- 172 We grouped the 969 false positives we observed in our data into seven categories, as shown in
- 173 Table 2. We then analyzed the false positives to inform improvements to our EMS firearm injury
- 174 definition.

175 Table 2. EMS Syndrome Definition False Positive Categories. Count (%) False Positive Category Other types of injuries or other types of false positives that could not be 222 (22.9%) categorized Pistol whips with no penetrating firearm discharge injury 185 (19.1%) Subsequent encounters for a previous firearm injury 173 (17.9%) Firearm injuries by bb guns, pellet guns, air guns, nail guns, etc. 121 (12.5%) Near misses, e.g., injuries sustained near gunshots but with no evidence of 113 (11.7%) direct injury by a bullet Mental health concerns without firearm injury 92 (9.5%) Stabbings 63 (6.5%)

176

177 The largest number of false positives, 22.9% (n = 222) fell into the other types of injuries category, 178 which included EMS encounters with no documented firearm discharge and other types of 179 assault. Pistol whips were also a common source of false positives; 173 (93.5%) of these had a 180 firearm injury cause code, typically Assault by unspecified firearm discharge (X95.9), even though 181 there was no documented firearm discharge injury in the EMS record. We often identified 182 subsequent encounters (n = 173, 17.9%) in the EMS narrative text; 80 (46.2%) of these 183 subsequent encounter records had a firearm discharge injury cause code. Injuries sustained near 184 gunshots but with no evidence of direct injury by a bullet, e.g., injuries sustained by falls running 185 away from gunshots, lacerations sustained from broken glass caused by bullets hitting 186 windshields and windows, etc., comprised 11.7% (n = 113) of false positives. Examples of the 92 187 encounters (9.5%) that fell into the mental health category included family members of firearm 188 injury victims who had a mental health crisis after the firearm injury event, suicidal patients with 189 firearms who did not suffer firearm injury, patients with mental health conditions claiming firearm 190 injury with no identified firearm injury, and patients with substance misuse claiming firearm injury 191 with no identified firearm injury. The 63 (6.5%) stabbing-related false positives resulted primarily 192 from including the Stab/Gunshot Wound/Penetrating Trauma dispatch complaint in our definition It is made available under a CC-BY 4.0 International license .

and this dispatch complaint text being repeated in the narrative text. Based on our false positive analysis, we hypothesize that the addition of chief complaint exclusion terms used in the CDCdeveloped ED firearm definition may improve our EMS firearm definition [9]. Additionally, exclusion terms that address false positives, e.g., pistol whips, even when the encounter has a firearm discharge injury cause code, will require logic that applies these exclusion terms to encounters with firearm discharge injury cause codes in addition to exclusion terms specific to the complaint and narrative free text fields.

200

201 EMS descriptive analyses

202 The primary cause of firearm related injury in 2021-2022 EMS encounter data was assault, which 203 encompassed 51.8% (n = 4446) of all recorded events, followed by undetermined (n = 1694; 204 19.7%), intentional self-harm (n = 1581; 18.4%), and unintentional (n = 862; 10.1%). The ICD-10-205 CM injury intent coding does not classify unintentional and undetermined firearm injury by 206 perpetrator, but our manual intent coding revealed that 7.3% (n = 634) of EMS firearm encounters 207 were for unintentional self-inflicted injury, e.g., persons shooting themselves while cleaning, 208 holstering, storing, retrieving, stepping on and/or dropping firearms. This finding presents an 209 opportunity to inform unintentional firearm injury prevention messaging with specific scenarios. 210 For example, public health messaging could remind firearm owners to clean their firearms while 211 sober, and to store their firearms unloaded.

212

Fig 1. EMS Firearm Injuries by shooter and intent

215

216 EMS demographic analysis

We categorized firearm injury intent by key demographics, revealing several important differences
by sex, age group, and race/ethnicity, as shown in Table 3. Notably, EMS firearm injuries were

It is made available under a CC-BY 4.0 International license .

219 more than 7.0 times more likely to occur among males in comparison to females. Males were also 220 4.7 times more likely to be involved in an assault-related firearm injury and 6.2 times more likely 221 to be involved in an intentional self-inflicted firearm injury compared to females. With respect to 222 age group, the highest rate of firearms related assaults was among those aged 19 to 24 (112.77 events per 100,000), followed by those aged 15 to 18 (92.90 events per 100,000). In contrast, 223 224 those aged 65+, followed closely by those aged 19 to 24 years of age, were more likely to be 225 involved in an intentional self-inflicted firearm injury, with rates of 23.31 per 100,000 and 19.28 226 per 100,000, respectively. For race/ethnicity, Black or African Americans were 13.3 times more 227 likely to be involved in a firearm injury related to assault in comparison to White persons, while 228 White persons had the highest rate of intentional self-inflicted firearm injury, highlighting the 229 disparities among these two demographic groups. Notably, American Indian/Alaskan Native 230 (AIAN) persons had the second highest assault firearm injury rate, which was nearly four times 231 higher than Whites, at 41.43 per 100,000.

232 These results highlight the importance of targeted interventions. For example, holistic 233 interventions that target middle-school and high-school age students could result in significant 234 reductions in assault and intentional self-harm events. Recent efforts focused on establishing 235 universal firearms risk injury screening in care settings for adolescents and youths [11], as well 236 as developing a better understanding of firearm access among this group [12], may help drive 237 preventative mechanisms in this area. Interventions that examine the underlying causes of the 238 increasing rates of intentional self-harm among those aged 65+ [13], might lead to interventions 239 such as increased health care provider communication and policies that address firearms access 240 to those with signs of dementia and/or depression [14, 15]. Finally, holistic efforts to address the 241 disparities that exist among different racial and ethnic groups with higher rates of firearm-related 242 injury events [16] would be an important step forward for reducing injuries among these groups.

It is made available under a CC-BY 4.0 International license .

244 Table 3. EMS Firearm Injury Demographics, by Count and Crude Rate.^a

Carolina. For the total row, the total population was used for the denominator.

| Demographic | Assault | Intentional Self-Harm | Undetermined | Unintentional | All Intents |
|--------------------|----------|--------------------------|--------------|---------------|----------------|
| Total | 4446 | 1581 | 1694 | 863 | 8584 |
| | (39.17) | (13.93) | (14.93) | (7.60) | (75.64) |
| | ••• | Se | X | | |
| Male | 3621 | 1342 | 1403 | 743 | 7109 |
| | (70.34) | (26.07) | (27.25) | (14.43) | (138.09) |
| Female | 808 | 230 | 270 | 112 | 1420 |
| | (14.77) | (4.21) | (4.90) | (2.05) | (25.99) |
| Unknown/Missing | 17 | 9 | 21 | 8 | 55 |
| | | Age G | roup | | |
| Infant/Preschool | 19 | 0 | 20 | 26 | 65 |
| (0-4) | (3.20) | (0.00) | (3.37) | (4.37) | (10.93) |
| Elementary School | 23 | 0 | 19 | 14 | 56 |
| (5-9) | (3.80) | (0.00) | (3.14) | (2.31) | (9.25) |
| Middle School | 86 | 22 | 36 | 32 | 176 |
| (10-14) | (13.23) | (3.38) | (5.54) | (4.92) | (27.07) |
| High School | 515 | 66 | 199 | 53 | 833 |
| (15-18) | (92.90) | (11.91) | (35.90) | (9.56) | (150.26) |
| College | 1012 | 173 | 354 | 175 | 1714 |
| (19-24) | (112.77) | (19.28) | (39.45) | (19.50) | (190.99) |
| Young Adult | 2074 | 434 | 751 | 311 | 3579 |
| (25-44) | (75.25) | (15.75) | (27.25) | (11.28) | (129.54) |
| Middle Aged | 580 | 443 | 229 | 155 | 1407 |
| (45-64) | (21.41) | (16.35) | (8.45) | (5.72) | (51.93) |
| Senior | 91 | 430 | 51 | 90 | 662 |
| (65+) | (4.93) | (23.31) | (2.76) | (4.88) | (35.88) |
| Unknown/Missing | 46 | 13 | 35 | 7 | 101 |
| | | Race/Et | hnicity | | |
| American | 74 | 6 | 61 | 12 | 153 |
| Indian/Alaskan | (41.43) | (3.36) | (34.15) | (6.72) | (85.67) |
| Native | | | | | |
| Asian/Pacific | 22 | 17 | 7 | 11 | 57 |
| Islander | (5.63) | (4.35) | (1.79) | (2.82) | (14.59) |
| Black or African | 3103 | 211 | 1020 | 266 | 4600 |
| American | (139.28) | (9.47) | (45.78) | (11.94) | (206.47) |
| Hispanic or Latino | 281 | 56 | 89 | 45 | 471 |
| | (24.24) | (4.83) | (7.68) | (3.88) | (40.63) |
| White | 801 | 1246 | 351 | 485 | 2883 |
| | (10.83) | (16.85) | (4.75) | (6.56) | (38.99) |
| Unknown/Missing | 165 | 45 | 166 | 44 | 420 |

245 246

247

.

248

249

250

251 Manual intent and injury cause code classification comparison

Next, we compared our manually assigned intent classifications to the EMS injury cause codes (ICD-10-CM) alone. The results are shown in Table 4. There were 2358 firearm injury encounters (27.5%) that were missing an injury cause code. Of the remaining 6226 EMS encounters, our manual intent coding was comparable to the intent identified solely by the injury cause code. Although our manual intent classifications referenced the injury cause codes during review, these results suggest that EMS injury cause codes can serve as a reliable indicator of firearm injury intent when available.

Table 4. Manual and Injury Cause Code Intent Classification Comparison, Count (Column
 %).

| Intent | Manual Injury Classification | EMS Injury Cause Code Classification* |
|------------------------------------|---------------------------------|--|
| Assault | 4446 (51.8%) | 3055 (49.1%) |
| Intentional Self - Harm/Suicide | 1581 (18.4%) | 1111 (17.8%) |
| Unintentional | 862 (10.1%) | 580 (9.3%) |
| Undetermined | 1694 (19.7%) | 1211 (19.5%) |
| Multiple Intent Codes | | 105 (1.7%) |
| Other (non-firearm) | | 164 (2.6%) |
| Total | 8584 (100%) | 6226 (100%) |

261

262

*Missing = 2,358

We further examined instances where our manual review did NOT match the intent as assigned by the EMS injury cause code. Our manual intent classifications did not agree with the assigned injury cause code intents for 773 EMS encounters, of which 641 (82.9%) were assigned an undetermined injury cause code. With manual review of the narrative text, we were able to assign 436 of these to assault, followed by 156 to intentional self-harm and 49 to unintentional. The manual intent assignments resulted in more specific intent coding rather than a change of intent

It is made available under a CC-BY 4.0 International license .

- for most EMS encounters. With an improvement in completeness, EMS injury cause codes alone
- 270 can be a reliable source of intent-specific firearm injury surveillance.

271 EMS disposition

272 We also grouped the EMS firearm injury encounters by EMS disposition to document both the

273 continuum of care for patients initially encountered by EMS, and to assess the potential for linking

each event to ED visit data (Table 5). Most events were recorded as '*Patient Treated, Transported*

by this EMS Unit' (n = 6054, 71%). A total of 1889 patients (22%) were dead at the scene from

firearm injury, 67% (n = 1062) of which were from intentionally self-inflicted gunshot wounds. For

- intents other than intentional self-harm, the majority of EMS encounters for firearm injury result in
- EMS transport.

| Intent | All Encounters | Patient treated, transported by this EMS Unit | Patient Dead at Scene (with and without transport) | Other Disposition |
|---------------|-------------------|---|--|----------------------|
| | | | | |
| Assault | 4446 | 3484 (78%) | 602 (14%) | 360 (8%) |
| Intentional | | | | |
| Self-Harm | 1581 | 467 (30%) | 1062 (67%) | 52 (3%) |
| Unintentional | 863 | 770 (89%) | 15 (2%) | 78 (9%) |
| Undetermined | 1694 | 1333 (79%) | 210 (12%) | 151 (9%) |
| All Intents | 8584 | 6054 (71%) | 1889 (22%) | 641 (7%) |

| (Row %). |
|----------|
| (|

280

281 EMS-ED linkage results

We then examined the linkage between the firearm injury events in our EMS analysis with ED data with the goal of comparing firearm injury intent across the two data sources. Of the 8854 EMS encounters included in our analysis, 6054 had a disposition of '*Patient Treated, Transported by this EMS Unit*' and, of those, 5558 had a destination type of '*Hospital-Emergency Department*', '*Freestanding Emergency Department*' or '*Missing*', and 5353 of those were transported to an ED in NC DETECT (i.e., all civilian, acute care EDs in NC), based on a review of the free text

It is made available under a CC-BY 4.0 International license .

288 destination name. We were able to link 3401 (63.5%) of these EMS encounters to a corresponding

- ED visit (Figure 2).
- 290 Fig 2. EMS-ED linkage results
- 291

292 For the EMS records that we successfully linked to an ED visit record (n = 3401), only 293 18.3% (n=355) of the assaults we identified in the EMS data (n=1945) were binned into the CDC 294 Firearm Assault v1 definition, and only 22.2% (n=38) of the intentional self-harm suicide 295 encounters (n=171) were binned into the CDC Firearm Intentional Self-Harm v1 definition. These 296 findings illustrate the challenges of accurate intent coding in the ED setting and the potential 297 impact of the CMS coding guidance that requires injury intent to default to unintentional if the 298 intent is not identified in the physician documentation [17]. They also suggest that EMS 299 encounters may serve as a better proximate for understanding intent for firearm injury than ED 300 data, given the current coding guidance. EMS encounters identified as firearm injuries that were 301 not identified as firearm injuries in the ED data often had trauma-related chief complaints and/or 302 had diagnosis codes for penetrating injuries without an external mechanism firearm ICD-10-CM 303 code.

Table 6. Count (Row %) of EMS encounters linked to ED visits and binned into ED
 Firearm Syndromes by EMS Manual Intent.^a

| Intent of EMS records assigned by manual review | EMS Firearm Encounters linked to ED visits | ED CDC Firearm All Intents V2 | ED CDC Firearm Assault V1 | ED CDC Firearm Intentional Self-Harm V1 | ED CDC Firearm Unintentional / Undetermined V1 | Not binned into any ED firearm injury- related syndrome |
|---|--|--|---------------------------------------|---|---|--|
| Assault | 1945 | 1794 (92.2%) | 355 (18.3%) | 6 (0.3%) | 1234 (63.4%) | 151 (7.8%) |
| Unintentional (self, other, or unknown shooter) | 509 | 472 (92.7%) | 9 (1.8%) | 3 (0.59%) | 378 (74.3%) | 37 (7.3%) |
| Intentional Self-Harm / Suicide | 171 | 136 (79.5%) | 3 (1.8%) | 38 (22.2%) | 77 (45.0%) | 35 (20.4%) |

It is made available under a CC-BY 4.0 International license .

| Undetermined | 776 | 721 | 126 | 6 | 482 | 55 |
|-----------------------------|-----|---------|---------|---------|---------|--------|
| (self, other, or unknown | | (92.9%) | (16.2%) | (0.77%) | (62.1%) | (7.1%) |
| shooter) | | | | | | |

306

^a An ED visit can be binned into 0,1 or more syndromes.

307

308 While this study did not include a comprehensive review of EMS encounters transported to an ED

309 that were not successfully linked to an ED visit record, our review of missed linkages revealed

310 that missing or inaccurate dates of birth represented a significant cause of missed linkages. For

311 serious events such as firearm injury, a reliable date of birth may be challenging for EMS to record

312 at the scene.

313 ED disposition for EMS-ED linked encounters

As shown in Table 7, 1721 (50.6%) of the ED visits linked to an EMS encounter had an ED disposition of discharged home, with 26.8% (n=913) admitted; 230 (6.8%) had a disposition of died in the ED. Of those who died in the ED, 76 (33%) were identified in the EMS encounter as intentional self-harm, and 103 (44.8%) as assault. These data indicate the overall severity of firearm injury and the lethality of firearms when used with intent to inflict injury. Only 2 (0.8%) deaths in the ED were for EMS identified unintentional firearm injuries.

320 Table 7. ED Disposition for EMS-ED linkage.

| ED Disposition for EMS-ED Linked Data | Count (%) |
|---|--------------|
| Discharged to home or self-care (routine discharge) | 1721 (50.6%) |
| Admitted to Hospital | 913 (26.8%) |
| Died | 230 (6.8%) |
| Transferred/discharged to another short-term general hospital | 249 (6.5%) |
| Transferred/discharged to another type of care setting | 65 (1.9%) |
| Transferred/discharged to prison or jail | 54 (1.6%) |
| Left AMA / Discontinued Care | 36 (1.1%) |
| Unknown/Missing | 133 (3.9%) |
| Total | 3401 (100%) |

321

322 Limitations

It is made available under a CC-BY 4.0 International license .

323 The analysis presented in this report has several limitations. When manually coding intent, we 324 relied on the EMS narrative but if the reporting of the incident intent to or by EMS was inaccurate, 325 then the resulting intent coding may be inaccurate. For example, if the patient stated that they 326 were shot in a drive-by shooting we coded this EMS encounter as an assault, but the patient may 327 have unintentionally shot themselves and reported it as a drive-by shooting. In addition, our 328 descriptive analysis included both nonfatal and fatal EMS encounters. While this reflects the 329 comprehensive EMS data for firearm injury, certain findings may be different when analyzing 330 nonfatal and fatal EMS firearm injury encounters separately. Finally, most of the EMS encounters 331 in this analysis were reviewed by only one reviewer; intent classifications may have been assigned differently by a different reviewer. 332

333

334 Conclusions

335 Timely and accurate injury cause codes are crucial for understanding the nature and scope of 336 firearm-related injuries. In this study, we manually assigned intent to 8584 eligible NC EMS 337 encounters from 2021 and 2022 to better understand the landscape of EMS-related firearm injury 338 calls and to assess the accuracy of firearm injury coding in the ED setting. Manual coding of the 339 data for shooter and intent revealed comparable results to those obtained from examining EMS 340 assigned injury cause code classifications, when available. Of the 63.5% encounters (n = 3401) 341 that could be linked to ED visit data, 1,945 of those were coded in the EMS data as assaults 342 (57.2%). However, only 18.3% (n = 355) of those encounters were classified as assaults based 343 on the ED CDC Assault v1 definition. Similarly, of the 171 encounters coded as intentional self-344 harm suicide encounters in the EMS, only 22.2% (n = 38) were binned to the CDC Firearm 345 Intentional Self-Harm v1 definition. The ED data likely reflects current CMS coding guidance that

It is made available under a CC-BY 4.0 International license .

requires injury intent to default to unintentional if the intent is not identified in the physiciandocumentation [17].

348 The outcomes of our study reveal that EMS encounter data may be useful for obtaining 349 more granular data from the scene of the injury incident. Such data might be important for 350 developing tailored approaches that reflect regional and demographic differences in firearms 351 injuries. In contrast, the lack of such detailed information in the ED visit data may limit such 352 interventions. For example, if an encounter is the result of an assault, but it is coded as 353 unintentional or undetermined, a patient may not be flagged for post-discharge follow-up that is 354 directed towards supporting victims of assault. Combined, the results of this work illustrate the importance of accurate intent coding for firearms injuries and provides a foundation for the 355 356 discussion of better coding guidance in the ED setting.

357

359

358 Acknowledgements

360 The North Carolina Disease Event Tracking and Epidemiologic Collection Tool (NC DETECT) is 361 an advanced, statewide public health surveillance system. NC DETECT is supported by the North 362 Carolina Division of Public Health through a federal Public Health Emergency Preparedness 363 Grant and is managed through a collaboration between NC DPH and the University of North 364 Carolina at Chapel Hill Department of Emergency Medicine's Carolina Center for Health 365 Informatics. The findings and conclusions in this publication are those of the author(s) and do not 366 necessarily represent the views of the North Carolina Department of Health and Human Services, 367 Division of Public Health.

368

The NC OEMS and the NC EMS Data System supports state, regional and local EMS and healthcare related service delivery from a patient care, resource allocation, and regulatory perspective. This manuscript was not prepared in collaboration with investigators of the NC EMS

18

- 372 Data System and does not necessarily reflect the opinions or views of the NC OEMS, ESO or the
- 373 study sites participating in the NC EMS Data System.

374

- 375 The authors also wish to thank Matthew Kerber for his work on the EMS-ED linkage components
- of this study.
- 377

378 **References**

379

- Barber C, Cook PJ, Parker ST. The emerging infrastructure of US firearms injury data. Preventive Medicine. 2022;165:107129.
- Keeping Families and Communities Safe: Public Health Approaches to Reduce
 Violence and Firearm Misuse Leading to Injury and Death [White Paper]. 2022.
- Cook PJ, Parker ST. Correcting Misinformation on Firearms Injuries. JAMA Netw
 Open. 2022;5(12):e2246434.
- 3864.NC-FASTER Reports. Carolina Center for Health Informatics in the Department of387Emergency Medicine, University of North Carolina at Chapel HIII & NC Division of388Health and Human Services, Division of Public Health. 2023. https://ncdetect.org/nc-389faster-firearm-quarterly-reports/.
- 3905.Miller M, Azrael D, Yenduri R, Barber C, Bowen A, MacPhaul E, et al. Assessment of391the Accuracy of Firearm Injury Intent Coding at 3 US Hospitals. JAMA Netw Open.3922022;5(12):e2246429.
- MacPhaul E, Zhou L, Mooney SJ, Azrael D, Bowen A, Rowhani-Rahbar A, et al.
 Classifying Firearm Injury Intent in Electronic Hospital Records Using Natural
 Language Processing. JAMA Netw Open. 2023;6(4):e235870.
- Hakenewerth AM, Waller AE, Ising AI, Tintinalli JE. North Carolina Disease Event
 Tracking and Epidemiologic Collection Tool (NC DETECT) and the National Hospital
 Ambulatory Medical Care Survey (NHAMCS): comparison of emergency department
 data. Academic emergency medicine. 2009;16(3):261-9.
- Fix J, Ising AI, Proescholdbell SK, Falls DM, Wolff CS, Fernandez AR, et al. Linking
 Emergency Medical Services and Emergency Department Data to Improve
 Overdose Surveillance in North Carolina. Public Health Rep.
 2021;136(1 suppl):54S-61S.
- 404
 9. Zwald ML, Holland KM, Bowen DA, Simon TR, Dahlberg LL, Stein Z, et al. Using the
 405
 406
 406
 407
 407
 408
 409
 409
 409
 409
 409
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400
 400</li
- 40810.Emergency Department Syndrome & Custom Event Definitions. Carolina Center for409Health Informatics in the Department of Emergency Medicine, University of North410Carolina at Chapel HIII & NC Division of Health and Human Services, Division of411Public Health. 2023. https://ncdetect.org/case-definitions/

| 412 413 414 | 11. | Ngo QM, Sigel E, Moon A, Stein SF, Massey LS, Rivara F, et al. State of the science: a scoping review of primary prevention of firearm injuries among children and adolescents. Journal of Behavioral Medicine. 2019;42(4):811-29. |
|--------------------------|-----|--|
| 415 416 | 12. | Fleegler EW. Our Limited Knowledge of Youth Suicide Risk and Firearm Access. JAMA Network Open. 2021;4(10):e2127965-e. |
| 417 418 419 | 13. | El Ibrahimi S, Xiao Y, Bergeron CD, Beckford NY, Virgen EM, Smith ML. Suicide Distribution and Trends Among Male Older Adults in the U.S., 1999-2018. Am J Prev Med. 2021;60(6):802-11. |
| 420 421 422 423 | 14. | Morgan ER, Gomez A, Rivara FP, Rowhani-Rahbar A. Household Firearm Ownership and Storage, Suicide Risk Factors, and Memory Loss Among Older Adults: Results From a Statewide Survey. Annals of Internal Medicine. 2019;171(3):220-2. |
| 424 425 426 | 15. | Mulcahy LB, Goyal MK, Cohen J. Documentation of Firearm Safety Screening During Emergency Department Visits for Assault-Related Injury. Health Promotion Practice. 2021;23(6):941-3. |
| 427 428 429 430 | 16. | Akinyemi OA, Weldeslase TA, Hughes K, Williams M, Cornwell EE. The Distressed Communities Index: A Measure of Community-Level Economic Deprivation and Rate of Firearm Injuries in Maryland. The American Surgeon [™] . 2023:00031348231191243. |
| 431 432 433 | 17. | ICD-10-CM Official Guidelines for Coding and Reporting: FY 2023 UPDATED April 1, 2023. 2023. https://www.cms.gov/files/document/fy-2023-icd-10-cm-coding-guidelines-updated-01/11/2023.pdf |
| 434 | | |




Fig 1

