

1 Acute intermittent hypoxia in people living with chronic stroke – a preliminary study to
2 examine safety and efficacy as a neurorehabilitation intervention.

3

4 Gregory EP Pearcey^{1,2}, Alexander J Barry³, Milap S Sandhu^{2,3}, Timothy Carroll⁴, Elliot J
5 Roth^{2,3}, and W Zev Rymer^{2,3}

6

7 1. Memorial University of Newfoundland

8 2. Northwestern University

9 3. Shirley Ryan Ability Lab

10 4. University of Chicago

11

12 Correspondence:

13 Gregory Pearcey: gearcey@mun.ca

14 Z Rymer: zrymer@sralab.org

15 **ABSTRACT**

16

17 **Background and Purpose:** Acute intermittent hypoxia (AIH) is a novel therapeutic
18 intervention that has the potential to facilitate recovery of function, but its safety and
19 efficacy have not been tested in people with stroke. The purpose here was to examine
20 whether AIH is safe and effective in people with stroke.

21

22 **Methods:** Participants (n=10) with a unilateral, ischemic, hemispheric stroke were
23 assessed before and following 4 sessions of AIH. Clinical tests and upper limb strength
24 were assessed before, ~15-30 minutes, and ~60 minutes after the intervention.

25

26 **Results:** AIH was well-tolerated and there were no adverse events observed. Although
27 no changes in strength were detected for the less-affected limb, grip strength and elbow
28 flexion force of the more-affected limb was increased after AIH.

29

30 **Conclusions:** AIH appears to be potentially safe and effective for improving strength in
31 the more-affected limb in people with stroke. Future work should explore the use of AIH
32 to enhance task-specific training-induced plasticity.

33 INTRODUCTION

34
35 Stroke is a leading cause of long-term disability worldwide. Despite some spontaneous
36 recovery following a stroke, many people with stroke show substantial residual functional
37 impairments that significantly affect their independence and quality of life. Novel
38 interventions to alleviate impairment in people with stroke are urgently needed. Several
39 therapies promote strength and endurance, but few capitalize on the unique potential of
40 the central nervous system to reorganize following injury.

41
42 Acute intermittent hypoxia (AIH), a protocol that involves brief respiratory exposure to
43 reduced oxygen concentrations, has been found to augment neuroplasticity and promote
44 functional recovery after spinal cord injury (SCI)¹. Although the pathology of stroke differs
45 from SCI, stroke is often accompanied by damage to white matter projections. Therefore,
46 there is great potential for similar therapeutic benefits arising from transient reductions in
47 oxygen on damaged cerebral projections. Published results in people with SCI have
48 shown enhanced strength and coordination rather quickly after AIH, which opens a
49 temporary window for enhanced neuroplasticity after a single intervention².

50
51 The purpose of this study was to investigate the safety and efficacy of AIH in people with
52 stroke. The objectives were two-fold, to examine whether brief reductions in inhaled
53 oxygen 1) are safely tolerated, and 2) improve upper limb strength. We hypothesized that
54 AIH would be safely tolerated and increase strength on the more affected side.

55 METHODS

56 *Participants*

57
58 Sixteen people were recruited and consented to complete the study, but only 10/16
59 completed all procedures. Data are thus reported on 10 individuals (58.2±5.8 years, 2
60 females) living with chronic stroke (10.4±7.1 years since injury). The remaining 6
61 participants were either withdrawn due to resting heart rates <50 bpm, dropped out due
62 to positive tests for Covid-19, or did not qualify following the screening session. The
63 experimental protocol was registered (NCT04019522) and participants provided informed
64 and written consent to procedures approved by the local ethics committee at
65 Northwestern University (IRB protocol STU00208610).

66 *Experimental sessions*

67
68
69 Participants attended a free-standing academic rehabilitation hospital for six visits. The
70 first visit was a baseline assessment, which included a structural magnetic resonance
71 imaging (MRI) scan; a medical history, demographics, and medications; a pregnancy test
72 for women of childbearing potential; performance of the outcome assessments; medical
73 monitoring; and a blood draw. The next four sessions had a consistent format, but the
74 level of inspired oxygen during AIH was modified from 21% (i.e., room air) in the first
75 session, reduced by 4% each session to reach 9% in the final experimental session. In
76 each experimental session, participants performed outcome assessments before and
77
78

79 after 15 sequences of brief oxygen reductions while vital functions were monitored. After
80 AIH, blood was drawn, and a neurological assessment was performed by a trained
81 physician. Each visit was separated by at least 48 hours. The final visit was a follow-up
82 assessment, which included a structural MRI scan; performance of the outcome
83 assessments; medical monitoring; and a blood draw.

84

85 ***AIH intervention***

86

87 Each session utilized a customized Hyp-123 device (Hypoxico, Gardiner, NY) to switch
88 between hypoxic air and normoxic air. Participants breathed lowered O₂ concentrations
89 for 30-60 seconds, followed immediately by breathing room air for 60-90 seconds, and
90 this was repeated for 15 cycles. The total duration of each intervention was approximately
91 30 minutes and O₂ concentrations were monitored with a Maxtec Handi+ oxygen analyzer
92 (Maxtec, Salt Lake City, UT).

93

94 Since this was the first study using the AIH protocol in people with stroke, a titration
95 approach was used for the O₂ concentrations across sessions and was decreased along
96 the above-mentioned schedule. This allowed the capacity for monitoring participants for
97 adverse events while slowly increasing the hypoxia levels to identify any potential issues.
98 A side effect questionnaire was given before, during, and after AIH where participants
99 were instructed to inform study staff if they experienced any notable changes.

100

101 ***Physiological monitoring***

102

103 Participants oxygen saturation levels (SpO₂) and heart rate were monitored with a Nonin
104 7500 SpO₂ monitor (Nonin Medical Inc., Plymouth, MN), and a 5-lead electrocardiogram
105 was read in real time by a registered nurse. Blood pressure was measured before, during,
106 and after the interventions.

107

108 ***Outcome Assessments***

109

110 ***Clinical assessments***

111

112 At their first visit, all participants underwent a battery of clinical assessments, prior to AIH
113 administration. A medical history was taken by the physician to confirm that the participant
114 met inclusion and exclusion criteria and that there were no other co-morbidities.
115 Participants underwent a structural brain MRI before the intervention to confirm size and
116 location of the stroke lesion, and after the intervention to determine whether the stroke
117 lesion size changed. Participants then received a 12-lead EKG assessment at rest, which
118 the study physician used to identify whether cardiac abnormalities were present prior to
119 intervention. A baseline blood draw was assessed for Troponin levels, and a second
120 blood draw was assessed for Troponin levels after each intervention session, before each
121 participant was discharged.

122

123 Prior to and following each intervention, a trained clinician performed: neurological
124 assessments to determine whether there were changes in cognition (NIH Stroke Scale,

125 a measure of stroke severity); cranial nerve assessment; a standard muscle strength test;
126 the Brunnstrom scale to assess level of motor recovery; sensory changes; reflexes;
127 coordination; assessment of heart and lung status; Fugl-Meyer test, a quantitative
128 measure of motor impairment; Chedoke McMaster Stroke Assessment, a measure of
129 impairment of the hand; Modified Ashworth Scale for Spasticity, and Delis Kaplan
130 Executive Function System Color-Word Interference Test, an assessment of verbal
131 processing and executive functioning.

132

133 *Strength testing*

134

135 Elbow flexion strength was assessed using a custom elbow dynamometer that utilizes a
136 TAS510 tension/compression load cell (HTC Sensors, Xi'An, China) to quantify elbow
137 flexion force. Grip strength was measured using a Jamar analog hand dynamometer
138 (Jamar Technologies, Hatfield, PA). For both elbow flexion and grip strength, the peak
139 force obtained during a three second maximal voluntary contraction was recorded. w60
140 seconds rest was provided before the next trial. A total of three trials of each test was
141 performed in each limb. The right limb was tested first, followed by the left limb and the
142 mean of three trials was used for further analysis.


143 RESULTS

144

145 Safety

146

147 Our primary objective was to determine if AIH is safely tolerated by people with stroke.
148 Across participants and sessions, there were no adverse consequences observed in
149 clinical assessments, neurological assessments, or troponin levels after AIH. Participants
150 tolerated breathing 9% O₂, which caused SpO₂ to drop to a nadir of $86.9 \pm 0.9\%$ ($P < 0.001$;
151 fig.1A) and heart rate to increase by 7.3 ± 4.8 beats per minute ($P < 0.001$; fig.1B).
152 Significant differences in apparent diffusion coefficient (ADC) values were observed in
153 the MRI scans but secondary analysis found that voxels with drops in ADC were disjoint
154 and located on the periphery of the brain. Thus, they were deemed to be a result of
155 imperfection in the image registration (i.e., noise), further excluding negative
156 consequences in brain morphology because of the AIH intervention.
157



158

159 **Figure 1: Physiological monitoring during acute intermittent hypoxia (AIH).** A) the
160 nadir oxygen saturation reached during each sequence of AIH. B) the maximum heart
161 rate reached during each sequence of AIH. In all plots, the 21% O₂ condition is in white,
162 while the 9% O₂ condition is in cyan. Error bars are 95% confidence limits. All timepoints
163 were significantly different between conditions in A and B.


164

165 Strength

166

167 The secondary objective was to determine if AIH improves strength in the upper limb of
168 stroke survivors. To account for within-participant variability, we computed the percent
169 change from baseline during the breathing of 21% and 9% O₂ (AIH). The change in elbow
170 flexion strength in the more affected limb during the 9% O₂ condition ($4.92 \pm 3.34\%$
171 increase from baseline) was significantly greater than the change during the 21% O₂
172 condition ($7.6 \pm 3.34\%$ decrease from baseline; $p = 0.003$; fig.2A). Similarly, in the more
173 affected limb, the change in grip strength during the 9% O₂ condition ($12.9 \pm 6.38\%$
174 increase from baseline) was significantly greater than the change during the 21% O₂
175 condition ($12 \pm 3.81\%$ decrease from baseline; $p = 0.012$, fig.2C). The change in grip

176 strength and elbow flexion strength from baseline was not different between conditions in
177 the less affected limb (fig.2B and 2D).
178



179
180

181 **Figure 2: Percent change in strength.** A) more-affected elbow flexion, B) less-affected
182 elbow flexion, C) more-affected grip, and D) less-affected grip. The 21% O₂ condition is
183 white, the 9% O₂ condition is cyan, error bars are 95% confidence limits, and * indicates
184 $p < 0.05$ for the effect of condition.

185 **DISCUSSION**

186
187 The primary purpose of this preliminary study was to investigate the safety of AIH in
188 chronic hemispheric stroke survivors. Our secondary objective was to determine if AIH
189 augments volitional strength after stroke. In support of our first objective, we showed that
190 AIH was tolerated without negative consequences. The observed ~5 and ~13% increase
191 in volitional strength of the hand and elbow musculature supports the assertion proposed
192 in our second objective, that AIH can improve volitional strength after stroke. The
193 mechanisms underlying the increases in strength remain speculative but might include
194 synaptic plasticity³, and/or improved neural drive to the motor pool⁴, which improves
195 motor unit output⁵.

196
197 Several limitations should be acknowledged: The sample size was small, and although
198 we did not observe any adverse events, medical monitoring will be important in future
199 trials. Additionally, we only administered a single bout of AIH to these individuals, and the
200 cumulative effects of hypoxia must be monitored. Repetitive AIH may enhance
201 neuroplasticity, and thus may provide greater therapeutic benefit than a single session².

202
203 **Conclusions**

204
205 Our findings suggest that AIH is potentially safe and effective for improving limb function
206 in people with stroke. Although AIH is a promising intervention, further investigation is
207 required to confirm safety and to maximize therapeutic outcomes.

208 **Acknowledgements**

209 Thanks to Leah O'Shea for assisting data collection and analysis, Deena Hasselballa for
210 help with neurological assessments, and the nurses for monitoring vitals.

211

212 **Sources of Funding**

213 This study was supported by the American Heart Association (AHA GRANT
214 #18IPA34170025).

215

216 **Disclosures**

217 The authors have nothing to disclose, financial or otherwise.

218 **References**

219

220 1. Vose AK, Welch JF, Nair J, Dale EA, Fox EJ, Muir GD, Trumbower RD, Mitchell GS.
221 Therapeutic acute intermittent hypoxia: A translational roadmap for spinal cord injury
222 and neuromuscular disease. *Exp Neurol*. 2022;347:113891.

223 2. Welch JF, Sutor TW, Vose AK, Perim RR, Fox EJ, Mitchell GS. Synergy between
224 Acute Intermittent Hypoxia and Task-Specific Training. *Exerc Sport Sci Rev*.
225 2020;48:125–132.

226 3. Christiansen L, Urbin MA, Mitchell GS, Perez MA. Acute intermittent hypoxia
227 enhances corticospinal synaptic plasticity in humans. *Elife*. 2018;7:e34304.

228 4. Afsharipour B, Pearcey GEP, Rymer WZ, Sandhu MS. Acute intermittent hypoxia
229 enhances strength, and modulates spatial distribution of muscle activation in
230 persons with chronic incomplete spinal cord injury. *Experimental Neurology*.
231 2023;367:114452.

232 5. Pearcey GEP, Afsharipour B, Holobar A, Sandhu MS, Rymer WZ. Acute intermittent
233 hypoxia-induced increases maximal motor unit discharge rates in people with
234 chronic incomplete spinal cord injury. *medRxiv*. 2023. Available from:
235 <https://www.medrxiv.org/content/10.1101/2023.05.22.23290235v1>

236