- 1 Acute intermittent hypoxia in people living with chronic stroke a preliminary study to
- 2 examine safety and efficacy as a neurorehabilitation intervention.
- 3
- Gregory EP Pearcey^{1,2}, Alexander J Barry³, Milap S Sandhu^{2,3}, Timothy Carroll⁴, Elliot J
 Roth^{2,3}, and W Zev Rymer^{2,3}
- 6
- 7 1. Memorial University of Newfoundland
- 8 2. Northwestern University
- 9 3. Shirley Ryan Ability Lab
- 10 4. University of Chicago
- 11
- 12 Correspondence:
- 13 Gregory Pearcey: <u>gearcey@mun.ca</u>
- 14 Z Rymer: <u>zrymer@sralab.org</u>

15 **ABSTRACT**

16

Background and Purpose: Acute intermittent hypoxia (AIH) is a novel therapeutic intervention that has the potential to facilitate recovery of function, but its safety and efficacy have not been tested in people with stroke. The purpose here was to examine whether AIH is safe and effective in people with stroke.

21

Methods: Participants (n=10) with a unilateral, ischemic, hemispheric stroke were assessed before and following 4 sessions of AIH. Clinical tests and upper limb strength were assessed before, ~15-30 minutes, and ~60 minutes after the intervention.

25

Results: AIH was well-tolerated and there were no adverse events observed. Although
 no changes in strength were detected for the less-affected limb, grip strength and elbow

- 28 flexion force of the more-affected limb was increased after AIH.
- 29

30 **Conclusions:** AlH appears to be potentially safe and effective for improving strength in

- the more-affected limb in people with stroke. Future work should explore the use of AIH
- 32 to enhance task-specific training-induced plasticity.

33 INTRODUCTION

34

Stroke is a leading cause of long-term disability worldwide. Despite some spontaneous recovery following a stroke, many people with stroke show substantial residual functional impairments that significantly affect their independence and quality of life. Novel interventions to alleviate impairment in people with stroke are urgently needed. Several therapies promote strength and endurance, but few capitalize on the unique potential of the central nervous system to reorganize following injury.

41

42 Acute intermittent hypoxia (AIH), a protocol that involves brief respiratory exposure to 43 reduced oxygen concentrations, has been found to augment neuroplasticity and promote 44 functional recovery after spinal cord injury (SCI)¹. Although the pathology of stroke differs 45 from SCI, stroke is often accompanied by damage to white matter projections. Therefore, there is great potential for similar therapeutic benefits arising from transient reductions in 46 oxygen on damaged cerebral projections. Published results in people with SCI have 47 shown enhanced strength and coordination rather guickly after AIH, which opens a 48 49 temporary window for enhanced neuroplasticity after a single intervention². 50

51 The purpose of this study was to investigate the safety and efficacy of AIH in people with 52 stroke. The objectives were two-fold, to examine whether brief reductions in inhaled 53 oxygen 1) are safely tolerated, and 2) improve upper limb strength. We hypothesized that 54 AIH would be safely tolerated and increase strength on the more affected side.

55 56 **METHODS**

57 58 *Participants*

59

60 Sixteen people were recruited and consented to complete the study, but only 10/16 completed all procedures. Data are thus reported on 10 individuals (58.2±5.8 years, 2 61 62 females) living with chronic stroke (10.4±7.1 years since injury). The remaining 6 participants were either withdrawn due to resting heart rates <50 bpm, dropped out due 63 to positive tests for Covid-19, or did not qualify following the screening session. The 64 65 experimental protocol was registered (NCT04019522) and participants provided informed and written consent to procedures approved by the local ethics committee at 66 67 Northwestern University (IRB protocol STU00208610).

68

69 Experimental sessions

70

71 Participants attended a free-standing academic rehabilitation hospital for six visits. The 72 first visit was a baseline assessment, which included a structural magnetic resonance imaging (MRI) scan; a medical history, demographics, and medications; a pregnancy test 73 74 for women of childbearing potential; performance of the outcome assessments; medical 75 monitoring; and a blood draw. The next four sessions had a consistent format, but the 76 level of inspired oxygen during AIH was modified from 21% (i.e., room air) in the first 77 session, reduced by 4% each session to reach 9% in the final experimental session. In 78 each experimental session, participants performed outcome assessments before and

after 15 sequences of brief oxygen reductions while vital functions were monitored. After 79 80 AIH, blood was drawn, and a neurological assessment was performed by a trained physician. Each visit was separated by at least 48 hours. The final visit was a follow-up 81 82 assessment, which included a structural MRI scan; performance of the outcome assessments; medical monitoring; and a blood draw. 83

84

85 AIH intervention

86

87 Each session utilized a customized Hyp-123 device (Hypoxico, Gardiner, NY) to switch between hypoxic air and normoxic air. Participants breathed lowered O₂ concentrations 88 89 for 30-60 seconds, followed immediately by breathing room air for 60-90 seconds, and 90 this was repeated for 15 cycles. The total duration of each intervention was approximately 91 30 minutes and O2 concentrations were monitored with a Maxtec Handi+ oxygen analyzer 92 (Maxtec, Salt Lake City, UT).

93

94 Since this was the first study using the AIH protocol in people with stroke, a titration 95 approach was used for the O2 concentrations across sessions and was decreased along 96 the above-mentioned schedule. This allowed the capacity for monitoring participants for 97 adverse events while slowly increasing the hypoxia levels to identify any potential issues. 98 A side effect questionnaire was given before, during, and after AIH where participants 99 were instructed to inform study staff if they experienced any notable changes.

100

Physiological monitoring 101

102

Participants oxygen saturation levels (SpO2) and heart rate were monitored with a Nonin 103 104 7500 SpO2 monitor (Nonin Medical Inc., Plymouth, MN), and a 5-lead electrocardiogram was read in real time by a registered nurse. Blood pressure was measured before, during, 105 106 and after the interventions.

107

108 **Outcome Assessments** 109

Clinical assessments

- 110
- 111

At their first visit, all participants underwent a battery of clinical assessments, prior to AIH 112 administration. A medical history was taken by the physician to confirm that the participant 113 met inclusion and exclusion criteria and that there were no other co-morbidities. 114 115 Participants underwent a structural brain MRI before the intervention to confirm size and location of the stroke lesion, and after the intervention to determine whether the stroke 116 117 lesion size changed. Participants then received a 12-lead EKG assessment at rest, which the study physician used to identify whether cardiac abnormalities were present prior to 118 intervention. A baseline blood draw was assessed for Troponin levels, and a second 119 120 blood draw was assessed for Troponin levels after each intervention session, before each 121 participant was discharged.

122

123 Prior to and following each intervention, a trained clinician performed: neurological 124 assessments to determine whether there were changes in cognition (NIH Stroke Scale,

a measure of stroke severity); cranial nerve assessment; a standard muscle strength test;
the Brunnstrom scale to assess level of motor recovery; sensory changes; reflexes;
coordination; assessment of heart and lung status; Fugl-Meyer test, a quantitative
measure of motor impairment; Chedoke McMaster Stroke Assessment, a measure of
impairment of the hand; Modified Ashworth Scale for Spasticity, and Delis Kaplan
Executive Function System Color-Word Interference Test, an assessment of verbal
processing and executive functioning.

- 132
- 133 Strength testing
- 134

135 Elbow flexion strength was assessed using a custom elbow dynamometer that utilizes a 136 TAS510 tension/compression load cell (HTC Sensors, Xi'An, China) to quantify elbow flexion force. Grip strength was measured using a Jamar analog hand dynamometer 137 (Jamar Technologies, Hatfield, PA). For both elbow flexion and grip strength, the peak 138 force obtained during a three second maximal voluntary contraction was recorded. w60 139 seconds rest was provided before the next trial. A total of three trials of each test was 140 141 performed in each limb. The right limb was tested first, followed by the left limb and the 142 mean of three trials was used for further analysis.

143 **RESULTS**

144

145 **Safety**

146

147 Our primary objective was to determine if AIH is safely tolerated by people with stroke. Across participants and sessions, there were no adverse consequences observed in 148 clinical assessments, neurological assessments, or troponin levels after AIH. Participants 149 tolerated breathing 9% O2, which caused SpO2 to drop to a nadir of 86.9±0.9% (P<0.001; 150 151 fig.1A) and heart rate to increase by 7.3±4.8 beats per minute (P<0.001; fig.1B). 152 Significant differences in apparent diffusion coefficient (ADC) values were observed in the MRI scans but secondary analysis found that voxels with drops in ADC were disjoint 153 154 and located on the periphery of the brain. Thus, they were deemed to be a result of 155 imperfection in the image registration (i.e., noise), further excluding negative 156 consequences in brain morphology because of the AIH intervention.





Figure 1: Physiological monitoring during acute intermittent hypoxia (AIH). A) the
nadir oxygen saturation reached during each sequence of AIH. B) the maximum heart
rate reached during each sequence of AIH. In all plots, the 21% O2 condition is in white,
while the 9% O2 condition is in cyan. Error bars are 95% confidence limits. All timepoints
were significantly different between conditions in A and B.

165 Strength

166

167 The secondary objective was to determine if AIH improves strength in the upper limb of 168 stroke survivors. To account for within-participant variability, we computed the percent 169 change from baseline during the breathing of 21% and 9% O2 (AIH). The change in elbow flexion strength in the more affected limb during the 9% O2 condition (4.92±3.34% 170 171 increase from baseline) was significantly greater than the change during the 21% O2 172 condition (7.6±3.34% decrease from baseline; p=0.003; fig.2A). Similarly, in the more 173 affected limb, the change in grip strength during the 9% O2 condition (12.9±6.38% increase from baseline) was significantly greater than the change during the 21% O2 174 condition (12±3.81% decrease from baseline; p=0.012, fig.2C). The change in grip 175

strength and elbow flexion strength from baseline was not different between conditions in

the less affected limb (fig.2B and 2D).

178



179 180

Figure 2: Percent change in strength. A) more-affected elbow flexion, B) less-affected
 elbow flexion, C) more-affected grip, and D) less-affected grip. The 21% O2 condition is
 white, the 9% O2 condition is cyan, error bars are 95% confidence limits, and * indicates

184 *p*<0.05 for the effect of condition.

185 **DISCUSSION**

186

The primary purpose of this preliminary study was to investigate the safety of AIH in 187 188 chronic hemispheric stroke survivors. Our secondary objective was to determine if AIH augments volitional strength after stroke. In support of our first objective, we showed that 189 190 AIH was tolerated without negative consequences. The observed ~5 and ~13% increase 191 in volitional strength of the hand and elbow musculature supports the assertion proposed 192 in our second objective, that AIH can improve volitional strength after stroke. The mechanisms underlying the increases in strength remain speculative but might include 193 194 synaptic plasticity³, and/or improved neural drive to the motor pool⁴, which improves 195 motor unit output⁵.

196

197 Several limitations should be acknowledged: The sample size was small, and although 198 we did not observe any adverse events, medical monitoring will be important in future 199 trials. Additionally, we only administered a single bout of AIH to these individuals, and the 200 cumulative effects of hypoxia must be monitored. Repetitive AIH may enhance 201 neuroplasticity, and thus may provide greater therapeutic benefit than a single session². 202

- 203 Conclusions
- 204

205 Our findings suggest that AIH is potentially safe and effective for improving limb function 206 in people with stroke. Although AIH is a promising intervention, further investigation \is 207 required to confirm safety and to maximize therapeutic outcomes.

208 Acknowledgements

Thanks to Leah O'Shea for assisting data collection and analysis, Deena Hasselballa for help with neurological assessments, and the nurses for monitoring vitals.

211

212 Sources of Funding

This study was supported by the American Heart Association (AHA GRANT #18IPA34170025).

215

216 Disclosures

217 The authors have nothing to disclose, financial or otherwise.

218 **References**

- 219
- Vose AK, Welch JF, Nair J, Dale EA, Fox EJ, Muir GD, Trumbower RD, Mitchell GS.
 Therapeutic acute intermittent hypoxia: A translational roadmap for spinal cord injury
 and neuromuscular disease. *Exp Neurol*. 2022;347:113891.
- Welch JF, Sutor TW, Vose AK, Perim RR, Fox EJ, Mitchell GS. Synergy between
 Acute Intermittent Hypoxia and Task-Specific Training. *Exerc Sport Sci Rev.* 2020;48:125–132.
- Christiansen L, Urbin MA, Mitchell GS, Perez MA. Acute intermittent hypoxia
 enhances corticospinal synaptic plasticity in humans. *Elife*. 2018;7:e34304.
- Afsharipour B, Pearcey GEP, Rymer WZ, Sandhu MS. Acute intermittent hypoxia enhances strength, and modulates spatial distribution of muscle activation in persons with chronic incomplete spinal cord injury. *Experimental Neurology*.
 2023;367:114452.
- Pearcey GEP, Afsharipour B, Holobar A, Sandhu MS, Rymer WZ. Acute intermittent hypoxia-induced increases maximal motor unit discharge rates in people with chronic incomplete spinal cord injury. *medRxiv*. 2023. Available from: https://www.medrxiv.org/content/10.1101/2023.05.22.23290235v1

236