It is made available under a CC-BY 4.0 International license .

## Geographical variation in dementia: systematic review with meta-analysis

Walesby K.E.<sup>1</sup>\*, Shenkin, S.D.<sup>2</sup>, Burton J.K.<sup>3</sup>, Dunlop C.<sup>4</sup>, Fenton C.<sup>5</sup>,

Gray D<sup>.6</sup>, McGillicuddy L.A.<sup>7</sup>, Starr J.M.<sup>1</sup>, Wilkinson T.<sup>8</sup>, Muniz Terrera G.<sup>9</sup>, Russ T.C.<sup>1</sup>

<sup>1</sup>Alzheimer Scotland Dementia Research Centre and Centre for Cognitive Ageing and Cognitive Epidemiology, 7 George Square, University of Edinburgh, UK

<sup>2.</sup> Ageing and Health Research Group, Centre for Population Health Sciences, and Advanced Care Research Centre, Usher Institute, University of Edinburgh

<sup>3.</sup> Academic Geriatric Medicine, School of Cardiovascular and Metabolic Health, College of Medical, Veterinary & Life Sciences, University of Glasgow

<sup>4.</sup> Old Age Psychiatry, Edinburgh Royal Infirmary, NHS Lothian, UK

<sup>5.</sup> NESSIE, Usher Institute, University of Edinburgh

<sup>6.</sup> Fairnington Centre, Hexham General Hospital, Cumbria, Northumberland, Tyne and Wear NHS Foundation Trust (CNTW), UK

<sup>7.</sup> Old Age Psychiatry, Royal Edinburgh Hospital, NHS Lothian, UK

<sup>8.</sup> Centre for Clinical Brain Sciences, University of Edinburgh, UK

<sup>9.</sup> Department of Social Medicine, Ohio University, Ohio University Heritage College of Osteopathic Medicine and Edinburgh Dementia Prevention, University of Edinburgh, UK

\*Corresponding author. Alzheimer Scotland Dementia Research Centre, 7 George Square, University of Edinburgh, EH8 9JZ. E-mail: <u>Katherine.walesby@ed.ac.uk</u>

## ABSTRACT

**Background:** Understanding geographical variation of dementia could highlight important modifiable socio-environmental risk factors. A previous systematic review (2012) identified an increased risk of Alzheimer dementia in in rural living in High-Income Countries (HICs), with a dearth of studies in Low to Middle-Income Countries (L-MICs). We updated this review to examine geographical variations in dementia, to encompass the growing number of studies in this field.

**Methods:** We systematically reviewed the literature for cross-sectional or longitudinal observational studies that compared dementia incidence or prevalence between two or more geographical areas including rural and urban settings.

We conducted a narrative synthesis of included papers. Where possible, we undertook metaanalysis, generating odds ratios for rural versus urban dementia prevalence and stratified the analysis by HICs and L-MICs.

**Results:** We identified 38 relevant papers, encompassing approximately 98,502,147 people. Twenty-seven papers were included in the quantitative synthesis. Study methodologies varied widely. Dementia rates varied geographically (0.43-38.5%). Overall, rural living was associated with small increased odds of dementia (OR, 1.20, 95% CI 1.03-1.40; *P* value = 0.0182). Stratification by HICs and L-MICs demonstrated further variation, with increased odds of dementia in rural areas in L-MICs but not HICs.

### Conclusions

There is some evidence of geographical variation of dementia. Rural living was associated with small increased odds of dementia, with stratification showing evidence in rural areas of L-MICs but not HICs. We believe this has not been reported previously. Future research

It is made available under a CC-BY 4.0 International license .

must consider life course geographical exposure and addressing heterogeneity in definitions of 'rural' and 'urban.'

Keywords: Dementia, Alzheimer disease, epidemiology, geography

#### What this study adds

We confirm that rural living (compared to urban living) is associated with a small increased odds of dementia (OR 1.20, 95%CI 1.03-1.40). We demonstrate for the first time that this is driven by increased odds of dementia in rural areas in Low to Middle-Income Countries (L-MICs) rather than High Income Countries (HICs), and that the odds of dementia were higher in urban areas in large studies in HICs. Future studies need to carefully consider study setting, method of dementia ascertainment, when exposures may occur, and risk of bias, to understand the role of environment and geography in dementia risk.

## **INTRODUCTION**

Dementia is a growing worldwide public health concern. The epidemiology of this complex syndrome is not yet fully understood.<sup>1</sup> Age and known genetic, health, and lifestyle risk factors play an important role in dementia incidence, but together they do not fully explain dementia aetiology. Life course geographical variation might influence dementia risk or protective factors.

A systematic review in 2012 found "evidence of geographical variation in rates of dementia in affluent countries at a variety of geographical scales."<sup>2</sup> Furthermore, an increased risk of Alzheimer's dementia was associated with rural living (Prevalence OR 1.50, 90% CI 1.33-1.69), with early life rural living further increasing this risk to approximately double (OR 2.22, 90% CI 1.19-4.16). However, these results were based on very few studies, and there

It is made available under a CC-BY 4.0 International license .

have been several subsequent studies. We therefore updated this systematic review<sup>2</sup> to reflect the developing body of evidence in this field. In our present systematic review, we report a comprehensive overview of the literature specifically examining the influence of rural and urban living on risk of dementia.

#### METHODS

## **Registration and reporting**

We prospectively registered the protocol with PROSPERO, reference CRD42016050323 (http://www.crd.york.ac.uk/) and report the review in accordance with the Preferred Reporting of Items in Systematic Reviews and Meta-Analyses (PRISMA) guidance.<sup>3,4</sup>

#### **Information sources**

We searched multiple databases on 18<sup>th</sup>-31st October 2018 for papers since the published review<sup>2</sup> search date of 15<sup>th</sup> April 2010: search information and indicative search strategies were developed with an Information Specialist (CF), (Appendix 1).

#### **Eligibility criteria**

Eligible studies included: cross-sectional or longitudinal studies of any duration comparing primary dementia rates (prevalence or incidence) between two or more different geographical sites. Our initial search was broad then focused to rural/urban; therefore, many studies (117) were excluded as 'wrong comparator.'

It is made available under a CC-BY 4.0 International license .

We excluded studies purely on young onset dementia (<60 years old) or mild cognitive impairment. We did not restrict to English language (Appendix 2).

#### Study selection and data collection

Two reviewers (from KEW, JKB, and TCR) independently screened all titles and abstracts using Covidence,<sup>5</sup> and two then reviewed full text with conflicts resolved by discussion. Google translate was used for data extraction from papers not in English.

Two of KEW, LAM, CD, or DG extracted data using a bespoke and piloted form with discrepancies resolved by discussion and reference to the original paper.

Where articles did not report odds ratios for dementia risk in rural and urban areas,<sup>6-16</sup> we computed these if the relevant raw data were reported or contacted authors<sup>9,17-20</sup> : data provided by two authors.<sup>18,19</sup> We contacted authors regarding full-text publications from abstracts identified, and overlap between studies. When both unadjusted and multivariable-adjusted odds ratios were reported, we report unadjusted odds ratio. When both DSM-IV and 10/66 method were used for dementia ascertainment, we report the 10/66 results<sup>21-23</sup> (as this has shown greater dementia ascertainment in LICs and LMICs<sup>24,25</sup>) except when the 10/66 data were incomplete.<sup>26</sup>

It is made available under a CC-BY 4.0 International license .

#### Strategy for data synthesis

Where sufficient data were available, we conducted a random effects meta-analysis of odds ratios. Post-hoc we decided to explore the data by stratifying by Higher Income Countries (HICs) and Low to Middle-Income Countries (L-MICs)<sup>27</sup> (Appendix 3), and further by study size. We recognise this classification of country income can change with time (Appendix 3). Between-study heterogeneity was assessed by the  $I^2$  statistic. Studies which could not be

combined quantitatively were reported in a narrative synthesis.

## Risk of bias assessment in individual studies

KEW, SDS, and TCR assessed the risk of bias using an adapted Risk of Bias Assessment Tool for Non-Randomized studies (RoBANS), (Appendix 4), with disagreements resolved by discussion.

It is made available under a CC-BY 4.0 International license .

#### RESULTS

The narrative synthesis includes 38 new papers (**Table 1**), 27 of which – including 33 cohorts - could be meta-analysed (**Figure 1**).

#### Geographical locations and sample sizes

Studies were predominantly in the Americas (N=11) and Western Pacific region (especially China, N=11), and also Europe (N=9) South-East Asia (N=4), Africa (N=3), and Eastern Mediterranean region (N=2) (one study covered three areas). Of the meta-analysis papers, 12 papers (13 studies) were based in HICs and 15 papers (20 studies) in L-MICs.

In the 35 papers that included a sample size, this ranged from 90 to 73,000,000 people, median 2170. Of 27 papers in the meta-analysis, study samples ranged from 300 to 21,624,228 people, median 2162.<sup>6-11,13,15,16,18,19,21,22,28-39</sup> The definition of 'rural' and 'urban' was heterogeneous. Too few papers included information on education, deprivation, or other socio-economic factors in relation to rural or urban settings and subsequent dementia risk to include in the meta-analysis. Three papers considered childhood and adulthood rural or urban residence and risk of dementia,<sup>18,40,41</sup> only one was included in the meta-analysis.<sup>18</sup>

It is made available under a CC-BY 4.0 International license .

#### **Dementia diagnosis**

There was considerable heterogeneity in dementia ascertainment methodologies (**Appendix 5**, **Supplementary Table 1**), mostly using surveys/interviews, clinical assessment and ICD codes. Dementia prevalence ranged from 0.43%-38.5% of the study populations in all studies, and 2.4-21% in the meta-analysis studies. Few identified dementia subtype. Most papers compared dementia against non-dementia or 'normal', with some including cognitive impairment/MCI in the 'no dementia' group, and others excluding those with MCI.

It is made available under a CC-BY 4.0 International license .



1

- 2 Figure 1: PRISMA flowchart detailing selection of studies for inclusion in the review:
- 3 rural/urban differences in dementia

4

5

6

	Year			Study Population		Dementia <i>n</i> (% of
Author	(data collection)	Country	Age (years)	(n) Urban n (% of total)		total)
Abner 2016* <sup>28</sup>	2007-2013	USA	Any	1,209,976	766,250 (63.3)	82,278 (6.8)
					Major cities = 139,071 (53.1%)	
Astell_Burt 2018 <sup>*29</sup>	2006-2009	Australia	>=45	261,669	Inner regional area = 92,734 (35.4)	3046 (1.2)
Chammartin 2016 <sup>42</sup>	2008-2012	Switzerland	Any	60,388	50,388 N/A	
Chen 2017 <sup>30</sup> *	2001-2013	Canada	55–85	2,066,639	1,674,591 (81.0)	257,816 (12.4)
Chen 2012 <sup>31</sup> Anhui						
study (a)*	2001-2003	China	>=65	2,917	1,736 (59.5)	210 (7.2)
Chen 2012 <sup>31</sup> Four						
provinces study						
(b)*	2008-2009	China	>=60	3,327	1,787 (53.7)	341 (10.2)
			>=65			
			(mean 72.9 +/-			
Contador 2015 <sup>41</sup>	1994-1998	Spain	6.1)	2,711	852 (31.4)	91 (3.4)

## 7 Table 1: Executive summary of studies meeting inclusion criteria: rural/urban differences in dementia

	Year			Study Population		Dementia <i>n</i> (% of
Author	(data collection)	Country	Age (years)	( <i>n</i> )	Urban <i>n</i> (% of total)	total)
Cornutiu 2010 <sup>43</sup>	1980-2006	Romania	Any	Unclear	N/A	N/A
Deng 2018* <sup>32</sup>	Oct-Dec 2015	China	≥60	1,781	464 (26.1)	195 (10.9)
			>=60			
Fronza da Silva			(mean 72.3 +/-			
2013 <sup>44</sup>	July-Oct 2007	Brazil	7.7)	229	131 (57.2)	11 (4.8)
Ganesh 2010 <sup>17</sup>	2003	USA	any	31,660	N/A	427 (1.3)
Goncalves-Pereira						
2017* <sup>21</sup>	N/A	Portugal	≥65	1,405	690 (49.5)	DSM IV: 51 (7.4)
Goodman 2017* <sup>33</sup>	2011-2013	USA	>=68	21,624,228	16,477,662 (76.2)	3,110,654 (18.9)
		Central				
Guerchet (a)		African				
2013* <sup>26</sup>	2011-2012	Republic	≥65	973	500 (51.4)	72 (7.4)
Guerchet (b)		Republic of				
2013* <sup>26</sup>	2011-2012	Congo	≥65	1,029	500 (48.5)	63 (6.1)

	Year			Study Population		Dementia <i>n</i> (% of
Author	(data collection)	Country	Age (years)	( <i>n</i> )	Urban <i>n</i> (% of total)	total)
Hendrie 2018 <sup>*18</sup>	1992-2009	USA	>=70	3,259	2326 (71.4)	285 (8.7)
			Any			
			(mean 58.7 +/-			
Huang 2014 <sup>45</sup>	1997-2007	Taiwan	14)	142,744	41,458 (29.0)	612 (0.43)
Jia 2014 <sup>38</sup> *	2008-2009	China	≥65	10,276	6,096 (59.3)	528 (5.1)
Khedr 2015 <sup>6</sup> *	2011-2013	Egypt	>=60	691		35 (5.1)
Kim 2011* <sup>39</sup>	2008	South Korea	>=65	1,673	N/A	351 (21.0)
Koller 2010 <sup>15</sup> *	2004-2006	Germany	≥65	9,216	6,578 (71.4)	1846 (20.0)
			>60			
Nadel 2014 <sup>40</sup>	2011	Costa Rica	(mean 76.8+8.1)	90	N/A	2 (2.3)
Nunes 2010* <sup>34</sup>	2003	Portugal	55-79	1,146	713 (62.2)	31 (2.7)
		Central				
		African				
Pilleron 2015 <sup>13</sup> (a)*	2011-2012	republic	≥65	973	452 (46.4)	72 (7.5)

	Year			Study Population		Dementia <i>n</i> (% of
Author	(data collection)	Country	Age (years)	(n)	Urban <i>n</i> (% of total)	total)
		Republic of				
Pilleron 2015 <sup>13</sup> (b)*	2011-2012	Congo	≥65	1,029	479 (46.6)	63(6.1)
Poddar 2011 <sup>12</sup> *	Unclear	India	≥50	2890	793 (27.4)	147 (5.1)
Prince 2012 <sup>22</sup> * (a)	2003-2010	Peru	>=65	1933	1381	10/66: 69 (3.6)
			(mean age			DSM-IV: 53 (2.7)
			follow-up 75.7-			
			78.6)			
Prince 2012 <sup>22</sup> * (b)	2003-2010	Mexico	>=65	2003	1003	10/66: 121 (6.0)
			(mean age			DSM-IV: 67 (3.3)
			follow-up 75.7-			
			78.6)			

	Year			Study Population		Dementia <i>n</i> (% of
Author	(data collection)	Country	Age (years)	( <i>n</i> )	Urban <i>n</i> (% of total)	total)
			>=65			
			(mean age			
			follow-up 75.7-			10/66: 161 (7.4)
Prince 2012 <sup>22</sup> * (c)	2003-2010	China	78.6)	2162	1160	DSM-IV: 106 (4.9)
						2
						2
Raina 2014 <sup>7</sup> *	unclear	India	>60	1000	500 (50)	3 (2.3)
Rodríguez-Agudelo						
2011 <sup>23</sup> *	2011	Mexico	>65	2003	1003 (50.1)	171 (8.5)
						After 2 years follow
		The Republic				up, dementia =23
Samba 2015 <sup>9</sup> *	2012-2014	of Congo	≥65	966	467 (48.3)	(2.38)

	Year			Study Population		Dementia <i>n</i> (% of
Author	(data collection)	Country	Age (years)	( <i>n</i> )	Urban <i>n</i> (% of total)	total)
			>=60			
			(mean age 69.2			
Sharifi 2016 <sup>35</sup> *	2012	Iran	years, 68.7-69.7)	1257	681 (54.2)	99 (7.9)
						1999= 44,536
Taylor 2017 <sup>46</sup>	1999-2014	USA	Any age	Unclear	Unclear	2014 = 93,541
			>=60			
Tengku Aizan			(average age			
2010 <sup>10</sup> *	2004-2005	Malaysia	70.5, SD = 7.22)	2980	1635 (54.9)	418 (14.0)
			>=65			
Tola-Arribas			(mean age 76.5			
2013 <sup>11</sup> *	2009-2010	Spain	+/- 7.8)	2170	1459 (67.2)	184 (8.5)
	Unknown					
	(published 2015					
Tornau 2015 <sup>20</sup>	as abstract)	Germany	>59	Unclear	Not specified	Not specified

	Year			Study Population		Dementia <i>n</i> (% of
Author	(data collection)	Country	Age (years) (n)		Urban <i>n</i> (% of total)	total)
Trinathi 2012 <sup>36</sup> *	2003-2005	India	44-85	300	190 (63 3)	150 (50)
	2000 2000	maid	11 05		150 (05.5)	150 (50)
27						
Weden 2018 <sup>37</sup> (a)*	2000	USA	>=55	16,386	12,096 (73.8)	958 (5.8)
Weden 2018 <sup>37</sup> (b)*	2010	USA	>=55	16,311	12,410 (76.1)	727 (4.5)
Wu 2017 <sup>16</sup> *	2008-2011	England	>=65	7505	5951 (79.3)	328 (4.4)
Yin 2016 <sup>47</sup>	2006-2012	China	>65	73 million approx.	Unclear	Unclear
Yuan 2016 <sup>8</sup> *	1997-2002	China	>=55	12,881	5687 (44.2)	311 (2.4)

		Year			Study Population		Dementia <i>n</i> (% of
	Author	(data collection)	Country	Age (years)	(n)	Urban <i>n</i> (% of total)	total)
					1235 completed		
	Zhao 2010 <sup>48</sup>	1997-1998	China	>=55	phase 3	Unclear	475 (38.5)
8	AD = Alzheimer's	Disease: ADRD =	Alzheimer's R	elated Dementia: N	N/A = not available (u)	(nable to extract). Grev = not in meta-	-analysis

9 Additional information: Studies separating geographical areas e.g. "100% urban" and "mixed urban", multiple different metropolitan sizes or

10 conurbation and urban, were combined together as "urban". One study<sup>35</sup> used rural areas as the referent, and we reversed the results using

11 *natural logarithms. One study*<sup>18</sup> *had two cohorts but only raw data from authors available for one cohort to calculate Odds ratios.* 

12

It is made available under a CC-BY 4.0 International license .

#### 13 Quantitative synthesis

14	Overall, rural living w	as associated with small	increased odds of	dementia, com	pared to urban
----	-------------------------	--------------------------	-------------------	---------------	----------------

15 living (OR, 1.20, 95% CI 1.03-1.40, P value = 0.0182; N=25,274,741, N<sub>Dementia</sub>=3,462,039;

- 16 33 studies), Figure 2. However, stratifying by HICs and L-MICs status revealed a difference,
- with HICs studies showing no effect of rural vs urban (1.01, 0.88 to 1.16, P value = 0.8842;
- 18 N=25,221,600, N<sub>Dementia</sub>=3,458,633; 13 studies) and L-MICs suggesting increased odds of
- 19 dementia in rural areas (1.33, 1.06 1.68, *P* value = 0.0141; N=53,141, N<sub>Dementia</sub>=3406 20
- studies). There was high heterogeneity:  $I^2$  value = 99.78% (for all studies), 99.72% (for only
- 21 HICs studies) and 87.64% (for only L-MICs studies).
- 22 Sensitivity analysis to explore the impact of potential overlap in L-MICs study
- 23 participants<sup>9,13,22,23,26</sup> did not significantly affect the results: rural living was still linked with a
- slight increase in dementia risk (1.35, 1.02-1.79, *P* value = 0.0354,  $I^2 = 90.69\%$ ). Including
- studies from the previous systematic review also showed a slight increased risk of dementia in
- 26 rural areas, (1.18, 1.01-1.37) (**Appendix 6**).
- 27 Further stratifying HICs and L-MICs by study size confirmed in larger L-MICs studies
- 28 (N>10,000) higher odds of dementia in rural areas (or lower in urban areas; 1.56, 1.20-2.02;
- 29 N=23,157, N<sub>Dementia</sub>=839; 2 studies), unchanged when moderate-sized studies (N>1000) were
- 30 included (1.50, 1.017-1.93; N=48,439, N<sub>Dementia</sub>=2856; 14 studies). However, larger HICs
- studies (N>1,000,000) showed *lower* odds of dementia in rural areas (0.88, 0.84-0.92;
- 32 N=24,900,843, N<sub>Dementia</sub>=3,450,748; 3 studies). This effect was reduced when moderate-sized
- 33 studies (N>10,000) were included (0.93, 0.76-1.13; N=25,195,209, N<sub>Dementia</sub>=3,455,479; six
- 34 studies). Studies in L-MICs were generally smaller than those in HICs.
- 35

It is made available under a CC-BY 4.0 International license .

oodman 2017 hen 2017 hen 2017 hen 2016 Yeden 2018 Yeden 2018a High Income Countries: N>10,000 High Income Countries: N>10,000 oler 2010 V 2017 endria 2018 Dia-Anthas 2013 im 2011 oncalves-Pereira 2017 unes 2010	USA Canada USA USA USA USA Cermany England USA Spain South Korea Portugal Portugal	21,624,228 2,066,839 1,209,976 261,669 16,386 16,311 9,216 7,505 3,276 2,170 1,673 1,405	3,110,654 257,816 82,278 3,046 958 727 1,848 328 285 184	0.84 (0.84, 0.85) 0.91 (0.90, 0.92) 0.88 (0.84, 0.92) 0.63 (0.56, 0.73) 1.34 (1.10, 1.63) 1.17 (0.92, 1.46) 0.93 (0.76, 1.13) 0.96 (0.86, 1.08) 0.92 (0.64, 1.33) 1.56 (1.21, 2.01) 0.99 (0.72, 1.37)			•		
hen 2017 High Income Countries: N>1,000,000 High Income Countries: N>1,000,000 Keden 2018a High Income Countries: N>10,000 Ider 2010 Iv 2017 endrie 2018 Jaa-Arribas 2013 im 2011 noralves-Pereira 2017 unes 2010	Canada USA Australia USA USA USA Cermany England USA Spain South Korea Portugal Portugal	2,066,639 1,209,976 261,669 16,386 16,311 9,216 7,505 3,276 2,170 1,673 1,405	257,816 82,278 3,046 958 727 1,846 328 285 184	0.91 (0.90, 0.92) 0.89 (0.88, 0.91) 0.88 (0.84, 0.92) 0.63 (0.56, 0.73) 1.34 (1.10, 1.63) 1.17 (0.92, 1.46) 0.93 (0.76, 1.13) 0.96 (0.86, 1.08) 0.92 (0.64, 1.33) 1.56 (1.21, 2.01) 0.99 (0.72, 1.37)		-			
hene 2016 High Income Countries: N>1,000,000 sell-Burt 2018 feden 2018a High Income Countries: N>10,000 olier 2010 u 2017 endrie 2018 bla-Arribas 2013 im 2011 orcalvise-Pereira 2017 unes 2010	USA Australia USA USA Cermany England USA Spain South Korea Portugal Portugal	1,209.976 261,669 16,386 16,311 9,216 7,505 3,276 2,170 1,673 1,405	82,278 3,046 958 727 1,848 328 285 184	0.88 (0.88, 0.91) 0.88 (0.84, 0.92) 0.63 (0.56, 0.73) 1.34 (1.10, 1.63) 1.17 (0.92, 1.48) 0.93 (0.76, 1.13) 0.96 (0.86, 1.08) 0.92 (0.84, 1.33) 1.56 (1.21, 2.01) 0.99 (0.72, 1.37)		-	•		
High Income Countries: N>1,000,000 stell-Burt 2018 (veden 2018a (veden 2018b High Income Countries: N>10,000 oller 2010 4, 2017 endrie 2018 ole-Antibas 2013 im 2011 oncalves-Pereira 2017 unes 2010	Australia USA USA Germany England USA Spain South Korea Portugal Portugal	261,669 16,386 16,311 9,216 7,505 3,276 2,170 1,673 1,405	3.046 958 727 1.846 328 285 184	0.88 (0.84, 0.92) 0.63 (0.56, 0.73) 1.34 (1.10, 1.63) 1.17 (0.92, 1.48) 0.93 (0.76, 1.13) 0.96 (0.86, 1.08) 0.92 (0.64, 1.33) 1.56 (1.21, 2.01) 0.99 (0.72, 1.37)			•		
stel-Burt 2018 (reden 2018a (weden 2018b High Income Countries: N>10,000 oller 2010 /u 2017 endride 2018 Dia-Anthas 2013 im 2011 Concalves-Pereira 2017 unes 2010	Australia USA USA England USA Spain South Korea Portugal Portugal	261,669 16,386 16,311 9,216 7,505 3,276 2,170 1,673 1,405	3,046 958 727 1,848 328 285 184	0.63 (0.56, 0.73) 1.34 (1.10, 1.63) 1.17 (0.92, 1.48) <b>0.93 (0.76, 1.13)</b> 0.95 (0.86, 1.08) 0.92 (0.64, 1.33) 1.56 (1.21, 2.01) 0.99 (0.72, 1.37)			+		
Veden 2018a Veden 2018b <b>High Income Countries: N&gt;10,000</b> oller 2010 Av 2017 endria: 2018 ola-Ambas 2013 im 2011 concalves-Pereira 2017 unes 2010	USA USA Germany England USA Spain South Korea Portugal Portugal	16.386 16.311 9.216 7.505 3.276 2.170 1.673 1.405	958 727 1,846 328 285 184	1.34 (1.10, 1.63) 1.17 (0.92, 1.48) <b>0.93 (0.76, 1.13)</b> 0.95 (0.86, 1.08) 0.92 (0.64, 1.33) 1.56 (1.21, 2.01) 0.99 (0.72, 1.37)			+	-	
Veden 2018b High Income Countries: N>10,000 Income Countries: N>10,000 Income 2018 Jola-Ambas 2013 Im 2011 Oncalves-Pereira 2017 unes 2010	USA Germany England USA Spain South Korea Portugal Portugal	9,216 7,505 3,276 2,170 1,673 1,405	727 1,846 328 285 184	1.17 (0.92, 1.48) <b>0.93 (0.76, 1.13)</b> 0.96 (0.86, 1.08) 0.92 (0.64, 1.33) 1.56 (1.21, 2.01) 0.99 (0.72, 1.37)			+		
High Income Countries: N>10,000 olier 2010 v 2017 endrie 2018 Dia-Arribas 2013 im 2011 orcalvise-Pereitra 2017 unes 2010	Germany England USA Spain South Korea Portugal Portugal	9,216 7,505 3,276 2,170 1,673 1,405	1.846 328 285 184	0.93 (0.76, 1.13) 0.96 (0.86, 1.08) 0.92 (0.64, 1.33) 1.56 (1.21, 2.01) 0.99 (0.72, 1.37)			-		
olier 2010 V 2017 endrie 2018 Dia-Anthas 2013 im 2011 moralvis-Pereitra 2017 unes 2010	Germany England USA Spain South Korea Portugal Portugal	9,216 7,505 3,276 2,170 1,673 1,405	1,846 328 285 184	0.96 (0.86, 1.08) 0.92 (0.64, 1.33) 1.56 (1.21, 2.01) 0.99 (0.72, 1.37)			-		
va 2017 endrie 2018 ole-Antibas 2013 im 2011 oncalves-Pereira 2017 unes 2010	England USA Spain South Korea Portugal Portugal	7,505 3,276 2,170 1,673	328 285 184	0.92 (0.64, 1.33) 1.56 (1.21, 2.01) 0.99 (0.72, 1.37)					
endre 2018 ola-Arribas 2013 im 2011 oncalves-Pereira 2017 unes 2010	USA Spain South Korea Portugal Portugal	3,276 2,170 1,673	184	1.56 (1.21, 2.01)			573) 		
im 2011 im 2011 oncalves-Pereira 2017 unes 2010	Spain South Korea Portugal Portugal	2,170	184	0.99 (0.72, 1.37)					
m 2011 oncalves-Pereira 2017 unes 2010	Portugal Portugal	1,673							
oncalves-Pereira 2017 unes 2010	Portugal Portugal	1.405	351	1,17 (1.00, 1.38)				-	
unes 2010	Portugal	1,400	129	0.99 (0.69, 1.42)					
		1,146	31	2,12 (0.91, 4.96)			-	2.5	
Il High Income Countries				1.01 (0.88, 1.16)			-		
uan 2016	China	12,881	311	1.32 (1.05, 1.66)					
a 2014	China	10,276	528	1.45 (1.22, 1.73)					
Low-to-Middle Income Countries: N>10,00	)			1.40 (1.22, 1.61)				-	
hen 2012b	China	3,327	341	2.26 (1.79, 2.86)					
engku Aizan 2010	Malaysia	2,980	418	1.87 (1.52, 2.31)					
hen 2012a	China	2,917	210	4.27 (3.13, 5.84)					-
oddar 2011	India	2,890	146	1.49 (0.99, 2.24)					
rince 2012c	China	2,162	161	0.93 (0.67, 1.29)					
rince 2012b	Mexico	2,003	121	2.24 (1.51, 3.31)					
odriguez-Agudelo 2011	Mexico	2,003	171	0.99 (0.72, 1.36)			-		
rince 2012a	Peru	1,933	69	1.36 (0.82, 2.23)					
eng 2018	China	1,781	195	1.56 (0.99, 2.45)					
harifi 2016	Iran	1,257	99	1.88 (1.23, 2.86)					
uerchet 2013b	ROC	1,029	63	0.85 (0.51, 1.42)					
aina 2014	India	1,000	23	0.43 (0.18, 1.05)					
Low-to-Middle Income Countries: N>1000				1.50 (1.17, 1.93)				-	
uerchet 2013a	CAR	973	72	1.35 (0.83, 2.19)					
amba 2015	ROC	966	23	1.78 (0.75, 4.23)					
illeron 2015b	ROC	912	135	1.01 (0.60, 1.68)					
illeron 2015a	CAR	860	135	1.43 (0.88, 2.32)					
hedr 2015	Egypt	691	35	0.44 (0.22, 0.90)					
ripathi 2012	India	300	150	0.56 (0.35, 0.90)				-	
Il Low-to-Middle Income Countries				1.33 (1.06, 1.68)			-		
II Studies				1.20 (1.03, 1.40)				•	
					0.05	1			 1

36 37

38 Figure 2: Forest plot showing meta-analysed rural/urban odds ratios of dementia

39 stratified by HICs/L-MICs status and size of study: rural/urban differences in dementia<sup>1</sup>

<sup>&</sup>lt;sup>1</sup> Guerchet 2013, Pilleron 2015, Prince 2012 (Mexico only) Rodríguez-Agudelo 2011 and Samba 2015: unclear if there is overlap in these L-MICs papers (as more than one similar study in the same country). Sensitivity analysis removing the smaller cohorts (Pilleron 2015, Rodríguez-Agudelo 2011 and Samba 2015) shows no significant change in result

It is made available under a CC-BY 4.0 International license .

# 40 Narrative synthesis of non-meta-analysis studies

41	The 11 heterogeneous papers not included in the meta-analysis, were undertaken in Europe
42	(Switzerland, Romania, Germany and Spain), <sup>20,41-43</sup> South-America (Brazil), <sup>44</sup> Central
43	America (Costa Rica), <sup>40</sup> North America (USA) <sup>17,46</sup> and Asia (Taiwan and China). <sup>45,47,48</sup>
44	between 1993 and 2012. One study investigated dementia in diabetic patients, <sup>45</sup> the other
45	studies investigated dementia in general. Five studies had no age restriction, <sup>17,42,43,45,46</sup> the
46	others included people $>55/59$ years or $>60/65$ years. <sup>20,40,41,44,47,48</sup> , with sample size from 90 to
47	approximately 73 million (median, 17,186).
48	The dementia ascertainment methodologies varied greatly: data linkage with ICD codes;
49	<sup>17,20,42,43,45-47</sup> cross sectional incidence surveys; questionnaires; structured interviews and/or
50	screening by clinical assessment. <sup>40,41,44,48</sup>
51	Just over half of these studies reflected the findings of the meta-analysis, with an increase of
52	dementia in rural areas. <sup>41,43,44,46-48</sup> The results were unclear in several of these studies, <sup>20,40</sup> and
53	showed no difference in one study investigating the variation of AD drug dispensing <sup>17</sup> . Of the
54	two studies suggesting an increase in urban areas, one was specifically investigating dementia
55	(AD) deaths, <sup>42</sup> and the other newly diagnosed diabetic patients. <sup>45</sup>
56	Most studies only assessed the relationship between residence at one time point and
57	subsequent dementia, but one study <sup>41</sup> investigated childhood and adulthood residence, finding
58	childhood rural residence was linked with a non-significant trend for risk of dementia (HR =
59	1.52, 95% CI = 0.93-2.49, p=0.08), which became significant in adulthood (HR 1.61, 95% CI
60	1.06-2.46, p<0.05). <sup>41</sup> , but there was no association in late-life (HR 0.78, 95% CI 0.49-1.23,
61	p=0.29). <sup>41</sup>

62

It is made available under a CC-BY 4.0 International license .

## 63 **Risk of Bias**

- 64 Figures 3 and Appendix 7 show the risk of bias assessments for included studies. There was
- great variation in the risk of bias and, in particular, high risk in measurement of exposure
- 66 (geographical location) in approximately half of the studies.

67



69 Figure 3. Summary of Risk of Bias assessment: rural/urban differences in dementia

It is made available under a CC-BY 4.0 International license .

#### DISCUSSION

Overall, rural living was associated with small increased odds of dementia compared to urban living (OR, 1.20, 95% CI 1.03-1.40, *P* value = 0.0182; N=25,274,741, N<sub>Dementia</sub>=3,462,039; 33 studies). Further stratification by HICs and L-MICs showed this difference was mostly driven by L-MICs (1.33, 1.06 -1.68, *P* value = 0.0141; N=53,141, N<sub>Dementia</sub>=3406 20 studies). In large HICs, odds of dementia were higher in urban areas (0.88, 0.84-0.92). To our knowledge, this difference has not been reported before, and has important implications for research and policy in this area. Studies were heterogeneous in the definition of 'rural' or 'urban' and identification of dementia, and there was high risk of bias in the exposure (geographical variation). Studies often did not report when in the life-course the geographical exposure was measured, and whether this changed across their lifetime, and the association between rurality and dementia prevalence and incidence may change with at different ages.

Increased risk of dementia in those living in rural areas may be due to specific aspects of living in these geographical locations, such as access to healthcare services/dementia diagnosis, air pollution or social connections, or may be confounded or mediated by education or socio-economic status.<sup>1</sup> There is a complex interplay between the environment and the individual living in it. For dementia risk, this includes direct and indirect associations between the environment, and cognition, and noting that both of these can change across the life course.<sup>49</sup>

Dementia prevalence can vary country to country, influenced by the drive to seek a dementia diagnosis as well as the different the methodologies used for dementia ascertainment. A recent systematic review recommends using multiple sources, including access to medical records, to ascertain dementia to improve consistency in methodologies between studies.<sup>50,51</sup>

23

For example, in another systematic review<sup>52</sup> which found a mixed picture between green space exposure and dementia risk, green space exposure was associated with higher incidence of dementia measured by anti-dementia medication and reversed if the other methods for detecting dementia were used.<sup>53</sup> It is therefore essential that the method(s) of dementia ascertainment is clearly reported and the potential limitations and biases of this acknowledged.

When comparing multiple studies it is important that the dementia ascertainment rate for the study population is calculated and compared with an expected rate.<sup>51</sup> Dementia diagnosis, particularly in L-MICs, can still be challenging, and there is ongoing debate about the best criteria for dementia diagnosis:<sup>54</sup> criteria (such as the 10/66 criteria) may have greater accuracy than DSM-IV in low levels of literacy.<sup>54</sup> Furthermore, in rural areas in general, or L-MICs, there may be limited access to diagnostic assessment, <sup>55</sup> especially if services are centralised.<sup>56</sup> Our systematic review did not show an increased risk of dementia in urban settings to suggest this bias, but this needs considered when data are included from more countries.

There may be increased cognitive reserve in people living in urban areas <sup>49,57,58</sup>, particularly during childhood, which influences dementia variation between rural and urban living. The impact from social deprivation may be higher than that from rural or urban location (e.g. the odds of a cognitive disorder were twice as high in those living in poor-quality rural environments, than those in rural areas).<sup>16</sup>

Education influences dementia risk, <sup>59</sup> but few studies investigated education in relation to the geographical area across time. Socio-economic status (SES) also influences dementia prevalence rates<sup>50,60</sup>. Our systematic review was unable to assess for SES in influencing the difference in rural versus urban dementia risk. Whilst education, SES and other variables for

overall dementia risk were often reported, these variables were rarely considered in relation to rural or urban areas. Further work is required to understand the complex interplay between geographical location, education, socioeconomic circumstances, prior cognitive ability and dementia risk: whether there are critical time points in a person's life where individual or combined factors influence dementia risk. Future studies should include more detailed information on participants' education, SES measures in relation to geographical area, as well as the different definitions of geographical categorisation.

## Comparison with other literature

This large and comprehensive systematic review builds on a previous systematic review in 2010 which showed, in 25 rural versus urban studies, that there was some evidence of greater all-cause dementia odds with rural living [prevalence OR = 1.11, 90% CI 0.79-1.57; incidence OR = 1.20, 90% CI 0.84-1.71], and that the association between rurality and dementia risk was stronger in studies only including people with Alzheimer Disease (OR=1.50, 90% CI 1.33-1.69), but weaker in studies only including people with vascular dementia (OR=1.09, 90% CI 0.65-1.83). The review also suggested that exposure to rural living early in life further increases the risk of subsequent dementia.

In comparison, our subsequent review which included 38 new papers was unable to investigate the dementia sub-type risk, or the life course effect of geographical location, as too few studies which reported adequate data on these predictors to undertake meaningful metaanalysis. Most studies used dementia as a 'blanket' term encompassing different sub-types of dementia, and did not provide adequate data for the life course effect of geographical location on dementia risk. The previous review investigated dementia prevalence and incidence at a number of geographical scales (country-to-country comparisons, rural to urban comparisons,

and regional comparisons), but there has been a rapid increase studies published on this topic, therefore our current review focussed on rural to urban comparisons.

### Limitations and strengths: of the systematic review

The literature search for this comprehensive systematic review was developed with an information specialist and used a validated comprehensive search strategy from our previous review, with a pre-registered protocol, and was reported according to PRISMA guidance. At least two reviewers independently reviewed titles and abstracts, and extracted and analysed the studies with high levels of agreement using proforma which was produced and trialled for data extraction, and risk of bias (adapting the ROBANS tool). Authors were contacted for additional information, to increase data available for meta-analysis. The language was not limited to English to minimise bias in selecting papers for review.

The limitations of this study include that despite a comprehensive search strategy including forward citation of included studies; it is possible that relevant papers may have been missed, as we did not search for grey literature. There is also significant variation within and between countries of the level of geography which is recorded.

The Risk of Bias assessment tool was based on ROBANS and whilst it did assess the quality of the method of dementia ascertainment, for the studies using big data and ICD-10 codes it was not possible to assess the quality of that dementia diagnosis.

## Limitations and strengths: of the included studies

We identified 38 new relevant papers in the last 5 years, and were able to extract, or calculate, odds ratios for 27 papers (33 studies) for the meta-analysis, including studies from a range of

It is made available under a CC-BY 4.0 International license .

L-MICs and HICs. The sample sizes were overall large (range 90 to 73,000,000, median 2170 for all papers, and 300 to 21,624,228, median 2162, for those in the meta-analysis) and the method of defining dementia was reported in the studies.

The individual studies were heterogeneous in their methodologies. In particular, there was significant heterogeneity in of the definition of 'rural' and 'urban', and how this translates at an international level for comparison. Too few studies reported the impact of geographical location at different time points in a person's life to include in meta-analysis. This means there could be confounding or bias influencing the risk of dementia.

There is a need for standard definitions of urbanisation and rurality, and a clearer definition of "place."<sup>49</sup> Despite proposed quantifiable measures of urbanisation through population density and sprawl and further evaluation of physical characteristics of proximal environments,<sup>49</sup> none of our included papers considered these factors.

It is also important to acknowledge that diagnostic coverage, those diagnosed with dementia rather than the true dementia prevalence, can be low<sup>61</sup> and therefore missed cases (particularly within studies using large big datasets of routinely collected data<sup>62</sup>) may introduce bias, potentially under-reporting, in interpreting geographical variation of dementia versus traditional face-to-face epidemiological studies.

#### Implications

There has been a rapid increase in interest in the geographical predictors of dementia, therefore this study, limited only to urban versus rural geographical variation, confirmed previous findings of increased odds of subsequent dementia for people living in rural areas. The heterogeneity of the studies means understanding rural versus urban risk of dementia is complicated. To ensure a future robust study to investigate the geographical variation of rural

versus urban living and dementia risk (given our finding of OR of 1.16 to 1.26), the sample size would need to be approximately 1000-4 000 people.

Further consensus on the term 'geographical location' with clear definitions of 'rural' and 'urban' that is translatable across, and within, different countries is important to understand the complex interplay between the environment and the individual in relation to dementia risk. Future large-scale studies must also include detailed information about the area (including a measure of deprivation, greenspace, access to services etc.) and a person's geographical location across time.

We know the incidence and prevalence of dementia is rising in L-MICs, but larger studies in this systematic review are in HICs, where they are more likely to have healthcare systems with routinely collected data allowing larger scale studies. Whereas, L-MICs are more likely to use traditional (smaller scale) face-to-face epidemiological studies. As these research methods are more costly than analysing nationally collected data, it results in fewer studies in L-MICs, with lower numbers of participants, but likely to provide more detailed analysis. Finally, the definition of LIC, HMIC and HIC has the potential to change with time, which is important to consider given the longitudinal aspect for dementia risk.

### Conclusions

Overall, rural living was associated with small-increased odds of dementia, compared to urban living (OR, 1.20, 95% CI 1.03-1.40, P value = 0.0182). Stratifying by HICs and L-MICs suggested increased odds of dementia in L-MICs rural areas, (1.33, 1.06 -1.68, P value = 0.0141). However, methodological differences complicate combining the extant literature. Therefore, better-designed studies with careful consideration of mediating and confounding variables are required to answer this important question definitively.

## Acknowledgements

KEW was supported by clinical research fellowship from Alzheimer Scotland and the University of Edinburgh Centre for Cognitive Ageing and Cognitive Epidemiology, part of the cross-council Lifelong Health and Wellbeing initiative (MR/L501530/1). Funding from the Biotechnology and Biological Sciences Research Council (BBSRC) and Medical Research Council (MRC) is gratefully acknowledged. KEW is also supported by Alzheimer Scotland Dementia Research Centre.

JKB is supported by an NHS Education for Scotland/Chief Scientists Office Postdoctoral Clinical Lectureship (PCL/21/01).

GMT is supported by the Osteopathic Heritage Foundation Ralph S. Licklider D.O.

Lastly, we would like to thank Professor John Starr (co-author) for his huge lifetime contribution to dementia research worldwide. We are deeply saddened by his sudden unexpected death on 9 December 2018 and miss our colleague greatly. His input as co-author was important to this research.

It is made available under a CC-BY 4.0 International license .

#### **REFERENCES**

- Livingston G, Huntley J, Sommerlad A, et al. Dementia prevention, intervention, and care:
   2020 report of the Lancet Commission. *Lancet* 2020;**396**(10248):413-446.
- Russ TC, Batty GD, Hearnshaw GF, Fenton C, Starr JM. Geographical variation in dementia: systematic review with meta-analysis. *International Journal of Epidemiology* 2012;**41**(4):1012-1032.
- 3. Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015;**4**(1):1.
- Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Bmj* 2021;**372**:n71.
- Covidence: Systematic Review Management. <u>https://www.covidence.org</u> Accessed 14th September 2023.
- Khedr E, Fawi G, Abbas MA, et al. Prevalence of mild cognitive impairment and dementia among the elderly population of Qena Governorate, Upper Egypt: a community-based study. *Journal of Alzheimer's Disease* 2015;45(1):117-26.
- Raina SK, Raina S, Chander V, et al. Is dementia differentially distributed? A study on the prevalence of dementia in migrant, urban, rural, and tribal elderly population of Himalayan region in Northern India. *North American Journal of Medical Sciences* 2014;6(4):172-177.
- Yuan J, Zhang Z, Wen H, et al. Incidence of dementia and subtypes: A cohort study in four regions in China. *Alzheimer's & Dementia: The Journal of the Alzheimer's Association* 2016;12(3):262-271.
- Samba H, Guerchet M, Bandzouzi BN, et al. Incidence of dementia among older adults in central africa: First results from the republic of congo in the EPIDEMCA-FU study. *Alzheimer's and Dementia* 2015;1):P221-P222.
- Tengku Aizan H, Krishnaswamy S, Siti Suhailah A, Yadollah Abolfathi M.
   Sociodemographic risk factors and correlates of dementia in older Malaysians. *Dementia and Geriatric Cognitive Disorders* 2010;**30**(6):533-539.

- Tola-Arribas MA, Yugueros MI, Garea MJ, et al. Prevalence of dementia and subtypes in Valladolid, northwestern Spain: The DEMINVALL Study. *PLoS ONE* 2013;8(10).
- Poddar K, Kant S, Singh A, Singh TB. An epidemiological study of dementia among the habitants of Eastern Uttar Pradesh, India. *Annals of the Indian Academy of Neurology* 2011;14(3):164-168.
- Pilleron S, Clement JP, Ndamba-Bandzouzi B, et al. Is dependent personality disorder associated with mild cognitive impairment and dementia in Central Africa? A result from the EPIDEMCA programme. *International Psychogeriatrics* 2015;27(2):279-288.
- 14. Kim KW, Park JH, Kim MH, et al. A nationwide survey on the prevalence of dementia and mild cognitive impairment in South Korea. *Journal of Alzheimer's Disease* 2011;23(2):281-291.
- Koller D, Eisele M, Kaduszkiewicz H, et al. Ambulatory health services utilization in patients with dementia - is there an urban-rural difference? *International Journal of Health Geographics* 2010;9(59).
- Wu YT, Prina AM, Jones A, et al. The Built Environment and Cognitive Disorders: Results From the Cognitive Function and Ageing Study II. *American Journal of Preventive Medicine* 2017;**53**(1):25-32.
- Ganesh C. The diffusion of prescription drugs for Alzheimer's disease among medicare beneficiaries [Ph.D.]. The Pennsylvania State University, 2010.
- Hendrie HC, Smith-Gamble V, Lane KA, et al. The Association of Early Life Factors and Declining Incidence Rates of Dementia in an Elderly Population of African Americans. *Journals of Gerontology Series B: Psychological Sciences & Social Sciences* 2018;73:S82-S89.
- Guerchet M, Bandzouzi-Ndamba B, Mbelesso P, et al. Prevalence of dementia in two countries of central africa: Comparison of rural and urban areas in the epidemca study. *Neuroepidemiology* 2013;41 (3-4):253-254.
- 20. Tornau F, Bruchmann G, Juckel G. Prevalence of psychiatric disorders in the elderly in rural and urban areas in Germany. *International Psychogeriatrics* 2015;**27**:S167.

- Goncalves-Pereira M, Cardoso A, Verdelho A, et al. The prevalence of dementia in a Portuguese community sample: a 10/66 Dementia Research Group study. *BMC Geriatrics* 2017;**17**(1):261.
- Prince M, Acosta D, Ferri CP, et al. Dementia incidence and mortality in middle-income countries, and associations with indicators of cognitive reserve: A 10/66 Dementia Research Group population-based cohort study. *The Lancet* 2012;**380**(9836):50-58.
- Rodriguez-Agudelo Y, Solis-Vivanco R, Acosta-Castillo I, et al. Neuropsychiatric symptoms in older adulte with and without dementia in urban and rural regions. Results of the 10/66 Dementia Research Group in Mexico. [Spanish]
- Sintomas neuropsiquiatricos en adultos mayores con y sin demencia de regiones urbana y rural. Resultados del Grupo de Investigacion en Demencia 10/66 en Mexico. *Revista de Investigacion Clinica* 2011;**63**(4):382-390.
- Prince M. Dementia in developing countries. A consensus statement from the 10/66 Dementia Research Group. *Int J Geriatr Psychiatry* 2000;15(1):14-20.
- Prina AM, Mayston R, Wu YT, Prince M. A review of the 10/66 dementia research group. Soc Psychiatry Psychiatr Epidemiol 2019;54(1):1-10.
- Guerchet M, Ndamba-Bandzouzi B, Mbelesso P, et al. Comparison of rural and urban dementia prevalences in two countries of central africa: The epidemca study. *Alzheimer's and Dementia* 2013;1):P688.
- Wellcome. Low-and middle-income countries. <u>https://wellcome.org/grant-</u> <u>funding/guidance/low-and-middle-income-countries</u> Accessed 8th September, 2023.
- Abner EL, Jicha GA, Christian WJ, Schreurs BG. Rural-urban differences in Alzheimer's disease and related disorders diagnostic prevalence in Kentucky and West Virginia. *Journal* of Rural Health 2016;32(3):314-320.
- 29. Astell-Burt T, Feng X. Is the risk of developing Alzheimer's disease really higher in rural areas? A multilevel longitudinal study of 261,669 Australians aged 45 years and older tracked over 11 years. *Health and Place* 2018;**54**:132-137.

- Chen H, Kwong JC, Copes R, et al. Exposure to ambient air pollution and the incidence of dementia: A population-based cohort study. *Environment International* 2017;108:271-277.
- 31. Chen R, Ma Y, Wilson K, et al. A multicentre community-based study of dementia cases and subcases in older people in China--the GMS-AGECAT prevalence and socio-economic correlates. *International Journal of Geriatric Psychiatry* 2012;27(7):692-702.
- 32. Deng J, Cao C, Jiang Y, et al. Prevalence and effect factors of dementia among the community elderly in Chongqing, China. *Psychogeriatrics* 2018;**18**(5):412-420.
- Goodman RA, Lochner KA, Thambisetty M, et al. Prevalence of dementia subtypes in United States Medicare fee-for-service beneficiaries, 2011-2013. *Alzheimer's & Dementia* 2017;13(1):28-37.
- Nunes B, Silva RD, Cruz VT, et al. Prevalence and pattern of cognitive impairment in rural and urban populations from Northern Portugal. *BMC Neurology Vol 10 2010, ArtID 42* 2010;10.
- 35. Sharifi F, Fakhrzadeh H, Varmaghani M, et al. Prevalence of Dementia and Associated Factors among Older Adults in Iran: National Elderly Health Survey (NEHS). Archives of Iranian Medicine 2016;19(12):838-844.
- Tripathi M, Vibha D, Gupta P, et al. Risk factors of dementia in North India: A case-control study. *Aging & Mental Health* 2012;16(2):228-235.
- Weden MM, Shih RA, Kabeto MU, Langa KM. Secular trends in dementia and cognitive impairment of U.S. rural and urban older adults. *American Journal of Preventive Medicine* 2018;54(2):164-172.
- 38. Jia J, Wang F, Wei C, et al. The prevalence of dementia in urban and rural areas of China. *Alzheimer's & Dementia: The Journal of the Alzheimer's Association* 2014;**10**(1):1-9.
- Kim K, Park J, Kim M, et al. A nationwide survey on the prevalence of dementia and mild cognitive impairment in South Korea. *JAD, Journal of Alzheimer's Disease* 2011;23(2):281-291.
- Nadel JL, Ulate D. Incidence and risk factors for cognitive impairment in rural elderly populations in Costa Rica^ien

It is made available under a CC-BY 4.0 International license .

Incidencia y factores de riesgo para la discapacidad cognitiva en poblaciones rurales de tercera edad en Costa Rica^ies. *Rev. biol. trop* 2014;**62**(3):869-876.

- Contador I, Bermejo-Pareja F, Puertas-Martin V, Benito-Leon J. Childhood and Adulthood Rural Residence Increases the Risk of Dementia: NEDICES Study. *Current Alzheimer Research* 2015;12(4):350-7.
- 42. Chammartin F, Probst-Hensch N, Utzinger J, Vounatsou P. Mortality atlas of the main causes of death in Switzerland, 2008-2012. *Swiss Medical Weekly* 2016;**146**:w14280.
- Cornutiu G. The incidence and prevalence of Alzheimer's disease. *Neurodegenerative Diseases* 2010;8(1-2):9-14.
- Silva EFd, Paniz VMV, Laste G, Torres ILdS. Prevalência de morbidades e sintomas em idosos: um estudo comparativo entre zonas rural e urbana. *Ciência & amp; Saúde Coletiva* 2013;18(4):1029-1040.
- 45. Huang C, Chung C, Leu H, et al. Diabetes mellitus and the risk of Alzheimer's disease: a nationwide population-based study. *PLoS ONE* 2014;**9**(1).
- 46. Taylor CA, Greenlund SF, McGuire LC, Lu H, Croft JB. Deaths from Alzheimer's Disease -United States, 1999-2014. *Mmwr* 2017; Morbidity and mortality weekly report.
  66(20):521-526.
- 47. Yin P, Feng X, Astell-Burt T, et al. Temporal trends and geographic variations in dementia mortality in China between 2006 and 2012: Multilevel evidence from a nationally representative sample. *Alzheimer Disease and Associated Disorders* 2016;**30**(4):348-353.
- Zhao Q, Zhou B, Ding D, Guo Q, Hong Z. Prevalence, mortality, and predictive factors on survival of dementia in Shanghai, China. *Alzheimer Disease and Associated Disorders* 2010;**24**(2):151-158.
- 49. Cassarino M, Setti A. Environment as 'Brain Training': A review of geographical and physical environmental influences on cognitive ageing. *Ageing Research Reviews* 2015;**Part B.**23:167-182.
- 50. Mullin DS, Stirland LE, Buchanan E, et al. Identifying dementia using medical data linkage in a longitudinal cohort study: Lothian Birth Cohort 1936. *BMC Psychiatry* 2023;**23**(1):303.

- Sibbett RA, Russ TC, Deary IJ, Starr JM. Dementia ascertainment using existing data in UK longitudinal and cohort studies: a systematic review of methodology. *BMC Psychiatry* 2017;17(1):239.
- Chen X, Lee C, Huang H. Neighborhood built environment associated with cognition and dementia risk among older adults: A systematic literature review. *Soc Sci Med* 2022;292:114560.
- 53. Astell-Burt T, Navakatikyan MA, Feng X. Urban green space, tree canopy and 11-year risk of dementia in a cohort of 109,688 Australians. *Environ Int* 2020;**145**:106102.
- 54. Ferri CP, Oliveira D. Harmonization of epidemiological studies on dementia in Latin America Why does it matter? *Dement Neuropsychol* 2019;13(4):363-366.
- Zakarias JK, Jensen-Dahm C, Nørgaard A, et al. Geographical Variation in the Diagnostic Rate and Quality of Dementia Diagnoses. *J Alzheimers Dis* 2019;69(2):513-520.
- 56. Jorgensen TS, Torp-Pedersen C, Gislason GH, Andersson C, Holm E. Time trend in Alzheimer diagnoses and the association between distance to an Alzheimer clinic and Alzheimer diagnosis. *European journal of public health* 2015;25(3):522-527.
- 57. Brayne C, Wu YT. Population-Based Studies in Dementia and Ageing Research: A Local and National Experience in Cambridgeshire and the UK. *Am J Alzheimers Dis Other Demen* 2022;**37**:15333175221104347.
- Cassarino M, O'Sullivan V, Kenny RA, Setti A. Environment and cognitive aging: A crosssectional study of place of residence and cognitive performance in the Irish longitudinal study on aging. *Neuropsychology* 2016;**30**(5):543-557.
- 59. Cadar D, Lassale C, Davies H, et al. Individual and area-based socioeconomic factors associated with dementia incidence in England: Evidence from a 12-year follow-up in the English longitudinal study of ageing. JAMA Psychiatry 2018;75(7):723-732.
- 60. Matthews FE, Arthur A, Barnes LE, et al. A two-decade comparison of prevalence of dementia in individuals aged 65 years and older from three geographical areas of England: results of the Cognitive Function and Ageing Study I and II. *Lancet* 2013;**382**(9902):1405-12.

It is made available under a CC-BY 4.0 International license .

61. Gauthier S, Rosa-Neto P, Morais JA, C W. World Alzheimer Report 2021: Journey through the diagnosis of dementia

London, England: Alzheimer's Disease International, 2021.

62. Wilkinson T, Schnier C, Bush K, et al. Identifying dementia outcomes in UK Biobank: a validation study of primary care, hospital admissions and mortality data. *Eur J Epidemiol* 2019;**34**(6):557-565.