

Detection of Noroviruses GII in wastewater samples of Bhopal, India, using Droplet Digital PCR

Nema R K^{1*}, Nagar J¹, Singh A K¹, Tandekar A¹, Singh S², Rahman A¹, Sharma O P¹, Tiwari R R³, Sarma D K¹, Mishra P K¹.

¹Division of Environmental Biotechnology, Genetics, and Molecular Biology, ICMR- National Institute for Research in Environmental Health, Bhopal - 462030 (MP), India

²Division of Environmental Monitoring and Exposure Assessment (Water & Soil), ICMR – National Institute for Research in Environmental Health, Bhopal – 462 030 (MP), India

³ICMR – National Institute for Research in Environmental Health, Bhopal – 462 030 (MP), India

Corresponding author's Email: ramkumar.nema@icmr.gov.in

Abstract: Noroviruses are a significant cause of global gastroenteritis outbreaks, underscoring the importance of effective surveillance. Wastewater-based epidemiology helps identify viral pathogens in communities. Recent advancements in wastewater-based molecular surveillance have shown that viruses can be detected in feces and urine early, making sewage monitoring an essential tool for tracking viral presence. We aimed to create and validate a new method for detecting and monitoring Noroviruses GII in Bhopal's wastewater using Automated Droplet Digital PCR (ddPCR) technology. In this study, a ddPCR assay targeting the ORF1-2 region of Norovirus GII was developed, allowing viral nucleic acid quantification without a standard curve. A total of 27 samples from five Sewage Treatment Plants located in Bhopal city were collected during the summer season (April and May 2023) at fortnightly intervals and analyzed for the presence of Norovirus using the novel ddPCR assay. Among the samples tested, 33% tested positive for Norovirus, with the highest detection rate observed as 72.72%, followed by 25%. The concentrations of Noroviruses GII in positive samples ranged from 0.06 to 6.60 copies/μl. These findings indicate a potentially higher patient population within Bansal Hospital's catchment area than the other STPs in the Bhopal region. The study underscores Norovirus's varying prevalence and distribution in wastewater across different STPs in Bhopal. Moreover, it demonstrates the utility of wastewater surveillance and digital PCR in accurately and specifically detecting Norovirus in wastewater. The practical application of this wastewater surveillance strategy could serve as an early warning system for communities, enabling timely preparedness for impending viral outbreaks, implementation of effective administrative containment measures, and intensified vaccination campaigns.

Keywords: Noroviruses, wastewater-based epidemiology, molecular surveillance, Droplet Digital PCR (ddPCR)

Introduction: Norovirus is responsible for around 20% of all diarrhea cases globally, affecting children and adults equally. It is estimated to cause over 200,000 deaths annually in developing countries (1). These highly contagious and widespread RNA viruses are responsible for a notable proportion of global cases of acute gastroenteritis. viral gastroenteritis cases worldwide. The major cause of acute viral gastroenteritis worldwide is rotavirus, with norovirus ranking second in prevalence (2). Noroviruses can cause severe gastroenteritis, with symptoms including nausea, vomiting, diarrhea, and stomach cramps. Norovirus infection can result in dehydration, especially in young children, older adults, and individuals with compromised immune systems. (3, 4). Noroviruses have the ability to infect not only humans but also a diverse range of other mammalian hosts. For the first time, two previously unidentified norovirus genomes were discovered in raccoon dogs (5). The viruses can survive in the environment for extended periods and are resistant to many common disinfectants, making them difficult to control (3). Infection control measures, such as improved sanitation and hand hygiene, are essential for preventing the spread of noroviruses. One of the ways in which noroviruses can spread is through the fecal-oral route, leading to the contamination of water sources and wastewater (6).

Noroviruses belong to the family Caliciviridae; considering the complete VP1 amino acid sequences, Noroviruses can be genetically classified into ten genogroups labelled as GI to GX. Genogroups I, II, IV, VIII and IX are known to infect humans, with genogroup II (GII) being the most prevalent. Currently, within these ten genogroups, there are 49 confirmed capsid genotypes based on the amino acids of the complete VP1 sequence. These include nine genotypes for GI, 27 for GII, 3 for GIII, 2 for GIV, 2 for GV, 2 for GVI, and one genotype each for GVII, GVIII, GLX, and GX. Moreover, there are 60 confirmed P-types based on partial nucleotide sequences of the RdRp regions, with 14 for GI, 37 for GII, 2 for GIII, 1 for GIV, 2 for GV, 2 for GVI, 1 for GVII, and 1 for GX (7, 8). The GII strain of Norovirus encompasses both porcine and human strains, while GIII consists solely of bovine strains, and GV exclusively contains murine strains (7). Diagnosing a norovirus infection can be challenging due to the presenceexistence of non-specific symptoms and the challenges in diagnosis. However, laboratory testing, particularly molecular assays, has proven to be the most reliable method for diagnosing norovirus infection (9). In addition to laboratory testing, clinical evaluation and epidemiological investigation play crucial roles in diagnosing norovirus infection, especially during outbreaks. These tools help in identifying patterns and sources of infection. Timely diagnosis and the enforcement of control measures are crucial to prevent the further spread of the viruses(10).

Traditional methods for detecting microbial pathogens in water and wastewater often rely on cultivating them in artificial media or cell cultures. However, these methods can be

time consuming and less sensitive. Emerging technologies like Polymerase Chain Reaction (PCR) offer a promising alternative. PCR can rapidly and precisely detect a wide range of pathogens from a single water sample, addressing the limitations of traditional methods. In recent years, molecular techniques such as PCR and RT-qPCR have been employed to detect noroviruses in wastewater samples. However, it is important to acknowledge that these methods may have limitations in terms of sensitivity, specificity, and the ability to detect low levels of the virus (11). Digital droplet PCR (ddPCR) represents a cutting-edge approach to nucleic acid detection, boasting several advantages over traditional PCR methods. This innovative technique offers heightened sensitivity, reduced variability, and the unique capability to quantify low levels of target nucleic acids. Beyond these advantages, one of the key features of ddPCR is its versatility in detecting multiple targets simultaneously within a single reaction, making it particularly invaluable for the surveillance of various pathogens, including noroviruses, in wastewater (12). The sensitivity and specificity inherent in ddPCR enable the detection of viruses even at markedly lower concentrations, providing a more profound understanding of viral epidemiology. This breakthrough has significantly impacted clinical research and the surveillance of enteric viruses in wastewater. Molecular surveillance plays a crucial role in understanding the evolution and spread of Norovirus, a highly contagious gastrointestinal pathogen (13). By analysing the genetic sequences of various strains, researchers can acquire insights into the epidemiology, evolution, and ecology of this significant human pathogen. Ultimately, this research will contribute to developing more effective strategies for preventing norovirus infections in the population.

Methods:

Waste water sample collection: Sewage samples, 2x500 ml, were collected fortnightly from each of the 5 Waste Water Treatment Plants (WWTP) in the Municipal Corporation Area of Bhopal, Madhya Pradesh. These plants were situated in Bansal Hospital, Charimli, Shirin River, Professor Colony, and Jamuniya Cheer area and had a well-defined catchment area. Samples collected from these areas were brought to the lab and processed for RNA isolation immediately or stored at 4°C and were processed within 24 hours to avoid further degradation of viral nucleic acid.

Virus concentration using Polyethylene Glycol Precipitation Method: Virus concentration was done using modified polyethylene glycol method (14-16). Initially, the sewage samples (50ml) were centrifuged at 4500 x g for 30 minutes, followed by filtering the supernatant using 0.22µm filters. The pH of the supernatant was then adjusted to 7.0-7.5. Subsequently, 10% PEG 8000 (polyethyleneglycol) and NaCl were introduced to the supernatant to achieve a final concentration of 0.3 Molar. The blend was subsequently left to

incubated overnight at 4°C followed by centrifugation at 4500x g for 30 minutes. The obtained supernatant was discarded, and the resulting pellets were resuspended in 500µL of phosphate buffer saline solution (PBS) to form a virus concentrate. The virus concentrate can either be processed for nucleic acid extraction or stored at -80°C for long-term storage until extraction.

RNA extraction: Total RNA was extracted from an aliquot of each viral concentrate by AllPrep Power Viral DNA/RNA Kit (Qiagen, Germany), following the manufacturer's instructions. This technique selectively bound viral RNA to a silica-gel membrane, while contaminants were efficiently eliminated during washing steps, removing PCR inhibitors such as divalent cations and proteins (17-19). Extracted RNA was eluted in 100 µL of the elution buffer and stored in 10 µL aliquots at -80°C until use.

cDNA Synthesis: The cDNA was synthesized using the cDNA Synthesis Kit GoScript™ Reverse Transcription System (Cat. No. A5000) (Promega), following the modified manufacturer's protocol (20, 21). Briefly, 4 µl of RNA and 1 µl of 0.2µg/µl random hexamer primer underwent denaturation at 70°C for 5 min. A 15 µl reverse transcription mix, including 5x reaction buffer, Ribonuclease Inhibitor, dNTP mix, and RT enzyme, was added to the RNA/primer mix. After proper mixing and brief centrifugation, the tube underwent incubation at 25°C for 5 min, followed by 42°C for 60 min for cDNA synthesis. Termination of RT reaction was achieved at 70°C for 10 min. The tubes were subsequently chilled on ice, and the cDNA was stored at -80°C for future use.

ddPCR Assay: Primers and TaqMan probe targeting the highly conserved ORF1, 2 regions of the Norovirus GII genotype were synthesized. The specificity of these selections was validated using the BLAST program from NCBI. Notably, the TaqMan probe was tagged with 5'-FAM for the fluorescent reporter dye and 3'-BHQ-1 as the quencher, ensuring accurate signal detection in the assay. The sequences of the primers and probes are shown in Table 1. The ddPCR reaction was prepared using 2µl (50 ng) of cDNA and a reaction mixture that included ddPCR Supermix for probes, forward and reverse primers, and the TaqMan probe. Droplet generation was then carried out using an Automatic Droplet Generator, QX200™ AutoDG™, in a 96-well polypropylene plate. The subsequent PCR cycling involved initial denaturation, followed by 45 cycles of denaturation (95°C for 15 seconds), annealing, and extension (55°C for 30 seconds), culminating in a final denaturation step (55°C for 7 minutes). After PCR, the plate containing the droplets was transferred to a QX200 droplet reader for analysis. Finally, data analysis was performed using QuantaSoft™ Analysis Pro software, which involved discriminating positive from negative droplets based on fluorescence amplitude, thereby identifying the positive droplets containing the amplification

products. This comprehensive approach ensured precise quantification and high-resolution analysis of the Norovirus genetic material.

Table 1: Primers and probe used in the development of ddPCR assay for Noroviruses.

S. No.	Primers	Sequences of primers	Region	Reference
1	Norovirus FP	5'CARGARBCNATGTTYAGRTGGATGAG 3'	ORF1,2	Kageyama T et.al., 2003 (28)
2	Norovirus RP	5'TCGACGCCATCTTCATTCACA3'	ORF1,2	
3	Norovirus Probe	5'FAM-'TGGGAGGGCGATCGCAATCT - BHQ 3'	ORF1,2	

Result: A ddPCR assay targeting the ORF1-2 region of Norovirus GII was developed, allowing viral nucleic acid quantification without a standard curve. A total of 27 sewage samples were collected from five Sewage Treatment Plants (STPs) located in Bhopal for the purpose of investigating the presence of Norovirus using the highly sensitive norovirus droplet digital polymerase chain reaction (ddPCR) assay. Among these samples, nine were confirmed to be positive for Norovirus using the ddPCR assay. In particular, within the Bansal Hospital STP subset, consisting of 11 samples, a significant majority of 8 samples (72.72%) exhibited a positive result for Norovirus when subjected to the ddPCR assay. Similarly, within the Char Imli STP subset, encompassing four samples, one sample (25%) was found to be positive for Norovirus through the ddPCR assay (Table 2). The concentrations of Noroviruses GII in positive samples ranged from 0.06 to 6.60 copies/ μ l (Table 3). However, the samples collected from Professor Colony, Jamunia Cheer, and Shirin River, representing distinct geographic locations, yielded negative results for Norovirus in the ddPCR assay. This suggests the absence of detectable Norovirus within the samples obtained from these areas.

Table 2: Distribution of Norovirus in wastewater treatment plants of Bhopal City.

S. No.	Name of WWTP	No. of Sample Collected	No. of norovirus Positive samples	% Positivity
1	Bansal Hospital	11	8	72.72%
2	Char Imli	4	1	25%
3	Professor Colony	4	0	0%
4	Shirin River	4	0	0%

5	Jamunia Cheer	4	0	0%
	Total	27	9	33.33%

Table 3: Concentration of Noroviruses GII in positive samples.

S. No.	Date of sample collection	Name of STPs	No. of DDPCR Positive Droplets	Concentration (Copies/ μ l)
1	19.04.2023	Bansal Hospital	96	6.60
2	22.04.2023	Char Imli	17	0.94
3	22.04.2023	Bansal Hospital	58	3.53
4	23.04.2023	Bansal Hospital	27	1.64
5	24.04.2023	Bansal Hospital	35	2.23
6	25.04.2023	Bansal Hospital	36	2.44
7	26.04.2023	Bansal Hospital	01	0.06
8	27.04.2023	Bansal Hospital	12	0.77
9	28.04.2023	Bansal Hospital	03	0.20

Discussion: The augmented Norovirus positivity observed in this study is hypothesized to be associated with the presence of a hospitalized patient population. This correlation could plausibly lead to an increased likelihood of Norovirus shedding in feces and urine among the affected individuals(22). As a result, the sewage system may experience an elevation in the viral load, potentially contributing to the heightened Norovirus prevalence in the studied area. Further investigation is warranted to validate and understand the implications of these findings on public health and environmental management.

The results of this study offer valuable insights into the prevalence of Norovirus in Sewage Treatment Plants (STPs) located in Bhopal, India. The detection and analysis of Norovirus using the droplet digital polymerase chain reaction (ddPCR) assay yielded significant results, highlighting the importance of understanding the presence and distribution of this pathogen in wastewater (23).

The overall norovirus detection rate of approximately 35.7% among the collected samples indicates a significant presence of Norovirus in the sewage samples analyzed. These findings are consistent with prior research that has recognized wastewater as a potential

reservoir for the transmission of norovirus (24). The high detection rate emphasizes the importance of wastewater surveillance for monitoring and managing potential health risks associated with norovirus contamination.

The Bansal Hospital STP demonstrated the highest norovirus detection rate, with 72% of the samples testing positive for the virus. This finding suggests a higher prevalence and potential concentration of Norovirus in the sewage originating from this specific treatment plant. The increased detection rate in this STP might be attributed to factors such as a higher number of infected patients admitted to Bansal Hospital, contributing to the viral load in the wastewater treatment plants. Further investigation is necessary to identify the specific factors influencing the higher detection rate in this particular STP. Conversely, the Char Imli STP exhibited a lower norovirus detection rate of approximately 25%. Although the overall detection rate is lower than the Bansal Hospital STP, it still signifies Norovirus's presence in the analyzed wastewater samples. The differences observed in detection rates between the two STPs may be attributed to variations in the sources of wastewater, treatment processes, or geographical factors influencing the viral load in the sewage. Understanding these variations is crucial for implementing effective control measures and optimizing treatment processes to minimize norovirus transmission. Interestingly, no norovirus was detected in the samples collected from Professor Colony, Jamunia Cheer, and Sirin River. This result suggests either the absence of norovirus contamination in these specific locations or the presence of low viral loads that fell below the detection threshold of the ddPCR assay. Further investigations using alternative detection methods, such as quantitative PCR or sequencing, could provide a more comprehensive understanding of norovirus presence in these areas (25).

The findings of this study underscore the importance of robust surveillance programs and the implementation of adequate wastewater treatment processes to mitigate norovirus transmission. Wastewater-based epidemiology, including the monitoring of sewage for the presence of pathogens like Norovirus, can serve as an early warning system for outbreaks and provide valuable information for public health interventions. Improvements in wastewater treatment processes, particularly in STPs with higher detection rates, should be considered to ensure the effective removal of Norovirus and minimize its potential spread through environmental contamination.

Furthermore, the identification of norovirus-positive samples in STPs highlights the importance of public awareness and hygiene practices to reduce the risk of infection. Proper hand hygiene, food handling, and the disinfection of contaminated surfaces are crucial in preventing norovirus outbreaks in communities(26). To strengthen the validity and applicability of research findings, future efforts should prioritize recruiting a larger and more diverse sample.. Additionally, sequencing and phylogenetic analysis of the norovirus strains

detected in the wastewater samples would provide insights into the genetic diversity and potential sources of the virus.

References:

1. Lopman BA, Steele D, Kirkwood CD, Parashar UD. The Vast and Varied Global Burden of Norovirus: Prospects for Prevention and Control. *PLoS Med.* 2016;13(4):e1001999.
2. Mathew S, Alansari K, M KS, Zaraket H, Al Thani AA, Yassine HM. Epidemiological, Molecular, and Clinical Features of Norovirus Infections among Pediatric Patients in Qatar. *Viruses.* 2019;11(5).
3. Barclay L, Park GW, Vega E, Hall A, Parashar U, Vinje J, et al. Infection control for norovirus. *Clin Microbiol Infect.* 2014;20(8):731-40.
4. Wang J, Gao Z, Yang ZR, Liu K, Zhang H. Global prevalence of asymptomatic norovirus infection in outbreaks: a systematic review and meta-analysis. *BMC Infect Dis.* 2023;23(1):595.
5. Li J, Liu W, Tian F, Tu Q, Xia X, Liu C, et al. First report of norovirus sequences isolated from raccoon dogs in mainland China. *Virus Res.* 2021;305:198546.
6. Prevention CfDca. How You Treat Norovirus: CDC; 2023 [Available from: (<https://www.cdc.gov/norovirus/about/treatment.html>)].
7. Chen L, Xu D, Wu X, Liu G, Ji L. An increasing prevalence of non-GII.4 norovirus genotypes in acute gastroenteritis outbreaks in Huzhou, China, 2014-2018. *Arch Virol.* 2020;165(5):1121-8.
8. Ao Y, Wang J, Ling H, He Y, Dong X, Wang X, et al. Norovirus GII.P16/GII.2-Associated Gastroenteritis, China, 2016. *Emerg Infect Dis.* 2017;23(7):1172-5.
9. Cardemil CV, Sherchand JB, Shrestha L, Sharma A, Gary HE, Estivariz CF, et al. Pathogen-Specific Burden of Outpatient Diarrhea in Infants in Nepal: A Multisite Prospective Case-Control Study. *J Pediatric Infect Dis Soc.* 2017;6(3):e75-e85.
10. Robilotti E, Deresinski S, Pinsky BA. Norovirus. *Clin Microbiol Rev.* 2015;28(1):134-64.
11. Gyawali P, Kc S, Beale DJ, Hewitt J. Current and Emerging Technologies for the Detection of Norovirus from Shellfish. *Foods.* 2019;8(6).
12. Jahne MA, Brinkman NE, Keely SP, Zimmerman BD, Wheaton EA, Garland JL. Droplet digital PCR quantification of norovirus and adenovirus in decentralized wastewater and graywater collections: Implications for onsite reuse. *Water Res.* 2020;169:115213.
13. Ogunsakin RE, Ebenezer O, Ginindza TG. A Bibliometric Analysis of the Literature on Norovirus Disease from 1991-2021. *Int J Environ Res Public Health.* 2022;19(5).
14. Thongprachum A, Fujimoto T, Takanashi S, Saito H, Okitsu S, Shimizu H, et al. Detection of nineteen enteric viruses in raw sewage in Japan. *Infect Genet Evol.* 2018;63:17-23.
15. Adachi Katayama Y, Hayase S, Ando Y, Kuroita T, Okada K, Iwamoto R, et al. COPMAN: A novel high-throughput and highly sensitive method to detect viral nucleic acids including SARS-CoV-2 RNA in wastewater. *Sci Total Environ.* 2023;856(Pt 1):158966.
16. Zheng X, Wang M, Deng Y, Xu X, Lin D, Zhang Y, et al. A rapid, high-throughput, and sensitive PEG-precipitation method for SARS-CoV-2 wastewater surveillance. *Water Res.* 2023;230:119560.
17. Anderson-Coughlin BL, Kniel KE. Recovery and Detection of Enteric Viruses from Non-Traditional Irrigation Water Sources. *Methods Protoc.* 2019;2(3).
18. Bhattarai B, Sahulka SQ, Podder A, Hong S, Li H, Gilcrease E, et al. Prevalence of SARS-CoV-2 genes in water reclamation facilities: From influent to anaerobic digester. *Sci Total Environ.* 2021;796:148905.
19. Kaya D, Niemeier D, Ahmed W, Kjellerup BV. Evaluation of multiple analytical methods for SARS-CoV-2 surveillance in wastewater samples. *Sci Total Environ.* 2022;808:152033.

20. Wang Y, Huang J, Sun Y, Stubbs D, He J, Li W, et al. SARS-CoV-2 suppresses mRNA expression of selenoproteins associated with ferroptosis, endoplasmic reticulum stress and DNA synthesis. *Food Chem Toxicol.* 2021;153:112286.
21. Sharif N, Ahmed SN, Sharif N, Alzahrani KJ, Alsuwat MA, Alzahrani FM, et al. High prevalence of norovirus GII.4 Sydney among children with acute gastroenteritis in Bangladesh, 2018-2021. *J Infect Public Health.* 2023;16(7):1015-22.
22. da Silva AK, Le Saux JC, Parnaudeau S, Pommepuy M, Elimelech M, Le Guyader FS. Evaluation of removal of noroviruses during wastewater treatment, using real-time reverse transcription-PCR: different behaviors of genogroups I and II. *Appl Environ Microbiol.* 2007;73(24):7891-7.
23. Song M, Hwang Y, Park J, Cha E, Jeong H, Kim M, et al. Quantitative differential analysis of norovirus outbreak samples using RT-ddPCR. *Lett Appl Microbiol.* 2022;75(1):29-35.
24. Han TH, Kim SC, Kim ST, Chung CH, Chung JY. Detection of norovirus genogroup IV, klassevirus, and pepper mild mottle virus in sewage samples in South Korea. *Arch Virol.* 2014;159(3):457-63.
25. Vinje J. Advances in laboratory methods for detection and typing of norovirus. *J Clin Microbiol.* 2015;53(2):373-81.
26. Prevention CfDCa. Norovirus Guidelines for Healthcare Settings: CDC; [updated November 5, 2015. Available from: <https://www.cdc.gov/infectioncontrol/guidelines/norovirus/index.html>.