

# 1 **Epigenetic scores indicate differences in the proteome of preterm infants**

2 Katie Mckinnon<sup>1</sup>, Eleanor L.S. Conole<sup>2</sup>, Kadi Vaher<sup>1,4</sup>, Robert F. Hillary<sup>3</sup>, Danni A. Gadd<sup>3</sup>, Justyna

3 Binkowska<sup>1</sup>, Gemma Sullivan<sup>4</sup>, Anna J. Stevenson<sup>3</sup>, Amy Corrigan<sup>1</sup>, Lee Murphy<sup>5</sup>, Heather C.

4 Whalley<sup>4</sup>, Hilary Richardson<sup>6</sup>, Riccardo E. Marioni<sup>3</sup>, Simon R. Cox<sup>2</sup>, James P. Boardman<sup>1,4</sup>

5

6 1. Centre for Reproductive Health, Institute for Regeneration and Repair, University of Edinburgh,  
7 Edinburgh, UK.

8 2. Lothian Birth Cohorts, Department of Psychology, University of Edinburgh, Edinburgh, UK.

9 3. Centre for Genomic and Experimental Medicine, Institute of Genetics and Cancer, University of  
10 Edinburgh, Edinburgh, UK.

11 4. Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh, UK.

12 5. Edinburgh Clinical Research Facility, University of Edinburgh, Edinburgh, UK.

13 6. School of Philosophy, Psychology, and Language Sciences, University of Edinburgh, Edinburgh,  
14 UK.

15

16 Corresponding author: Professor James P Boardman

17 Institute for Regenerative Medicine, 4-5 Little France Drive, Edinburgh, EH16 4UU

18 [James.Boardman@ed.ac.uk](mailto:James.Boardman@ed.ac.uk)

19

20

21

22

23

24

25

26

27

28

29

30

# Abstract

**Background:** Epigenetic scores (EpiScores), reflecting DNA methylation (DNAm)-based surrogates for complex traits, have been developed for multiple circulating proteins. EpiScores for pro-inflammatory proteins, such as C-reactive protein (DNAm CRP), are associated with brain health and cognition in adults and with inflammatory comorbidities of preterm birth in neonates. Social disadvantage can become embedded in child development through inflammation, and deprivation is over-represented in preterm infants. We tested the hypotheses that preterm birth and socioeconomic status (SES) are associated with alterations in a set of EpiScores enriched for inflammation-associated proteins.

**Results:** 104 protein EpiScores were derived from saliva samples of 332 neonates born at gestational age (GA) 22.14 to 42.14 weeks. Saliva sampling was between 36.57 and 47.14 weeks. Forty-three (41%) EpiScores were associated with low GA at birth (standardised estimates [0.14 to 0.88], Bonferroni-adjusted  $p$ -value  $<8.3 \times 10^{-3}$ ). These included EpiScores for chemokines, growth factors, proteins involved in neurogenesis and vascular development, cell membrane proteins and receptors, and other immune proteins. Three EpiScores were associated with SES, or the interaction between birth GA and SES: afamin, intercellular adhesion molecule 5 and hepatocyte growth factor-like protein (standardised estimates [0.06 to 0.13], Bonferroni-adjusted  $p$ -value  $<8.3 \times 10^{-3}$ ). In a preterm sub-group ( $n=217$ , median [range] GA 29.29 weeks [22.14 to 33.0 weeks]), SES-EpiScore associations did not remain statistically significant after adjustment for sepsis, bronchopulmonary dysplasia, necrotising enterocolitis, and histological chorioamnionitis.

**Conclusions:** Low birth GA is substantially associated with a set of EpiScores. The set was enriched for inflammatory proteins, providing new insights into immune dysregulation in preterm infants. SES had fewer associations with EpiScores; these tended to have small effect sizes and were not statistically significant after adjusting for inflammatory comorbidities. This suggests that inflammation is unlikely to be the primary axis through which SES becomes embedded in the development of preterm infants in the neonatal period.

**Keywords:** Epigenetic, Inflammation, Neonatal, Preterm birth, Socioeconomic status.

## 1.0 Background

Preterm birth (delivery at less than 37 weeks of gestation) affects around 10% of births worldwide and is closely associated with increased likelihood of cerebral palsy, neurocognitive impairment, behavioural, social and communication difficulties, and mental and cardiometabolic health diagnoses across the life course [1–5]. These adverse outcomes can be explained, in part, by deleterious effects of early exposure to extrauterine life on brain and cardiac development, and they are often accompanied by changes in blood proteins, including those reflecting the perinatal innate and adaptive immune response [6–8].

Socioeconomic status (SES) is also associated with the adverse neurodevelopmental and health outcomes listed above [9–12], and social deprivation is consistently over-represented among preterm children and their families [13,14]. In a meta-analysis of 43 studies (n=111,156 individuals), low SES associated with increased inflammatory markers of disease risk (C-reactive protein [CRP] and interleukin-6 [IL6]), which suggests that pro-inflammatory pathways may be important mechanisms for translating social inequalities into health disparities [15]. However, only four studies included participants under 10 years of age, leaving uncertainty about SES-inflammation correlations in early life [16–19].

Although protein levels are commonly used as biomarkers of exposure and disease risk, they are limited because they are often phasic in the systemic circulation, rely on venepuncture, and may not capture baseline status or chronicity. For example, inflammation is often measured using acute-phase inflammatory proteins such as CRP [20,21], but it is not always reliable [22], particularly in neonates, and a single-time-point measure may not reflect baseline inflammation or capture chronic inflammation [23]. These challenges have been addressed by the development of DNA methylation (DNAm) markers of protein expression (EpiScores), which are derived from a linear weighted sum of DNAm sites that are correlated with protein levels. Several EpiScores are associated with magnetic resonance imaging (MRI) measures of brain health, cognition, child mental health, stroke, ischaemic heart disease Alzheimer's disease, lung cancer [24–31]. In neonates, DNAm CRP is associated with birth gestational age (GA), perinatal inflammatory processes, and MRI features of encephalopathy of prematurity [32]. Childhood SES is associated with differential DNAm in inflammation-related genes

[33,34] and at CpG sites that correlate with an inflammation index [35]. Adult SES and social mobility are associated with variations in DNAm in inflammation-related genes [33,36]. Importantly, SES-related DNAm variations are associated with differences in gene expression so may have functional consequences [33,36].

We investigated relationships between preterm birth, SES, and 104 EpiScores enriched for inflammation-related proteins [26–28,31,37]. We tested the following hypotheses: first, low GA is associated with differences in EpiScores; second, SES is correlated with EpiScores, and interacts with birth GA, but the relationship is attenuated by inflammatory disease burden in preterm infants.

## 2.0 Methods

### 2.1 Participants

Participants were preterm infants (born  $\leq 33$  weeks' gestation) and term-born infants born at the Royal Infirmary of Edinburgh, UK. These infants were recruited to a longitudinal cohort study designed to investigate the effect of preterm birth on brain development and outcomes with multimodal data collection [38]. Infants were recruited between February 2012 and December 2021.

Exclusion criteria were congenital malformation, chromosomal abnormality, congenital infection, cystic periventricular leukomalacia, haemorrhagic parenchymal infarction, and post-haemorrhagic ventricular dilatation. These criteria mean the cohort is representative of the majority of survivors of modern intensive care practices [38].

Final participants included were 217 preterm infants (born  $\leq 33$  weeks' GA) and 115 term-born infants, with median birth GA of 29.29 weeks and 39.71 weeks respectively. Their demographic characteristics are shown in Table 1. The three SES measures (Scottish Index of Multiple Deprivation (SIMD 2016) [39], maternal education and maternal occupation) differed between the preterm and term groups (Cohen's  $d$  effect sizes 0.52-0.68). Ethnicity did not differ between groups and is representative of the Edinburgh area [40].

[Table 1 near here]

1

## 2 *2.2 DNA methylation*

3 Saliva samples for DNAm were collected at term-equivalent age using Oragene OG-575 Assisted  
4 Collection kits (DNA Genotek, ON, Canada) and DNA was extracted using prepIT.L2P reagent (DNA  
5 Genotek, ON, Canada). Saliva sampling was used due to accessibility and the non-invasiveness of  
6 the method; DNAm patterns measured via saliva samples correlate with brain and other tissue DNAm  
7 patterns [41,42]. We chose to sample at the term-corrected gestation timepoint to include the  
8 allostatic load of both prenatal and early postnatal exposures.

9

10 DNA was bisulphite converted and methylation levels were measured using Illumina  
11 HumanMethylationEPIC BeadChip (Illumina, San Diego, CA, USA) at the Edinburgh Clinical  
12 Research Facility (Edinburgh, UK). The arrays were imaged on the Illumina iScan or HiScan platform,  
13 and genotypes were called automatically using GenomeStudio Analysis software version 2011.1  
14 (Illumina). DNAm was processed in four batches.

15

16 Raw intensity (.idat) files were read into the R environment using minfi. watermelon and minfi were  
17 used for preprocessing, quality control and normalisation [43]. The pfilter function in watermelon was  
18 used to exclude samples with 1% of sites with a detection  $p$ -value  $>0.05$ , sites with beadcount  $<3$  in  
19 5% of samples, and sites with 1% of samples with detection  $p$ -value  $>0.05$ . Cross hybridising probes,  
20 probes targeting single nucleotide polymorphisms with overall minor allele frequency  $\geq 0.05$ , and  
21 control probes were also removed. Samples were removed if there was a mismatch between  
22 predicted sex (minfi) and recorded sex ( $n=3$ ), or if samples did not meet preprocessing quality control  
23 criteria ( $n=29$ ). Data were danet normalised, which includes background correction and dye bias  
24 correction [43]. Saliva contains different cells types, including buccal epithelial cells. Epithelial cell  
25 proportions were estimated with epigenetic dissection of intra-sample heterogeneity with the reduced  
26 partial correlation method implemented in the R package EpiDISH [44]. Probes located on sex  
27 chromosomes were removed before analysis. The cohort includes twins ( $n=32$ ); these were randomly  
28 removed leaving one participant per twin pair. This left a final sample size of  $n=332$ .

29

## 30 *2.3 EpiScore calculation*

1 The 104 protein EpiScores included 100 EpiScores from Gadd *et al* [26], a study enriched for  
2 inflammatory-related proteins, excluding those where the required CpGs were not available, owing to  
3 differences in assay platform CpG coverage relative to Gadd *et al* [26]. For duplicate proteins, those  
4 developed using Olink platform-identified proteins (antibody-based assays) were prioritised over those  
5 from SOMAscan platforms (aptamer-based assays), due to specificity and reproducibility, and the  
6 variable correlation between the two methods [45–48] (for details see Supplementary eMethods,  
7 Additional File 1). In addition, we included EpiScores for growth and differentiation factor 15 (GDF15)  
8 and N-terminal-pro B-type natriuretic peptide (NTproBNP) [37], and CRP. The CRP EpiScore used  
9 was Barker *et al*'s seven-CpG variation of Ligthart *et al*'s CRP EpiScore [27,31], as this is known to  
10 correlate with birth GA, perinatal pro-inflammatory exposures, and neonatal brain development [32].

11

12 For each individual, EpiScores were obtained by multiplying the methylation proportion at a given  
13 CpG by the effect size from previous studies. This was performed using the MethylDetectR platform  
14 [49] for those inflammatory proteins currently included and using R for those not currently included  
15 (CRP, GDF15, IL6, NTproBNP). All CpG sites and coefficients required to calculate the 104  
16 EpiScores are in Supplementary Table 1 (Additional File 2).

17

## 18 2.4 Statistics

19 The predictor variables were SES and birth GA. SES was operationalised in three ways:  
20 neighbourhood-level SES using the Scottish Index of Multiple Deprivation (SIMD) [39], and two  
21 measures of family-level SES, which were maternal education (highest educational qualification) and  
22 maternal occupation (current or most recent occupation). For further details, see Supplementary  
23 eMethods, Additional File 1. Birth GA was a continuous variable to maximise statistical power [50,51].  
24 We adjusted for GA at saliva sampling, DNAm batch, infant sex, and birthweight z-score.

25

26 All statistical analyses were performed in R (version 4.3.1) and were preregistered [52].

27

28 Principal component analysis (PCA) was used to determine the significance threshold for controlling  
29 type 1 error in analyses of multiple EpiScores [53]. We began with a correlation analysis, which  
30 showed correlation coefficients between EpiScores of |0.01 to 0.93| (Figure 1A). To determine the

number of statistical “families” among the 104 EpiScores, PCA was performed. This yielded two principal components with eigenvalues >1, our pre-specified threshold, which explained 59.5% and 17.2% of variance, respectively (Figure 1B). Standardised component loadings are provided in Supplementary Table 3 (Additional File 1). In all subsequent analyses, we corrected for multiple comparisons across EpiScores and SES measures using a Bonferroni-adjusted  $p$ -value threshold of  $8.3 \times 10^{-3}$ . This is  $0.05/(2 \times 3)$ , with two reflecting the two principal components for EpiScores and three reflecting the number of SES measures used.

*[Figure 1 near here]*

We constructed generalised linear regression models for each EpiScore as outcome measure to assess associations between GA, each of the three SES measures (separate models for each of SIMD, maternal education, and maternal occupation), and the product interaction term SES\*birth GA (removing the term if not significant), and adjusting for GA at sampling, sex, and batch.

For the preterm sub-group, we then additionally adjusted for perinatal inflammatory exposures known to be associated with the CRP EpiScore as, to our knowledge, this is the only DNAm proxy of an inflammatory protein that has been studied in this context [32]. These were histological chorioamnionitis (HCA), sepsis, bronchopulmonary dysplasia (BPD) and necrotising enterocolitis (NEC). For definitions see Supplementary eMethods (Additional File 1) and for frequencies see Supplementary Table 2 (Additional File 1).

### **3.0 Results**

#### **3.1 Associations between gestational age and EpiScores**

Gestational age associated with 43 of the 104 EpiScores after adjustment for SIMD, maternal education, or maternal occupation (Figure 2A-C). The proteins represented by the 43 EpiScores are listed in Table 2 categorised by functional annotation adapted from the STRING database[54], and their broader roles in immune processes and inflammation, and the pathogenesis of neonatal diseases, where known, are described in Supplementary Table 4 (Additional File 1).

29 (67%) EpiScores negatively associated with birth GA (standardised estimates |0.14-0.76|, adjusted  $p$ -value  $<8.3 \times 10^{-3}$ ), and 14 (33%) EpiScores positively associated with birth GA (standardised estimates 0.14-0.88, adjusted  $p$ -value  $<8.3 \times 10^{-3}$ ).

33 EpiScores associated with low GA irrespective of SES measure used in the model. The results for all 104 EpiScores are provided in Supplementary Figures 1-9 (Additional File 1).

[Table 2 near here]

[Figure 2 near here]

### 3.2 Associations between SES, EpiScores, and the effect of inflammatory comorbidities of preterm birth.

Three out of 104 EpiScores associated with SES measures or the interaction between SES and GA (Figure 3). There was a small effect size association between higher afamin EpiScore and higher maternal occupation (standardised  $\beta=0.06$ , 95% confidence interval (CI) 0.02-0.11,  $p=0.0082$ ), and DNAm afamin associated with the birth GA\*maternal education interaction term such that afamin positively correlated with birth GA among babies with mothers without university education, and negatively correlated with birth GA among babies with mothers with university education (undergraduate or postgraduate) (standardised  $\beta=-0.12$ , 95% CI -0.20 to -0.04,  $p=0.0041$ , Supplementary Figure 10, Additional File 1).

Higher intercellular adhesion molecule 5 (ICAM5) EpiScore associated with higher SIMD (standardised  $\beta=0.13$ , 95% CI 0.03-0.23,  $p=0.0079$ ). Hepatocyte growth factor-like protein (HGFI) associated with the birth GA\*maternal occupation interaction term (standardised  $\beta=-0.10$ , 95% CI -0.16- -0.04,  $p=0.0021$ , Supplementary Figure 11, Additional File 1), such that HGFI EpiScore positively correlated with birth GA among babies with mothers who were unemployed, homemakers, in full time education, or in unskilled or manual occupations, but negatively correlated with birth GA among babies with mothers in partly skilled, non-manual skilled or professional occupations.

In the planned analysis of preterm-born babies only, when controlling for inflammatory exposures (sepsis, HCA, NEC, and BPD), a greater proportion of  $R^2$  was explained ( $R^2=0.053$ -0.28 in unadjusted



models,  $R^2=0.029-0.437$  in adjusted models), but the EpiScores no longer met our statistical threshold ( $p$ -values  $>0.045$  with adjusted  $p$ -value threshold  $<8.3 \times 10^{-3}$ , see Supplementary Table 5, Additional File 1).

[Figure 3 near here]

## 4.0 Discussion

In this study, we identified several associations between a set of EpiScores enriched for inflammatory proteins and low GA at birth. Few EpiScores associated with SES within the whole sample and these associations were partially attenuated in preterm infants who experienced inflammatory comorbidities. This is the first study to assess the impact of preterm birth and social status using epigenetic signatures designed to reflect the circulating proteome.

### 4.1 Associations between birth GA and EpiScores

43 EpiScores associated with preterm birth when sampled at term equivalent age. The EpiScores reflect chemokines, growth factors, proteins required for neurogenesis and vascular development, cell membrane proteins and receptors, and immune response proteins (Table 2). As well as having specific immunoregulatory roles, in the neonatal setting or relevant animal models, these proteins are associated with several comorbidities and developmental consequences of preterm birth. These include lung development and disease such as BPD [55–67], in utero and postnatal growth failure [68–71], HCA [7,72,73], patent ductus arteriosus [74–77], retinopathy of prematurity [78–82], NEC [83–85], hyperglycaemia [86], sepsis [85,87–90], brain injury [32,91–94], and neurodevelopmental outcomes [95–99].

### 4.2 Associations between SES measures and EpiScores

SES appears to play a much smaller role in the patterning of EpiScores compared to birth GA. SES measures, or interactions between birth GA and SES measures, were associated with only three of the 104 EpiScores studied: afamin, ICAM5 and HGFI. Afamin and ICAM5 positively associated with maternal occupation and SIMD, respectively. Afamin and HGFI both associated with the interaction term between SES and birth GA. Afamin is involved in vitamin E transport [100], ICAM5 has a role in

microglial regulation [101], and HGFI is a macrophage-stimulating protein [102]. The relationships between these proteins and SES have not previously been investigated, although afamin is associated with the development of metabolic syndrome [103], which varies with SES [104].

Among preterm infants, no SES-EpiScore associations survived adjustment for inflammatory exposures, which suggests that the weak effects of SES on the neonatal proteome that we observed in a small number of EpiScores are at least partially accounted for by inflammatory pathologies in early life. Taken together, the results suggest immune dysregulation, proxied by EpiScores, may not be the primary axis through which SES becomes embedded in the development of preterm infants during neonatal intensive care.

SES has been consistently associated with inflammation in adulthood, including in relation to childhood deprivation [15,105], but less is known about the relationship between SES and inflammation in the neonatal period. A longitudinal study by Leviton *et al* [106], with five sampling timepoints during the first month of life after preterm birth, showed that maternal eligibility for Medicaid associated with levels of 14 inflammatory proteins (IL6R, TNFR1, TNFR2, IL8, ICAM1, VCAM1, TSH, EPO, bFGF, IGF1, VEGF, PIGF, Ang-1, Ang-2). However, only three were significant at more than one timepoint during the month after preterm birth (TSH, bFGF, Ang-1), and none associated at all five measurement timepoints. These studies, taken together with our results, suggest that the impact of SES on immune regulation is relatively modest and inconsistent in the newborn period but accrues through to adulthood. Further research is required to understand how and when SES becomes embedded in child development and whether early life events such as preterm birth modify that process; EpiScores could be a powerful tool for investigating the temporal dynamics of social determinants of child health.

#### 4.3 Strengths and limitations

Strengths of this study include the large sample of term and preterm neonates; to the best of our knowledge, this is the first examination of multiple DNA methylation-based estimators of circulating proteins in a neonatal sample. We derived EpiScores from minimally-invasive sampling (buccal cells from saliva) which overcomes the ethical challenge of venepuncture for research in children. The

EpiScores, serving as proxies of inflammatory proteins and sampled at term-equivalent age in preterm infants, were selected for their potential to capture chronic, cumulative inflammation associated with preterm birth and neonatal intensive care exposures [23–25]. We adjusted for variables associated with DNAm, and additionally for inflammatory exposures to increase the clinical validity of our results.

The study has some limitations. The EpiScores used were developed in adult cohorts [26–28,37] and have not been validated in neonatal populations with neonatal protein levels, although we have previously established that neonatal DNAmCRP scores correlate with cumulative inflammatory exposures [32]. Further studies that evaluate serial blood protein levels with EpiScores could be informative but would need to rely on small volume samples taken during time of venepuncture for clinical reasons, given the practical and ethical challenges of serial venepuncture for research purposes in neonates. The EpiScores were also trained using blood samples [26–28,37], whereas we have projected these scores into saliva samples. However, previous studies have successfully used similar cross-tissue techniques [32,107,108], and in neonates saliva provides a non-invasive and accessible sample method. Not all inflammatory-related proteins are represented, as we were limited by available EpiScores, so we may have underestimated the full complexity of the relationship between birth GA, SES, and inflammation. Longitudinal investigations are imperative for elucidating whether the DNA methylation signatures associated with gestational age identified in this study exert a causal influence on the inflammation-associated mechanisms in preterm birth. It remains crucial to discern whether these signatures represent a direct downstream consequence of GA itself or are induced by specific factors correlated with GA, yet not necessarily driven by chronic inflammation. Mendelian randomization studies, integrating genomic and epigenomic determinants, are a promising methodological approach to disentangle the directionality of these intricate relationships. The study population is comparable to other neonatal populations in high-income, majority-white settings, but these results may not generalise to settings with different socioeconomic or ethnicity profiles. We studied several measures of SES but were not able to include all that could be relevant, such as household income.

## 5.0 Conclusion

We identified 43 EpiScores enriched for inflammatory proteins that associated with low birth GA. These 43 proteins offer novel insights into the physiological response to preterm birth and warrant further study to explore their role in the relationship between preterm birth, inflammation, and longer-term outcomes. We found only three EpiScores associated with SES in the neonatal period, none of which survived adjustment for perinatal pro-inflammatory exposures, suggesting that inflammation is unlikely to be the primary axis through which SES becomes embedded in the development of preterm infants in the neonatal period.

## 6.0 Declarations

### 6.1 Ethics approval and consent to participate

Ethical approval for the study was obtained from the National Research Ethics Service, South-East Scotland Research Ethics Committee (REC 11/55/0061, 13/SS/0143, 16/SS/0154), and NHS Lothian Research and Development (2016/0255). Written informed consent was obtained from parents.

### 6.2 Consent for publication

Not applicable.

### 6.3 Availability of data and materials

DNA methylation data are available to researchers subject to the terms of the Data Access Policy: <https://www.ed.ac.uk/centre-reproductive-health/tebc/about-tebc/for-researchers/data-access-collaboration>.

### 6.4 Competing interests

LM has received speaker and consultancy fees from Illumina. REM is a scientific advisor to the Epigenetic Clock Development Foundation and to Optima Partners. All other authors declare that they have no competing interests.

### 6.5 Funding

This work was supported by Theirworld ([www.theirworld.org](http://www.theirworld.org)) and a UKRI MRC Programme Grant MR/X003434/1. KM receives salary from NHS Scotland. GS is funded by an MRC Clinician Scientist

Fellowship MR/X019535/1. SRC is supported by a Sir Henry Dale Fellowship jointly funded by the Wellcome Trust and the Royal Society (221890/Z/20/Z).

#### *6.6 Authors' contributions*

KM: Conceptualisation, methodology, formal analysis, writing (original draft), writing (review/editing), visualization. ELSC: Formal analysis, writing (review/editing). KV: Formal analysis, writing (review/editing), visualization. RH: Methodology, formal analysis, writing (review/editing). DG: Methodology, formal analysis, writing (review/editing). JB: Formal analysis, writing (review/editing). GS: Investigation, writing (review/editing). AJS: Methodology, writing (review/editing). AC: Investigation, resources, data curation, writing (review/editing). LM: Investigation, resources, writing (review/editing). HCW: Methodology, writing (review/editing). HR: Conceptualisation, writing (original draft), writing (review/editing), supervision. REM: Conceptualisation, methodology, writing (original draft), writing (review/editing), supervision. SRC: Conceptualisation, methodology, writing (original draft), writing (review/editing), supervision. JPB: Conceptualisation, methodology, writing (original draft), writing (review/editing), supervision, project administration, funding acquisition.

#### *6.7 Acknowledgements*

The authors are grateful to the families who consented to take part in the study.

# Figure legends

## Figure 1. Determining significance threshold

Principal component analysis was used to determine the adjusted statistical significance threshold, given multiple statistical comparisons. Figure 1A shows a correlation matrix of 104 EpiScores, showing correlation coefficient as red for positive and blue negative associations when significant ( $p < 0.05$ ). Figure 1B shows a scree plot of principal components, with the eigenvalues for each component. Standardised component loadings for principal components one and two are provided in Supplementary Table 3 (Additional File 1).

## Figure 2. EpiScores associated with gestational age in regression models adjusted for socioeconomic status

EpiScores associated with gestational age in regression models with (A) Scottish Index of Multiple Deprivation, (B) maternal education, and (C) maternal occupation. Figure 2A ( $n=331$ ) shows 39 associations, Figure 2B ( $n=323$ ) shows 35 associations, and Figure 2C ( $n=328$ ) shows 39 associations. Points and bars represent standardised beta and 95% confidence intervals, with red indicating positive and blue negative associations. Covariates included in all models: age at sample, birthweight z-score, sex, and methylation processing batch. Bonferroni-adjusted  $p$ -value  $< 8.3 \times 10^{-3}$ . CCL11 – C-C chemokine 11, CCL18 – C-C chemokine 18, CCL21 – C-C chemokine 21, CCL22 – C-C chemokine 22, CCL25 – C-C chemokine 25, CD5L – CD5 antigen-like protein, CD6 – T-cell differentiation antigen, CD163 – scavenger receptor cysteine-rich type 1 protein M130, CI – confidence interval, CRP – C-reactive protein, CRTAM – Cytotoxic and regulatory T-cell molecule, CXCL9 – C-X-C motif chemokine 9, CXCL10 – C-X-C motif chemokine 10, FAP – Fibroblast Activation Protein alpha, FCGR3B – Low affinity immunoglobulin gamma Fc region receptor III-B, FcRL2 – Fc receptor-like protein 2, FGF21 – Fibroblast growth factor 21, GDF15 – Growth/differentiation factor 15, GHR – Growth hormone receptor, HCII – Heparin cofactor II, HGF – Hepatocyte growth factor alpha chain, IGFBP4 – Insulin-like growth factor-binding protein 4, MMP9 – Matrix metalloproteinase-9, NCAM1 – Neural cell adhesion molecule 1, PIGR – Polymeric immunoglobulin receptor, SCGF – stem cell growth factor, SIMD – Scottish Index of Multiple Deprivation, SKR3 – Serine/threonine-protein kinase receptor R3, SLITRK5 – SLIT and NTRK-like

protein 5, VCAM1 – Vascular cell adhesion protein 1, VEGFA – Vascular endothelial growth factor A, WFIKKN2 – WAP Kazal immunoglobulin Kunitz and NTR domain-containing protein 2.

*Figure 3. EpiScores associated with socioeconomic status or an interaction between socioeconomic status and birth gestational age*

EpiScores associated with socioeconomic status (Scottish Index of Multiple Deprivation, maternal education, or maternal occupation), or with an interaction between socioeconomic status and birth gestational age. Regression models with gestational age at birth, gestational age at sample, birthweight z-score, sex, and methylation processing batch. Sample sizes: for the Scottish Index of Multiple Deprivation n=331, for maternal education n=323, and for maternal occupation n=328. 3/104 EpiScores were significant (Bonferroni-adjusted  $p$ -value  $<8.3 \times 10^{-3}$ ). Points and bars represent standardised beta and 95% confidence intervals, with red indicating positive and blue negative associations.

CI – confidence interval, GA – gestational age, HGFI – Hepatocyte growth factor-like protein alpha chain, ICAM5 – Intercellular adhesion molecule 5, SIMD – Scottish Index of Multiple Deprivation.

# 1 Tables

## 2 Table 1. Participant characteristics

Demographic measure		Preterm (n=217)	Term (n=115)
Sex	Male	114/217 (52.5%)	64/115 (55.7%)
	Female	103/217 (47.5%)	51/115 (44.3%)
Birth GA (weeks) - median (range)		29.29 (22.14-33.0)	39.71 (37.0-42.14)
Birthweight (g) - median (range)		1200 (370-2510)	3450 (2346-4670)
Birthweight z-score - median (range)		0.10 (-3.13-2.07)	0.43 (-2.3-2.96)
SIMD rank - median (range) <sup>A</sup>		3720 (6-6966)	5344 (267-6967)
Maternal ethnicity	African	1/217 (0.5%)	0/115 (0%)
	Bangladeshi	0/217 (0%)	1/115 (0.9%)
	Caribbean	0/217 (0%)	0/115 (0%)
	Chinese	0/217 (0%)	1/115 (0.9%)
	Indian	3/217 (2.4%)	1/115 (0.9%)
	Pakistani	4/214 (1.8%)	1/115 (0.9%)
	White	195/217 (89.9%)	106/115 (92.2%)
	White/Asian	2/217 (0.9%)	0/115 (0%)
	White/Black African	0/217 (0%)	1/115 (0.9%)
	White/Black Caribbean	1/217 (0.5%)	1/115 (0.9%)
	Other Asian <sup>B</sup>	1/217 (0.5%)	2/115 (1.7%)
	Other ethnic group <sup>C</sup>	6/217 (2.8%)	0/115 (0%)
	Other mixed ethnic background <sup>D</sup>	4/217 (1.8%)	1/115 (0.9%)
Maternal education	None	7/208 (3.4%)	0/115 (0%)
	Basic high school qualification (1-4)	5/208 (2.4%)	2/115 (1.7%)
	Basic high school qualification (>=5)	8/208 (3.8%)	1/115 (0.9%)
	Advanced high school	32/208 (15.4%)	3/115 (2.6%)



	qualification		
	College qualification	46/208 (22.1%)	8/115 (7.0%)
	University undergraduate	61/208 (29.3%)	50/115 (43.5%)
	University postgraduate	49/208 (23.6%)	51/115 (44.3%)
Maternal occupation	Unemployed	12/213 (5.6%)	1/115 (0.9%)
	Homemaker	10/213 (4.7%)	1/115 (0.9%)
	Still in full time education	9/213 (4.2%)	1/115 (0.9%)
	Sheltered employment	1/213 (0.5%)	0/115 (0%)
	Unskilled	11/213 (5.2%)	3/115 (2.6%)
	Partly skilled	7/213 (3.3%)	2/115 (1.7%)
	Manual skilled	25/213 (11.7%)	9/115 (7.8%)
	Non-manual skilled	44/213 (20.7%)	11/115 (9.6%)
	Professional	94/213 (44.1%)	87/115 (75.7%)
Sample GA (weeks) - median (range)		40.57 (36.57-45.86)	42.0 (39.86-47.14)
Batch	1	92/217 (42.4%)	46/115 (40.0%)
	2	62/217 (28.6%)	56/115 (48.7%)
	3	32/217 (14.7%)	1/115 (0.9%)
	4	31/217 (14.3%)	12/115 (10.4%)

1 A: Preterm n=216, term n=115

2 B: "Other Asian" includes Sri Lankan (n=1), Malay (n=1)

3 C: "Other ethnic group" includes Arab (n=1), Iraqi (n=1), white Bulgarian (n=1), Fijian (n=1), Japanese  
4 (n=1), Hong Kong (n=1).

5 D: "Other mixed ethnic background" includes Pakistani/Scottish (n=2), British/Arab (n=1), Sri  
6 Lankan/Black (n=1), Sri Lankan/Indian (n=1).

7 GA = gestational age, SIMD = Scottish index of multiple deprivation.

8

9

10

11

1 *Table 2. Protein EpiScores associated with birth gestational age*

Chemokines	Growth factors	Neurogenesis	Vascular development	Cell membrane proteins / receptors	Other immune response
CCL11	FGF21	NCAM1	SKR3	Afamin	Complement C9
CCL18	GDF15	Semaphorin 3E	VCAM1	CD5L	CRP
CCL21	GHR	SLITRK5	VEGFA	CD6	CRTAM
CCL22	HGF			CD48	FAP
CCL25	IGFBP4			CD163	HCII
CXCL9	SCGF alpha			CD209	L-selectin
CXCL10	SCGF beta			Contactin-4	Lactotransferrin
	WFIKKN2			FcRL2	MMP9
				FcGR3B	S100A9
				PIGR	Sialoadhesin
				Thrombopoietin receptor	Trypsin-2

2 43 EpiScores associated with birth gestational age in regression models adjusted for socioeconomic status. Roles adapted from the STRING database[54].

3 See Supplementary Table 4, Additional File 1, for further details of the functional roles of each protein, including roles in immunity and inflammation, and in  
4 preterm infants specifically.

5 CCL11 – C-C chemokine 11, CCL18 – C-C chemokine 18, CCL21 – C-C chemokine 21, CCL22 – C-C chemokine 22, CCL25 – C-C chemokine 25, CD5L –  
6 CD5 antigen-like protein, CD6 – T-cell differentiation antigen, CD163 – scavenger receptor cysteine-rich type 1 protein M130, CRP – C-reactive protein,  
7 CRTAM – Cytotoxic and regulatory T-cell molecule, CXCL9 – C-X-C motif chemokine 9, CXCL10 – C-X-C motif chemokine 10, FAP – Fibroblast Activation  
8 Protein alpha, FCGR3B – Low affinity immunoglobulin gamma Fc region receptor III-B, FcRL2 – Fc receptor-like protein 2, FGF21 – Fibroblast growth factor

21, GDF15 – Growth/differentiation factor 15, GHR – Growth hormone receptor, HCII – Heparin cofactor II, HGF – Hepatocyte growth factor alpha chain, IGFBP4 – Insulin-like growth factor-binding protein 4, MMP9 – Matrix metalloproteinase-9, NCAM1 – Neural cell adhesion molecule 1, PIGR – Polymeric immunoglobulin receptor, SCGF – stem cell growth factor, SIMD – Scottish Index of Multiple Deprivation, SKR3 – Serine/threonine-protein kinase receptor R3, SLITRK5 – SLIT and NTRK-like protein 5, VCAM1 – Vascular cell adhesion protein 1, VEGFA – Vascular endothelial growth factor A, WFIKKN2 – WAP Kazal immunoglobulin Kunitz and NTR domain-containing protein 2.

# References

1. Ohuma EO, Moller A-B, Bradley E, Chakwera S, Hussain-Alkhateeb L, Lewin A, et al. National, regional, and global estimates of preterm birth in 2020, with trends from 2010: a systematic analysis. *Lancet*. 2023;402:1261–71.
2. Johnson S, Marlow N. Early and long-term outcome of infants born extremely preterm. *Archives of Disease in Childhood*. 2017;102:97–102.
3. Agrawal S, Rao SC, Bulsara MK, Patole SK. Prevalence of Autism Spectrum Disorder in Preterm Infants: A Meta-analysis. *Pediatrics*. 2018;142:1–14.
4. Twilhaar ES, Wade RM, deKieviet JF, vanGoudoever JB, vanElburg RM, Oosterlaan J. Cognitive Outcomes of Children Born Extremely or Very Preterm Since the 1990s and Associated Risk Factors: A Meta-analysis and Meta-regression. *Jama Pediatr*. 2018;172:361.
5. Crump C. An overview of adult health outcomes after preterm birth. *Early Hum Dev*. 2020;150:105187.
6. Hagberg H, Mallard C, Ferriero DM, Vannucci SJ, Levison SW, Vexler ZS, et al. The role of inflammation in perinatal brain injury. *Nat Rev Neurol*. 2015;11:192–208.
7. Sullivan G, Galdi P, Blesa-Cábez M, Borbye-Lorenzen N, Stoye DQ, Lamb GJ, et al. Interleukin-8 dysregulation is implicated in brain dysmaturation following preterm birth. *Brain Behav Immun*. 2020;90:311–8.
8. Raisi-Estabragh Z, Cooper J, Bethell MS, McCracken C, Lewandowski AJ, Leeson P, et al. Lower birth weight is linked to poorer cardiovascular health in middle-aged population-based adults. *Heart*. 2023;109:535–41.
9. Farah MJ. The Neuroscience of Socioeconomic Status: Correlates, Causes, and Consequences. *Neuron*. 2017;96:56–71.
10. Kivimäki M, Batty GD, Pentti J, Shipley MJ, Sipilä PN, Nyberg ST, et al. Association between socioeconomic status and the development of mental and physical health conditions in adulthood: a multi-cohort study. *Lancet Public Heal*. 2020;5:e140–9.
11. Duncan GJ, Ziol-Guest KM, Kalil A. Early-Childhood Poverty and Adult Attainment, Behavior, and Health. *Child Dev*. 2010;81:306–25.
12. Pillas D, Marmot M, Naicker K, Goldblatt P, Morrison J, Pikhart H. Social inequalities in early childhood health and development: a European-wide systematic review. *Pediatr Res*. 2014;76:418–24.
13. Thomson K, Moffat M, Arisa O, Jesurasa A, Richmond C, Odeniyi A, et al. Socioeconomic inequalities and adverse pregnancy outcomes in the UK and Republic of Ireland: a systematic review and meta-analysis. *Bmj Open*. 2021;11:e042753.
14. Ruiz M, Goldblatt P, Morrison J, Kukla L, Švancara J, Riitta-Järvelin M, et al. Mother's education and the risk of preterm and small for gestational age birth: a DRIVERS meta-analysis of 12 European cohorts. *J Epidemiol Commun H*. 2015;69:826–33.
15. Muscatell KA, Brosso SN, Humphreys KL. Socioeconomic status and inflammation: a meta-analysis. *Mol Psychiatr*. 2020;25:2189–99.

- 1 16. Broyles ST, Staiano AE, Drazba KT, Gupta AK, Sothorn M, Katzmarzyk PT. Elevated C-Reactive  
2 Protein in Children from Risky Neighborhoods: Evidence for a Stress Pathway Linking Neighborhoods  
3 and Inflammation in Children. *Plos One*. 2012;7:e45419.
- 4 17. Dowd JB, Zajacova A, Aiello AE. Predictors of Inflammation in U.S. Children Aged 3–16 Years.  
5 *Am J Prev Med*. 2010;39:314–20.
- 6 18. Schmeer KK, Yoon AJ. Home sweet home? Home physical environment and inflammation in  
7 children. *Soc Sci Res*. 2016;60:236–48.
- 8 19. Schmeer KK, Yoon A. Socioeconomic status inequalities in low-grade inflammation during  
9 childhood. *Arch Dis Child*. 2016;101:1043.
- 10 20. Chiesa C, Natale F, Pascone R, Osborn JF, Pacifico L, Bonci E, et al. C reactive protein and  
11 procalcitonin: Reference intervals for preterm and term newborns during the early neonatal period.  
12 *Clin Chim Acta*. 2011;412:1053–9.
- 13 21. Brown JVE, Meader N, Cleminson J, McGuire W. C-reactive protein for diagnosing late-onset  
14 infection in newborn infants. Group CN, editor. *Cochrane Database of Systematic Reviews*.  
15 2019;10:16–78.
- 16 22. Bower JK, Lazo M, Juraschek SP, Selvin E. Within-Person Variability in High-Sensitivity C-  
17 Reactive Protein. *Arch Intern Med*. 2012;172:1519–21.
- 18 23. Dammann O, Leviton A. Intermittent or sustained systemic inflammation and the preterm brain.  
19 *Pediatr Res*. 2014;75:376–80.
- 20 24. Stevenson AJ, McCartney DL, Harris SE, Taylor AM, Redmond P, Starr JM, et al. Trajectories of  
21 inflammatory biomarkers over the eighth decade and their associations with immune cell profiles and  
22 epigenetic ageing. *Clin Epigenetics*. 2018;10:159.
- 23 25. Stevenson AJ, McCartney DL, Hillary RF, Campbell A, Morris SW, Bermingham ML, et al.  
24 Characterisation of an inflammation-related epigenetic score and its association with cognitive ability.  
25 *Clin Epigenetics*. 2020;12:113.
- 26 26. Gadd DA, Hillary RF, McCartney DL, Zaghloul SB, Stevenson AJ, Cheng Y, et al. Epigenetic  
27 scores for the circulating proteome as tools for disease prediction. *Elife*. 2022;11:e71802.
- 28 27. Ligthart S, Marzi C, Aslibekyan S, Mendelson MM, Conneely KN, Tanaka T, et al. DNA  
29 methylation signatures of chronic low-grade inflammation are associated with complex diseases.  
30 *Genome Biol*. 2016;17:255.
- 31 28. Stevenson AJ, Gadd DA, Hillary RF, McCartney DL, Campbell A, Walker RM, et al. Creating and  
32 Validating a DNA Methylation-Based Proxy for Interleukin-6. *Journals Gerontology Ser Biological Sci*  
33 *Medical Sci*. 2021;76:2284–92.
- 34 29. Conole ELS, Stevenson AJ, Maniega SM, Harris SE, Green C, Hernández M del CV, et al. DNA  
35 Methylation and Protein Markers of Chronic Inflammation and Their Associations With Brain and  
36 Cognitive Aging. *Neurology*. 2021;97:e2340–52.
- 37 30. Green C, Shen X, Stevenson AJ, Conole ELS, Harris MA, Barbu MC, et al. Structural brain  
38 correlates of serum and epigenetic markers of inflammation in major depressive disorder. *Brain*  
39 *Behav Immun*. 2021;92:39–48.
- 40 31. Barker ED, Cecil CAM, Walton E, Houtepen LC, O'Connor TG, Danese A, et al. Inflammation-  
41 related epigenetic risk and child and adolescent mental health: A prospective study from pregnancy to  
42 middle adolescence. *Dev Psychopathol*. 2018;30:1145–56.

32. Conole ELS, Vaher K, Blesa-Cábez M, Sullivan G, Stevenson AJ, Hall J, et al. Immuno-epigenetic signature derived in saliva associates with the encephalopathy of prematurity and perinatal inflammatory disorders. *Brain Behav Immun*. 2023;110:322–38.
33. Needham BL, Smith JA, Zhao W, Wang X, Mukherjee B, Kardia SLR, et al. Life course socioeconomic status and DNA methylation in genes related to stress reactivity and inflammation: The multi-ethnic study of atherosclerosis. *Epigenetics*. 2015;10:958–69.
34. Stringhini S, Polidoro S, Sacerdote C, Kelly RS, Veldhoven K van, Agnoli C, et al. Life-course socioeconomic status and DNA methylation of genes regulating inflammation. *Int J Epidemiol*. 2015;44:1320–30.
35. McDade TW, Ryan C, Jones MJ, MacIsaac JL, Morin AM, Meyer JM, et al. Social and physical environments early in development predict DNA methylation of inflammatory genes in young adulthood. *Proc National Acad Sci*. 2017;114:7611–6.
36. Smith JA, Zhao W, Wang X, Ratliff SM, Mukherjee B, Kardia SLR, et al. Neighborhood characteristics influence DNA methylation of genes involved in stress response and inflammation: The Multi-Ethnic Study of Atherosclerosis. *Epigenetics*. 2017;12:662–73.
37. Gadd DA, Smith HM, Mullin D, Chybowska O, Hillary RF, Kimenai DM, et al. DNAm scores for serum GDF15 and NT-proBNP levels associate with a range of traits affecting the body and brain. *medRxiv*. 2023;2023.10.18.23297200.
38. Boardman JP, Hall J, Thrippleton MJ, Reynolds RM, Bogaert D, Davidson DJ, et al. Impact of preterm birth on brain development and long-term outcome: protocol for a cohort study in Scotland. *Bmj Open*. 2020;10:e035854.
39. Scottish National Statistics. SIMD - Scottish Index of Multiple Deprivation: SIMD16 technical notes. Edinburgh: Scottish National Statistics; 2016 p. 1–69.
40. National Records of Scotland. Scotland's Census [Internet]. 2020 [cited 2023 Nov 23]. Available from: <https://www.scotlandscensus.gov.uk/search-the-census/#/>
41. Lowe R, Gemma C, Beyan H, Hawa MI, Bazeos A, Leslie RD, et al. Buccals are likely to be a more informative surrogate tissue than blood for epigenome-wide association studies. *Epigenetics*. 2013;8:445–54.
42. Braun PR, Han S, Hing B, Nagahama Y, Gaul LN, Heinzman JT, et al. Genome-wide DNA methylation comparison between live human brain and peripheral tissues within individuals. *Transl Psychiat*. 2019;9:47.
43. Pidsley R, Wong CCY, Volta M, Lunnon K, Mill J, Schalkwyk LC. A data-driven approach to preprocessing Illumina 450K methylation array data. *Bmc Genomics*. 2013;14:293.
44. Zheng SC, Webster AP, Dong D, Feber A, Graham DG, Sullivan R, et al. A novel cell-type deconvolution algorithm reveals substantial contamination by immune cells in saliva, buccal and cervix. *Epigenomics-uk*. 2018;10:925–40.
45. Katz DH, Robbins JM, Deng S, Tahir UA, Bick AG, Pampana A, et al. Proteomic profiling platforms head to head: Leveraging genetics and clinical traits to compare aptamer- and antibody-based methods. *Sci Adv*. 2022;8:eabm5164.
46. Haslam DE, Li J, Dillon ST, Gu X, Cao Y, Zeleznik OA, et al. Stability and reproducibility of proteomic profiles in epidemiological studies: comparing the Olink and SOMAscan platforms. *PROTEOMICS*. 2022;22:e2100170.

- 1 47. Raffield LM, Dang H, Pratte KA, Jacobson S, Gillenwater LA, Ampleford E, et al. Comparison of  
2 Proteomic Assessment Methods in Multiple Cohort Studies. *PROTEOMICS*. 2020;20:1900278.
- 3 48. Joshi A, Mayr M. In Aptamers They Trust: The caveats of the SOMAscan biomarker discovery  
4 platform from SomaLogic. *Circulation*. 2018;138:2482–5.
- 5 49. Hillary RF, Marioni RE. MethylDetectR: a software for methylation-based health profiling.  
6 *Wellcome Open Res*. 2021;5:283.
- 7 50. Preacher KJ. Extreme Groups Designs. *The Encyclopedia of Clinical Psychology*. 2015;1–4.
- 8 51. Fisher JE, Guha A, Heller W, Miller GA. Extreme-Groups Designs in Studies of Dimensional  
9 Phenomena: Advantages, Caveats, and Recommendations. *J Abnorm Psychol*. 2020;129:14–20.
- 10 52. Mckinnon K, Conole ELS, Vaher K, Binkowska J, Sullivan G, Hillary R, et al. The relationship  
11 between socioeconomic status, preterm birth and systemic inflammation using DNA methylation  
12 proxies. *OSF*. 2023;
- 13 53. Gao X, Becker LC, Becker DM, Starmer JD, Province MA. Avoiding the high Bonferroni penalty in  
14 genome-wide association studies. *Genet Epidemiol*. 2010;34:100–5.
- 15 54. Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, et al. The STRING  
16 database in 2023: protein–protein association networks and functional enrichment analyses for any  
17 sequenced genome of interest. *Nucleic Acids Res*. 2022;51:D638–46.
- 18 55. Kandasamy J, Roane C, Szalai A, Ambalavanan N. Serum eotaxin-1 is increased in extremely-  
19 low-birth-weight infants with bronchopulmonary dysplasia or death. *Pediatr Res*. 2015;78:498–504.
- 20 56. Almudares F, Hagan J, Chen X, Devaraj S, Moorthy B, Lingappan K. Growth and differentiation  
21 factor 15 (GDF15) levels predict adverse respiratory outcomes in premature neonates. *Pediatr*  
22 *Pulmonol*. 2023;58:271–8.
- 23 57. Al-Mudares F, Reddick S, Ren J, Venkatesh A, Zhao C, Lingappan K. Role of Growth  
24 Differentiation Factor 15 in Lung Disease and Senescence: Potential Role Across the Lifespan.  
25 *Frontiers Medicine*. 2020;7:594137.
- 26 58. Wan Y, Fu J. GDF15 as a key disease target and biomarker: linking chronic lung diseases and  
27 ageing. *Mol Cell Biochem*. 2023;1–14.
- 28 59. Zhang Y, Jiang W, Wang L, Lingappan K. Sex-specific differences in the modulation of Growth  
29 Differentiation Factor 15 (GDF15) by hyperoxia in vivo and in vitro: Role of Hif-1 $\alpha$ . *Toxicol Appl*  
30 *Pharm*. 2017;332:8–14.
- 31 60. Wang G, Wen B, Deng Z, Zhang Y, Kolesnichenko OA, Ustiyany V, et al. Endothelial progenitor  
32 cells stimulate neonatal lung angiogenesis through FOXF1-mediated activation of BMP9/ACVRL1  
33 signaling. *Nat Commun*. 2022;13:2080.
- 34 61. Collaco JM, McGrath-Morrow SA, Griffiths M, Chavez-Valdez R, Parkinson C, Zhu J, et al.  
35 Perinatal Inflammatory Biomarkers and Respiratory Disease in Preterm Infants. *J Pediatrics*.  
36 2022;246:34-39.e3.
- 37 62. Oak P, Hilgendorff A. The BPD trio? Interaction of dysregulated PDGF, VEGF, and TGF signaling  
38 in neonatal chronic lung disease. *Mol Cell Pediatrics*. 2017;4:11.
- 39 63. Course CW, Lewis PA, Kotecha SJ, Cousins M, Hart K, Watkins WJ, et al. Characterizing the  
40 urinary proteome of prematurity-associated lung disease in school-aged children. *Respir Res*.  
41 2023;24:191.



- 1 64. Bose CL, Dammann CEL, Laughon MM. Bronchopulmonary dysplasia and inflammatory  
2 biomarkers in the premature neonate. *Archives Dis Child - Fetal Neonatal Ed.* 2008;93:F455.
- 3 65. Zasada M, Suski M, Bokinić R, Szwarc-Duma M, Borszewska-Kornacka MK, Madej J, et al.  
4 Comparative two time-point proteome analysis of the plasma from preterm infants with and without  
5 bronchopulmonary dysplasia. *Ital J Pediatr.* 2019;45:112.
- 6 66. Lassus P, Heikkilä P, Andersson LC, Boguslawski K von, Andersson S. Lower concentration of  
7 pulmonary hepatocyte growth factor is associated with more severe lung disease in preterm infants. *J*  
8 *Pediatr.* 2003;143:199–202.
- 9 67. Wagner BD, Babinec AE, Carpenter C, Gonzalez S, O'Brien G, Rollock K, et al. Proteomic  
10 Profiles Associated with Early Echocardiogram Evidence of Pulmonary Vascular Disease in Preterm  
11 Infants. *Am J Respir Crit Care Med.* 2018;197:394–7.
- 12 68. Guasti L, Silvennoinen S, Bulstrode NW, Ferretti P, Sankilampi U, Dunkel L. Elevated FGF21  
13 Leads to Attenuated Postnatal Linear Growth in Preterm Infants Through GH Resistance in  
14 Chondrocytes. *J Clin Endocrinol Metab.* 2014;99:E2198–206.
- 15 69. Spencer R, Maksym K, Hecher K, Maršál K, Figueras F, Ambler G, et al. Ultrasound and  
16 biochemical predictors of pregnancy outcome at diagnosis of early-onset fetal growth restriction.  
17 *Medrxiv.* 2023;2023.01.27.23285087.
- 18 70. Qiu Q, Bell M, Lu X, Yan X, Rodger M, Walker M, et al. Significance of IGFBP-4 in the  
19 Development of Fetal Growth Restriction. *J Clin Endocrinol Metab.* 2012;97:E1429–39.
- 20 71. Voller SB, Chock S, Ernst LM, Su E, Liu X, Farrow KN, et al. Cord blood biomarkers of vascular  
21 endothelial growth (VEGF and sFlt-1) and postnatal growth: A preterm birth cohort study. *Early Hum*  
22 *Dev.* 2014;90:195–200.
- 23 72. Kim MJ, Romero R, Kim CJ, Tarca AL, Chhauy S, LaJeunesse C, et al. Villitis of Unknown  
24 Etiology Is Associated with a Distinct Pattern of Chemokine Up-Regulation in the Feto-Maternal and  
25 Placental Compartments: Implications for Conjoint Maternal Allograft Rejection and Maternal Anti-  
26 Fetal Graft-versus-Host Disease. *J Immunol.* 2009;182:3919–27.
- 27 73. Romero R, Chaemsathong P, Chaiyasit N, Docheva N, Dong Z, Kim CJ, et al. CXCL10 and IL-6:  
28 Markers of two different forms of intra-amniotic inflammation in preterm labor. *Am J Reprod Immunol.*  
29 2017;78:e12685.
- 30 74. Sellmer A, Henriksen TB, Palmfeldt J, Bech BH, Astono J, Bennike TB, et al. The Patent Ductus  
31 Arteriosus in Extremely Preterm Neonates Is More than a Hemodynamic Challenge: New Molecular  
32 Insights. *Biomol.* 2022;12:1179.
- 33 75. Olsson KW, Larsson A, Jonzon A, Sindelar R. Exploration of potential biochemical markers for  
34 persistence of patent ductus arteriosus in preterm infants at 22–27 weeks' gestation. *Pediatr Res.*  
35 2019;86:333–8.
- 36 76. Waleh N, Seidner S, McCurnin D, Giavedoni L, Hodara V, Goelz S, et al. Anatomic Closure of the  
37 Premature Patent Ductus Arteriosus: The Role of CD14+/CD163+ Mononuclear Cells and VEGF in  
38 Neointimal Mound Formation. *Pediatr Res.* 2011;70:332–8.
- 39 77. Xu C, Su X, Chen Y, Xu Y, Wang Z, Mo X. Proteomics analysis of plasma protein changes in  
40 patent ductus arteriosus patients. *Ital J Pediatr.* 2020;46:64.
- 41 78. Cheng Y, Zhu X, Linghu D, Xu Y, Liang J. Serum levels of cytokines in infants treated with  
42 conbercept for retinopathy of prematurity. *Sci Rep-uk.* 2020;10:12695.



- 1 79. Sato T, Kusaka S, Shimojo H, Fujikado T. Simultaneous Analyses of Vitreous Levels of 27  
2 Cytokines in Eyes with Retinopathy of Prematurity. *Ophthalmology*. 2009;116:2165–9.
- 3 80. Rathi S, Jalali S, Patnaik S, Shahulhameed S, Musada GR, Balakrishnan D, et al. Abnormal  
4 Complement Activation and Inflammation in the Pathogenesis of Retinopathy of Prematurity. *Front*  
5 *Immunol*. 2017;8:1868.
- 6 81. The Royal College of Ophthalmologists. Treating Retinopathy of Prematurity in the UK. The Royal  
7 College of Ophthalmologists; 2022.
- 8 82. Rivera JC, Holm M, Austeng D, Morken TS, Zhou T (Ellen), Beaudry-Richard A, et al. Retinopathy  
9 of prematurity: inflammation, choroidal degeneration, and novel promising therapeutic strategies. *J*  
10 *Neuroinflamm*. 2017;14:165.
- 11 83. Klerk DH, Plösch T, Verkaik-Schakel RN, Hulscher JBF, Kooi EMW, Bos AF. DNA Methylation of  
12 TLR4, VEGFA, and DEFA5 Is Associated With Necrotizing Enterocolitis in Preterm Infants. *Frontiers*  
13 *Pediatrics*. 2021;9:630817.
- 14 84. Olaloye OO, Liu P, Toothaker JM, McCourt BT, McCourt CC, Xiao J, et al. CD16+CD163+  
15 monocytes traffic to sites of inflammation during necrotizing enterocolitis in premature infants. *J Exp*  
16 *Med*. 2021;218:e20200344.
- 17 85. Pammi M, Suresh G. Enteral lactoferrin supplementation for prevention of sepsis and necrotizing  
18 enterocolitis in preterm infants. *Cochrane Db Syst Rev*. 2020;3:CD007137.
- 19 86. Satrom KM, Ennis K, Sweis BM, Matveeva TM, Chen J, Hanson L, et al. Neonatal hyperglycemia  
20 induces CXCL10/CXCR3 signaling and microglial activation and impairs long-term synaptogenesis in  
21 the hippocampus and alters behavior in rats. *J Neuroinflamm*. 2018;15:82.
- 22 87. Groselj-Grenc M, Ihan A, Derganc M. Neutrophil and Monocyte CD64 and CD163 Expression in  
23 Critically Ill Neonates and Children with Sepsis: Comparison of Fluorescence Intensities and  
24 Calculated Indexes. *Mediat Inflamm*. 2008;2008:202646.
- 25 88. Kingsmore SF, Kennedy N, Halliday HL, Velkinburgh JCV, Zhong S, Gabriel V, et al. Identification  
26 of Diagnostic Biomarkers for Infection in Premature Neonates. *Mol Cell Proteomics*. 2008;7:1863–75.
- 27 89. Zilow EP, Hauck W, Linderkamp O, Zilow G. Alternative Pathway Activation of the Complement  
28 System in Preterm Infants with Early Onset Infection. *Pediatr Res*. 1997;41:334–9.
- 29 90. Mohammed A, Okwor I, Shan L, Onyilagha C, Uzonna JE, Gounni AS. Semaphorin 3E Regulates  
30 the Response of Macrophages to Lipopolysaccharide-Induced Systemic Inflammation. *J Immunol*.  
31 2020;204:128–36.
- 32 91. Leviton A, Allred EN, Fichorova RN, O'Shea TM, Fordham LA, Kuban KKC, et al. Circulating  
33 biomarkers in extremely preterm infants associated with ultrasound indicators of brain damage. *Eur J*  
34 *Paediatr Neuro*. 2018;22:440–50.
- 35 92. Schultz SJ, Aly H, Hasanen BM, Khashaba MT, Lear SC, Bendon RW, et al. Complement  
36 component 9 activation, consumption, and neuronal deposition in the post-hypoxic–ischemic central  
37 nervous system of human newborn infants. *Neurosci Lett*. 2005;378:1–6.
- 38 93. Morales DM, Townsend RR, Malone JP, Ewersmann CA, Macy EM, Inder TE, et al. Alterations in  
39 Protein Regulators of Neurodevelopment in the Cerebrospinal Fluid of Infants with Posthemorrhagic  
40 Hydrocephalus of Prematurity. *Mol Cell Proteom*. 2012;11:M111.011973.
- 41 94. Ochoa TJ, Sizonenko SV. Lactoferrin and prematurity: a promising milk protein? *Biochem Cell*  
42 *Biol*. 2017;95:22–30.

- 1 95. Che X, Hornig M, Bresnahan M, Stoltenberg C, Magnus P, Surén P, et al. Maternal mid-  
2 gestational and child cord blood immune signatures are strongly associated with offspring risk of ASD.  
3 *Mol Psychiatry*. 2022;27:1527–41.
- 4 96. Leviton A, Joseph RM, Fichorova RN, Allred EN, Taylor HG, O'Shea TM, et al. Executive  
5 Dysfunction Early Postnatal Biomarkers among Children Born Extremely Preterm. *J Neuroimmune*  
6 *Pharm*. 2019;14:188–99.
- 7 97. Allred EN, Dammann O, Fichorova RN, Hooper SR, Hunter SJ, Joseph RM, et al. Systemic  
8 Inflammation during the First Postnatal Month and the Risk of Attention Deficit Hyperactivity Disorder  
9 Characteristics among 10 year-old Children Born Extremely Preterm. *J Neuroimmune Pharm*.  
10 2017;12:531–43.
- 11 98. Aly H, Khashaba M, Nada A, Hasanen B, McCarter R, Schultz S, et al. The Role of Complement  
12 in Neurodevelopmental Impairment following Neonatal Hypoxic-Ischemic Encephalopathy. *Am J*  
13 *Perinatol*. 2009;26:659–65.
- 14 99. Limbrick DD, Morales DM, Shannon CN, Wellons JC, Kulkarni AV, Alvey JS, et al. Cerebrospinal  
15 fluid NCAM-1 concentration is associated with neurodevelopmental outcome in post-hemorrhagic  
16 hydrocephalus of prematurity. *PLoS ONE*. 2021;16:e0247749.
- 17 100. Voegelé AF, Jerković L, Wellenzohn B, Eller P, Kronenberg F, Liedl KR, et al. Characterization  
18 of the Vitamin E-Binding Properties of Human Plasma Afamin. *Biochemistry*. 2002;41:14532–8.
- 19 101. Paetau S, Rolova T, Ning L, Gahmberg CG. Neuronal ICAM-5 Inhibits Microglia Adhesion and  
20 Phagocytosis and Promotes an Anti-inflammatory Response in LPS Stimulated Microglia. *Front Mol*  
21 *Neurosci*. 2017;10:431.
- 22 102. Skeel A, Yoshimura T, Showalter SD, Tanaka S, Appella E, Leonard EJ. Macrophage stimulating  
23 protein: purification, partial amino acid sequence, and cellular activity. *J Exp Med*. 1991;173:1227–34.
- 24 103. Kronenberg F, Kollerits B, Kiechl S, Lamina C, Kedenko L, Meisinger C, et al. Plasma  
25 Concentrations of Afamin Are Associated With the Prevalence and Development of Metabolic  
26 Syndrome. *Circ: Cardiovasc Genet*. 2018;7:822–9.
- 27 104. Montez JK, Bromberger JT, Harlow SD, Kravitz HM, Matthews KA. Life-Course Socioeconomic  
28 Status and Metabolic Syndrome Among Midlife Women. *J Gerontol: Ser B*. 2016;71:1097–107.
- 29 105. Castagné R, Delpierre C, Kelly-Irving M, Campanella G, Guida F, Krogh V, et al. A life course  
30 approach to explore the biological embedding of socioeconomic position and social mobility through  
31 circulating inflammatory markers. *Sci Rep-uk*. 2016;6:25170.
- 32 106. Leviton A, Allred EN, Dammann O, Joseph RM, Fichorova RN, O'Shea TM, et al. Socioeconomic  
33 status and early blood concentrations of inflammation-related and neurotrophic proteins among  
34 extremely preterm newborns. *Plos One*. 2019;14:e0214154.
- 35 107. Suarez A, Lahti J, Lahti-Pulkkinen M, Girchenko P, Czamara D, Arloth J, et al. A polyepigenetic  
36 glucocorticoid exposure score at birth and childhood mental and behavioral disorders. *Neurobiology*  
37 *Stress*. 2020;13:100275.
- 38 108. Blostein FA, Fisher J, Dou J, Schnepel L, Ware EB, Notterman DA, et al. Polymethylation scores  
39 for prenatal maternal smoke exposure persist until age 15 and are detected in saliva in the Fragile  
40 Families and Child Wellbeing cohort. *Epigenetics*. 2022;17:2223–40.

41

Component



