Effect of nutrition assessment, counselling and support integration on mother- infant nutritional status, practices and health in Tororo and Butaleja districts, Uganda: A comparative non equivalent quasi experimental study

5					
6	Samalie Namukose ¹ ,	Gakenia Wamuy	u Maina², Suzanne N Ki	wanuka ¹ Fredrick E	dward Makumbi ³

- 8 1 Department of Health Policy Planning and Management, School of Public Health, College
- 9 of Health Sciences, Makerere University, Kampala Uganda
- 10 2 Department of Community Health and Behavioural Sciences, School of Public Health,
- 11 College of Health Sciences, Makerere University, Kampala Uganda
- 12 3 Department of Epidemiology and Biostatistics, School of Public Health, College of Health
- 13 Sciences, Makerere University, Kampala Uganda
- 14 * <u>snamukoseb@gmail.com</u>
- 15

7

16 Abstract

Background: Malnutrition remains a health challenge for women aged 15 to 49 years and their
infants. While Nutrition Assessment Counselling and Support (NACS) is considered a
promising strategy, evidence on its effectiveness remains scanty. This study assessed the effect
of comprehensive NACS package on the mother-infant practices, health and nutrition outcomes
in two districts in Eastern Uganda.

Methods; Andrewick and Antiparticipation and Routine NACS (Butaleja). Pregnant mothers
 were enrolled spanning various trimesters and followed through the antenatal periods and post-

delivery for health and nutrition status. Infants were followed for feeding practices, health and
nutritional status at birth and weeks 6, 10, 14 and at month 6, 9 and 12 post-delivery.

Propensity score matching ensured study group comparability. The NACS effect was estimated
by nearest neighbour matching and the logistic regression methods. Statistical analysis utilised
STATA version 15 and R version 4.1.1.

Results: A total of 666/784 (85%) with complete data and were analysed (routine: 412,
comprehensive: 254). Both groups were comparable by mothers' age, MUAC, prior antenatal
visits, meal frequency, micronutrient supplementation and instances of maternal headache,
depression and diarrhoea. However, differences existed in gestation age, income, family size,
education and other living conditions.

Comprehensive NACS infants exhibited higher infant birth weights, weight- for- age z-scores at the 3^{rd} -6th visits (p<0.001), length- for- age z scores at the 4th -7th visits (p<0.001) and weight-for-length z-scores at the 3^{rd} - 5th (p<=0.001) visits. Despite fewer episodes of diarrhoea and fever, upper respiration infections were higher.

39 Conclusion: The comprehensive NACS demonstrated improved mother-infant nutritional and
40 other health outcomes suggesting the need for integrated and holistic care for better maternal,
41 infant and child health.

42 Keywords: Effect, nutrition assessment counselling and support, practices, health and nutrition
43 outcomes, mothers, infants.

44

46 Introduction

47

Maternal and infant malnutrition is a significant global health concern with significant 48 implications on the overall health and well-being of both the mothers and their infants. The 49 Global Nutrition report of 2022 [1] indicated that 29.9% of women of reproductive age suffer 50 from anaemia while 9.1% of all women were underweight. The prevalence of low birth weights 51 was 14.6% among the newborns while 22%, 6.7% and 5.7% of children under 5 years were 52 stunted, wasted and overweight respectively. In the same report, Sub-saharan Africa was noted 53 54 to contribute to the highest burden of malnutrition with 32.6% of children under 5 year stunted, 5.2% wasted and 4% overweight while anaemia among the women of reproductive age was 55 31.9%. According to the Uganda Demographic Health Survey (UDHS) of 2016 [2], the 56 57 prevalence of stunting among children under 5 years was 29% while underweight and wasting 58 was 11% and 4% respectively. Additionally, aneamia affects 32% of the women of reproductive age. These surveys and reports indicate the persistent challenge of malnutrition 59 among the women of reproductive health and children calling for urgent need for intervention 60 and improvement. 61

Maternal nutrition is vital for the health and well being of both the mother and her developing infant. Maternal interventions aimed at improving nutrition practices before pregnancy, during pregnancy and lactation have been extensively studied for their potential to enhance maternal and infant health down the line [3–7]. Well nourished and healthy mothers are more likely to give birth to health babies, experience a healthy pregnancy and are less likely to experience life-threatening complications during pregnancy [8,9].

68

Several studies have demonstrated the positive impact of maternal interventions on the nutrition practices and growth of infants particularly when implemented in a multi-sectoral approach. For instance, implementation of a comprehensive range of interventions such as; breastfeeding promotion, education and counselling, maternal mental health, women empowerment, family planning, water, hygiene and sanitation, agricultural interventions has

shown promising results in reducing stunting rates [10]. Notable studies by Olutayo et al [11], 74 Nadia et al [12], Bhutta et al [13] emphasize the importance of these holistic interventions in 75 reducing stunting. However, it is important to consider the perspective put forth by 76 77 USAID/Advancing Nutrition [14] which argues against using stunting as a primary indicator of success of short term or single interventions at individual. Instead, stunting should be 78 interpreted as a reflection of the population's well-being. This view suggests a more 79 80 comprehensive assessment of interventions success focusing of multiple short and long term causal factors to malnutrition instead of looking at the immediate outcomes. 81

Additionally, numerous studies have shown that Nutrition Counselling and education during 82 pregnancy significantly improve maternal-infant nutrition practices as well as the overall health 83 and nutritional status of both mothers and infants. Dearden et al [15] demonstrated a positive 84 effect of nutrition counselling and education on maternal meal frequency and diet 85 diversification. These findings align with a quasi- experimental study conducted by Kaleem et 86 al [16] which indicated that Antenatal Counselling improved the maternal dietary practices 87 88 and nutritional status. Similarly, a study by Perez-Escamilla et al [17] revealed a positive effect of maternal counselling on maternal and infant health and nutrition outcomes including 89 birth weights and prevention of pre-term births. In contrast, Ghosh-Jereth et al [18] found that 90 91 the maternal dietary intake remained low and anaemia rates were high despite targeted antenatal care including counselling at every visit. The authors attributed this lack of 92 improvement to the poor quality of counselling, a finding also echoed by Nsiah-Asamoah et al 93 [19] in their study on nutrition counselling interactions between the health workers and 94 caregivers. While some studies showed positive effect of maternal nutrition counselling on the 95 maternal-infant health and nutrition outcomes, contrasting findings indicate that the quality of 96 97 the counselling and how it is delivered can have negative impact on these outcomes. More research is needed to bridge this gap and provide a clearer understanding on the effect of 98 delivery of a comprehensive package including counselling on the mother-infant health and 99 100 nutrition outcomes.

101 Therefore, providing high-quality health services, including preventive health services, antenatal, maternity and postnatal services as well as early diagnosis and treatment of medical 102 conditions such as anaemia is crucial for improving women's health. The World Health 103 104 Organisation (WHO) and the Ministry of Health, Uganda recommends a comprehensive package of nutrition interventions to pregnant women for a positive outcome, including 105 counselling on healthy eating and physical activity, guidance on infant and young child feeding, 106 107 nutrition education on energy and protein intake, and daily iron and folic acid supplements. The package also includes energy and protein dietary supplements and high-protein 108 supplements for the undernourished populations [20,21]. 109

Even before the release of the WHO guidelines in 2020, the Ministry of Health in Uganda had 110 been implementing NACS initiative, aiming to integrate nutrition into the health system and 111 112 consequently improving the health and nutrition practices and outcomes of the beneficiaries. The NACS intervention package was tailored to the specific nutrition needs of the clients and 113 was in line with WHO's recommendations on maternal nutrition care. Support was provided 114 to mothers and their children, covering aspects such as optimal maternal nutrition, diversified 115 diets, iron/folic acid supplementation, iodated salt consumption, deworming, malaria 116 prevention, and provision of antenatal care package and encouragement to attend all the 8 117 visits. Breastfeeding education emphasized early initiation, exclusive breastfeeding for 6 118 months, and extended breastfeeding. Mothers of older infants received guidance on 119 120 complementary feeding practices. Caregivers of sick children were advised on continued breastfeeding. The community system offered ongoing health and nutrition care, including 121 livelihood and economic support to improve the health and nutrition outcomes. The 122 123 malnourished mothers received therapeutic foods [22].

While existing literature has assessed the impact of vertical maternal interventions on the health and nutritional status of mothers and infants, there is limited body of research on the effect of broad integrated interventions, such as NACS on the health and nutrition outcomes of beneficiaries. This study therefore, sought to assess the effect of comprehensive NACS package on the health and nutrition practices and status of mothers and their infants in Tororo

and Butaleja districts in Eastern Uganda. We tested the hypotheses that there was no difference in the maternal-infant health, nutrition practices and outcomes between the facilities which integrated comprehensive NACS, verses those with routine NACS. The findings of this study contribute to the growing body of evidence on the effectiveness of broad integrated interventions on the health and nutrition outcomes of the beneficiaries and provide insights and recommendations for scaling up the NACS approach.

Pathways on the effect of NACS integration in the health system on maternal and infant health, nutrition practices, and outcomes

Integration of comprehensive NACS into the health system will result into an integrated 137 nutrition service delivery system which aims to foster a productive interaction between the 138 139 service providers and mothers. Based on the health belief model, service providers were 140 expected be motivated to impart knowledge and skills to mothers, enabling them take charge of their own health and nutrition. Based on the health belief model, these empowered mothers 141 would then embrace optimal nutrition practices resulting in enhanced health and nutrition well-142 being. Consequently, this positive change would improve the health and nutrition outcomes of 143 their infants, as illustrated in S1 Fig 144

145 Insert S1 Fig

146

147 Material and Methods

148 Study Design

149 The study used a comparative non-equivalent quasi-experimental design with two groups;

150 comprehensive NACS integration compared to routine NACS integration.

151

152 Study Setting and population

The study involved pregnant and lactating mothers, along with their respective infants. The 153 two hospitals selected for the study were Tororo Hospital as the comprehensive NACS and 154 Busolwe Hospital as the routine NACS. The hospitals were similar by level of facility, 155 ownership, funding, staffing norms, services provided and client load. Pregnant mothers in 156 various trimesters were enrolled and their health and nutrition status monitored at the antenatal 157 visits and post-delivery. Only women accessing antenatal care and residing in Tororo and 158 159 Butaleja were included. During post-delivery, infants were monitored for their feeding 160 practices, health and nutrition status till 12 months.

Mothers and infants were followed through the scheduled visits at their respective health service points, which included, antenatal, labour suite/maternity, postnatal, children wards, young child and ART clinics.

164

165 Comprehensive NACS versus routine service delivery

The comprehensive NACS package targeted both the health workers and mothers with their 166 infants. The support to the health workers included: five-day training in NACS and Health 167 Management Information System (HMIS) for nutrition in-service courses for staff at key health 168 contact points, such as antenatal clinics, maternity, postnatal clinics, young child and HIV 169 clinics; provision of anthropometric equipment, policy guidelines, job aides, information, 170 education and communication materials for mothers; mentorships/support supervision of 171 172 health staff to ensure quality service delivery; monitoring and reporting of nutrition interventions; employing quality improvement support to address gaps in NACS 173 implementation; linking study subjects to community support structures for continuous 174 175 nutrition care and support; and collaborating with key stakeholders and the district health management team to establish supervisory and support mechanisms for the intervention. 176

To the mothers and their infants, the package included: nutrition assessment and categorization of the nutritional status; health and nutrition education on a diversified diet, recommended antenatal clinic visits, iron/folic acid supplementation, water hygiene, and sanitation; maternal-

180 infant nutrition counselling; provision of therapeutic feeds to identified malnourished cases;

181 active follow-up of mother-baby pairs to ensure they receive the necessary nutrition services.

182

183 Routine service delivery

In the routine NACS setting, some elements of NACS were integrated into the regular health 184 care services provided such as growth monitoring and promotion for children, iron/folic acid 185 supplementation. To ensure comparability, staff at both study settings were trained in NACS 186 and HMIS for nutrition. They were provided with information, education and communication 187 materials to enhance their capacity in nutrition education and counselling. Subsequently, the 188 staff carried on with their services as per usual. The nutrition counselling placed a strong 189 emphasis on promoting the consumption of locally available foods for the management of 190 191 malnutrition. [23]

We determined the level of exposure to comprehensive verses routine NACS by closely supervising the data collection process and enhancing documentation of both the services rendered and the frequency with which the respondents accessed these services.

195 Sampling

The study employed purposive sampling approach, enrolling subjects who had given their consent on a continuous basis until the desired sample size was attained. In both study settings, the antenatal care clinic served as the entry point and the ANC register as the sampling frame. The enrolment of the study participants took 8 months from starting from 23rd October 2018 to 25th May, 2019. The mothers were followed up till they gave birth, and the mother-infant pairs followed up for 12 months. Data was collected from 23rd October, 2018 to 25th July 2021.

203

204 Sample size calculation

205 The sample size was calculated using the formular by V. Kasiulevicius et al [24] based on

206 infant underweight as an outcome variable.

 $n = \frac{\left| z_{\alpha} \sqrt{(1+\frac{1}{m}) \bar{p}(1-\bar{p})} + z_{\beta} \sqrt{\frac{po(1-p0)}{m} + p1(1-p1)} \right|^2}{(p0-p1)^2}$

208

Where
$$\overline{p} = \frac{p1 + mp0}{m+1}$$

209 Where,

- 210 P_0 was the probability of underweight infants in the control group -0.113.
- P₁was the probability of underweight infants in the intervention group -0.07
- P₀ was based on the Uganda Demographic Health Survey 2011 burden of malnutrition in the
- eastern region while P1 was an estimated reduction in underweight with the intervention.
- 214 If α (alpha) = 0.05 then $z_{\alpha} = 1.96$
- 215 If β (beta) = 0.80, then $z_{\beta} = 0.845$
- 216 m was the number of control subjects per experimental subject = 2
- 217 $\overline{p} = 0.0987$
- n = 652 with the inclusion of 20% loss to follow up of mother-baby pairs. A sample size of 652
- 219 (217 in the intervention group and 435 in the control group) was estimated to detect a 4.3%
- reduction in the underweight infants at 80% power and 5% level of significance.
- 221

222 Data collection

223 We pre-tested the data collection tools among the mothers and feedback was used to finalize

the tools. The tool was designed in excel to facilitate the tracking of the mother- baby variables

for their scheduled visits.

Our research assistants underwent training on the data collection tools and data capture methods at a minimum of four points: recruitment/baseline, antenatal clinic visits, delivery and postnatal care clinic, and immunization scheduled visits. We encouraged mothers to deliver at the health facility where they were provided with a package of both routine and comprehensive

230 package of services. To ensure data quality, we conducted regular supervision and spot checks.

At baseline/recruitment, we collected data on various aspects, including socio-economic and demographic characteristics, maternal health and nutrition practices, and maternal nutritional status. Throughout each antenatal visits, we monitored the mothers' anthropometric data, health status and nutrition practices. Following delivery, we collected data on infant's anthropometric measurement such as birth weight, length, head circumference as well as details about their feeding practices. Subsequently, we continued to track the infants' anthropometric data, health and nutrition practices and status during the scheduled immunization visits.

Anthropometric data and feeding practices for both the mother and her infant were collected using standard procedures. Mother's weight was taken to the nearest 0.1gm using a digital Uniscale. The infant weight was measured to nearest 0.1 gm using the neonatal weighing scales at birth and thereafter a digital uniscale. Infant length was measured to nearest 0.1cm using an infantometer at birth and a height board for the subsequent visits.

We measured the head circumference and Mid Upper Arm Circumference (MUAC) of infants using specialized tapes, with measurements recorded to the nearest 0.1cm. MUAC was measured for infants above 6 months and mothers. Additionally, we conducted health assessment for mothers, including evaluation for illnesses such as headaches, depression, diarrhea, fever and cough. For infants, we assessed episodes of diarrhea, fever and Upper Respiratory infections.

In total, there were 15 scheduled appointments from the time of mother's enrolment until the baby made 12 months of age. Mothers were encouraged to continue attending health facilities for continuous health care as well as participating in the informative health and nutrition education sessions.

255

Data management

We used excel for data capture, STATA version 15 for data, cleaning and performing bivariate tests on all confounding background variables for both study settings. The variables included; weeks of gestation, age of the mothers, mothers' education, marital status, mothers' occupation,

260 mothers' income, spouses' income, spouses' education level, previous ANC visits, distance to 261 the health facility, type of transport used, total numbers of children, number of children alive, 262 number of family members, number of children under 5 years, fuel for cooking, water source 263 and faecal matter disposal. We cleaned data by synchronising the variable codes for the two 264 data sets, checked for missing data, and excluded the variables with insignificant data.

We characterised variables as continuous, binary, categorical and generated new variables. We checked the data set for normal distribution for the continuous variables. Descriptive analysis was conducted to compare mothers' background characteristics in the 2 study arms. Continuous variables were compared using a 2 sample t-test while the categorical variables were compared using the chi square test.

270 Data analysis

Because these groups were not randomly assigned and this being a quasi non- equivalent experimental study, we conducted propensity score matching to minimise potential imbalance and also create reasonable comparable groups, before assessment of the effectiveness of the NACS intervention.

By creating more comparable intervention and control groups, propensity score matching can result into a more precise estimates of intervention effects and reducing confounders. On the other hand, matching reduces the sample size, because not all individuals may find suitable matches resulting in a loss of statistical power and precision [25–27].

The propensity score matching process involved; defining the intervention (comprehensive NACS) verses control (routine NACS) groups, identification of the variables before administration of the intervention, estimating the propensity scores, checking the initial balance of the variables for both groups using mean differences, using the nearest-neighbor matching method to pair individuals in intervention and control group based on their propensity scores, assessing the quality of the matches, and thereafter estimating the effect of comprehensive NACS on maternal-infant practices, health and nutritional status [27].

Using the R software version 4.1.1, we estimated the NACS effect on maternal-infant nutrition 286 practices, health and nutritional status by comparing various methods such as nearest 287 neighbour, null and full matching methods. The nearest neighbour matching using the logistic 288 regression propensity score model provided the best balance compared to other matching 289 methods such as; full matching using a probit regression propensity score, nearest neighbour 290 matching using a probit regression propensity score, null probit, full matching using a logistic 291 regression propensity score as determined by the lower standardised mean difference statistics. 292 The enrolment and data analysis flow chart is illustrated in S2 Fig. 293

294 Insert S2 Fig

295 Ethical approval

This study was approved by: the Higher Degrees, Research, and Ethics Committee -296 297 Institutional Review Board at Makerere University School of Public Health (MaKSPH HDREC 24/01/2017), the Uganda National Council of Science and Technology (SS 4251), and 298 the Office of the President in Uganda (ADM/194/212/01). An official letter from the Ministry 299 of Health was written to the Tororo District Health Officer to seek for permission to conduct 300 the study. The Principal Investigator informed both the District Health Officer and the District 301 Resident Commissioner about the study plan before its execution. Participating mothers were 302 asked to sign informed consent form. The mothers who were unable to read and write provided 303 their informed consent using their thumbprint. 304

306 **Results**

307 A total of 784 mothers were enrolled in the study; 423 from the Routine NACS setting and 361 308 from the Comprehensive NACS setting. One hundred (118) mothers were excluded from the 309 analysis due to missing data while 666 mothers were considered in the final analysis, majority 310 from the routine (412) compared to comprehensive (254) NACS groups.

311 The mothers' characteristics at enrollment were compared in the two study arms and the results 312 are shown in the Tables 1 and 2. The findings indicated no significant difference between the 313 mother's age (p= 0.466), prior antenatal visits for this pregnancy (p=0.316), number of family 314 members (p=0.007) between the two groups at enrollment. However, there was a significant 315 difference in the weeks of gestation (p=0.023), mothers' income (p=0.000), spouses' income 316 (p=0.000), number of children (p=0.000), number of children alive (p=0.000), number of children 317 < 5years (p=0.001), distance to health facility (p<0.001), mothers' education (p<0.001), marital 318 status (p<0.001), mothers' occupation (p<0.001), spouses' education (p<0.001), type of transport 319 (p<0.001), cooking method (p<0.001), water source (p<0.001), and fecal matter disposal 320 (p<0.001).

321	Table 1. Mothers'	characteristics at	enrolment in the	routine verses	comprehensive	study a	rms for
322	continuous variabl	es					

Variable name (Comprehensive NACS=254, Routine NACS = 412)	t-test	p-value
Mothers' age	-0.730	0.466
Weeks of gestation	2.277	0.023
Mothers' income	-4.682	0.000
Spouse income	-4.997	0.000
Prio antennal visits for this pregnancy	1.003	0.316
Number of children	11.400	0.000
Number of children alive	5.725	0.000
Number of family members	2.669	0.007
Number of children less than 5 years	6.842	0.000

325 Table 2. Mothers' characteristics at enrolment in the routine verses comprehensive NACS study arms for categorical variables before propensity score matching 326

Variable nome	Douting NACS	Comprehensive NACS N=254	Chi-square Test (p-Value)
v ar lable frame	N=412	IN-234	
Distance to health facility	06 10/	(0, (0/	
<5 km	3.9%	39.4%	< 0.001
Mothers Education		/	
No Education	5.8%	1.2%	
Fillinal y Secondary	26.9%	44.9%	<0.001
Higher	4.4%	13.4%	0.001
Marital Status			
Married	94.4%	99.6%	
Not married	5.6%	0.4%	<0.001
Mothers occupation			
Formal	6.8%	18.1%	<0.001
informat	95.2%	81.9%	<0.001
Spouse's education	4.20/	1 20/	
No Education Primary	4.2%	1.2%	<0.001
Secondary	38.7%	55.1%	<0.001
Higher	7.8%	18.1%	
Type of transport			
Motorised	50.7%	92.1%	
Walking	49.3%	7.9%	<0.001
Cooking method	70.00/	50.00/	-0.001
Firewood	/9.9%	52.8% 46.1%	<0.001
Gas	0.2%	1.2%	
XX7 4			
Water source	0.2%	16.1%	
Borehole	96.8%	44.1%	< 0.001
Tap water	2.9%	39.8%	
Faecal matter Disposal			
Latrine	99.3%	89.0%	.0.001
loilet	0.7%	11.0%	< 0.001

327

329 Propensity score matching was used to create comparability between the study groups. The Null 330 model was used to check the initial imbalance in the two groups that the matching methods 331 eliminated step wise. Table 3 shows severe imbalances as reflected by the standard mean 332 differences computed by the R software. All values close to zero in the standard mean differences 333 reflected better matches while those away from zero reflected severe imbalances. The variable 334 'number of children' had the highest difference (-4.7263) indicating severe imbalance while the 335 variable 'nutrition status by MUAC' had the lowest difference (0.0019) indicating that mothers in 336 the both groups had comparable nutritional status, which conclusions resonate with exiting 337 literature [28].

338	Table	3:	Propensity	score	matching	null	model	for	checking	initial	imbalance	between	the
339	compr	ehe	ensive and re	outine	NACS stu	dy ar	ms						

Variables	Means Treated	Means Control	Mean difference
Distance	0.8513	0.1053	3.1896
Mother age			
15-24	0.4762	0.5140	-0.0758
25-49	0.5238	0.4860	0.0758
Mothers' education			
1. No Education	0.0119	0.0562	-0.4082
2. Primary	0.4444	0.6433	-0.4001
3. Secondary	0.4087	0.2556	0.3115
4. Higher	0.1349	0.0449	0.2634
Marital status			
1. Married	0.9960	0.9438	0.8305
2. Not Married	0.0040	0.0562	-0.8305
Mothers' occupation			
1. Formal	0.1825	0.0674	0.2980
2. Informal	0.8175	0.9326	-0.2980
Spouse education			
1. No Education	0.0119	0.0365	-0.2269
2. Primary	0.2579	0.5056	-0.5661
3. Secondary	0.5476	0.3820	0.3327
4. Higher	0.1825	0.0758	0.2762
Prior ANC visits	2.0397	2.1011	-0.0626
Distance to hospital			
1. Less than 5km	0.6111	0.9635	-0.7228
$2. \geq 5 \text{km}$	0.3889	0.0365	0.7228
Type of transport			
1. Walking	0.0794	0.5028	-1.5665
2. Motorized	0.9206	0.4972	1.5665
No. of children	1.0198	2.6236	-4.7263
No. of children alive	1.5437	2.4129	-0.5786
No of chn < 5years	0.7540	1.2809	-0.6142
Type of fuel used			
1. Firewood	0.5278	0.7978	-0.5408
2. Charcoal	0.4603	0.2022	0.5178
3. Gas	0.0119	0.0000	0.1098

Water source			
1. Well	0.1627	0.0028	0.4332
2. Borehole	0.4405	0.9691	-1.0648
3. Tap Water	0.3968	0.0281	0.7537
Faecal matter disposal			
1. Latrine	0.8889	0.9916	-0.3267
2. Toilet	0.1111	0.0084	0.3267
Mother wtg (kg) 1 st visit	64.2496	60.7542	0.3124
Nutritional status by MUAC			
1. Normal (Green)	0.9722	0.9719	0.0019
2 Malnourished (Yellow/Red)	0.0278	0.0281	0.0019
No. of meals	3 0992	2 8792	0 2940
Iron folic acid supplement	0.0772	2.0772	
No	0.0238	0.0028	0 1377
Vag	0.0238	0.0028	0.1377
	0.9702	0.9972	-0.1377
History of headache	1 0000	0.0024	0.6465
No	1.0000	0.8034	0.6465
Yes	0.0000	0.1966	-0.6465
History of depression			
No	1.0000	0.9803	0.1851
Yes	0.0000	0.0197	-0.1851
History of diarrhoea			
No	1.0000	0.9888	0.1393
Yes	0.0000	0.0112	0.1393
History of fever			
No	0.9960	0.9298	1.0539
Yes	0.0040	0.0702	-1.0539
History of cough			
No	1 0000	0 9522	0 2927
Yes	0.0000	0.0478	-0 2927
1.00	0.0000	0.0170	0.2/2/

340

341 Effect NACS integration on the mothers-infant health, nutrition practices and

342 status

343 The study assessed the effect of NACS integration on maternal-infant nutrition practices as well 344 as its effects on health and nutritional status. This assessment employed the nearest neighbour 345 matching method along with a logistic regression propensity score model and the results are shown 346 in Table 4.

347

349 Table 4. The effect of NACS integration on the mother-infant nutrition practices, health and 350 nutrition status using nearest neighbour matching logistic regression propensity score model

Variable name	Contrast		Estimate	SE	P value	CI
	1-(Comp NACS)	0-(Routine NACS)				
Mother variables	N=252	N=	252			
Meal frequency at 2 nd visit	1	0	0.008	0.070	0.911	-0.13, -0.146
Diet Diversity Score						
2 nd visit 3 rd visit	1	0	-3.110	0.263	<0.001 <0.001	-3.630, - 2.600
4 th visit	1	0	-2 690	0.149	<0.001	2 990 -2 400
Iron/ folic acid supplementation	1	0	-0.029	0.010	0.006	-0.049 -0.008
3 rd visit	1	0	-0.029	0.010	0.000	-0.043 , -0.005
4 th visit	1	0	-0.035	0.015	0.017	-0.064 , -0.006
Weight at						
2 nd visit	1	0	1.040	0.272	< 0.001	0.507 -1.570
3 rd visit	1	0	2.690	0.422	< 0.001	1.860 - 3.520
4 th visit	1	0	5.860	1.990	0.003	1.950 - 9.760
Nut.status by (MUAC)	1	0	0.039	0.019	0.022	0.002 0.072
3 rd visit	1	0	0.038	0.018	0.032	0.008 -0.044
4 th visit	1	0	0.025	0.015	0.091	0.004- 0.054
History of headache	1	0	0.020	0.000	0.020	0.002 0.027
2 rd visit	1	0	0.020	0.009	<0.020	0.003 - 0.037
4 th visit	1	0	0.078	0.022	< 0.001	0.034 - 0.121
History of depression						
2 nd visit	1	0	0.013	0.007	0.070	-0.001, -0.026
3 rd VISIt	1	0	0.008	0.005	0.160	-0.003, -0.018
Diarrnoea at 3 rd visit	1	0	0.018	0.008	0.030	0.002 -0.034
History of fever	1	0	0.020	0.000	0.010	0.002 0.027
2 rd VISIt	1	0	0.020	0.009	0.019	0.003 - 0.037
3 rd VISIt	1	0	0.049	0.014	< 0.001	0.023 - 0.076
4 ^m visit	1	0	0.030	0.013	0.021	0.005 - 0.055
Infant variables						
Infant birth weight	1	0	0.191	0.100	0.056	-0.005, -0.387
Infant weight at						
3 rd visit	1	0	1.050	0.111	< 0.001	0.835 - 1.270
4 th Visit	1	0	1.130	0.106	< 0.001	0.923 - 1.340
5 th visit	1	0	1.930	0.107	< 0.001	1.720 - 2.140

Infant head cirm	1	0	2 000	0.040	-0.001	2 410 4 200
3 rd v1s1t 4 th visit	1	0	3.900	0.248	<0.001 <0.001	3.410 - 4.380
5 th visit	1	0	0.020	0.240	0.001	0.003 0.037
J'' visit	1	0	0.020	0.009	0.019	0.003 -0.037
within Age 2-scores						
3 rd visit	1	0	1.800	0.233	< 0.001	1.350-2.260
4 th visit	l	0	1.700	0.208	< 0.001	1.290 -2.110
5 th visit	1	0	2.730	0.202	< 0.001	2.330 - 3.130
6 th visit	1	0	0.265	0.159	0.096	-0.047, -0.577
Length for Age Z- scores						
4 th visit	1	0	0.634	0.246	0.010	0.151 - 1.120
5 th Visit	1	0	0.761	0.222	< 0.001	0.326 - 1.200
6 th visit	1	0	3.990	0.281	< 0.001	3.440 - 4.540
7 th visit	1	0	4.630	1.140	< 0.001	2.470 - 6.780
Wt. for Length Z- scores at						
3 rd visit	1	0	6.250	0.475	< 0.001	5.320 - 7.180
4 th visit	1	0	1.460	0.356	< 0.001	0.763 – 2.160
5 th visit	1	0	2.770	0.342	< 0.001	2.100 - 3.440
History of Upper Respiratory Infection	1	0	0.043	0.013	<0.001	0.018 - 0.069
5th Visit						
6th visit	1	0	0.155	0.023	< 0.001	0.110 - 0.199
7th visit	1	0	0.294	0.029	< 0.001	0.237 - 0.350
8th visit	1	0	0.315	0.029	< 0.001	0.258 - 0.372
History of infant diarrhoea 5th visit	1	0	-0.168	0.080	0.036	-0.324, -0.011
6th visit	1	0	-0.146	0.080	0.067	-0.302- 0.010
7th visit	1	0	-0.033	0.049	0.506	-0.130- 0.064
8th visit	1	0	-0.126	0.067	0.061	-0.130- 0.064
History of infant fever 1st visit	1	0	-0.054	0.030	0.072	-0.113- 0.005
3rd visit	1	0	-0.005	0.012	0.694	-0.029- 0.019
4th visit	1	0	-0.409	0.086	< 0.001	-0.576, -0.241
5th Visit	1	0	-0.111	0.076	0.145	-0.260- 0.038
6th visit	1	0	-0.023	0.062	0.712	-0.144 - 0.099
7th visit	1	0	0.028	0.050	0.569	-0.070 - 0.126
8th visit	1	0	-0.027	0.060	0.657	-0.143- 0.090

352

353

Mothers in both groups were similar in terms of; meal frequency (p=0.911), iron/folic acid 354 supplementation at the 2nd -4th visits ($\beta \le -0.035$), maternal nutritional status by MUAC at the 355 $2^{nd} - 4^{th}$ visits ($\beta \le 0.038$). Additionally there was no significant difference in maternal 356

instances of; headache at the 2nd -4th visits ($\beta \le 0.078$), depression at the 2nd -3rd visit ($\beta \le 0.013$) and diarrhoea at the 2nd -4th ($\beta \le 0.049$) visits.

Whereas mothers in the routine NACS group had a significantly higher diversity score at the $2^{nd} - 4^{th}$ visits (p<0.001), the comprehensive group had higher weights at the $2^{nd} - 4^{th}$ (p<=0.003) visits. The difference in weights increased with number of visits right from the time of mothers' enrolment.

Compared to routine, infants born to mothers in the comprehensive group had a significantly higher; birth weights at 10% level of significance (p=0.056, CI -0.005 – 0.387), weight-forage at the 3^{rd} -6th visits (p<0.001) with 20% reduction in underweight on average per visit. Furthermore, their length-for-age was significantly higher at the 4th -7th visits (p<0.001). The difference widened with the increasing number of visits. Similarly, the weight-for-length of the comprehensive NACS group was significantly much higher at the 3rd -5th visits (p<0.001). The difference remained constant throughout the subsequent visits.

Unlike the routine group, infants in the comprehensive NACS group had significantly higher episodes of upper respiratory infections at the 5th -8th (p<0.001, β <=0.315). On the other hand, the routine NACS infants experienced a significantly higher episodes of; diarrhoea at the 5th -8th visits (p<=0.061) and fever at the 1st (p=0.072, β =-0.054) and 4th visits (p<0.001, β =-0.409).

374

375

376

378 **Discussion**

The study aimed to assess the effect of NACS integration on the maternal-infant nutrition practices, health and nutritional status. The findings provide insights into the potential benefits of nutrition integration on the health system on the wellbeing of the mothers and their infants.

382 The key findings in light of existing evidence and their implication are discussed.

The study found no significant difference in meal frequency among the mothers in both study 383 groups, suggesting similar dietary habits. Compared to the comprehensive group, mothers in 384 the routine NACS setting had a significantly higher diversity scores on all visits. This disparity 385 may be attributed to the close proximity to the rural settings offering more natural and diverse 386 food choices. Maternal nutritional status by MUAC estimates exhibited no significant 387 differences across the visits in the two study settings. In contrast to the routine group, mothers 388 in the comprehensive displayed significantly higher weights at the 2nd-4th visits. This implied 389 that integration of comprehensive NACS had positive progressive impact on maternal weight 390 gain from the time of enrolment. The findings suggest potential program implication and 391 highlight the need to consider environmental context when implementing nutrition programs. 392 Future nutrition interventions should therefore be tailored to the specific needs and context of 393 the target population. Furthermore, the study re-enforces, existing body of evidence indicating 394 395 that maternal focused interventions particularly those with a multi-sectoral nature contribute to improved maternal diet diversity, micronutrient intake and overall nutritional status [29-31]. 396

Additionally, there were minimal difference in iron/folic acid supplementation between the two 397 groups at the 2nd - 4th visits implying consistent adherence to the Ministry of Health guidance 398 on routine iron/folic acid supplementation among pregnant mothers in both settings. However, 399 it is worth noting that the effect of iron/folic acid supplementation on haemoglobin levels in 400 both settings could not be assessed in both settings due to lack of equipment and supplies. 401 Studies by Michael Habtu et al [32], Sunita Taneja et al [7], Melesse Kuma et al [33] revealed 402 elevated haemoglobin levels among women in the intervention group, findings that our study 403 was unable to replicate due to the constraints related to equipment and supplies. 404

The estimates showed no difference between the two settings for maternal episodes of headache, depression and diarrhoea across the various visits. This implies that these health concerns are common and not influenced by the study settings. These need to be addressed in both setting for the well being of mothers.

Our investigation into the nutritional status of the infants revealed that integration of comprehensive NACS increased infant birth weights, reduced instances of underweight, stunted and wasted infants. This implies that nutrition integration had a potential benefit of on the foetal and infant growth and development. Our findings concur with; Veeena et al [34] M Barker et al [8], Von Salmuth et al [10], Olutayo et al [11], Micheal Habtu etal [35] on the effectiveness of a holistic approach to improving the nutritional status of children.

Routine NACS infants experienced significantly more episodes of diarrhea and fever at the various visits than the comprehensive NACS group. The findings concur with Gonzalenz-Fernandez et al [36] in their study in which implementation of the multisectoral approach lowered the risk of diarrhoea and respiratory infections. This implies that the health facilities in the routine NACS settings did not comprehensively address these health concerns hence the need for more interventions for better health and nutrition outcomes.

421 One of the strengths of this study lies in its comparison of two separate groups; routine versus comprehensive and its close monitoring of the practices and outcomes of the study participants. 422 This approach bolstered the study's findings, providing a clear and robust insight into the 423 effectiveness of the integrated intervention package. Moreover the study places emphasis on 424 favourable outcomes of comprehensive NACS highlighting the potential benefits of such 425 comprehensive interventions, which findings are also consistent with the existing literature. On 426 the other hand, the study lacked the ability to assess the impact of iron/folic acid 427 supplementation on haemoglobin levels due to lack of equipment and supplies, which is a 428 limitation in understanding the complete maternal health and nutrition outcomes. 429

431 Conclusions

The findings add to the existing body of evidence supporting improved maternal-infant health and nutrition practices and status with integrated nutrition services. While meal frequency, iron/folic acid supplementation were similar in the both groups, integration of comprehensive NACS intervention improved; maternal weights, infant birth weights, infant growth in light of weight-for-age, length-for-age and weight-for-length. This emphasises the potential benefits of integrated nutrition interventions in promoting the overall being of the mothers and their infants.

439 **Recommendations**

Based on the above findings, the Ministry of Health should consider: investing in acquiring the necessary equipment and supplies to assess the impact of iron/folic acid supplementation on haemoglobin levels for comprehensive evaluation of the women; scaling up integration of the comprehensive NACS in the health system as it has positive effect on the maternal-infant nutrition practices, health and nutrition outcomes; investing in digitization to ease monitoring and tracking trends in the health and nutrition status of the mother-infant pairs.

Future research can focus on implementation and effectiveness of digitization in monitoring
and tracking of the maternal-infant health and nutritional status in an integrated health system.
Secondly, it will be important to investigate the experiences of the women and caregivers
receiving these services.

451 **Supporting information**

- 452 S1 File Dataset for Tororo and Butaleja districts
- 453 S1 Fig. Pathways on the effect of NACS integration into the health system on maternal and
- 454 infant health, nutrition practices, and outcomes adapted from the Chronic Care Model
- 455 S2 Fig. Enrolment and data analysis flow chart

456 Acknowledgements

We extend our appreciation to the political and administrative authorities of Tororo and Butaleja districts for granting us the necessary permission to conduct this study. Our gratitude go to the health workers as well as the mothers and their infants from Tororo and Butaleja districts for the dedicated participation in advancing this research undertaking. Brian Wakoli is appreciated for the statistical support rendered during the analysis.

462 Author contribution

- 463 Conceptualization: SN FEM GWM SNK
- 464 **Data Curation:** SN FEM SNK
- 465 **Formal analysis:** SN SNK FEM
- 466 **Funding acquisition:** SN
- 467 Investigation: SN
- 468 Methodology: SN SNK FEM
- 469 **Project administration:** SN
- 470 **Resources:** SN
- 471 Software: SN
- 472 **Supervision:** SN FEM GWM SNK
- 473 Validation: SN FEM GWM SNK
- 474 Visualization: SN FEM
- 475 Writing- Original draft: SN

476 Writing – Review and editing: SN FEM GWM SNK

477

478 **References**

479 480 481	1.	Development Initiatives Poverty Research Ltd. Global Nutrition Report: Stronger commitments for greater action. Global Nutrition Report. Bristol. UK; 2022. Available: http://www.segenlan.gob.gt/2.0/index.php?ontion=com_content&view=article&id=47
483		2&Itemid=472
484 485	2.	UBOS. Uganda Demographic and Health Survey 2016. Udhs. 2016. Available: www.DHSprogram.com
486	3.	Marshall NE, Abrams B, Barbour LA, Catalano P, Christian P, Friedman JE, et al. The
487		importance of nutrition in pregnancy and lactation: lifelong consequences. Am J
488		Obstet Gynecol. 2022;226: 607-632. doi:10.1016/j.ajog.2021.12.035
489	4.	Brink LR, Bender TM, Davies R, Luo H, Miketinas D, Shah N, et al. Optimizing
490		Maternal Nutrition: The Importance of a Tailored Approach. Curr Dev Nutr. 2022;6:
491		1–14. doi:10.1093/cdn/nzac118
492	5.	Girard AW, Olude O. Nutrition education and counselling provided during pregnancy:
493		Effects on maternal, neonatal and child health outcomes. Paediatr Perinat Epidemiol.
494		2012;26: 191–204. doi:10.1111/j.1365-3016.2012.01278.x
495	6.	Taneja S. Impact of an integrated nutrition, health, water sanitation and hygiene,
496		psychosocialcare and support intervention package delivered. Bmc. 2020;8: 1-18.
497	7.	Taneja S, Upadhyay RP, Chowdhury R, Kurpad A V., Bhardwaj H, Kumar T, et al.
498		Impact of nutritional interventions among lactating mothers on the growth of their
499		infants in the first 6 months of life: A randomized controlled trial in Delhi, India. Am J
500		Clin Nutr. 2021;113: 884-894. doi:10.1093/ajcn/nqaa383
501	8.	M Barker, SU Dombrowski, T Colbourn, CHD Fall, NM Kriznik, W Lawrence, SA
502		Norris, G Ngaiza, D Patel, J Skordis-Worrall, FF Sniehotta, R Steegers-Theunissen, C
503		Vogel, K Woods Townsend and JS. Intervention strategies to improve nutrition and
504		health behaviours before conception. Lancet. 2018;391: 1853-1864.
505		doi:10.1016/S0140-6736(18)30313-1.Intervention
506	9.	Harriet Torlesse, Nita Dalmiya VT and VA. Counselling to improve maternal

507 508		nutrition: Considerations for programming with quality, equity and scale. Unicef. 2021; 1–16.
509 510 511 512	10.	Von Salmuth V, Brennan E, Kerac M, McGrath M, Frison S, Lelijveld N. Maternal- focused interventions to improve infant growth and nutritional status in lowmiddle income countries: A systematic review of reviews. PLoS One. 2021;16: 1–30. doi:10.1371/journal.pone.0256188
513 514 515	11.	Adeyemi O, Toure M, Covic N, van den Bold M, Nisbett N, Headey D. Understanding drivers of stunting reduction in Nigeria from 2003 to 2018: a regression analysis. Food Secur. 2022;14: 995–1011. doi:10.1007/s12571-022-01279-8
516 517 518	12.	Akseer N, Vaivada T, Rothschild O, Ho K, Bhutta ZA. Understanding multifactorial drivers of child stunting reduction in exemplar countries: A mixed-methods approach. Am J Clin Nutr. 2020;112: 792S-805S. doi:10.1093/ajcn/nqaa152
519 520 521	13.	Bhutta ZA, Akseer N, Keats EC, Vaivada T, Baker S, Horton SE, et al. How countries can reduce child stunting at scale: Lessons from exemplar countries. Am J Clin Nutr. 2020;112: 894S-904S. doi:10.1093/ajcn/nqaa153
522 523 524 525 526	14.	USAID Advancing Nutrition. Stunting: considerations for use as an indicator in nutrition projects. Arlington, VA: USAID Advancing Nutrition. USAID Advancing Nutrition; 2020. Available: https://www.advancingnutrition.org/sites/default/files/2021- 10/usaid_an_stunting_literature_review_2021.pdf
527 528 529 530	15.	Dearden KA, Bishwakarma R, Crookston BT, Masau BT, Mulokozi GI. Health facility-based counselling and community outreach are associated with maternal dietary practices in a cross-sectional study from Tanzania. BMC Nutr. 2021;7: 1–11. doi:10.1186/s40795-021-00447-x
531 532 533 534	16.	Kaleem R, Adnan M, Nasir M, Rahat T. Effects of antenatal nutrition counselling on dietary practices and nutritional status of pregnant women: A quasi-experimental hospital based study. Pakistan J Med Sci. 2020;36: 632–636. doi:10.12669/pjms.36.4.1919
535 536 537	17.	Pérez-Escamilla R, Segura-Pérez S, Hall Moran V. Dietary guidelines for children under 2 years of age in the context of nurturing care. Matern Child Nutr. 2019;15: 2–4. doi:10.1111/mcn.12855
538	18.	Ghosh-Jerath S, Devasenapathy N, Singh A, Shankar A, Zodpey S. Ante natal care

539		(ANC) utilization, dietary practices and nutritional outcomes in pregnant and recently
540		delivered women in urban slums of Delhi, India: an exploratory cross-sectional study.
541		Reprod Health. 2015;12. doi:10.1186/s12978-015-0008-9
542	19.	Nsiah-Asamoah C, Pereko KKA, Intiful FD. Nutritional counselling interactions
543		between health workers and caregivers of children under two years: Observations at
544		selected child welfare clinics in Ghana. BMC Health Serv Res. 2019;19: 1-15.
545		doi:10.1186/s12913-019-4692-y
546	20.	MINISTRY OF HEALTH. Guidelines on Maternal, Infant, Young Child and
547		Adolescent Nutrition. Ministry of Health, Uganda; 2021. Available:
548		https://www.health.go.ug/cause/guidelines-on-maternal-infant-young-child-and-
549		adolescent-nutrition/
550	21.	World Health Organization. WHO recommendations on antenatal care for a positive
551		pregnancy experience. WHO recommendations on antenatal care for a positive
552		pregnancy experience. Geneva: World Health Organization; 2020. Available:
553		www.who.int/maternal_child_adolescent
554	22.	Ministry of Health U. INTEGRATING NUTRITION ASSESSMENT,
555		COUNSELLING, AND SUPPORT INTO Training Course for Facility-Based Health
556		Providers. 2016.
557	23.	Levin CE, Self JL, Kedera E, Wamalwa M, Hu J, Grant F, et al. What is the cost of
558		integration? Evidence from an integrated health and agriculture project to improve
559		nutrition outcomes in Western Kenya. Health Policy Plan. 2019;34: 646-655.
560		doi:10.1093/heapol/czz083
561	24.	Kasiulevičius V, Šapoka V, Filipavičiūtė R. Sample size calculation in
562		epidemiological studies. Gerontologija. 2006;7: 225-231. doi:013165/AIM.0010
563	25.	Beal SJ, Kupzyk KA. An Introduction to Propensity Scores: What, When, and How. J
564		Early Adolesc. 2014;34: 66–92. doi:10.1177/0272431613503215
565	26.	Valojerdi AE, Janani L. A brief guide to propensity score analysis. Med J Islam Repub
566		Iran. 2018;32: 1–4. doi:10.14196/mjiri.32.122
567	27.	Garrido MM, Kelley AS, Paris J, Roza K, Meier DE, Morrison RS, et al. Methods for
568		constructing and assessing propensity scores. Health Serv Res. 2014;49: 1701-1720.
569		doi:10.1111/1475-6773.12182
570	28.	Dehejia RH, Wahba S. Propensity score-matching methods for nonexperimental causal

571		studies. Rev Econ Stat. 2002;84: 151–161. doi:10.1162/003465302317331982
572	29.	Taylor RM, Wolfson JA, Lavelle F, Dean M, Frawley J, Hutchesson MJ, et al. Impact
573		of preconception, pregnancy, and postpartum culinary nutrition education
574		interventions: A systematic review. Nutr Rev. 2021;79: 1186-1203.
575		doi:10.1093/nutrit/nuaa124
576	30.	Kim C, Mansoor GF, Paya PM, Ludin MH, Ahrar MJ, Mashal MO, et al. Review of
577		policies, data, and interventions to improve maternal nutrition in Afghanistan. Matern
578		Child Nutr. 2020;16. doi:10.1111/mcn.13003
579	31.	Tsegaye D, Tamiru D, Belachew T. Theory-based nutrition education intervention
580		through male involvement improves the dietary diversity practice and nutritional status
581		of pregnant women in rural Illu Aba Bor Zone, Southwest Ethiopia: A quasi-
582		experimental study. Matern Child Nutr. 2022;18. doi:10.1111/mcn.13350
583	32.	Habtu M, Agena AG, Umugwaneza M, Mochama M, Munyanshongore C. Effect of
584		integrated nutrition-sensitive and nutrition-specific intervention package on maternal
585		malnutrition among pregnant women in Rwanda. Matern Child Nutr. 2022;18.
586		doi:10.1111/mcn.13367
587	33.	Kuma MN, Tamiru D, Tefera B. Effects of nutrition education and home gardening
588		interventions on feto-maternal outcomes among pregnant women in Jimma Zone,
589		Soutwest Ethiopia: A cluster randomized control trial. Nutr Health. 2023;29: 97–105.
590		doi:10.1177/02601060211056745
591	34.	Singh V, Ahmed S, Dreyfuss ML, Kiran U, Chaudhery DN, Srivastava VK, et al. An
592		integrated nutrition and health program package on IYCN improves breastfeeding but
593		not complementary feeding and nutritional status in rural northern India: A quasi-
594		experimental randomized longitudinal study. PLoS One. 2017;12: 1-20.
595		doi:10.1371/journal.pone.0185030
596	35.	Habtu M, Agena AG, Umugwaneza M, Mochama M, Munyanshongore C.
597		Effectiveness of Integrated Maternal Nutrition Intervention Package on Birth Weight
598		in Rwanda. Front Nutr. 2022;9. doi:10.3389/fnut.2022.874714
599	36.	González-Fernández D, Mazzini Salom AS, Herrera Bendezu F, Huamán S, Rojas
600		Hernández B, Pevec I, et al. A Multi-Sectoral Approach Improves Early Child
601		Development in a Disadvantaged Community in Peru: Role of Community Gardens,
602		Nutrition Workshops and Enhanced Caregiver-Child Interaction: Project "Wawa
603		Illari." Front Public Heal. 2020;8. doi:10.3389/fpubh.2020.567900